

# STAT 510 Homework 11

Yifan Zhu

April 19, 2017

1. (a) For this model, we have the model matrix

$$\mathbf{X} = \left[ \mathbf{1}_{n_{11}+n_{12}+n_{21}+n_{22}}, \begin{bmatrix} \mathbf{1}_{n_{11}+n_{12}} \\ \mathbf{0}_{n_{21}+n_{22}} \end{bmatrix}, \begin{bmatrix} \mathbf{0}_{n_{11}+n_{12}} \\ \mathbf{1}_{n_{21}+n_{22}} \end{bmatrix}, \begin{bmatrix} \mathbf{1}_{n_{11}} \\ \mathbf{0}_{n_{12}} \\ \mathbf{1}_{n_{21}} \\ \mathbf{0}_{n_{22}} \end{bmatrix}, \begin{bmatrix} \mathbf{0}_{n_{11}} \\ \mathbf{1}_{n_{12}} \\ \mathbf{0}_{n_{21}} \\ \mathbf{1}_{n_{22}} \end{bmatrix} \right]$$

To make it full column rank with the same column space, then

$$\mathbf{X} = [\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3], \text{ where } \mathbf{x}_1 = \begin{bmatrix} \mathbf{1}_{n_{11}+n_{12}} \\ \mathbf{0}_{n_{21}+n_{22}} \end{bmatrix}, \mathbf{x}_2 = \begin{bmatrix} \mathbf{0}_{n_{11}+n_{12}} \\ \mathbf{1}_{n_{21}+n_{22}} \end{bmatrix}, \mathbf{x}_3 = \begin{bmatrix} \mathbf{1}_{n_{11}} \\ \mathbf{0}_{n_{12}} \\ \mathbf{1}_{n_{21}} \\ \mathbf{0}_{n_{22}} \end{bmatrix}$$

- (b) We already have  $\mathbf{x}_1 \perp \mathbf{x}_2$ , thus let  $\mathbf{w}_1 = \mathbf{x}_1$ ,  $\mathbf{w}_2 = \mathbf{x}_2$ . For  $\mathbf{w}_3$ , first we know

$$\mathbf{P}_{[\mathbf{w}_1, \mathbf{w}_2]} = \mathbf{P}_{[\mathbf{x}_1, \mathbf{x}_2]} = \begin{bmatrix} \frac{1}{n_{11}+n_{12}} \mathbf{1}\mathbf{1}^T & \mathbf{0} \\ \mathbf{0} & \frac{1}{n_{21}+n_{22}} \mathbf{1}\mathbf{1}^T \end{bmatrix}$$

Thus we have

$$\mathbf{w}_3 = (\mathbf{I} - \mathbf{P}_{[\mathbf{x}_1, \mathbf{x}_2]})\mathbf{x}_3 = \begin{bmatrix} \mathbf{1}_{n_{11}} \\ \mathbf{0}_{n_{12}} \\ \mathbf{1}_{n_{21}} \\ \mathbf{0}_{n_{22}} \end{bmatrix} - \begin{bmatrix} \frac{n_{11}}{n_{11}+n_{12}} \mathbf{1}_{n_{11}+n_{12}} \\ \frac{n_{21}}{n_{21}+n_{22}} \mathbf{1}_{n_{21}+n_{22}} \end{bmatrix} = \begin{bmatrix} \frac{n_{12}}{n_{11}+n_{12}} \mathbf{1}_{n_{11}} \\ -\frac{n_{11}}{n_{11}+n_{12}} \mathbf{1}_{n_{12}} \\ \frac{n_{22}}{n_{21}+n_{22}} \mathbf{1}_{n_{21}} \\ -\frac{n_{21}}{n_{21}+n_{22}} \mathbf{1}_{n_{22}} \end{bmatrix} = \begin{bmatrix} \frac{4}{5} \mathbf{1}_2 \\ -\frac{1}{5} \mathbf{1}_8 \\ \frac{2}{5} \mathbf{1}_6 \\ -\frac{3}{5} \mathbf{1}_4 \end{bmatrix}$$

$$\mathbf{W}^T \mathbf{W} = \begin{bmatrix} \mathbf{w}_1^T \\ \mathbf{w}_2^T \\ \mathbf{w}_3^T \end{bmatrix} [\mathbf{w}_1 \quad \mathbf{w}_2 \quad \mathbf{w}_3] = \begin{bmatrix} \mathbf{w}_1^T \mathbf{w}_1 & 0 & 0 \\ 0 & \mathbf{w}_2^T \mathbf{w}_2 & 0 \\ 0 & 0 & \mathbf{w}_3^T \mathbf{w}_3 \end{bmatrix}$$

$$\mathbf{w}_1^T \mathbf{w}_1 = n_{11} + n_{12} = 10$$

$$\mathbf{w}_2^T \mathbf{w}_2 = n_{21} + n_{22} = 10$$

$$\begin{aligned} \mathbf{w}_3^T \mathbf{w}_3 &= \left( \frac{n_{12}}{n_{11}+n_{12}} \right)^2 n_{11} + \left( \frac{n_{11}}{n_{11}+n_{12}} \right)^2 n_{12} + \left( \frac{n_{22}}{n_{21}+n_{22}} \right)^2 n_{21} + \left( \frac{n_{21}}{n_{21}+n_{22}} \right)^2 n_{22} \\ &= \left( \frac{4}{5} \right)^2 \times 2 + \left( \frac{1}{5} \right)^2 \times 8 + \left( \frac{2}{5} \right)^2 \times 6 + \left( \frac{3}{5} \right)^2 \times 4 = 4 \end{aligned}$$

$$\mathbf{W}^T \mathbf{y} = \begin{bmatrix} \mathbf{w}_1^T \\ \mathbf{w}_2^T \\ \mathbf{w}_3^T \end{bmatrix} \mathbf{y} = \begin{bmatrix} \mathbf{w}_1^T \mathbf{y} \\ \mathbf{w}_2^T \mathbf{y} \\ \mathbf{w}_3^T \mathbf{y} \end{bmatrix}$$

$$\mathbf{w}_1^T \mathbf{y} = n_{11} \bar{y}_{11\cdot} + n_{12} \bar{y}_{12\cdot} = 2 \cdot 3 + 8 \cdot 5 = 46$$

$$\mathbf{w}_2^T \mathbf{y} = n_{21} \bar{y}_{21\cdot} + n_{22} \bar{y}_{22\cdot} = 6 \cdot 7 + 4 \cdot 3 = 54$$

$$\mathbf{w}_3^T \mathbf{y} = \frac{n_{12}n_{11}}{n_{11}+n_{12}} \bar{y}_{11\cdot} - \frac{n_{11}n_{12}}{n_{11}+n_{12}} \bar{y}_{12\cdot} + \frac{n_{22}n_{21}}{n_{21}+n_{22}} \bar{y}_{21\cdot} - \frac{n_{21}n_{22}}{n_{21}+n_{22}} \bar{y}_{22\cdot} = 6.4$$

Then

$$(W^T W)^{-1} W^T \mathbf{y} = \begin{bmatrix} 1/w_1^T w_1 & 0 & 0 \\ 0 & 1/w_2^T w_2 & 0 \\ 0 & 0 & 1/w_3^T w_3 \end{bmatrix} \begin{bmatrix} w_1^T \mathbf{y} \\ w_2^T \mathbf{y} \\ w_3^T \mathbf{y} \end{bmatrix} = \begin{bmatrix} \frac{w_1^T \mathbf{y}}{w_1^T w_1} \\ \frac{w_2^T \mathbf{y}}{w_2^T w_2} \\ \frac{w_3^T \mathbf{y}}{w_3^T w_3} \end{bmatrix} = \begin{bmatrix} 4.6 \\ 5.4 \\ 1.6 \end{bmatrix}$$

$$\begin{aligned} W(W^T W)^{-1} W^T \mathbf{y} &= \begin{bmatrix} w_1 & w_2 & w_3 \end{bmatrix} \begin{bmatrix} \frac{w_1^T \mathbf{y}}{w_1^T w_1} \\ \frac{w_2^T \mathbf{y}}{w_2^T w_2} \\ \frac{w_3^T \mathbf{y}}{w_3^T w_3} \end{bmatrix} \\ &= \frac{w_1^T \mathbf{y}}{w_1^T w_1} w_1 + \frac{w_2^T \mathbf{y}}{w_2^T w_2} w_2 + \frac{w_3^T \mathbf{y}}{w_3^T w_3} w_3 \\ &= \begin{bmatrix} \left( \frac{w_1^T \mathbf{y}}{w_1^T w_1} + \frac{n_{12}}{n_{11}+n_{12}} \frac{w_3^T \mathbf{y}}{w_3^T w_3} \right) \mathbf{1}_{n_{11}} \\ \left( \frac{w_1^T \mathbf{y}}{w_1^T w_1} - \frac{n_{11}}{n_{11}-n_{12}} \frac{w_3^T \mathbf{y}}{w_3^T w_3} \right) \mathbf{1}_{n_{12}} \\ \left( \frac{w_2^T \mathbf{y}}{w_2^T w_2} + \frac{n_{22}}{n_{21}+n_{22}} \frac{w_3^T \mathbf{y}}{w_3^T w_3} \right) \mathbf{1}_{n_{21}} \\ \left( \frac{w_2^T \mathbf{y}}{w_2^T w_2} - \frac{n_{21}}{n_{21}-n_{22}} \frac{w_3^T \mathbf{y}}{w_3^T w_3} \right) \mathbf{1}_{n_{22}} \end{bmatrix} \\ &= \begin{bmatrix} (4.6 + 0.8 \times 1.6) \mathbf{1}_2 \\ (4.6 - 0.2 \times 1.6) \mathbf{1}_8 \\ (5.4 + 0.4 \times 1.6) \mathbf{1}_6 \\ (5.4 - 0.6 \times 1.6) \mathbf{1}_4 \end{bmatrix} \\ &= \begin{bmatrix} 5.88 \mathbf{1}_2 \\ 4.28 \mathbf{1}_8 \\ 6.04 \mathbf{1}_6 \\ 4.44 \mathbf{1}_4 \end{bmatrix} = P_W \mathbf{y} = P_X \mathbf{y} \end{aligned}$$

(c)

$$P_{[x_1, x_3]} \mathbf{y} = \begin{bmatrix} \frac{n_{11} + \bar{y}_{11} + n_{12} \bar{y}_{12}}{n_{21} + n_{22}} \mathbf{1}_{n_{11} + n_{12}} \\ \frac{n_{21} + \bar{y}_{21} + n_{22} \bar{y}_{22}}{n_{21} + n_{22}} \mathbf{1}_{n_{21} + n_{22}} \end{bmatrix} = \begin{bmatrix} 4.6 \mathbf{1}_{10} \\ 5.4 \mathbf{1}_{10} \end{bmatrix}$$

Then

$$\begin{aligned} SS(B|1, A) &= \mathbf{y}^T (P_X - P_{[x_1, x_2]}) \mathbf{y} \\ &= \mathbf{y}^T P_X \mathbf{y} - \mathbf{y}^T P_{[x_1, x_2]} \mathbf{y} \\ &= (P_X \mathbf{y})^T (P_X \mathbf{y}) - (P_{[x_1, x_2]} \mathbf{y})^T (P_{[x_1, x_2]} \mathbf{y}) \\ &= (5.88^2 \times 2 + 4.28^2 \times 8 + 6.04^2 \times 6 + 4.44^2 \times 4) - (4.6^2 \times 10 + 5.4^2 \times 10) \\ &= 10.24 \end{aligned}$$

2. (a)  $\boldsymbol{\mu} = [\mu_{ik}]$ ,  $i = 1, 2, 3$ ,  $k = 1, \dots, 4$ .  $\mathbf{w} = [w_{ij}]$   $i = 1, 2, 3$ ,  $j = 1, \dots, 5$ .  $\mathbf{bme} = [e_{ijk}]$ ,  $i = 1, 2, 3$ ,  $j = 1, \dots, 5$ ,  $k = 1, \dots, 4$ . Then

$$\mathbf{y} = (\mathbf{I}_{3 \times 3} \otimes \mathbf{1}_5 \otimes \mathbf{I}_{4 \times 4}) \boldsymbol{\mu} + (\mathbf{I}_{15 \times 15} \otimes \mathbf{1}_4) \mathbf{w} + \mathbf{I}_{60 \times 60} \mathbf{e}$$

Then

$$Var(\mathbf{y}) = (\mathbf{I}_{15 \times 15} \otimes \mathbf{1}_4) Var(\mathbf{w}) (\mathbf{I}_{15 \times 15} \otimes \mathbf{1}_4)^T + Var(\mathbf{e}) = \sigma_w^2 (\mathbf{I}_{15 \times 15} \otimes \mathbf{1}_4 \mathbf{1}_4^T) + \sigma_e^2 \mathbf{I}_{60 \times 60}$$

(b) Test statistic  $F = 7.1152$ .

Degree of freedom are (6, 36).

p-value  $< 0.0001$ .

Since  $p < 0.0001$  is a very small number less than 0.05. We reject the null hypothesis and conclude that there is significant evidence that there is drug-by-time interactions.

(c)  $H_0 : \mu_{24} - \mu_{14} = \mu_{34} - \mu_{14} = 0$

We have

$$\mathbf{C}\hat{\boldsymbol{\beta}} = \begin{bmatrix} \hat{\mu}_{24} - \hat{\mu}_{14} \\ \hat{\mu}_{34} - \hat{\mu}_{14} \end{bmatrix} = \begin{bmatrix} \hat{\alpha}_2 + \hat{\gamma}_{24} \\ \hat{\alpha}_3 + \hat{\gamma}_{14} \end{bmatrix} = \begin{bmatrix} 5.6 \\ 1.6 \end{bmatrix}$$

Also

$$\begin{aligned} \text{Var}(\mathbf{C}\hat{\boldsymbol{\beta}}) &= \text{Var}\left(\begin{bmatrix} \bar{y}_{2.4} - \bar{y}_{1.4} \\ \bar{y}_{3.4} - \bar{y}_{1.4} \end{bmatrix}\right) \\ &= \begin{bmatrix} \text{Var}(\bar{y}_{2.4}) + \text{Var}(\bar{y}_{1.4}) & \text{Cov}(\bar{y}_{2.4}, \bar{y}_{1.4}) \\ \text{Cov}(\bar{y}_{3.4}, \bar{y}_{1.4}) & \text{Var}(\bar{y}_{3.4}) + \text{Var}(\bar{y}_{1.4}) \end{bmatrix} \\ &= \begin{bmatrix} \frac{2}{5}\sigma_w^2 + \frac{2}{5}\sigma_e^2 & \frac{1}{5}\sigma_w^2 + \frac{1}{5}\sigma_e^2 \\ \frac{1}{5}\sigma_w^2 + \frac{1}{5}\sigma_e^2 & \frac{2}{5}\sigma_w^2 + \frac{2}{5}\sigma_e^2 \end{bmatrix} \\ &= \frac{1}{5}(\sigma_w^2 + \sigma_e^2) \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix} \end{aligned}$$

Hence

$$(\widehat{\text{Var}}(\mathbf{C}\hat{\boldsymbol{\beta}}))^{-1} = \frac{5}{\hat{\sigma}_w^2 + \hat{\sigma}_e^2} \begin{bmatrix} \frac{2}{3} & -\frac{1}{3} \\ -\frac{1}{3} & \frac{2}{3} \end{bmatrix}$$

Then the test statistic

$$F = \frac{(\mathbf{C}\hat{\boldsymbol{\beta}})^T (\widehat{\text{Var}}(\mathbf{C}\hat{\boldsymbol{\beta}}))^{-1} (\mathbf{C}\hat{\boldsymbol{\beta}})}{q} = \frac{5}{2(\hat{\sigma}_e^2 + \hat{\sigma}_w^2)} [5.6 \quad 1.6] \begin{bmatrix} 2/3 & -1/3 \\ -1/3 & 2/3 \end{bmatrix} \begin{bmatrix} 5.6 \\ 1.6 \end{bmatrix} = 1.109$$

For degrees of freedom, because

$$\hat{\sigma}_w^2 + \hat{\sigma}_e^2 = \frac{1}{4}(4\hat{\sigma}_w^2 + \hat{\sigma}_e^2) + \frac{3}{4}\hat{\sigma}_e^2 = \frac{1}{4}MS_{\text{woman}(\text{drug})} + \frac{3}{4}MS_{\text{error}}$$

Thus

$$df = \frac{(\frac{1}{4}MS_{\text{woman}(\text{drug})} + \frac{3}{4}MS_{\text{error}})^2}{\frac{(\frac{1}{4})^2 MS_{\text{woman}(\text{drug})}^2}{12} + \frac{(\frac{3}{4})^2 MS_{\text{error}}^2}{36}} = 17.06$$

Then p-value

$$p = P(F_{2,17.06} > 1.109) = 0.35$$

The p-value  $> 0.05$ . We fail to reject the null hypothesis and conclude that there is no significant evidence that the mean heart rate 15 minutes after treatment are not the same for all three drugs.

(d) We know

$$\begin{aligned} \hat{\mu}_{14} - \hat{\mu}_{24} &= -5.6 \\ SE(\hat{\mu}_{14} - \hat{\mu}_{24}) &= \sqrt{\frac{2}{5}(\hat{\sigma}_e^2 + \hat{\sigma}_w^2)} = 3.87 \end{aligned}$$

Then

$$CI = (\hat{\mu}_{14} - \hat{\mu}_{24} - t_{17.06,0.975} \cdot SE, \hat{\mu}_{14} - \hat{\mu}_{24} + t_{17.06,0.975} \cdot SE) = (-13.766, 2.566)$$

3. (a)  $\sigma = 6.12, \rho = 0.7769$ .  
 (b)  $AIC = 317.92, BIC = 344.12$ .  
 (c)  $\sigma = 6.00, \rho = 0.8278$ .  
 (d)  $AIC = 313.94, BIC = 340.14$   
 (e)  $\sigma = 6.10, \delta_2 = 1.085, \delta_3 = 0.995, \delta_4 = 0.928, \rho_{12} = 0.850, \rho_{13} = 0.889, \rho_{14} = 0.625, \rho_{23} = 0.870, \rho_{24} = 0.631, \rho_{34} = 0.794$ .  
 (f)  $AIC = 322.85, BIC = 364.01$   
 (g) AR1 is preferred. The model has smaller AIC and BIC.  
 (h)  $\hat{\mu}_{14} - \hat{\mu}_{24} = -5.6, SE(\hat{\mu}_{14} - \hat{\mu}_{24}) = \sqrt{\frac{2}{5}6^2} = 3.79$ .

$$CI = (-5.6 - z_{0.975} \times 3.79, -5.6 + z_{0.975} \times 3.79) = (-13.03, 1.83)$$

4. (a)

$$\mathbf{y} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 1 & 0 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} \mu_1 \\ \mu_2 \end{bmatrix} + \begin{bmatrix} 1 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} u_1 \\ u_2 \\ u_3 \end{bmatrix} + \begin{bmatrix} e_{11} \\ e_{12} \\ e_{21} \\ e_{31} \end{bmatrix}$$

$$\Sigma = \begin{bmatrix} 4 & 2 & 0 & 0 \\ 2 & 4 & 0 & 0 \\ 0 & 0 & 4 & 0 \\ 0 & 0 & 0 & 4 \end{bmatrix} \Rightarrow \Sigma^{-1} = \begin{bmatrix} 1/3 & -1/6 & 0 & 0 \\ -1/6 & 1/3 & 0 & 0 \\ 0 & 0 & 1/4 & 0 \\ 0 & 0 & 0 & 1/4 \end{bmatrix}, \boldsymbol{\beta} = (\mathbf{X}^T \Sigma^{-1} \mathbf{X})^{-1} \mathbf{X}^T \Sigma^{-1} \mathbf{y} = \begin{bmatrix} 50.33 \\ 53.67 \end{bmatrix} \Rightarrow \hat{\mu}_1 + \hat{\mu}_2 = 50.33 + 53.67 = 104.$$

- (b) From  $Var(y_{i1}) = Var(y_{i2}) = 4$  and  $Corr(y_{i1}, y_{i2}) = 0.5 \Rightarrow \sigma_u^2 = 2, \sigma_e^2 = 2$ .

$$\text{Thus } \mathbf{G} = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix}, \mathbf{Z}^T = \begin{bmatrix} 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}, \mathbf{y} - \mathbf{X}\boldsymbol{\beta} = \begin{bmatrix} 0.67 \\ 0.33 \\ -2.33 \\ 1.67 \end{bmatrix}. \text{ Hence}$$

$$BLUP(\mathbf{u}) = \mathbf{G}\mathbf{Z}^T \Sigma^{-1}(\mathbf{y} - \mathbf{X}\boldsymbol{\beta}) = \begin{bmatrix} 0.333 \\ -1.167 \\ 0.833 \end{bmatrix}$$

$$\text{Hence } y_{22} = 53.67 - 1.167 = 52.5, y_{23} = 53.67 + 0.833 = 54.5.$$