

Descriptive Statistics: Part 2/2 (Ch 3)

Yifan Zhu

Iowa State University

Outline

Descriptive
Statistics: Part
2/2 (Ch 3)

Yifan Zhu

Boxplots

Boxplots

Quantile-Quantile
(QQ) Plots

Quantile-Quantile (QQ) Plots

Theoretical
Quantile-Quantile
Plots

Theoretical Quantile-Quantile Plots


Numerical
Summaries

Numerical Summaries

Parameters

Parameters

Generic Boxplot



Boxplots

Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters

Example: bullet data

Quantiles of the Bullet Penetration Depth Distributions

$i^r = 20 \times 0.25 + 0.5 = 5.5$

P_i	i	i th Smallest 230 Grain Data Point = $Q(\frac{i-.5}{20})$	i th Smallest 200 Grain Data Point = $Q(\frac{i-.5}{20})$
	1	27.75	58.00
	2	37.35	58.65
	3	38.35	59.10
	4	38.35	59.50
	5	38.75	59.80
	6	39.75	60.70
	7	40.50	61.30
	8	41.00	61.50
	9	41.15	62.30
	10	42.55	62.65
	11	42.90	62.95
	12	43.60	63.30
	13	43.85	63.55
	14	47.30	63.80
	15	47.90	64.05
	16	48.15	64.65
	17	49.85	65.00
	18	51.25	67.75
	19	51.60	70.40
	20	56.00	71.70

Boxplots

Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters

Example: bullet data (230-grain bullets)

Step 1:

$$Q(.25) = .5Q(.225) + .5Q(.275) = .5(38.75) + .5(39.75) = 39.25 \text{ mm}$$

$$Q(.5) = .5Q(.475) + .5Q(.525) = .5(42.55) + .5(42.90) = 42.725 \text{ mm}$$

$$Q(.75) = .5Q(.725) + .5Q(.775) = .5(47.90) + .5(48.15) = 48.025 \text{ mm}$$

So Step 2:

$$IQR = 48.025 - 39.25 = 8.775 \text{ mm}$$

$$1.5IQR = 13.163 \text{ mm}$$

$$Q(.75) + 1.5IQR = 61.188 \text{ mm}$$

$$Q(.25) - 1.5IQR = 26.087 \text{ mm}$$

Boxplots


Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters

Example: bullet data



Outline

Descriptive
Statistics: Part
2/2 (Ch 3)

Yifan Zhu

Boxplots

Boxplots

Quantile-Quantile (QQ) Plots

Quantile-Quantile
(QQ) Plots

Theoretical Quantile-Quantile Plots

Theoretical
Quantile-Quantile
Plots

Numerical Summaries

Numerical
Summaries

Parameters

Parameters

- quantiles*
- ▶ **Quantile-quantile (QQ) plot:** a scatterplot of the sorted values of one dataset on the sorted values of another dataset.
 - ▶ This plot is used to tell if the distributional shapes of the datasets are the same or different.
 - ▶ If the points in the plot lie in a straight line, the distributional shapes are the same.
 - ▶ Otherwise, the shapes are different.
 - ▶ The datasets must be univariate, numerical, and of the same size.

- ▶ **Quantile-quantile (QQ) plot:** a scatterplot of the sorted values of one dataset on the sorted values of another dataset.
 - ▶ This plot is used to tell if the distributional shapes of the datasets are the same or different.
 - ▶ If the points in the plot lie in a straight line, the distributional shapes are the same.
 - ▶ Otherwise, the shapes are different.
 - ▶ The datasets must be univariate, numerical, and of the same size.

- ▶ **Quantile-quantile (QQ) plot:** a scatterplot of the sorted values of one dataset on the sorted values of another dataset.
 - ▶ This plot is used to tell if the distributional shapes of the datasets are the same or different.
 - ▶ If the points in the plot lie in a straight line, the distributional shapes are the same.
 - ▶ Otherwise, the shapes are different.
 - ▶ The datasets must be univariate, numerical, and of the same size.

- ▶ **Quantile-quantile (QQ) plot:** a scatterplot of the sorted values of one dataset on the sorted values of another dataset.
 - ▶ This plot is used to tell if the distributional shapes of the datasets are the same or different.
 - ▶ If the points in the plot lie in a straight line, the distributional shapes are the same.
 - ▶ Otherwise, the shapes are different.
 - ▶ The datasets must be univariate, numerical, and of the same size.

- ▶ **Quantile-quantile (QQ) plot:** a scatterplot of the sorted values of one dataset on the sorted values of another dataset.
 - ▶ This plot is used to tell if the distributional shapes of the datasets are the same or different.
 - ▶ If the points in the plot lie in a straight line, the distributional shapes are the same.
 - ▶ Otherwise, the shapes are different.
 - ▶ The datasets must be univariate, numerical, and of the same size.

x_1, \dots, x_n dataset 1
 y_1, \dots, y_n dataset 2

(x_i, y_i)

Example: bullet data

Quantiles of the Bullet Penetration Depth Distributions

i	$\frac{i-.5}{20}$	i th Smallest 230 Grain Data Point = $Q(\frac{i-.5}{20})$	i th Smallest 200 Grain Data Point = $Q(\frac{i-.5}{20})$
1	.025	27.75	58.00
2	.075	37.35	58.65
3	.125	38.35	59.10
4	.175	38.35	59.50
5	.225	38.75	59.80
6	.275	39.75	60.70
7	.325	40.50	61.30
8	.375	41.00	61.50
9	.425	41.15	62.30
10	.475	42.55	62.65
11	.525	42.90	62.95
12	.575	43.60	63.30
13	.625	43.85	63.55
14	.675	47.30	63.80
15	.725	47.90	64.05
16	.775	48.15	64.65
17	.825	49.85	65.00
18	.875	51.25	67.75
19	.925	51.60	70.40
20	.975	56.00	71.70

Boxplots

Quantile-Quantile
(QQ) Plots


Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters


Example: bullet data

- ▶ I can make a QQ plot of the bullet data by plotting the sorted 200-grain depths against the sorted 230-grain depths.
- ▶ The points lie in approximately a straight line, so the 200-grain depths are similarly shaped in distribution to the 230-grain depths.



Example: bullet data

- ▶ I can make a QQ plot of the bullet data by plotting the sorted 200-grain depths against the sorted 230-grain depths.
- ▶ The points lie in approximately a straight line, so the 200-grain depths are similarly shaped in distribution to the 230-grain depths.



Outline

Descriptive
Statistics: Part
2/2 (Ch 3)

Yifan Zhu

Boxplots

Boxplots

Quantile-Quantile
(QQ) Plots

Quantile-Quantile (QQ) Plots

Theoretical
Quantile-Quantile
Plots

Theoretical Quantile-Quantile Plots



Numerical
Summaries

Numerical Summaries

Parameters

Parameters

Theoretical quantile-quantile (QQ) plots

► Theoretical quantile-quantile (QQ) plot: a scatterplot with:

- The sorted values x_1, x_2, \dots, x_n of some real data set on the x axis.
- $Q(\frac{1-.5}{n}), Q(\frac{2-.5}{n}), \dots, Q(\frac{n-.5}{n})$ on the y axis.
 - Q is some **theoretical quantile function**: the quantile function we would *expect* from a dataset if that dataset had a certain shape.

► Example theoretical quantile functions:

- “Standard” bell-shaped (or normally-distributed) data should have:

$$Q(p) \approx 4.9(p^{0.14} - (1-p)^{0.14})$$

- “Exponentially distributed” data (a kind of highly right-skewed data) should have:

$$Q(p) \approx -\lambda^{-1} \log(1-p)$$

where λ is some constant.

Boxplots

Quantile-Quantile
(QQ) PlotsTheoretical
Quantile-Quantile
PlotsNumerical
Summaries

Parameters

Theoretical quantile-quantile (QQ) plots

- ▶ **Theoretical quantile-quantile (QQ) plot:** a scatterplot with:

- ▶ The sorted values x_1, x_2, \dots, x_n of some real data set on the x axis.

- ▶ $Q(\frac{1-.5}{n}), Q(\frac{2-.5}{n}), \dots, Q(\frac{n-.5}{n})$ on the y axis.

- ▶ Q is some **theoretical quantile function**: the quantile function we would *expect* from a dataset if that dataset had a certain shape.

- ▶ Example theoretical quantile functions:

- ▶ “Standard” bell-shaped (or normally-distributed) data should have:

$$Q(p) \approx 4.9(p^{0.14} - (1-p)^{0.14})$$

- ▶ “Exponentially distributed” data (a kind of highly right-skewed data) should have:

$$Q(p) \approx -\lambda^{-1} \log(1-p)$$

where λ is some constant.

Boxplots

Quantile-Quantile
(QQ) PlotsTheoretical
Quantile-Quantile
PlotsNumerical
Summaries

Parameters

Theoretical quantile-quantile (QQ) plots

- ▶ **Theoretical quantile-quantile (QQ) plot:** a scatterplot with:

- ▶ The sorted values x_1, x_2, \dots, x_n of some real data set on the x axis.
- ▶ $Q(\frac{1-.5}{n}), Q(\frac{2-.5}{n}), \dots, Q(\frac{n-.5}{n})$ on the y axis.

- ▶ Q is some **theoretical quantile function**: the quantile function we would *expect* from a dataset if that dataset had a certain shape.

- ▶ Example theoretical quantile functions:

- ▶ “Standard” bell-shaped (or normally-distributed) data should have:

$$Q(p) \approx 4.9(p^{0.14} - (1-p)^{0.14})$$

- ▶ “Exponentially distributed” data (a kind of highly right-skewed data) should have:

$$Q(p) \approx -\lambda^{-1} \log(1-p)$$

where λ is some constant.

Boxplots

Quantile-Quantile
(QQ) PlotsTheoretical
Quantile-Quantile
PlotsNumerical
Summaries

Parameters

Theoretical quantile-quantile (QQ) plots

- **Theoretical quantile-quantile (QQ) plot:** a scatterplot with:

- The sorted values x_1, x_2, \dots, x_n of some real data set on the x axis.
- $Q(\frac{1-.5}{n}), Q(\frac{2-.5}{n}), \dots, Q(\frac{n-.5}{n})$ on the y axis.
 - Q is some **theoretical quantile function**: the quantile function we would expect from a dataset if that dataset had a certain shape.

- Example theoretical quantile functions:

- “Standard” bell-shaped (or normally-distributed) data should have:

$$Q(p) \approx 4.9(p^{0.14} - (1-p)^{0.14})$$

- “Exponentially distributed” data (a kind of highly right-skewed data) should have:

$$Q(p) \approx -\lambda^{-1} \log(1-p)$$

where λ is some constant.

Boxplots

Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters

Theoretical quantile-quantile (QQ) plots

- ▶ **Theoretical quantile-quantile (QQ) plot:** a scatterplot with:

- ▶ The sorted values x_1, x_2, \dots, x_n of some real data set on the x axis.
- ▶ $Q(\frac{1-.5}{n}), Q(\frac{2-.5}{n}), \dots, Q(\frac{n-.5}{n})$ on the y axis.
 - ▶ Q is some **theoretical quantile function**: the quantile function we would *expect* from a dataset if that dataset had a certain shape.

- ▶ Example theoretical quantile functions:

- ▶ “Standard” bell-shaped (or normally-distributed) data should have:

$$Q(p) \approx 4.9(p^{0.14} - (1-p)^{0.14})$$

- ▶ “Exponentially distributed” data (a kind of highly right-skewed data) should have:

$$Q(p) \approx -\lambda^{-1} \log(1-p)$$

where λ is some constant.


Theoretical quantile-quantile (QQ) plots

- ▶ **Theoretical quantile-quantile (QQ) plot:** a scatterplot with:

- ▶ The sorted values x_1, x_2, \dots, x_n of some real data set on the x axis.
- ▶ $Q(\frac{1-.5}{n}), Q(\frac{2-.5}{n}), \dots, Q(\frac{n-.5}{n})$ on the y axis.
 - ▶ Q is some **theoretical quantile function**: the quantile function we would *expect* from a dataset if that dataset had a certain shape.

- ▶ Example theoretical quantile functions:

- ▶ “Standard” bell-shaped (or normally-distributed) data should have:



$$Q(p) \approx 4.9(p^{0.14} - (1-p)^{0.14})$$

$\leftarrow N(0,1)$

- ▶ “Exponentially distributed” data (a kind of highly right-skewed data) should have:

$$Q(p) \approx -\lambda^{-1} \log(1-p)$$

where λ is some constant.

Boxplots

Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters

Theoretical quantile-quantile (QQ) plots


- **Theoretical quantile-quantile (QQ) plot:** a scatterplot with:

- The sorted values x_1, x_2, \dots, x_n of some real data set on the x axis.
- $Q(\frac{1-.5}{n}), Q(\frac{2-.5}{n}), \dots, Q(\frac{n-.5}{n})$ on the y axis.
 - Q is some **theoretical quantile function**: the quantile function we would *expect* from a dataset if that dataset had a certain shape.

- Example theoretical quantile functions:

- “Standard” bell-shaped (or normally-distributed) data should have:

$$Q(p) \approx 4.9(p^{0.14} - (1-p)^{0.14})$$

- 
- “Exponentially distributed” data (a kind of highly right-skewed data) should have:

$$Q(p) \approx -\lambda^{-1} \log(1-p)$$

where λ is some constant.

$\text{Exp}(\lambda)$

$\lambda = 1$

Normal quantile-quantile (QQ) Plots

Descriptive
Statistics: Part
2/2 (Ch 3)

Yifan Zhu

Boxplots

Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters


- ▶ **Normal quantile-quantile (QQ) plot:** a theoretical QQ plot where the quantile function, Q , is the quantile function for “standard” bell-shaped *N(0,1)* (normally-distributed) data.
- ▶ If the points in a normal QQ plot are in a straight line, the dataset in question is normally-distributed and bell-shaped. Otherwise, the data is not normally-distributed (but still could be bell-shaped).

Normal quantile-quantile (QQ) Plots


- ▶ **Normal quantile-quantile (QQ) plot:** a theoretical QQ plot where the quantile function, Q , is the quantile function for “standard” bell-shaped (normally-distributed) data.
- ▶ If the points in a normal QQ plot are in a straight line, the dataset in question is normally-distributed and bell-shaped. Otherwise, the data is not normally-distributed (but still could be bell-shaped).

↑
bell-shaped.

Histogram of t-distribution with 1 df



Normal Q-Q plot of t-distribution with df 1



Example: towel breaking strength data

Breaking Strength and Standard Normal Quantiles

i	$\frac{i-.5}{10}$	$\frac{i-.5}{10}$ Breaking Strength	$\frac{i-.5}{10}$ Standard Normal Quantile
1	.05	7,583	-1.65
2	.15	8,527	-1.04
3	.25	8,572	-.67
4	.35	8,577	-.39
5	.45	9,011	-.13
6	.55	9,165	.13
7	.65	9,471	.39
8	.75	9,614	.67
9	.85	9,614	1.04
10	.95	10,688	1.65

Boxplots


Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters

Example: towel breaking strength data



- The points are roughly straight-line-shaped, so the breaking strength data is roughly bell-shaped and normally-distributed.

Boxplots


Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters

Normal QQ plot: 200-grain bullet penetration



- ▶ Since the points in the normal QQ plot are not quite arranged in a straight line, the 200-grain penetration depths are not quite normally-distributed. However, the departure from normality is not severe.
- ▶ The QQ plot of the bullet data from before revealed that the 200-grain depths had the same distributional shape as the 200-grain bullet depths. Thus, the 230-grain bullet data is not quite normally-distributed either.

- ▶ Since the points in the normal QQ plot are not quite arranged in a straight line, the 200-grain penetration depths are not quite normally-distributed. However, the departure from normality is not severe.
- ▶ The QQ plot of the bullet data from before revealed that the 200-grain depths had the same distributional shape as the 200-grain bullet depths. Thus, the 230-grain bullet data is not quite normally-distributed either.

↑ also make QQ plot.

Outline

Descriptive
Statistics: Part
2/2 (Ch 3)

Yifan Zhu

Boxplots

Boxplots

Quantile-Quantile
(QQ) Plots

Quantile-Quantile (QQ) Plots

Theoretical
Quantile-Quantile
Plots

Theoretical Quantile-Quantile Plots

Numerical
Summaries

Numerical Summaries

Parameters

Parameters

Numerical summaries

▶ Numerical summary (statistic)

- ▶ A number or list of numbers calculated using the data (and only the data).
- ▶ Numerical summaries highlight important features of the data (shape, center, spread, outliers).

▶ Examples:

- ▶ Measures of center:
 - ▶ Arithmetic mean
 - ▶ Median
 - ▶ Mode
- ▶ Measures of spread:
 - ▶ Sample variance
 - ▶ Sample standard deviation
 - ▶ Range
 - ▶ IQR
- ▶ Measures of shape:
 - ▶ All the quantiles together
 - ▶ Skew (beyond the scope of the class)
 - ▶ Kurtosis (beyond the scope of the class)

Numerical summaries

► Numerical summary (statistic)

- A number or list of numbers calculated using the data (and only the data).
- Numerical summaries highlight important features of the data (shape, center, spread, outliers).

► Examples:

► Measures of center:

- Arithmetic mean
- Median
- Mode

► Measures of spread:

- Sample variance
- Sample standard deviation
- Range
- IQR

► Measures of shape:

- All the quantiles together
- Skew (beyond the scope of the class)
- Kurtosis (beyond the scope of the class)

Boxplots

Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters

Numerical summaries

► Numerical summary (statistic)

- A number or list of numbers calculated using the data (and only the data).
- Numerical summaries highlight important features of the data (shape, center, spread, outliers).

► Examples:

► Measures of center:

- Arithmetic mean
- Median
- Mode

► Measures of spread:

- Sample variance
- Sample standard deviation
- Range
- IQR

► Measures of shape:

- All the quantiles together
- Skew (beyond the scope of the class)
- Kurtosis (beyond the scope of the class)

Boxplots

Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters

Numerical summaries

► Numerical summary (statistic)

- A number or list of numbers calculated using the data (and only the data).
- Numerical summaries highlight important features of the data (shape, center, spread, outliers).

► Examples:

► Measures of center:

- Arithmetic mean
- Median
- Mode

► Measures of spread:

- Sample variance
- Sample standard deviation
- Range
- IQR

► Measures of shape:

- All the quantiles together
- Skew (beyond the scope of the class)
- Kurtosis (beyond the scope of the class)

Boxplots

Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters

Numerical summaries

► Numerical summary (statistic)

- A number or list of numbers calculated using the data (and only the data).
- Numerical summaries highlight important features of the data (shape, center, spread, outliers).

► Examples:

► Measures of center:

- Arithmetic mean
- Median
- Mode

► Measures of spread:

- Sample variance
- Sample standard deviation
- Range
- IQR

► Measures of shape:

- All the quantiles together
- Skew (beyond the scope of the class)
- Kurtosis (beyond the scope of the class)

Boxplots

Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters

Numerical summaries

▶ Numerical summary (statistic)

- ▶ A number or list of numbers calculated using the data (and only the data).
- ▶ Numerical summaries highlight important features of the data (shape, center, spread, outliers).

▶ Examples:

▶ Measures of center:

- ▶ Arithmetic mean ← Sample mean
- ▶ Median
- ▶ Mode

▶ Measures of spread:

- ▶ Sample variance
- ▶ Sample standard deviation
- ▶ Range
- ▶ IQR

▶ Measures of shape:

- ▶ All the quantiles together
- ▶ Skew (beyond the scope of the class)
- ▶ Kurtosis (beyond the scope of the class)

Boxplots

Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters

Numerical summaries

► Numerical summary (statistic)

- A number or list of numbers calculated using the data (and only the data).
- Numerical summaries highlight important features of the data (shape, center, spread, outliers).

► Examples:

- Measures of center:
 - Arithmetic mean
 - Median
 - Mode
- Measures of spread:
 - Sample variance
 - Sample standard deviation
 - Range
 - IQR
- Measures of shape:
 - All the quantiles together
 - Skew (beyond the scope of the class)
 - Kurtosis (beyond the scope of the class)

Boxplots

Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters

Numerical summaries

► Numerical summary (statistic)

- A number or list of numbers calculated using the data (and only the data).
- Numerical summaries highlight important features of the data (shape, center, spread, outliers).

► Examples:

- Measures of center:
 - Arithmetic mean
 - Median
 - Mode
- Measures of spread:
 - Sample variance
 - Sample standard deviation
 - Range
 - IQR
- Measures of shape:
 - All the quantiles together
 - Skew (beyond the scope of the class)
 - Kurtosis (beyond the scope of the class)

Boxplots

Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters

Numerical summaries

▶ Numerical summary (statistic)

- ▶ A number or list of numbers calculated using the data (and only the data).
- ▶ Numerical summaries highlight important features of the data (shape, center, spread, outliers).

▶ Examples:

- ▶ Measures of center:
 - ▶ Arithmetic mean
 - ▶ Median
 - ▶ Mode
- ▶ Measures of spread:
 - ▶ Sample variance
 - ▶ Sample standard deviation
 - ▶ Range
 - ▶ IQR
- ▶ Measures of shape:
 - ▶ All the quantiles together
 - ▶ Skew (beyond the scope of the class)
 - ▶ Kurtosis (beyond the scope of the class)

Numerical summaries

▶ Numerical summary (statistic)

- ▶ A number or list of numbers calculated using the data (and only the data).
- ▶ Numerical summaries highlight important features of the data (shape, center, spread, outliers).

▶ Examples:

- ▶ Measures of center:
 - ▶ Arithmetic mean
 - ▶ Median
 - ▶ Mode
- ▶ Measures of spread:
 - ▶ Sample variance
 - ▶ Sample standard deviation
 - ▶ Range
 - ▶ IQR
- ▶ Measures of shape:
 - ▶ All the quantiles together
 - ▶ Skew (beyond the scope of the class)
 - ▶ Kurtosis (beyond the scope of the class)

Boxplots

Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters

▶ Numerical summary (statistic)

- ▶ A number or list of numbers calculated using the data (and only the data).
- ▶ Numerical summaries highlight important features of the data (shape, center, spread, outliers).

▶ Examples:

- ▶ Measures of center:
 - ▶ Arithmetic mean
 - ▶ Median
 - ▶ Mode
- ▶ Measures of spread:
 - ▶ Sample variance
 - ▶ Sample standard deviation
 - ▶ Range
 - ▶ IQR
- ▶ Measures of shape:
 - ▶ All the quantiles together
 - ▶ Skew (beyond the scope of the class)
 - ▶ Kurtosis (beyond the scope of the class)

▶ Numerical summary (statistic)

- ▶ A number or list of numbers calculated using the data (and only the data).
- ▶ Numerical summaries highlight important features of the data (shape, center, spread, outliers).

▶ Examples:

- ▶ Measures of center:
 - ▶ Arithmetic mean
 - ▶ Median
 - ▶ Mode
- ▶ Measures of spread:
 - ▶ Sample variance
 - ▶ Sample standard deviation
 - ▶ Range
 - ▶ IQR
- ▶ Measures of shape:
 - ▶ All the quantiles together
 - ▶ Skew (beyond the scope of the class)
 - ▶ Kurtosis (beyond the scope of the class)

Measures of center

x_1	x_2	x_3	x_4	x_5	x_6
0	1	1	2	3	5

- ▶ Arithmetic mean:
 - ▶ $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$
 - ▶ Here, $\bar{x} = \frac{1}{6}(0 + 1 + 1 + 2 + 3 + 5) = 2$
- ▶ Median: $Q(0.5)$.
 - ▶ A shortcut to calculating $Q(0.5)$ is:
 - ▶ $Q(0.5) = x_{[n/2]}$ if n is odd
 - ▶ $Q(0.5) = (x_{n/2} + x_{n/2+1})/2$ if n is even.
 - ▶ Here, $Q(0.5) = (1 + 2)/2 = 1.5$
- ▶ Mode (of a discrete or categorical dataset)
 - ▶ the most frequently-occurring value
 - ▶ Here, mode = 1.

Boxplots

Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters

Measures of center

x_1	x_2	x_3	x_4	x_5	x_6
0	1	1	2	3	5

► Arithmetic mean:

► $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$

► Here, $\bar{x} = \frac{1}{6}(0 + 1 + 1 + 2 + 3 + 5) = 2$

► Median: $Q(0.5)$.

► A shortcut to calculating $Q(0.5)$ is:

► $Q(0.5) = x_{[n/2]}$ if n is odd

► $Q(0.5) = (x_{n/2} + x_{n/2+1})/2$ if n is even.

► Here, $Q(0.5) = (1 + 2)/2 = 1.5$

► Mode (of a discrete or categorical dataset)

► the most frequently-occurring value

► Here, mode = 1.

Boxplots

Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters

Measures of center

x_1	x_2	x_3	x_4	x_5	x_6
0	1	1	2	3	5

► Arithmetic mean:

$$\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$$

$$\text{► Here, } \bar{x} = \frac{1}{6}(0 + 1 + 1 + 2 + 3 + 5) = 2$$

► Median: $Q(0.5)$.

► A shortcut to calculating $Q(0.5)$ is:

► $Q(0.5) = x_{[n/2]}$ if n is odd

► $Q(0.5) = (x_{n/2} + x_{n/2+1})/2$ if n is even.

► Here, $Q(0.5) = (1 + 2)/2 = 1.5$

► Mode (of a discrete or categorical dataset)

► the most frequently-occurring value

► Here, mode = 1.

Boxplots

Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters

Measures of center $n = 6$ (sample size) .

x_1	x_2	x_3	x_4	x_5	x_6
0	1	1	2	3	5

sample mean

► Arithmetic mean:

$$\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$$

$$\text{Here, } \bar{x} = \frac{1}{6}(0 + 1 + 1 + 2 + 3 + 5) = 2$$

► Median: $Q(0.5)$.

► A shortcut to calculating $Q(0.5)$ is:

► $Q(0.5) = x_{[n/2]}$ if n is odd

► $Q(0.5) = (x_{n/2} + x_{n/2+1})/2$ if n is even.

► Here, $Q(0.5) = (1 + 2)/2 = 1.5$

► Mode (of a discrete or categorical dataset)

► the most frequently-occurring value

► Here, mode = 1.

Measures of center

x_1	x_2	x_3	x_4	x_5	x_6
0	1	1	2	3	5

► Arithmetic mean:

► $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$

► Here, $\bar{x} = \frac{1}{6}(0 + 1 + 1 + 2 + 3 + 5) = 2$

► Median: $Q(0.5)$.

► A shortcut to calculating $Q(0.5)$ is:

► $Q(0.5) = x_{\lceil n/2 \rceil}$ if n is odd

► $Q(0.5) = (x_{n/2} + x_{n/2+1})/2$ if n is even.

► Here, $Q(0.5) = (1 + 2)/2 = 1.5$

► Mode (of a discrete or categorical dataset)

► the most frequently-occurring value

► Here, mode = 1.

Boxplots

Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters

Measures of center

x_1	x_2	x_3	x_4	x_5	x_6
0	1	1	2	3	5

► Arithmetic mean:

► $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$

► Here, $\bar{x} = \frac{1}{6}(0 + 1 + 1 + 2 + 3 + 5) = 2$

► Median: $Q(0.5)$.

► A shortcut to calculating $Q(0.5)$ is:

► $Q(0.5) = x_{\lceil n/2 \rceil}$ if n is odd

► $Q(0.5) = (x_{n/2} + x_{n/2+1})/2$ if n is even.

► Here, $Q(0.5) = (1 + 2)/2 = 1.5$

► Mode (of a discrete or categorical dataset)

► the most frequently-occurring value

► Here, mode = 1.

Boxplots

Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters

Measures of center

x_1	x_2	x_3	x_4	x_5	x_6
0	1	1	2	3	5

- ▶ Arithmetic mean:
 - ▶ $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$
 - ▶ Here, $\bar{x} = \frac{1}{6}(0 + 1 + 1 + 2 + 3 + 5) = 2$
- ▶ Median: $Q(0.5)$.
 - ▶ A shortcut to calculating $Q(0.5)$ is:
 - ▶ $Q(0.5) = x_{\lceil n/2 \rceil}$ if n is odd
 - ▶ $Q(0.5) = (x_{n/2} + x_{n/2+1})/2$ if n is even.
 - ▶ Here, $Q(0.5) = (1 + 2)/2 = 1.5$
- ▶ Mode (of a discrete or categorical dataset)
 - ▶ the most frequently-occurring value
 - ▶ Here, mode = 1.

Boxplots

Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters

Measures of center

x_1	x_2	x_3	x_4	x_5	x_6
0	1	1	2	3	5

► Arithmetic mean:

► $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$

► Here, $\bar{x} = \frac{1}{6}(0 + 1 + 1 + 2 + 3 + 5) = 2$

► Median: $Q(0.5)$.

► A shortcut to calculating $Q(0.5)$ is:

► $Q(0.5) = x_{\lceil n/2 \rceil}$ if n is odd

► $Q(0.5) = (x_{n/2} + x_{n/2+1})/2$ if n is even.

► Here, $Q(0.5) = (1 + 2)/2 = 1.5$

$n = 5, \lceil 5/2 \rceil$

$= \lceil 2.5 \rceil$

$= 3$

$n = 6, n/2 = 3$

► Mode (of a discrete or categorical dataset) $(x_3 + x_4)/2$

► the most frequently-occurring value

► Here, mode = 1.

Boxplots

Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters

x_1	x_2	x_3	x_4	x_5	x_6
0	1	1	2	3	5

- ▶ Arithmetic mean:
 - ▶ $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$
 - ▶ Here, $\bar{x} = \frac{1}{6}(0 + 1 + 1 + 2 + 3 + 5) = 2$
- ▶ Median: $Q(0.5)$.
 - ▶ A shortcut to calculating $Q(0.5)$ is:
 - ▶ $Q(0.5) = x_{\lceil n/2 \rceil}$ if n is odd
 - ▶ $Q(0.5) = (x_{n/2} + x_{n/2+1})/2$ if n is even.
 - ▶ Here, $Q(0.5) = (1 + 2)/2 = 1.5$
- ▶ Mode (of a discrete or categorical dataset)
 - ▶ the most frequently-occurring value
 - ▶ Here, mode = 1.

Boxplots

Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters

Measures of center

x_1	x_2	x_3	x_4	x_5	x_6
0	1	1	2	3	5

- ▶ Arithmetic mean:
 - ▶ $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$
 - ▶ Here, $\bar{x} = \frac{1}{6}(0 + 1 + 1 + 2 + 3 + 5) = 2$
- ▶ Median: $Q(0.5)$.
 - ▶ A shortcut to calculating $Q(0.5)$ is:
 - ▶ $Q(0.5) = x_{\lceil n/2 \rceil}$ if n is odd
 - ▶ $Q(0.5) = (x_{n/2} + x_{n/2+1})/2$ if n is even.
 - ▶ Here, $Q(0.5) = (1 + 2)/2 = 1.5$
- ▶ Mode (of a discrete or categorical dataset)
 - ▶ the most frequently-occurring value
 - ▶ Here, mode = 1.

Boxplots

Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters

Measures of center

x_1	x_2	x_3	x_4	x_5	x_6
0	1	1	2	3	5

- ▶ Arithmetic mean:
 - ▶ $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$
 - ▶ Here, $\bar{x} = \frac{1}{6}(0 + 1 + 1 + 2 + 3 + 5) = 2$
- ▶ Median: $Q(0.5)$.
 - ▶ A shortcut to calculating $Q(0.5)$ is:
 - ▶ $Q(0.5) = x_{\lceil n/2 \rceil}$ if n is odd
 - ▶ $Q(0.5) = (x_{n/2} + x_{n/2+1})/2$ if n is even.
 - ▶ Here, $Q(0.5) = (1 + 2)/2 = 1.5$
- ▶ Mode (of a discrete or categorical dataset)
 - ▶ the most frequently-occurring value
 - ▶ Here, mode = 1.

Boxplots

Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters

Measures of center

x_1	x_2	x_3	x_4	x_5	x_6
0	1	1	2	3	5

► Arithmetic mean:

► $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$

► Here, $\bar{x} = \frac{1}{6}(0 + 1 + 1 + 2 + 3 + 5) = 2$

► Median: $Q(0.5)$.

► A shortcut to calculating $Q(0.5)$ is:

► $Q(0.5) = x_{\lceil n/2 \rceil}$ if n is odd

► $Q(0.5) = (x_{n/2} + x_{n/2+1})/2$ if n is even.

► Here, $Q(0.5) = (1 + 2)/2 = 1.5$

► Mode (of a discrete or categorical dataset)

► the most frequently-occurring value

► Here, mode = 1.

Boxplots

Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters

Measures of spread

	x_1	x_2	x_3	x_4	x_5	x_6
x_i	0	1	1	2	3	5
$\frac{i-.5}{n}$.083	0.25	0.417	0.583	0.75	0.917

▶ Sample variance

$$\bullet s^2 = \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2$$

← unbiased estimator of variance

$$\bullet \text{Here, } s^2 = \frac{1}{6-1} [(0-2)^2 + (1-2)^2 + (1-2)^2 + (2-2)^2 + (3-2)^2 + (5-2)^2] = 3.2$$

▶ Sample standard deviation

$$\bullet s = \sqrt{s^2} = \sqrt{\frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2}$$

$$\bullet \text{Here, } s = \sqrt{3.2} = 1.78885438199983$$

▶ Range

$$\bullet \text{Range} = \text{Maximum} - \text{Minimum}$$

$$\bullet \text{Here, Range} = 5 - 0 = 5$$

▶ Interquartile range

$$\bullet \text{IQR} = Q(0.75) - Q(0.25)$$

$$\bullet \text{Here, IQR} = 3 - 1 = 2.$$

Boxplots

Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters

Your turn: sensitivity to outliers



Compare:

	x_1	x_2	x_3	x_4	x_5	x_6
x_i	0	1	1	2	3	5
$\frac{i-.5}{n}$.083	0.25	0.417	0.583	0.75	0.917

to:

	y_1	y_2	y_3	y_4	y_5	y_6
x_i	0	1	1	2	3	817263489
$\frac{i-.5}{n}$.083	0.25	0.417	0.583	0.75	0.917

outlier

extreme case of right skewed

which measures of center and spread differ drastically between the x_i 's and the y_i 's? Which ones are about the same?

Boxplots

Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters

Answers: sensitivity to outliers

Data	x_i	y_i
Mean	2	136210582.666667
Median	1.5	1.5
Mode	1	1
Sample Variance	3.2	111319934694019232
Sample Std. Dev.	1.78885438199983	333646421.671235
Range	5	817263489
IQR	2	2

Sensitivity of numerical summaries

- ▶ Numerical summaries sensitive to outliers *and* skewness:
 - ▶ Mean *Sample mean*
 - ▶ Sample variance
 - ▶ Sample standard deviation
 - ▶ Range
- ▶ Less sensitive numerical summaries:
 - ▶ Median
 - ▶ Mode
 - ▶ IQR

Outline

Descriptive
Statistics: Part
2/2 (Ch 3)

Yifan Zhu

Boxplots

Boxplots

Quantile-Quantile (QQ) Plots

Quantile-Quantile
(QQ) Plots

Theoretical Quantile-Quantile Plots

Theoretical
Quantile-Quantile
Plots

Numerical Summaries

Numerical
Summaries

Parameters

Parameters

- ▶ **Statistic:** numerical summary of data on the *sample*
- ▶ **Parameter:** numerical summary of a theoretical distribution or data on an entire *population*.
 - ▶ Population mean ("true" mean):
 - ▶ $\mu = \frac{1}{N} \sum_{i=1}^N x_i$ if N the *finite* population size.
 - ▶ $\bar{x} \approx \mu$.
 - ▶ Population variance ("true" variance):
 - ▶ $\sigma^2 = \frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2$ if N the *finite* population size.
 - ▶ $s^2 \approx \sigma^2$.
 - ▶ Population standard deviation ("true" standard deviation):
 - ▶ $\sigma = \sqrt{\frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2}$ if N is the *finite* population size.
 - ▶ $s \approx \sigma$.

Boxplots

Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters

Statistics and parameters

approx

- ▶ **Statistic:** numerical summary of data on the ~~sample~~
- ▶ **Parameter:** numerical ~~summary~~ *characteristic* of a theoretical distribution or data on an entire *population*. *unknown fixed*

- ▶ Population mean ("true" mean):

- ▶ $\mu = \frac{1}{N} \sum_{i=1}^N x_i$ if N the *finite* population size.
- ▶ $\bar{x} \approx \mu$.

- ▶ Population variance ("true" variance):

- ▶ $\sigma^2 = \frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2$ if N the *finite* population size.
- ▶ $s^2 \approx \sigma^2$.

- ▶ Population standard deviation ("true" standard deviation):

- ▶ $\sigma = \sqrt{\frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2}$ if N is the *finite* population size.
- ▶ $s \approx \sigma$.

Boxplots

Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters

- ▶ **Statistic:** numerical summary of data on the *sample*
- ▶ **Parameter:** numerical summary of a theoretical distribution or data on an entire *population*.
 - ▶ Population mean ("true" mean):
 - ▶ $\mu = \frac{1}{N} \sum_{i=1}^N x_i$ if N the *finite* population size.
 - ▶ $\bar{x} \approx \mu$.
 - ▶ Population variance ("true" variance):
 - ▶ $\sigma^2 = \frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2$ if N the *finite* population size.
 - ▶ $s^2 \approx \sigma^2$.
 - ▶ Population standard deviation ("true" standard deviation):
 - ▶ $\sigma = \sqrt{\frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2}$ if N is the *finite* population size.
 - ▶ $s \approx \sigma$.

Boxplots

Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters

- ▶ **Statistic:** numerical summary of data on the *sample*
- ▶ **Parameter:** numerical summary of a theoretical distribution or data on an entire *population*.
 - ▶ Population mean ("true" mean):
 - ▶ $\mu = \frac{1}{N} \sum_{i=1}^N x_i$ if N the *finite* population size.
 - ▶ $\bar{x} \approx \mu$.
 - ▶ Population variance ("true" variance):
 - ▶ $\sigma^2 = \frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2$ if N the *finite* population size.
 - ▶ $s^2 \approx \sigma^2$.
 - ▶ Population standard deviation ("true" standard deviation):
 - ▶ $\sigma = \sqrt{\frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2}$ if N is the *finite* population size.
 - ▶ $s \approx \sigma$.

Boxplots

Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters

- ▶ **Statistic:** numerical summary of data on the *sample*
- ▶ **Parameter:** numerical summary of a theoretical distribution or data on an entire *population*.

- ▶ Population mean ("true" mean): $N = \infty, E(X)$

- ▶ $\mu = \frac{1}{N} \sum_{i=1}^N x_i$ if N the *finite* population size.

- ▶ $\bar{x} \approx \mu$. $n \ll N, \bar{x} \rightarrow \mu, n \rightarrow \infty$

- ▶ Population variance ("true" variance):

- ▶ $\sigma^2 = \frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2$ if N the *finite* population size.

- ▶ $s^2 \approx \sigma^2$.

- ▶ Population standard deviation ("true" standard deviation):

- ▶ $\sigma = \sqrt{\frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2}$ if N is the *finite* population size.

- ▶ $s \approx \sigma$.

Boxplots

Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters

- ▶ **Statistic:** numerical summary of data on the *sample*
- ▶ **Parameter:** numerical summary of a theoretical distribution or data on an entire *population*.
 - ▶ Population mean (“true” mean):
 - ▶ $\mu = \frac{1}{N} \sum_{i=1}^N x_i$ if N the *finite* population size.
 - ▶ $\bar{x} \approx \mu$.
 - ▶ Population variance (“true” variance):
 - ▶ $\sigma^2 = \frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2$ if N the *finite* population size.
 - ▶ $s^2 \approx \sigma^2$.
 - ▶ Population standard deviation (“true” standard deviation):
 - ▶ $\sigma = \sqrt{\frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2}$ if N is the *finite* population size.
 - ▶ $s \approx \sigma$.

Boxplots

Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters

- ▶ **Statistic:** numerical summary of data on the *sample*
- ▶ **Parameter:** numerical summary of a theoretical distribution or data on an entire *population*.
 - ▶ Population mean (“true” mean):
 - ▶ $\mu = \frac{1}{N} \sum_{i=1}^N x_i$ if N the *finite* population size.
 - ▶ $\bar{x} \approx \mu$.
 - ▶ Population variance (“true” variance):
 - ▶ $\sigma^2 = \frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2$ if N the *finite* population size.
 - ▶ $s^2 \approx \sigma^2$.
 - ▶ Population standard deviation (“true” standard deviation):
 - ▶ $\sigma = \sqrt{\frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2}$ if N is the *finite* population size.
 - ▶ $s \approx \sigma$.

Boxplots

Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters

- ▶ **Statistic:** numerical summary of data on the *sample*
- ▶ **Parameter:** numerical summary of a theoretical distribution or data on an entire *population*.
 - ▶ Population mean ("true" mean):
 - ▶ $\mu = \frac{1}{N} \sum_{i=1}^N x_i$ if N the *finite* population size.
 - ▶ $\bar{x} \approx \mu$.
 - ▶ Population variance ("true" variance):
 - ▶ $\sigma^2 = \frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2$ if N the *finite* population size.
 - ▶ $s^2 \approx \sigma^2$. Var(X)
 - ▶ Population standard deviation ("true" standard deviation):
 - ▶ $\sigma = \sqrt{\frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2}$ if N is the *finite* population size.
 - ▶ $s \approx \sigma$.

Boxplots

Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters

- ▶ **Statistic:** numerical summary of data on the *sample*
- ▶ **Parameter:** numerical summary of a theoretical distribution or data on an entire *population*.
 - ▶ Population mean (“true” mean):
 - ▶ $\mu = \frac{1}{N} \sum_{i=1}^N x_i$ if N the *finite* population size.
 - ▶ $\bar{x} \approx \mu$.
 - ▶ Population variance (“true” variance):
 - ▶ $\sigma^2 = \frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2$ if N the *finite* population size.
 - ▶ $s^2 \approx \sigma^2$.
 - ▶ Population standard deviation (“true” standard deviation):
 - ▶ $\sigma = \sqrt{\frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2}$ if N is the *finite* population size.
 - ▶ $s \approx \sigma$.

Boxplots

Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters

- ▶ **Statistic:** numerical summary of data on the *sample*
- ▶ **Parameter:** numerical summary of a theoretical distribution or data on an entire *population*.
 - ▶ Population mean (“true” mean):
 - ▶ $\mu = \frac{1}{N} \sum_{i=1}^N x_i$ if N the *finite* population size.
 - ▶ $\bar{x} \approx \mu$.
 - ▶ Population variance (“true” variance):
 - ▶ $\sigma^2 = \frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2$ if N the *finite* population size.
 - ▶ $s^2 \approx \sigma^2$.
 - ▶ Population standard deviation (“true” standard deviation):
 - ▶ $\sigma = \sqrt{\frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2}$ if N is the *finite* population size.
 - ▶ $s \approx \sigma$.

Boxplots

Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters

- ▶ **Statistic:** numerical summary of data on the *sample*
- ▶ **Parameter:** numerical summary of a theoretical distribution or data on an entire *population*.
 - ▶ Population mean (“true” mean):
 - ▶ $\mu = \frac{1}{N} \sum_{i=1}^N x_i$ if N the *finite* population size.
 - ▶ $\bar{x} \approx \mu$.
 - ▶ Population variance (“true” variance):
 - ▶ $\sigma^2 = \frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2$ if N the *finite* population size.
 - ▶ $s^2 \approx \sigma^2$.
 - ▶ Population standard deviation (“true” standard deviation):
 - ▶ $\sigma = \sqrt{\frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2}$ if N is the *finite* population size.
 - ▶ $s \approx \sigma$.

Boxplots

Quantile-Quantile
(QQ) Plots

Theoretical
Quantile-Quantile
Plots

Numerical
Summaries

Parameters