

Discrete Random Variables (Ch. 5.1)

Yifan Zhu

Iowa State University

Outline

What is a random variable?

What is a random
variable?

Probability

Probability

Probability Mass Functions (pmf)

Probability Mass
Functions (pmf)

Cumulative Distribution Functions (cdf)

Cumulative
Distribution
Functions (cdf)

Expected Value

Expected Value

Variance and Standard Deviation

Variance and
Standard Deviation

What is a random variable?

- ▶ **Random variable**; a quantity that can be thought of as dependent on chance phenomena.
 - ▶ X = the value of a coin toss (heads or tails). ¹ ⁰
 - ▶ Z = the amount of torque required to loosen the next bolt.
 - ▶ T = the time you'll have to wait for the next bus home.
 - ▶ N = the number of defective widgets in manufacturing process in a day.
 - ▶ S = the number of provoked shark attacks off the coast of Florida next year. ^{0, 1, 2, ...}
- ▶ Two types:
 - ▶ **Discrete random variable**: one that can only take on a set of isolated points (X , N , and S). ^{integers}
 - ▶ **Continuous random variable**: one that can fall in an interval of real numbers (T and Z).

Discrete random variables

Support

- ▶ A discrete random variable has a list of possible values:
 - ▶ X = roll of a 6-sided fair die = 1, 2, 3, 4, 5, or 6. }
 - ▶ Y = roll of a 6-sided *unfair* die = 1, 2, 3, 4, 5, or 6. }
- ▶ But how do you distinguish between X and Y ?

Outline

What is a random variable?

What is a random
variable?

Probability

Probability

Probability Mass Functions (pmf)

Probability Mass
Functions (pmf)

Cumulative Distribution Functions (cdf)

Cumulative
Distribution
Functions (cdf)

Expected Value

Expected Value

Variance and Standard Deviation

Variance and
Standard Deviation

Probability

repeat tossing n times.

$$\frac{\text{\# of 3's out of } n \text{ times}}{n} \rightarrow \frac{1}{6}, \quad n \rightarrow \infty$$

► $P(X = x)$, the **probability** that X equals x , is the fraction of times that X will land on x

1. We expect a fair die to land the number 3 roughly one out of every 6 tosses. Thus, $P(X = 3) = 1/6$
2. Suppose the unfair die is weighted so that the number 3 only lands one out of every 22 tosses. Then, $P(Y = 3) = 1/22$.

What is a random variable?

Probability

Probability Mass Functions (pmf)

Cumulative Distribution Functions (cdf)

Expected Value


Variance and Standard Deviation

► X has the following probabilities:

Support x \rightarrow *distribution of X*


x	1	2	3	4	5	6
$P(X = x)$	$1/6$	$1/6$	$1/6$	$1/6$	$1/6$	$1/6$

prob.



► Say Y has the probabilities:

y	1	2	3	4	5	6
$P(Y = y)$	$5/22$	$7/44$	$1/22$	$7/44$	$2/11$	$5/22$



What is a random variable?

Probability

Probability Mass Functions (pmf)

Cumulative Distribution Functions (cdf)

Expected Value

Variance and Standard Deviation

What is a random variable?

Probability

Probability Mass Functions (pmf)


Cumulative Distribution Functions (cdf)

Expected Value

Variance and Standard Deviation

- S , the number of provoked shark attacks off FL next year, has infinite number of possible values. Here is one possible (made up) distribution:

s	1	2	3	...	k	...
$P(S = s)$	$\frac{6}{\pi^2}$	$\frac{1}{2} \frac{6}{\pi^2}$	$\frac{1}{4} \frac{6}{\pi^2}$...	$\frac{1}{2^k} \frac{6}{\pi^2}$...



 →
 mean ↑
 — > median

Outline

What is a random variable?

What is a random
variable?

Probability

Probability

Probability Mass Functions (pmf)

Probability Mass
Functions (pmf)

Cumulative Distribution Functions (cdf)

Cumulative
Distribution
Functions (cdf)

Expected Value

Expected Value

Variance and Standard Deviation

Variance and
Standard Deviation

Probability mass functions (pmf)


- ▶ The **probability mass function (pmf)** $f(x)$ of a random variable X is just $P(X = x)$ + support of X .
 - ▶ X has $f(x) = 1/6$, $x = 1, 2, \dots, 6$.
 - ▶ S has $f(s) = \frac{1}{2^s \pi^2}$, $s = 1, 2, 3, \dots$.
- ▶ We could also write f_X for the pmf of X and f_S for the pmf of S .
- ▶ Rules of the pmf f :
 - ▶ $f(x) \geq 0$ for all x . $f(x) > 0$, x in the support.
 - ▶ $\sum_x f(x) = 1$.

Your turn: calculating probabilities

- Let Z = the torque, rounded to the nearest integer, required to loosen the next bolt on an apparatus.

z	11	12	13	14	15
$f(z) = P(Z = z)$	0.03	0.03	0.03	0.06	0.26

z	16	17	18	19	20
$f(z) = P(Z = z)$	0.09	0.12	0.20	0.15	0.03



- Calculate:

1. $P(Z \leq 14)$
2. $P(Z > 16)$
3. $P(Z \text{ is an even number})$
4. $P(Z \text{ in } \{15, 16, 18\})$

Answers: calculating probabilities

What is a random
variable?

Probability

Probability Mass
Functions (pmf)

Cumulative
Distribution
Functions (cdf)

Expected Value

Variance and
Standard Deviation

1.

event



$$\begin{aligned} P(Z \leq 14) &= P(Z = 11 \text{ or } Z = 12 \text{ or } Z = 13 \text{ or } Z = 14) \\ &= P(Z = 11) + P(Z = 12) + P(Z = 13) + P(Z = 14) \\ &= f(11) + f(12) + f(13) + f(14) \\ &= 0.03 + 0.03 + 0.03 + 0.06 \\ &= 0.15 \end{aligned}$$

2.

$$\begin{aligned} P(Z > 16) &= P(Z = 17 \text{ or } Z = 18 \text{ or } Z = 19 \text{ or } Z = 20) \\ &= P(Z = 17) + P(Z = 18) + P(Z = 19) + P(Z = 20) \\ &= f(17) + f(18) + f(19) + f(20) \\ &= 0.12 + 0.20 + 0.15 + 0.03 \\ &= 0.5 \end{aligned}$$

Answers: calculating probabilities

3.

$$\begin{aligned}P(\underline{Z \text{ even}}) &= P(Z = 12 \text{ or } Z = 14 \text{ or } Z = 16 \text{ or } Z = 18 \text{ or } Z = 20) \\&= P(\underline{Z = 12}) + P(\underline{Z = 14}) + P(\underline{Z = 16}) + P(\underline{Z = 18}) \\&\quad + P(\underline{Z = 20}) \\&= f(12) + f(\underline{14}) + f(16) + f(18) + f(20) \\&= 0.03 + 0.06 + 0.09 + 0.20 + 0.03 \\&= \underline{0.41}\end{aligned}$$

4.

$$\begin{aligned}P(Z \text{ in } \{15, 16, 18\}) &= P(Z = 15 \text{ or } Z = 16 \text{ or } Z = 18) \\&= P(Z = 15) + P(Z = 16) + P(Z = 18) \\&= f(15) + f(16) + f(18) \\&= 0.26 + 0.09 + 0.02 \\&= 0.37\end{aligned}$$

Outline

What is a random variable?

Probability

Probability Mass Functions (pmf)

Cumulative Distribution Functions (cdf)

Expected Value

Variance and Standard Deviation

The cumulative distribution function (cdf)

- **Cumulative distribution function (cdf):** a function, F , defined by:

$$f(x) = P(X = x)$$

$$F(x) = P(X \leq x)$$

$$= \sum_{z \leq x} f(z)$$

- F has the following properties:

- $F(x) \geq 0$ for all real numbers x .
- F is monotonically increasing. $x_1 < x_2, F(x_1) \leq F(x_2)$.
not strictly monotonically increasing
- $\lim_{x \rightarrow -\infty} F(x) = 0$
- $\lim_{x \rightarrow \infty} F(x) = 1$

$$x \rightarrow -\infty, \sum_{\emptyset} f(z) = 0$$

$$x \rightarrow \infty, \sum_z f(z) = 1$$

Example: torque random variable, Z

z , Torque	$f(z) = P[Z = z]$	$F(z) = P[Z \leq z]$
11	.03	.03
<u>12</u>	.03	<u>.06</u>
13	.03	<u>.09</u>
14	.06	.15
<u>15</u>	.26	<u>.41</u>
16	.09	<u>.50</u>
17	.12	.62
18	.20	.82
19	.15	.97
20	.03	1.00

What is a random
variable?

Probability

Probability Mass
Functions (pmf)

Cumulative
Distribution
Functions (cdf)

Expected Value

Variance and
Standard Deviation

Example: torque random variable, Z

What is a random
variable?


Probability

Probability Mass
Functions (pmf)

Cumulative
Distribution
Functions (cdf)

Expected Value

Variance and
Standard Deviation



$$\varepsilon = 10^{-10}, \quad F(15 - \varepsilon) = F(14) = 0.15.$$
$$P(Z \leq 15 - \varepsilon) = P(Z \leq 14) = F(14)$$

Your turn: calculating probabilities

What is a random
variable?

Probability


Probability Mass
Functions (pmf)

Cumulative
Distribution
Functions (cdf)

Expected Value

Variance and
Standard Deviation

z	11	12	13	14	15
$F(z) = P(Z \leq z)$	0.03	0.06	0.09	0.15	0.41
z	16	17	18	19	20
$F(z) = P(Z \leq z)$	0.50	0.62	0.82	0.97	1



► Using the cdf only, calculate:

1. $F(10.7)$
2. $P(Z \leq 15.5)$
3. $P(12.1 < Z \leq 14)$
4. $P(15 \leq Z < 18)$

Answers: calculating probabilities

What is a random
variable?

Probability

Probability Mass
Functions (pmf)

Cumulative
Distribution
Functions (cdf)

Expected Value

Variance and
Standard Deviation

1. $F(10.7) = P(Z \leq 10.7) = 0$
2. $P(Z \leq 15.5) = P(Z \leq 15) = 0.41$
- 3.

$F(15)$

$$P(12.1 < Z \leq 14) = P(Z = 13 \text{ or } 14)$$

$$= f(14) + f(13)$$

$$= [f(14) + f(13) + f(12) + f(11)]$$

$$- [f(12) + f(11)]$$

$$= P(Z \leq 14) - P(Z \leq 12)$$

$$= F(14) - F(12)$$

$$= 0.15 - 0.06$$

$$= 0.09$$

$$P(Z \leq 14) - P(Z \leq 12.1) = P(Z \leq 12)$$

Answers: calculating probabilities

$$P(Z < 18)$$

$$= P(\underline{Z < 15}) + P(\underline{15 \leq Z < 18})$$

4.

$$\begin{aligned} P(15 \leq Z < 18) &= P(Z = 15, 16, \text{ or } 17) \\ &= \underline{P(Z \leq 17)} - \underline{P(Z \leq 14)} \\ &= F(17) - F(14) \\ &= 0.62 - 0.15 \\ &= 0.47 \end{aligned}$$

$$\begin{aligned} &\underline{P(Z < 18) - P(Z < 15)} \\ &\quad \parallel \qquad \qquad \parallel \\ &P(Z \leq 17) \qquad P(Z \leq 14) \end{aligned}$$

Your turn: drawing the cdf

- Say we have a random variable Q with pmf:

q	1	2	3	7
$f(q)$	0.34	0.1	0.22	0.34

- Draw the cdf.

Support of Q : $\{1, 2, 3, 7\}$.

1. break the real numbers up into 5 parts:


$$(-\infty, 1): F(q) = P(Q \leq q) = 0$$

$$[1, 2): F(q) = P(Q \leq q) = P(Q=1) = 0.34$$

$$[2, 3): F(q) = P(Q \leq q) = P(Q=1) + P(Q=2) = 0.57$$

$$[3, 7): \dots = P(Q=1) + P(Q=2) + P(Q=3) = 0.57$$

$$[7, +\infty): \dots = P(Q=1) + P(Q=2) + P(Q=3) + P(Q=7) = 1$$



Answer: drawing the cdf

What is a random
variable?


Probability

Probability Mass
Functions (pmf)

Cumulative
Distribution
Functions (cdf)

Expected Value

Variance and
Standard Deviation



Outline

What is a random variable?

Probability

Probability Mass Functions (pmf)

Cumulative Distribution Functions (cdf)

Expected Value

Variance and Standard Deviation

What is a random
variable?

Probability

Probability Mass
Functions (pmf)

Cumulative
Distribution
Functions (cdf)

Expected Value

Variance and
Standard Deviation

Expected Value

- ▶ The **expected value** $E(X)$ (also called μ) of a random variable X is given by:

$$\sum_x x \cdot f(x)$$

- ▶ When X is the roll of a fair die,

$$\begin{aligned} E(X) &= 1f(1) + 2f(2) + 3f(3) + 4f(4) + 5f(5) + 6f(6) \\ &= 1(1/6) + 2(1/6) + 3(1/6) + 4(1/6) + 5(1/6) + 6(1/6) \\ &= \frac{1 + 2 + 3 + 4 + 5 + 6}{6} \\ &= \underline{3.5} \end{aligned}$$

- ▶ $E(X)$ is a *weighted average* of the possible values of X , weighted by their probabilities.
- ▶ $E(X)$ is the **mean of the distribution** of X

Property of Expected Value

- ▶ Expected value of the function of a discrete random variable:

$$E(g(X)) = \sum_x g(x) \cdot f(x)$$


- ▶ Linearity (true for both discrete and continuous random variables):

$$E(X + Y) = E(X) + E(Y)$$

$$E(\underbrace{aX}_{\text{constants}} + \underbrace{b}_{\text{constants}}) = aE(X) + b$$

Your turn: expected value

y	1	2	3	4	5	6
$P(Y = y)$	5/22	7/44	1/22	7/44	2/11	5/22



- Calculate $E(Y)$, the expected value of a toss of the *unfair* die.

What is a random variable?

Probability

Probability Mass Functions (pmf)

Cumulative Distribution Functions (cdf)

Expected Value


Variance and Standard Deviation

Answer: expected value



$$\begin{aligned} E(Y) &= 1(5/22) + 2(7/44) + 3(1/22) \\ &\quad + 4(7/44) + 5(2/11) + 6(5/22) \\ &= \underline{3.5909091} \end{aligned}$$

- ▶ The average roll of the unfair die is 3.5909.
- ▶ $E(Y)$ is the mean of the distribution of Y .



Your turn: expected value

- Calculate $E(2X + Y)$, the expected value of the sum of two tosses of a fair die and a toss of the unfair die.

$$E(2X + Y)$$

$$= 2E(X) + E(Y)$$

$$= 2 \times 3.5 + 3.5909$$

$$= 10.5909.$$

What is a random variable?

Probability

Probability Mass Functions (pmf)

Cumulative Distribution Functions (cdf)

Expected Value

Variance and Standard Deviation

Answer: expected value

$$E(2X + Y) = 2E(X) + Y = 2 \times 3.5 + 3.5909 = 10.5909$$

- The average of the sum of two tosses of a fair die and a toss of the unfair die is 10.5909.

Your turn: expected value

z	11	12	13	14	15
$f(z) = P(Z = z)$	0.03	0.03	0.03	0.06	0.26
z	16	17	18	19	20
$f(z) = P(Z = z)$	0.09	0.12	0.20	0.15	0.03

- Calculate $E(Z)$, the expected value of the torque required to loosen the next bolt.

Answer: expected value

$$\begin{aligned}E(Z) &= 11(0.03) + 12(0.03) + 13(0.03) + 14(0.06) + 15(0.26) \\&= 16(0.09) + 17(0.12) + 18(0.20) + 19(0.15) + 20(0.03) \\&= \underline{16.35}\end{aligned}$$

- The average torque required to loosen the next bolt is 16.35 units.

Outline

What is a random variable?

What is a random
variable?

Probability

Probability

Probability Mass Functions (pmf)

Probability Mass
Functions (pmf)

Cumulative Distribution Functions (cdf)

Cumulative
Distribution
Functions (cdf)

Expected Value

Expected Value

Variance and Standard Deviation

Variance and
Standard Deviation

Variance

- **Variance:** the variance $\text{Var}(X)$ (also called σ^2) of a random variable X is given by:

$$\begin{aligned}\text{Var}(X) &= E((X - E(X))^2) \\ &= \sum_x (x - E(X))^2 f(x)\end{aligned}$$

$$\text{Var}(X) = E((X - E(X))^2)$$

$$\begin{aligned}&= E(X^2 - 2E(X) \cdot X + (E(X))^2) = E(X^2) - 2E(X)E(X) + (E(X))^2 \\ &= E(X^2) - (E(X))^2\end{aligned}$$

- Shortcut formulas:

$$\begin{aligned}\text{Var}(X) &= \left[\sum_x x^2 f(x) \right] - (E(X))^2 \\ &= E(X^2) - E^2(X)\end{aligned}$$

- The variance is the average squared deviation of random variable from its mean

Example: calculating the variance

q	1	2	3	7
$f(q)$	0.34	0.1	0.22	0.34

► Long way:

$$\begin{aligned} E(Q) &= 1(0.34) + 2(0.1) + 3(0.22) + 7(0.34) \\ &= 3.58 \end{aligned}$$

$$\begin{aligned} \text{Var}(Q) &= (1 - 3.58)^2 0.34 + (2 - 3.58)^2 0.1 \\ &\quad + (3 - 3.58)^2 0.22 + (7 - 3.58)^2 0.34 \\ &= 6.56 \end{aligned}$$

► Short way:

$$\begin{aligned} E(Q^2) &= \sum_q q^2 f(q) \\ &= 1(0.34) + 4(0.1) + 9(0.22) + 49(0.34) \\ &= 19.38 \end{aligned}$$

$$\begin{aligned} \text{Var}(Q) &= E(Q^2) - E^2(Q) \\ &= 19.38 - 3.58^2 \\ &= 6.56 \end{aligned}$$

What is a random variable?

Probability

Probability Mass Functions (pmf)

Cumulative Distribution Functions (cdf)

Expected Value

Variance and Standard Deviation

Your turn: calculating the variance

x	1	2	3	4	5	6
$f(x)$	1/6	1/6	1/6	1/6	1/6	1/6

- Calculate $\text{Var}(X)$
- Calculate $\text{SD}(X)$

$$\text{SD} = \sqrt{\text{Var}(X)}$$

Your turn: answers

What is a random
variable?

Probability

Probability Mass
Functions (pmf)

Cumulative
Distribution
Functions (cdf)

Expected Value

Variance and
Standard Deviation



$$\begin{aligned} E(X) &= 1(1/6) + 2(1/6) + 3(1/6) + 4(1/6) + 5(1/6) + 6(1/6) \\ &= 3.5 \end{aligned}$$

$$\begin{aligned} E(X^2) &= \sum_{x=1}^6 x^2 f(x) \\ &= 1^2(1/6) + 2^2(1/6) + 3^2(1/6) + 4^2(1/6) + 5^2(1/6) + 6^2(1/6) \\ &= 15.17 \end{aligned}$$

$$\begin{aligned} \text{Var}(X) &= E(X^2) - E^2(X) \\ &= 15.17 - 3.5^2 \\ &= 2.92 \end{aligned}$$

► $SD(X) = \sqrt{2.92} = 1.7088007$