

Functions of Several Random Variables (Ch. 5.5)

Yifan Zhu

Iowa State University

Outline

Functions of
Several Random
Variables (Ch. 5.5)

Yifan Zhu

Functions of
Several Random
Variables

Expectations and
variances of linear
combinations

Approximating the
Mean and Variance
of a Function

The Central Limit
Theorem

Functions of Several Random Variables

Expectations and variances of linear combinations

Approximating the Mean and Variance of a Function

The Central Limit Theorem

Functions of several random variables

- ▶ We often consider functions of random variables of the form:

$$U = g(X, Y, \dots, Z)$$

where X, Y, \dots, Z are random variables.

- ▶ U is itself a random variable.

Example: connecting steel parts

- ▶ Suppose that a steel plate with nominal thickness .15 in. is to rest in a groove of nominal width .155 in., machined on the surface of a steel block.

Relative Frequency Distribution of Plate Thicknesses

Plate Thickness (in.)	Relative Frequency
.148	.4
.149	.3
.150	.3

Relative Frequency Distribution of Slot Widths

Slot Width (in.)	Relative Frequency
.153	.2
.154	.2
.155	.4
.156	.2

- ▶ X = plate thickness
- ▶ Y = slot width
- ▶ $U = Y - X$, the “wiggle room” of the plate

The distributions of X , Y , and U

The Probability Function for the
Clearance $U = Y - X$

Marginal and Joint Probabilities for X and Y

$y \backslash x$.148	.149	.150	$f_Y(y)$
.156	.08	.06	.06	.2
.155	.16	.12	.12	.4
.154	.08	.06	.06	.2
.153	.08	.06	.06	.2
$f_X(x)$.4	.3	.3	

u	$f(u)$
.003	.06
.004	.12 = .06 + .06
.005	.26 = .08 + .06 + .12
.006	.26 = .08 + .12 + .06
.007	.22 = .16 + .06
.008	.08

- Determining the distribution of U is difficult in the continuous case.

Outline

Functions of
Several Random
Variables (Ch. 5.5)

Yifan Zhu

Functions of
Several Random
Variables

Expectations and
variances of linear
combinations

Approximating the
Mean and Variance
of a Function

The Central Limit
Theorem

Functions of Several Random Variables

Expectations and variances of linear combinations

Approximating the Mean and Variance of a Function

The Central Limit Theorem

Expectations and variances of linear combinations

Functions of
Several Random
Variables (Ch. 5.5)

Yifan Zhu

Functions of
Several Random
Variables

Expectations and
variances of linear
combinations

Approximating the
Mean and Variance
of a Function

The Central Limit
Theorem

- ▶ X_1, X_2, \dots, X_n are independent random variables and

$$Y = a_0 + a_1X_1 + a_2X_2 + \cdots + a_nX_n$$

then:

$$\begin{aligned} E(Y) &= E(a_0 + a_1X_1 + a_2X_2 + \cdots + a_nX_n) \\ &= a_0 + a_1E(X_1) + a_2E(X_2) + \cdots + a_nE(X_n) \end{aligned}$$

$$\begin{aligned} \text{Var}(Y) &= \text{Var}(a_0 + a_1X_1 + a_2X_2 + \cdots + a_nX_n) \\ &= a_1^2 \cdot \text{Var}(X_1) + a_2^2 \cdot \text{Var}(X_2) + \cdots + a_n^2 \cdot \text{Var}(X_n) \end{aligned}$$

Your turn: linear combinations

- Say we have two independent random variables X and Y with $E(X) = 3.3$, $Var(X) = 1.91$, $E(Y) = 25$, and $Var(Y) = 65$.
- Find:

$$E(3 + 2X - 3Y)$$

$$E(-4X + 3Y)$$

$$E(-4X - 6Y)$$

$$Var(3 + 2X - 3Y)$$

$$Var(2X - 5Y)$$

$$Var(-4X - 6Y)$$

Answers: linear combinations

Functions of
Several Random
Variables (Ch. 5.5)

Yifan Zhu

Functions of
Several Random
Variables

Expectations and
variances of linear
combinations

Approximating the
Mean and Variance
of a Function

The Central Limit
Theorem

$$\begin{aligned}E(3 + 2X - 3Y) &= 3 + 2E(X) - 3E(Y) \\&= 3 + 2 \cdot 3.3 - 3 \cdot 25 \\&= -65.4\end{aligned}$$

$$\begin{aligned}E(-4X + 3Y) &= -4E(X) + 3E(Y) \\&= -4 \cdot 3.3 + 3 \cdot 25 \\&= 61.8\end{aligned}$$

$$\begin{aligned}E(-4X - 6Y) &= -4 \cdot E(X) - 6 \cdot E(Y) \\&= -4 \cdot 3.3 - 6 \cdot 25 \\&= -163.2\end{aligned}$$

Answers: linear combinations

$$\begin{aligned}\text{Var}(3 + 2X - 3Y) &= 2^2 \cdot \text{Var}(X) + (-3)^2 \text{Var}(Y) \\ &= 4 \cdot 1.91 + 9 \cdot 65 \\ &= 592.64\end{aligned}$$

$$\begin{aligned}\text{Var}(2X - 5Y) &= 2^2 \cdot \text{Var}(X) + (-5)^2 \text{Var}(Y) \\ &= 4 \cdot 1.91 + 25 \cdot 65 \\ &= 1632.64\end{aligned}$$

$$\begin{aligned}\text{Var}(-4X - 6Y) &= (-4)^2 \cdot \text{Var}(X) + (-6)^2 \text{Var}(Y) \\ &= 16 \cdot 1.91 + 36 \cdot 65 \\ &= 2370.56\end{aligned}$$

Your turn: more linear combinations

Functions of
Several Random
Variables (Ch. 5.5)

Yifan Zhu

Functions of
Several Random
Variables

Expectations and
variances of linear
combinations

Approximating the
Mean and Variance
of a Function

The Central Limit
Theorem

- ▶ Say $X \sim \text{Binomial}(n = 10, p = 0.5)$ and $Y \sim \text{Poisson}(\lambda = 3)$.
- ▶ Calculate:

$$E(5 + 2X - 7Y)$$

$$\text{Var}(5 + 2X - 7Y)$$

Answer: more linear combinations

- First, note that:

$$E(X) = np = 10 \cdot 0.5 = 5$$

$$E(Y) = \lambda = 3$$

$$\text{Var}(X) = np(1 - p) = 10(0.5)(1 - 0.5) = 2.5$$

$$\text{Var}(Y) = \lambda = 3$$

Now, we can calculate:

$$\begin{aligned} E(5 + 2X - 7Y) &= 5 + 2E(X) - 7E(Y) \\ &= 5 + 2 \cdot 5 - 7 \cdot 3 \\ &= -6 \end{aligned}$$

$$\begin{aligned} \text{Var}(5 + 2X - 7Y) &= 2^2 \cdot \text{Var}(X) + (-7)^2 \cdot \text{Var}(Y) \\ &= 4 \cdot 2.5 + 49 \cdot 3 \\ &= 157 \end{aligned}$$

Outline

Functions of
Several Random
Variables (Ch. 5.5)

Yifan Zhu

Functions of
Several Random
Variables

Expectations and
variances of linear
combinations

Approximating the
Mean and Variance
of a Function

The Central Limit
Theorem

Functions of Several Random Variables

Expectations and variances of linear combinations

Approximating the Mean and Variance of a Function

The Central Limit Theorem

Approximating $E(U)$ and $Var(U)$ when determining $f_U(u)$ is too hard

Functions of
Several Random
Variables (Ch. 5.5)

Yifan Zhu

Functions of
Several Random
Variables

Expectations and
variances of linear
combinations

Approximating the
Mean and Variance
of a Function

The Central Limit
Theorem


- ▶ If X, Y, \dots, Z are independent, g is well-behaved, and the variances $Var(X), Var(Y), \dots, Var(Z)$ are small enough, then $U = g(X, Y, \dots, Z)$ has:

$$E(U) \approx g(E(X), E(Y), \dots, E(Z))$$

$$Var(U) \approx \left(\frac{\partial g}{\partial x}\right)^2 Var(X) + \left(\frac{\partial g}{\partial y}\right)^2 Var(Y) + \dots + \left(\frac{\partial g}{\partial z}\right)^2 Var(Z)$$

- ▶ These formulas are often called the **propagation of error formulas**.

Example: an electric circuit



- ▶ R is the total resistance of the circuit.
- ▶ R_1 , R_2 , and R_3 are the resistances of resistors 1, 2, and 3, respectively.
- ▶ $E(R_i) = 100$, $\text{Var}(R_i) = 2$, $i = 1, 2, 3$.

$$R = g(R_1, R_2, R_3) = R_1 + \frac{R_2 R_3}{R_2 + R_3}$$

Example: an electric circuit

$$E(R) \approx g(100, 100, 100) = 100 + \frac{(100)(100)}{100 + 100} = 150\Omega$$

$$\frac{\partial g}{\partial r_1} = 1$$

$$\frac{\partial g}{\partial r_2} = \frac{(r_2 + r_3)r_3 - r_2r_3}{(r_2 + r_3)^2} = \frac{r_3^2}{(r_2 + r_3)^2}$$

$$\frac{\partial g}{\partial r_3} = \frac{(r_2 + r_3)r_2 - r_2r_3}{(r_2 + r_3)^2} = \frac{r_2^2}{(r_2 + r_3)^2}$$

$$\begin{aligned}\text{Var}(R) &\approx (1)^2(2)^2 + \left(\frac{(100)^2}{(100 + 100)^2}\right)^2 (2)^2 + \left(\frac{(100)^2}{(100 + 100)^2}\right)^2 (2)^2 \\ &= 4.5\end{aligned}$$

$$\text{SD}(R)\sqrt{4.5} \approx 2.12\Omega$$

Outline

Functions of
Several Random
Variables (Ch. 5.5)

Yifan Zhu

Functions of
Several Random
Variables

Expectations and
variances of linear
combinations

Approximating the
Mean and Variance
of a Function

The Central Limit
Theorem

Functions of Several Random Variables

Expectations and variances of linear combinations

Approximating the Mean and Variance of a Function

The Central Limit Theorem

iid random variables.

- ▶ **Identically Distributed:** Random variables X_1, X_2, \dots, X_n are identically distributed if they have the same probability distribution.
- ▶ **“iid”:** Random variables X_1, X_2, \dots, X_n are iid if they are **I**ndependent and **I**dentically **D**istributed.

Averages of iid random variables

- ▶ X_1, X_2, \dots, X_n are iid with expectation μ and variance σ^2 .
- ▶ Derive:

$$E(\bar{X})$$

$$\text{Var}(\bar{X})$$

where:

$$\bar{X} = \frac{X_1 + X_2 + \dots + X_n}{n}$$

the mean of the X_i 's.

Averages of iid random variables

Functions of
Several Random
Variables (Ch. 5.5)

Yifan Zhu

Functions of
Several Random
Variables

Expectations and
variances of linear
combinations

Approximating the
Mean and Variance
of a Function

The Central Limit
Theorem

$$\begin{aligned} E(\bar{X}) &= E\left(\frac{X_1 + X_2 + \cdots + X_n}{n}\right) \\ &= \frac{1}{n}E(X_1) + \frac{1}{n}E(X_2) + \cdots + \frac{1}{n}E(X_n) \\ &= \underbrace{\frac{1}{n}\mu + \frac{1}{n}\mu + \cdots + \frac{1}{n}\mu}_{n \text{ times}} \\ &= n \cdot \frac{1}{n}\mu \\ &= \boxed{\mu} \end{aligned}$$

- Remember $E(\bar{X}) = \mu$: it's an important result.

Answers: averages of iid random variables

$$\begin{aligned} \text{Var}(\bar{X}) &= \text{Var}\left(\frac{X_1 + X_2 + \cdots + X_n}{n}\right) \\ &= \left(\frac{1}{n}\right)^2 \text{Var}(X_1) + \left(\frac{1}{n}\right)^2 \text{Var}(X_2) + \cdots + \left(\frac{1}{n}\right)^2 \cdot \text{Var}(X_n) \\ &= \underbrace{\frac{1}{n^2}\sigma^2 + \frac{1}{n^2}\sigma^2 + \cdots + \frac{1}{n^2}\sigma^2}_{n \text{ times}} \\ &= n \cdot \frac{1}{n^2}\sigma^2 \\ &= \boxed{\frac{\sigma^2}{n}} \end{aligned}$$

- Remember $\text{Var}(\bar{X}) = \frac{\sigma^2}{n}$: it's another important result.

Example: length of seeds

- ▶ A botanist has collected a sample of 10 seeds and measures the length of each.
- ▶ The seed lengths X_1, X_2, \dots, X_{10} are supposed to be iid with mean $\mu = 5$ mm and variance $\sigma^2 = 2$ mm².

$$E(\bar{X}) = \mu = 5$$

$$\text{Var}(\bar{X}) = \sigma^2/n = 2/10 = 0.2$$

The Central Limit Theorem

Functions of
Several Random
Variables (Ch. 5.5)

Yifan Zhu

Functions of
Several Random
Variables

Expectations and
variances of linear
combinations

Approximating the
Mean and Variance
of a Function

The Central Limit
Theorem

- ▶ If X_1, X_2, \dots, X_n are *any* iid random variables with mean μ and variance $\sigma^2 < \infty$, and if $n \geq 25$,

$$\bar{X} \approx \text{Normal} \left(\mu, \frac{\sigma^2}{n} \right)$$

- ▶ The Central Limit Theorem (CLT) one of the most important and useful results in statistics.

Example: tool serial numbers

- ▶ W_1 = last digit of the serial number observed next Monday at 9 AM
- ▶ W_2 = last digit of the serial number the Monday after at 9 AM
- ▶ W_1 and W_2 are independent with pmf:



$$f(w) = \begin{cases} 0.1 & w = 0, 1, 2, \dots, 9 \\ 0 & \text{otherwise} \end{cases}$$

- ▶ $\bar{W} = \frac{1}{2}(W_1 + W_2)$ has the pmf:


The Probability Function for \bar{W} for $n = 2$

\bar{w}	$f(\bar{w})$	\bar{w}	$f(\bar{w})$	\bar{w}	$f(\bar{w})$	\bar{w}	$f(\bar{w})$	\bar{w}	$f(\bar{w})$
0.0	.01	2.0	.05	4.0	.09	6.0	.07	8.0	.03
0.5	.02	2.5	.06	4.5	.10	6.5	.06	8.5	.02
1.0	.03	3.0	.07	5.0	.09	7.0	.05	9.0	.01
1.5	.04	3.5	.08	5.5	.08	7.5	.04		

Example: tool serial numbers




- What if $\bar{W} = \frac{1}{8}(W_1 + W_2 + \cdots + W_8)$, the average of 8 days of initial serial numbers?



Example: excess sale time

- ▶ \bar{S} = sample mean excess sale time (over a 7.5 s threshold) for 100 stamp sales.
- ▶ Each individual excess sale time should have an $\text{Exp}(\alpha = 16.5 \text{ s})$ distribution. That means:
 - ▶ $E(\bar{S}) = \alpha = 16.5 \text{ s}$
 - ▶ $SD(\bar{S}) = \sqrt{\text{Var}(\bar{S})} = \sqrt{\frac{\alpha^2}{100}} = 1.65 \text{ s}$
 - ▶ By the Central Limit Theorem, $\bar{S} \approx N(16.5, 1.65^2)$
- ▶ We want to approximate $P(\bar{S} > 17)$.

The approximate probability distribution of \bar{S} is normal with mean 16.5 and standard deviation 1.65



Example: excess sale time

$$\begin{aligned}P(\bar{S} > 17) &= P\left(\frac{\bar{S} - 16.5}{1.65} > \frac{17 - 16.5}{1.65}\right) \\&\approx P(Z > 0.303) \quad (Z \sim N(0, 1)) \\&= 1 - P(Z \leq 0.303) \\&= 1 - \Phi(0.303) \\&= 1 - 0.62 \quad \text{from the standard normal table} \\&= 0.38\end{aligned}$$

Example: net weight of baby food jars

Functions of
Several Random
Variables (Ch. 5.5)

Yifan Zhu

Functions of
Several Random
Variables

Expectations and
variances of linear
combinations

Approximating the
Mean and Variance
of a Function

The Central Limit
Theorem

- ▶ Individual jar weights are iid with unknown mean μ and standard deviation $\sigma = 1.6$ g
- ▶ \bar{V} = sample mean weight of n jars $\approx N\left(\mu, \frac{1.6^2}{n}\right)$.
- ▶ We want to find μ . One way to hone in on μ is to find n such that:

$$P(\mu - 0.3 < \bar{V} < \mu + 0.3) = 0.8$$

That way, our measured value of \bar{V} is likely to be close to μ .

Example: net weight of baby food jars

Functions of
Several Random
Variables (Ch. 5.5)

Yifan Zhu

Functions of
Several Random
Variables

Expectations and
variances of linear
combinations

Approximating the
Mean and Variance of
a Function

The Central Limit
Theorem

$$\begin{aligned}0.8 &= P(\mu - 0.3 < \bar{V} < \mu + 0.3) \\&= P\left(\frac{-0.3}{1.6/\sqrt{n}} < \frac{\bar{V} - \mu}{1.6/\sqrt{n}} < \frac{0.3}{1.6/\sqrt{n}}\right) \\&\approx P(-0.19\sqrt{n} < Z < 0.19\sqrt{n}) \quad (\text{by CLT}) \\&= 1 - 2\Phi(-0.19\sqrt{n}) \quad (\text{look at the } N(0,1) \text{ pdf})\end{aligned}$$

$$\begin{aligned}\Phi^{-1}(0.1) &= -0.19\sqrt{n} \\n &= \frac{\Phi^{-1}(0.1)^2}{(-0.19)^2} \\&= \frac{(-1.28)^2}{(-0.19)^2} \quad (\text{standard normal table}) \\&= 46.10\end{aligned}$$

► Hence, we'll need a sample size of $n = 47$.

Example: cars

- ▶ Suppose a bunch of cars pass through certain stretch of road. Whenever a car comes, you look at your watch and record the time.
- ▶ Let X_i be the time (in hours) between when the i 'th car comes and the $(i+1)$ 'th car comes, $i = 1, \dots, 44$. Suppose you know:

$$X_1, X_2, \dots, X_{44} \sim \text{iid } f(x) = e^{-x} \quad x \geq 0$$

- ▶ Find the probability that the average time gap between cars exceeds 1.05 hours.

Example: cars

$$\mu = E(X_1)$$

$$= \int_{-\infty}^{\infty} xf(x)dx$$

$$= \int_0^{\infty} xe^{-x}dx$$

$$= -e^{-x}(x+1)|_0^{\infty} \quad \text{integration by parts}$$

$$= 1$$

Example: cars

$$\begin{aligned}E(X_1^2) &= \int_{-\infty}^{\infty} x^2 f(x) dx \\&= \int_0^{\infty} x^2 e^{-x} dx \\&= -e^{-x}(x^2 + 2x + 2)|_0^{\infty} \quad \text{integration by parts} \\&= 2 \\ \sigma^2 &= \text{Var}(X_1) \\&= E(X_1^2) - E^2(X_1) \\&= 2 - 1^2 \\&= 1\end{aligned}$$

Example: cars

$$\begin{aligned}\bar{X} &\sim \text{approx. } N(\mu, \sigma^2/n) \\ &= N(1, 1/44)\end{aligned}$$

Thus:


$$\frac{\bar{X} - 1}{\sqrt{1/44}} \sim N(0, 1)$$

Example: cars


Now, we're ready to approximate:

$$\begin{aligned}P(\bar{X} > 1.05) &= P\left(\frac{\bar{X} - 1}{\sqrt{1/44}} > \frac{1.05 - 1}{\sqrt{1/44}}\right) \\&= P\left(\frac{\bar{X} - 1}{\sqrt{1/44}} > 0.332\right) \\&\approx P(Z > 0.332) \\&= 1 - P(Z \leq 0.332) \\&= 1 - \Phi(0.332) \\&= 1 - 0.630 = 0.370\end{aligned}$$

Example: cars



Example: cars



Example: cars

