

# Game-theoretic Foundations of Multi-agent Systems

## Lecture 6: Repeated Games

Seyed Majid Zahedi

UNIVERSITY OF  
**WATERLOO**



# Outline

1. Finitely Repeated Games
2. Infinitely Repeated Games
3. Folk Theorem
4. Repeated Games with Imperfect Monitoring



# Repeated Games

- In a (typical) repeated game:
  - Agents play a given game (aka. **stage game**)
  - Then, they get their utilities
  - And, they play again ...



# Repeated Games

- In a (typical) repeated game:
  - Agents play a given game (aka. **stage game**)
  - Then, they get their utilities
  - And, they play again ...
- Can be repeated **finitely** or **infinitely** many times



# Repeated Games

- In a (typical) repeated game:
  - Agents play a given game (aka. **stage game**)
  - Then, they get their utilities
  - And, they play again ...
- Can be repeated **finitely** or **infinitely** many times
- Really, an extensive form game
  - Would like to find subgame-perfect equilibria

## Repeated Games (cont.)

- One subgame-perfect equilibrium:
  - Keep repeating some Nash equilibrium of the stage game
  - **Memoryless** strategy, called a **stationary strategy**



## Repeated Games (cont.)

- One subgame-perfect equilibrium:
  - Keep repeating some Nash equilibrium of the stage game
  - **Memoryless** strategy, called a **stationary strategy**
- But are there other equilibria?
  - Strategy space of repeated game is much richer than that of stage game

# Key Questions

- Do agents see what the other agents played earlier?
- Do they remember what they knew?
- Given utility of each stage game, what is the utility of the entire repeated game?





# Finitely Repeated Games (with Perfect Monitoring)

- Agents play stage game  $G$  for  $R$  rounds



# Finitely Repeated Games (with Perfect Monitoring)

- Agents play stage game  $G$  for  $R$  rounds
- At each round, outcomes of all past rounds are observed by all agents



# Finitely Repeated Games (with Perfect Monitoring)

- Agents play stage game  $G$  for  $R$  rounds
- At each round, outcomes of all past rounds are observed by all agents
- Agents' overall utility is sum of **discounted utilities** at each round
  - Discount factor is  $0 \leq \delta \leq 1$
  - Game is denoted by  $G^R(\delta)$



## Finitely Repeated Games (with Perfect Monitoring)

- Agents play stage game  $G$  for  $R$  rounds
- At each round, outcomes of all past rounds are observed by all agents
- Agents' overall utility is sum of **discounted utilities** at each round
  - Discount factor is  $0 \leq \delta \leq 1$
  - Game is denoted by  $G^R(\delta)$
- Given sequence of utilities  $u_i^{(1)}, \dots, u_i^{(R)}$ ,  $u_i = \sum_{r=1}^R \delta^{r-1} u_i^{(r)}$

## Example: Finitely Repeated Prisoner's Dilemma

- Two agents play Prisoner's Dilemma for  $R$  rounds ( $\delta = 1$ )

	D	C
D	-2, -2	-4, -1
C	-1, -4	-3, -3



## Example: Finitely Repeated Prisoner's Dilemma

- Two agents play Prisoner's Dilemma for  $R$  rounds ( $\delta = 1$ )

	D	C
D	-2, -2	-4, -1
C	-1, -4	-3, -3

- Starting from last round, (C, C) is dominant strategy



## Example: Finitely Repeated Prisoner's Dilemma

- Two agents play Prisoner's Dilemma for  $R$  rounds ( $\delta = 1$ )

	D	C
D	-2, -2	-4, -1
C	-1, -4	-3, -3

- Starting from last round, (C, C) is dominant strategy
- Hence, in second-to-last round, there is no way to influence what will happen



## Example: Finitely Repeated Prisoner's Dilemma

- Two agents play Prisoner's Dilemma for  $R$  rounds ( $\delta = 1$ )

	D	C
D	-2, -2	-4, -1
C	-1, -4	-3, -3

- Starting from last round, (C, C) is dominant strategy
- Hence, in second-to-last round, there is no way to influence what will happen
- So, (C, C) is dominant strategy at this round as well





## Example: Finitely Repeated Prisoner's Dilemma

- Two agents play Prisoner's Dilemma for  $R$  rounds ( $\delta = 1$ )

	D	C
D	-2, -2	-4, -1
C	-1, -4	-3, -3

- Starting from last round, (C, C) is dominant strategy
- Hence, in second-to-last round, there is no way to influence what will happen
- So, (C, C) is dominant strategy at this round as well
- The unique SPE is (C, C) at each round

# SPE in Finitely Repeated Games

## [Theorem]

- If stage game  $G$  has unique strategy equilibrium  $s^*$ , then  $G^R(\delta)$  has unique SPE in which  $s^{(r)} = s^*$  for all  $r = 1, \dots, R$ , regardless of history

## [Proof]

- By backward induction, at round  $R$ , we have  $s^{(R)} = s^*$
- Given this, then we have  $s^{(R-1)} = s^*$ , and continuing inductively,  $s^{(r)} = s^*$  for all  $r = 1, \dots, R$ , regardless of history



## SPE: Example I

- Two agents play the following game for 2 rounds ( $\delta = 1$ )

	D1	D2	C
D1	4, 4	1, 1	6, 0
D2	1, 1	2, 2	6, 0
C	0, 6	0, 6	5, 5



## SPE: Example I

- Two agents play the following game for 2 rounds ( $\delta = 1$ )

	D1	D2	C
D1	4, 4	1, 1	6, 0
D2	1, 1	2, 2	6, 0
C	0, 6	0, 6	5, 5

- Consider the following strategy:
  - In round 1, cooperate;
  - In round 2, if someone defected in round 1, play D2; otherwise, play D1



## SPE: Example I

- Two agents play the following game for 2 rounds ( $\delta = 1$ )

	D1	D2	C
D1	4, 4	1, 1	6, 0
D2	1, 1	2, 2	6, 0
C	0, 6	0, 6	5, 5

- Consider the following strategy:
  - In round 1, cooperate;
  - In round 2, if someone defected in round 1, play D2; otherwise, play D1
- If both agents play this, is that SPE?

## SPE: Example II

- Two agents play the following game for 2 rounds ( $\delta = 1$ )

	D	Crazy	C
D	4, 4	1, 0	6, 0
Crazy	0, 1	0, 0	0, 1
C	0, 6	1, 0	5, 5



## SPE: Example II

- Two agents play the following game for 2 rounds ( $\delta = 1$ )

	D	Crazy	C
D	4, 4	1, 0	6, 0
Crazy	0, 1	0, 0	0, 1
C	0, 6	1, 0	5, 5

- What are the subgame perfect equilibria?



## SPE: Example II

- Two agents play the following game for 2 rounds ( $\delta = 1$ )

	D	Crazy	C
D	4, 4	1, 0	6, 0
Crazy	0, 1	0, 0	0, 1
C	0, 6	1, 0	5, 5

- What are the subgame perfect equilibria?
- Consider the following strategy:
  - In round 1, cooperate;
  - In round 2, if someone played D or Crazy in round 1, play Crazy; otherwise, play D





## SPE: Example II

- Two agents play the following game for 2 rounds ( $\delta = 1$ )

	D	Crazy	C
D	4, 4	1, 0	6, 0
Crazy	0, 1	0, 0	0, 1
C	0, 6	1, 0	5, 5

- What are the subgame perfect equilibria?
- Consider the following strategy:
  - In round 1, cooperate;
  - In round 2, if someone played D or Crazy in round 1, play Crazy; otherwise, play D
- If both agents play this, is that NE (not SPE)?

## TSPE: Example III

- If  $G$  has multiple equilibria, then  $G^R(\delta)$  does not have unique SPE
- Consider following example

	x	y	z
x	3, 3	0, 4	-2, 0
y	4, 0	1, 1	-2, 0
z	0, -2	0, -2	-1, -1

- Stage game has two pure NE:  $(y, y)$  and  $(z, z)$
- Consider the following policy:
  - Play x in first round
  - Play y in second round if opponent played x; otherwise, play z
- Is both agents playing this SPE?



# Outline

1. Finitely Repeated Games
2. Infinitely Repeated Games
3. Folk Theorem
4. Repeated Games with Imperfect Monitoring



# Utilities in Infinitely Repeated Games

- Limit-average utility:

$$u_i = \lim_{R \rightarrow \infty} \frac{\sum_{r=1}^R u_i^{(r)}}{R}$$

- Future-discounted utility:

$$u_i = (1 - \delta) \sum_{r=1}^{\infty} \delta^{r-1} u_i^{(r)},$$

for some  $0 \leq \delta < 1$



# Subgame Perfection in Infinitely Repeated Games

- **One-shot deviation** from strategy  $s$  means deviating from  $s$  in single stage and conforming to it thereafter
- Strategy profile  $s^*$  is SPE **if and only if** there are no **profitable** one-shot deviation for **each subgame** and **every agent**
- This follows from principle of optimality of **dynamic programming**
- This applies to finitely repeated games as well



# Trigger Strategies (TS)

- Agents get **punished** if they deviate from agreed profile
- In **non-forgiving** TS (or grim TS), punishment continues forever

$$s_i^{(t)} = \begin{cases} s_i^* & \text{if } s^{(r)} = s^* \quad \forall r < t, \\ \underline{s}_i^j & \text{otherwise} \end{cases}$$

- Here,  $s^*$  is agreed profile, and  $\underline{s}_i^j$  is punishment strategy of  $i$  against agent  $j$
- Single deviation by  $j$  triggers agent  $i$  to switch to  $\underline{s}_i^j$  **forever**



## Example: Infinitely Repeated Prisoner's Dilemma

- Consider **trigger** strategy:
  - Deny as long as everyone denies
  - Once a player confesses, confess **forever**

	D	C
D	-2, -2	-4, -1
C	-1, -4	-3, -3



## Example: Infinitely Repeated Prisoner's Dilemma

- Consider **trigger** strategy:
  - Deny as long as everyone denies
  - Once a player confesses, confess **forever**
- Is both agents playing this SPE?

	D	C
D	-2, -2	-4, -1
C	-1, -4	-3, -3





## Example: Infinitely Repeated Prisoner's Dilemma

- Consider **trigger** strategy:
  - Deny as long as everyone denies
  - Once a player confesses, confess **forever**
- Is both agents playing this SPE?
- Does it depend on  $\delta$ ?

	D	C
D	-2, -2	-4, -1
C	-1, -4	-3, -3

# Trigger Strategy for Infinitely Repeated Prisoners' Dilemma

- We can use one-stage deviation principle



# Trigger Strategy for Infinitely Repeated Prisoners' Dilemma

- We can use one-stage deviation principle
- There are two types of subgames:
  - Type 1: Both agents denied so far
  - Type 2: At least one agent confessed in the past



# Trigger Strategy for Infinitely Repeated Prisoners' Dilemma

- We can use one-stage deviation principle
- There are two types of subgames:
  - Type 1: Both agents denied so far
  - Type 2: At least one agent confessed in the past
- Type-1 subgames: (D is best response to D)



# Trigger Strategy for Infinitely Repeated Prisoners' Dilemma

- We can use one-stage deviation principle
- There are two types of subgames:
  - Type 1: Both agents denied so far
  - Type 2: At least one agent confessed in the past
- Type-1 subgames: (D is best response to D)
  - Utility from no deviation:  $(1 - \delta)(-2 - 2\delta - 2\delta^2 + \dots) = -2$



# Trigger Strategy for Infinitely Repeated Prisoners' Dilemma

- We can use one-stage deviation principle
- There are two types of subgames:
  - Type 1: Both agents denied so far
  - Type 2: At least one agent confessed in the past
- Type-1 subgames: (D is best response to D)
  - Utility from no deviation:  $(1 - \delta)(-2 - 2\delta - 2\delta^2 + \dots) = -2$
  - Utility from on-shot deviation:  $(1 - \delta)(-1 + (-3\delta - 3\delta^2 + \dots)) = -(1 - \delta) - 3\delta$



# Trigger Strategy for Infinitely Repeated Prisoners' Dilemma

- We can use one-stage deviation principle
- There are two types of subgames:
  - Type 1: Both agents denied so far
  - Type 2: At least one agent confessed in the past
- Type-1 subgames: (D is best response to D)
  - Utility from no deviation:  $(1 - \delta)(-2 - 2\delta - 2\delta^2 + \dots) = -2$
  - Utility from on-shot deviation:  $(1 - \delta)(-1 + (-3\delta - 3\delta^2 + \dots)) = -(1 - \delta) - 3\delta$
  - Deviation is not beneficial if  $\delta \geq 1/2$



# Trigger Strategy for Infinitely Repeated Prisoners' Dilemma

- We can use one-stage deviation principle
- There are two types of subgames:
  - Type 1: Both agents denied so far
  - Type 2: At least one agent confessed in the past
- Type-1 subgames: (D is best response to D)
  - Utility from no deviation:  $(1 - \delta)(-2 - 2\delta - 2\delta^2 + \dots) = -2$
  - Utility from on-shot deviation:  $(1 - \delta)(-1 + (-3\delta - 3\delta^2 + \dots)) = -(1 - \delta) - 3\delta$
  - Deviation is not beneficial if  $\delta \geq 1/2$
- Type-2 subgames: (C is best response to C)





# Trigger Strategy for Infinitely Repeated Prisoners' Dilemma

- We can use one-stage deviation principle
- There are two types of subgames:
  - Type 1: Both agents denied so far
  - Type 2: At least one agent confessed in the past
- Type-1 subgames: (D is best response to D)
  - Utility from no deviation:  $(1 - \delta)(-2 - 2\delta - 2\delta^2 + \dots) = -2$
  - Utility from on-shot deviation:  $(1 - \delta)(-1 + (-3\delta - 3\delta^2 + \dots)) = -(1 - \delta) - 3\delta$
  - Deviation is not beneficial if  $\delta \geq 1/2$
- Type-2 subgames: (C is best response to C)
  - Other agents will always play C, thus C is best response

# Tit-for-tat Strategy

- Consider **tit-for-tat** strategy:
  - Deny in 1st round
  - Then, do whatever other agent did in previous round



# Tit-for-tat Strategy

- Consider **tit-for-tat** strategy:
  - Deny in 1st round
  - Then, do whatever other agent did in previous round
- Is both agents playing this NE?



# Tit-for-tat Strategy

- Consider **tit-for-tat** strategy:
  - Deny in 1st round
  - Then, do whatever other agent did in previous round
- Is both agents playing this NE?
- Is both agents playing this SPE?



# Tit-for-tat Strategy

- Consider **tit-for-tat** strategy:
  - Deny in 1st round
  - Then, do whatever other agent did in previous round
- Is both agents playing this NE?
- Is both agents playing this SPE?
- What about one playing TFT and other trigger?

## Remarks

- If  $s^*$  is NE of  $G$ , then “each agent plays  $s_i^*$ ” is SPE of  $G^R(\delta)$ 
  - Future play of other agents is independent of how each agent plays
  - Optimal play is to maximize current utility, i.e., play static best response
- Sets of equilibria for finite and infinite horizon versions can be **different**
  - Multiplicity of equilibria in repeated prisoner's dilemma only occurs at  $R = \infty$
  - For any finite  $R$  (thus for  $R \rightarrow \infty$ ), repeated prisoners' dilemma has unique SPE



## Repetition Could Lead to Bad Outcomes

- Consider the following game

	x	y	z
x	2, 2	2, 1	0, 0
y	1, 2	1, 1	-1, 0
z	0, 0	0, -1	-1, -1

- Strategy x strictly dominates y and z for both agents
- Unique NE of stage game is (x, x)
- If  $\delta \geq 1/2$ , this game has SPE in which (y, y) is played in every round
- It is supported by slightly more complicated strategy than grim trigger
  - I. Play y in every round unless someone deviates, then go to II
  - II. Play z. If no one deviates go to I. If someone deviates stay in II



# Outline

1. Finitely Repeated Games
2. Infinitely Repeated Games
3. Folk Theorem
4. Repeated Games with Imperfect Monitoring





# Characterizing NE of Infinitely Repeated Games

- Characterizing all equilibrium strategy profiles might be challenging



# Characterizing NE of Infinitely Repeated Games

- Characterizing all equilibrium strategy profiles might be challenging
- Instead, we can characterize utilities obtained in them



# Characterizing NE of Infinitely Repeated Games

- Characterizing all equilibrium strategy profiles might be challenging
- Instead, we can characterize utilities obtained in them
- Such utilities must be **feasible**
  - There must be outcomes of game such that agents, on average, get these utilities



# Characterizing NE of Infinitely Repeated Games

- Characterizing all equilibrium strategy profiles might be challenging
- Instead, we can characterize utilities obtained in them
- Such utilities must be **feasible**
  - There must be outcomes of game such that agents, on average, get these utilities
- They must also be **enforceable**
  - Deviation should lead to punishment that outweighs benefits of deviation



# Characterizing NE of Infinitely Repeated Games

- Characterizing all equilibrium strategy profiles might be challenging
- Instead, we can characterize utilities obtained in them
- Such utilities must be **feasible**
  - There must be outcomes of game such that agents, on average, get these utilities
- They must also be **enforceable**
  - Deviation should lead to punishment that outweighs benefits of deviation
- **Folk theorem** states that utility vector can be realized by some NE iff it is both feasible and enforceable

## Feasible Utilities: Formal Definition

- **Utility profile**  $u = (u_1, u_2, \dots, u_n)$  is **feasible** if there exist **rational, non-negative** values  $\{\alpha_a\}$  such that for all  $i$ ,  $u_i = \sum_{a \in A} \alpha_a u_i(a)$ , with  $\sum_{a \in A} \alpha_a = 1$



## Feasible Utilities: Formal Definition

- **Utility profile**  $u = (u_1, u_2, \dots, u_n)$  is **feasible** if there exist **rational, non-negative** values  $\{\alpha_a\}$  such that for all  $i$ ,  $u_i = \sum_{a \in A} \alpha_a u_i(a)$ , with  $\sum_{a \in A} \alpha_a = 1$
- You could think of feasible utilities as **convex hull** of possible outcomes:

$$U = \text{Conv}\{u \in \mathbb{R}^{|N|} \mid \text{there exists } a \in A \text{ such that } u(a) = u\}$$



## Feasible Utilities: Formal Definition

- **Utility profile**  $u = (u_1, u_2, \dots, u_n)$  is **feasible** if there exist **rational, non-negative** values  $\{\alpha_a\}$  such that for all  $i$ ,  $u_i = \sum_{a \in A} \alpha_a u_i(a)$ , with  $\sum_{a \in A} \alpha_a = 1$
- You could think of feasible utilities as **convex hull** of possible outcomes:

$$U = \text{Conv}\{u \in \mathbb{R}^{|N|} \mid \text{there exists } a \in A \text{ such that } u(a) = u\}$$

- Note that  $U \neq \{u \in \mathbb{R}^{|N|} \mid \text{there exists } s \in S \text{ such that } u(s) = u\}$



## Feasibility: Example

	Left	Right
Left	2, 2	0, 3
Right	3, 0	1, 1

- Utility vector  $(2, 2)$  is feasible as it is one of outcomes of game



## Feasibility: Example

	Left	Right
Left	2, 2	0, 3
Right	3, 0	1, 1

- Utility vector  $(2, 2)$  is feasible as it is one of outcomes of game
- Utility vector  $(1, 2.5)$  is feasible as agents can alternate between  $(2, 2)$  and  $(0, 3)$



## Feasibility: Example

	Left	Right
Left	2, 2	0, 3
Right	3, 0	1, 1

- Utility vector  $(2, 2)$  is feasible as it is one of outcomes of game
- Utility vector  $(1, 2.5)$  is feasible as agents can alternate between  $(2, 2)$  and  $(0, 3)$
- What about  $(0.5, 2.75)$ ?



## Feasibility: Example

	Left	Right
Left	2, 2	0, 3
Right	3, 0	1, 1

- Utility vector  $(2, 2)$  is feasible as it is one of outcomes of game
- Utility vector  $(1, 2.5)$  is feasible as agents can alternate between  $(2, 2)$  and  $(0, 3)$
- What about  $(0.5, 2.75)$ ?
- What about  $(3, 0.1)$ ?

# Enforceable and Individually Rational Utilities

- Recall **minmax value** of agent  $i$ :

$$\underline{v}_i = \min_{s_{-i}} \max_{s_i} u_i(s_i, s_{-i})$$



# Enforceable and Individually Rational Utilities

- Recall **minmax value** of agent  $i$ :

$$\underline{v}_i = \min_{s_{-i}} \max_{s_i} u_i(s_i, s_{-i})$$

- Utility profile  $u \in \mathbb{R}^{|N|}$  is **individually rational** if  $u_i \geq \underline{v}_i$  for all  $i$



# Enforceable and Individually Rational Utilities

- Recall **minmax value** of agent  $i$ :

$$\underline{v}_i = \min_{s_{-i}} \max_{s_i} u_i(s_i, s_{-i})$$

- Utility profile  $u \in \mathbb{R}^{|N|}$  is **individually rational** if  $u_i \geq \underline{v}_i$  for all  $i$
- Utility profile  $u = (u_1, u_2, \dots, u_n)$  is **enforceable** if it is individually rational

# Nash Folk Theorem

- Consider infinitely repeated game  $G$  played by agents with **average utilities**





# Nash Folk Theorem

- Consider infinitely repeated game  $G$  played by agents with **average utilities**
- If  $u$  is utility profile for any NE of repeated  $G$ , then  $u_i$  is enforceable for all  $i$



# Nash Folk Theorem

- Consider infinitely repeated game  $G$  played by agents with **average utilities**
- If  $u$  is utility profile for any NE of repeated  $G$ , then  $u_i$  is enforceable for all  $i$
- If  $u$  is both feasible and enforceable, then  $u$  is utility profile for some NE of  $G$



# Nash Folk Theorem

- Consider infinitely repeated game  $G$  played by agents with **average utilities**
- If  $u$  is utility profile for any NE of repeated  $G$ , then  $u_i$  is enforceable for all  $i$
- If  $u$  is both feasible and enforceable, then  $u$  is utility profile for some NE of  $G$
- Folk theorem can be stated for agents with discounted utilities as well

# Problems with Nash Folk Theorem

- Any feasible and enforceable utility can be achieved (for **patient enough** agents)



# Problems with Nash Folk Theorem

- Any feasible and enforceable utility can be achieved (for **patient enough** agents)
- Enforcement is often done by grim trigger strategy



# Problems with Nash Folk Theorem

- Any feasible and enforceable utility can be achieved (for **patient enough** agents)
- Enforcement is often done by grim trigger strategy
  - Play certain strategy as long as no one deviates



# Problems with Nash Folk Theorem

- Any feasible and enforceable utility can be achieved (for **patient enough** agents)
- Enforcement is often done by grim trigger strategy
  - Play certain strategy as long as no one deviates
  - If some agent  $j$  deviates, then play minmax strategy against that agent thereafter



# Problems with Nash Folk Theorem

- Any feasible and enforceable utility can be achieved (for **patient enough** agents)
- Enforcement is often done by grim trigger strategy
  - Play certain strategy as long as no one deviates
  - If some agent  $j$  deviates, then play minmax strategy against that agent thereafter
- NE involves non-forgiving TS which may be costly for punishers





# Problems with Nash Folk Theorem

- Any feasible and enforceable utility can be achieved (for **patient enough** agents)
- Enforcement is often done by grim trigger strategy
  - Play certain strategy as long as no one deviates
  - If some agent  $j$  deviates, then play minmax strategy against that agent thereafter
- NE involves non-forgiving TS which may be costly for punishers
- NE may include **non-credible threats**



# Problems with Nash Folk Theorem

- Any feasible and enforceable utility can be achieved (for **patient enough** agents)
- Enforcement is often done by grim trigger strategy
  - Play certain strategy as long as no one deviates
  - If some agent  $j$  deviates, then play minmax strategy against that agent thereafter
- NE involves non-forgiving TS which may be costly for punishers
- NE may include **non-credible threats**
- NE may not be subgame perfect

## Example

	L	R
U	6, 6	0, -100
D	7, 1	0, -100

- Unique NE in this game is (D, L)
- Minmax values are given by  $\underline{v}_1 = 0$  and  $\underline{v}_2 = 1$
- Minmax strategy against agent 1 requires agent 2 to play R
- R is strictly dominated by L for agent 2



# Outline

1. Finitely Repeated Games
2. Infinitely Repeated Games
3. Folk Theorem
4. Repeated Games with Imperfect Monitoring



# Motivation

- So far, we assumed that agents observe actions of others at each round of game
- Next, we consider games where agents' actions may not be directly observable
- We assume that agents observe only an **imperfect signal** of stage game actions



# Example: Cournot Competition with Noisy Demand

[Green and Porter, Non-cooperative Collusion under Imperfect Price Information, 1984]

- Firms set production levels  $q_1^{(r)}, \dots, q_n^{(r)}$  **privately** at round  $r$
- Firms do not observe each others' output levels
- Market demand is **stochastic**
- Market price depends on total production and market demand
- Low price could be due to high production or low demand
- Firms utility depends on their own production and market price



# Model

- We focus on game with **public information**
- At each round, **all agents** observe some **public outcome**
- Let  $y^{(r)} \in Y$  denote publicly observed outcome at round  $r$
- Each action profile  $a$  induces **probability distribution** over  $y$
- Let  $\pi(y, a)$  denote probability distribution of  $y$  under action profile  $a$
- Public information at round  $r$  is  $h^{(r)} = (y^{(1)}, \dots, y^{(r-1)})$
- Strategy of agent  $i$  is **sequence of maps**  $s_i^{(r)} : h^{(r)} \rightarrow S_i$



## Model (cont.)

- Agents utility depends **only** on their own action and public outcome





## Model (cont.)

- Agents utility depends **only** on their own action and public outcome
- Dependence on actions of others is through their effect on distribution of  $y$



## Model (cont.)

- Agents utility depends **only** on their own action and public outcome
- Dependence on actions of others is through their effect on distribution of  $y$
- Agent  $i$ 's **realized** utility at round  $r$  is  $u_i(a_i^{(r)}, y^{(r)})$



## Model (cont.)

- Agents utility depends **only** on their own action and public outcome
- Dependence on actions of others is through their effect on distribution of  $y$
- Agent  $i$ 's **realized** utility at round  $r$  is  $u_i(a_i^{(r)}, y^{(r)})$
- Agent  $i$ 's expected stage utility is

$$u_i(a) = \sum_{y \in Y} \pi(y, a) u_i(a_i, y)$$



## Model (cont.)

- Agents utility depends **only** on their own action and public outcome
- Dependence on actions of others is through their effect on distribution of  $y$
- Agent  $i$ 's **realized** utility at round  $r$  is  $u_i(a_i^{(r)}, y^{(r)})$
- Agent  $i$ 's expected stage utility is

$$u_i(a) = \sum_{y \in Y} \pi(y, a) u_i(a_i, y)$$

- Agent  $i$ 's average discounted utility when sequence  $\{a^{(t)}\}$  is played is

$$(1 - \delta) \sum_{r=1}^{\infty} \delta^{r-1} u_i(a^{(r)})$$

## Simpler Example: Noisy Prisoner's Dilemma

- Prisoners do not observe each others actions, instead, they observe signal  $y$



## Simpler Example: Noisy Prisoner's Dilemma

- Prisoners do not observe each others actions, instead, they observe signal  $y$ 
  - $u_1(D, y) = 1 + y$        $u_1(C, y) = 4 + y$



## Simpler Example: Noisy Prisoner's Dilemma

- Prisoners do not observe each others actions, instead, they observe signal  $y$ 
  - $u_1(D, y) = 1 + y$                        $u_1(C, y) = 4 + y$
  - $u_2(D, y) = 1 + y$                        $u_2(C, y) = 4 + y$



## Simpler Example: Noisy Prisoner's Dilemma

- Prisoners do not observe each others actions, instead, they observe signal  $y$ 
  - $u_1(D, y) = 1 + y$                        $u_1(C, y) = 4 + y$
  - $u_2(D, y) = 1 + y$                        $u_2(C, y) = 4 + y$
- Signal  $y$  is defined by cont. random variable  $X$  with CDF  $F(x)$  and  $\mathbb{E}[X] = 0$





## Simpler Example: Noisy Prisoner's Dilemma

- Prisoners do not observe each others actions, instead, they observe signal  $y$ 
  - $u_1(D, y) = 1 + y$                        $u_1(C, y) = 4 + y$
  - $u_2(D, y) = 1 + y$                        $u_2(C, y) = 4 + y$
- Signal  $y$  is defined by cont. random variable  $X$  with CDF  $F(x)$  and  $\mathbb{E}[X] = 0$ 
  - If  $a = (D, D)$ , then  $y = X$



## Simpler Example: Noisy Prisoner's Dilemma

- Prisoners do not observe each others actions, instead, they observe signal  $y$ 
  - $u_1(D, y) = 1 + y$                        $u_1(C, y) = 4 + y$
  - $u_2(D, y) = 1 + y$                        $u_2(C, y) = 4 + y$
- Signal  $y$  is defined by cont. random variable  $X$  with CDF  $F(x)$  and  $\mathbb{E}[X] = 0$ 
  - If  $a = (D, D)$ , then  $y = X$
  - If  $a = (D, C)$  or  $(C, D)$ , then  $y = X - 2$



## Simpler Example: Noisy Prisoner's Dilemma

- Prisoners do not observe each others actions, instead, they observe signal  $y$ 
  - $u_1(D, y) = 1 + y$                        $u_1(C, y) = 4 + y$
  - $u_2(D, y) = 1 + y$                        $u_2(C, y) = 4 + y$
- Signal  $y$  is defined by cont. random variable  $X$  with CDF  $F(x)$  and  $\mathbb{E}[X] = 0$ 
  - If  $a = (D, D)$ , then  $y = X$
  - If  $a = (D, C)$  or  $(C, D)$ , then  $y = X - 2$
  - If  $a = (C, C)$ , then  $y = X - 4$



## Simpler Example: Noisy Prisoner's Dilemma

- Prisoners do not observe each others actions, instead, they observe signal  $y$ 
  - $u_1(D, y) = 1 + y$        $u_1(C, y) = 4 + y$
  - $u_2(D, y) = 1 + y$        $u_2(C, y) = 4 + y$
- Signal  $y$  is defined by cont. random variable  $X$  with CDF  $F(x)$  and  $\mathbb{E}[X] = 0$ 
  - If  $a = (D, D)$ , then  $y = X$
  - If  $a = (D, C)$  or  $(C, D)$ , then  $y = X - 2$
  - If  $a = (C, C)$ , then  $y = X - 4$
- Normal-form stage game is

	D	C
D	$1 + X, 1 + X$	$-1 + X, 2 + X$
C	$2 + X, -1 + X$	$X, X$

# Trigger-price Strategy

- Consider following trigger strategy
  - (I) - Play  $(D, D)$  until  $y \leq y^*$ , then go to (II)
  - (II) - Play  $(C, C)$  for  $R$  rounds, then go back to (I)
- Notice that strategy is stationary and symmetric
- Also notice that punishment uses **NE of stage game**
- We can choose  $y^*$  and  $R$  such that this strategy profile is SPE



## Trigger-price Strategy (cont.)

- We use one-shot deviation principle
- Deviation in (II) is obviously not beneficial
- In (I), if agents do not deviate, their **expected utility** is

$$v = (1 - \delta) \left( (1 + 0) + \delta \left( F(y^*) \delta^R v + (1 - F(y^*)) v \right) \right)$$

- From this, we obtain

$$v = \frac{1 - \delta}{1 - \delta(1 - \delta)(1 - F(y^*)(1 - \delta^R))}$$



## Trigger-price Strategy (cont.)

- If some agent deviates in (1), then her expected utility is

$$v_d = (1 - \delta) \left( (2 + 0) + \delta \left( F(y^* + 2) \delta^R v + (1 - F(y^* + 2)) v \right) \right)$$

- Deviation provides immediate utility, but increases probability of entering (II)
- To have SPE, we must have  $v \geq v_d$  which means

$$v \geq \frac{2(1 - \delta)}{1 - \delta(1 - \delta)(1 - F(y^* + 2)(1 - \delta^R))}$$
$$\Rightarrow F(y^* + 2) - 2F(y^*) \geq \frac{1 - \delta(1 - \delta)}{\delta(1 - \delta)(1 - \delta^R)}$$

- Any  $R$  and  $y^*$  that satisfy this constraint construct SPE
- **Best trigger-price strategy** can be found by maximizing  $v$  s.t. this constraint



# Acknowledgment

- This lecture is a slightly modified version of ones prepared by
  - Asu Ozdaglar [MIT 6.254]
  - Vincent Conitzer [Duke CPS 590.4]
- Elly Khodaie helped with importing slides from PowerPoint to L<sup>A</sup>T<sub>E</sub>X

