




Supplementary Figure 3 | Example genes with significant isoform switching due to switched promoter and TSS usage.

a, Genome browser view of the *PRKCZ* locus. Tracks include the finalized SALA transcriptome alongside GENCODE v39 gene and transcript annotations. Switched isoform track displays only transcript models with significant changes in expression fraction between any two of the three cell types (as identified by isoformSwitchAnalyzerR). Color scale indicates CPM values derived from *bambu*, normalized within each cell type. CFC-seq tracks show the aggregated signal for 5' ends, 3' ends, and exons across all samples. Additional tracks show annotations from ATAC-seq, cCRE promoter and enhancer regions, and SCAFE-defined tCREs. **b**, Isoform expression fractions of *PRKCZ* in the three cell types, estimated by isoformSwitchAnalyzerR. **c**, Chromatin activity of corresponding ex5_clusters derived from ATAC-seq signal normalized by library size. **d**, Predicted ORF alignment of the switched isoforms of *PRKCZ*. **e**, Genome browser view of the *CA14* locus, track layout as in (a). **f**, Isoform expression fractions of the *CA14* in the three cell types. **g**, Chromatin activity of ex5_clusters derived from ATAC-seq signal. **h**, Predicted ORF alignment of the switched isoforms of *CA14*.