

WEYL-CHAMBER GEOMETRY, POINCARÉ INEQUALITY MEETS SCHATTEN CLASS PROPERTIES OF DUNKL-RIESZ TRANSFORM COMMUTATORS

ZHIJIE FAN, JI LI, AND DMITRIY ZANIN

ABSTRACT.

CONTENTS

1. Preliminaries	1
1.1. Harmonic analysis in the Dunkl setting	1
1.2. Systems of dyadic cubes	5
1.3. Nearly weakly orthonormal sequences	6
2. Analytic tools for Weyl chamber geometry	6
3. Spectral asymptotic formula	14
4. First Rochberg-Semmes based estimate	31
4.1. Proof of Theorem 4.1	31
4.2. Proof of Theorem 4.2	36
5. Second Rochberg-Semmes based estimate	40
6. Appendix	41
References	42

1. PRELIMINARIES

1.1. Harmonic analysis in the Dunkl setting. Let $\mathbb{R}^N$ be the Euclidean space equipped with the usual scalar product $\langle \cdot, \cdot \rangle$ and the induced norm $\| \cdot \|$. For a nonzero vector $v \in \mathbb{R}^N$, the reflection σ_v with respect to the hyperplane $v^\perp$ orthogonal to v is defined by

$$\sigma_v x := x - 2 \frac{\langle x, v \rangle}{\|v\|^2} v.$$

A finite set $R \subset \mathbb{R}^N \setminus \{0\}$ is called a root system if $\sigma_v(R) = R$ for every $v \in R$. A root system R is called a normalized reduced root system if $\|v\|^2 = 2$ for every $v \in R$. The finite group G generated by the reflections $\{\sigma_v\}_{v \in R}$ is called the Weyl group (reflection group) of the root system R . Let $k : R \rightarrow [0, \infty)$ be a multiplicity function, which is a G -invariant function fixed throughout the paper. We will denote by $\mathcal{O}(x) = \{\sigma(x) : \sigma \in G\}$ the G -orbit of a point $x \in \mathbb{R}^N$, and $\mathcal{O}(E) = \bigcup_{x \in E} \mathcal{O}(x)$ the G -orbit of $E \subset \mathbb{R}^N$.

2010 *Mathematics Subject Classification.* 47B10, 42B20, 43A85.

Key words and phrases. Schatten class, Riesz transform commutators, Dunkl operator, Besov space, Nearly weakly orthogonal sequence.

Define the Dunkl measure dw on $\mathbb{R}^N$ by

$$dw(x) := \prod_{v \in R} |\langle x, v \rangle|^{k(v)} dx,$$

where dx stands for the Lebesgue measure on $\mathbb{R}^N$. Let $B(x, r) := \{y \in \mathbb{R}^N : \|x - y\| < r\}$ be the Euclidean ball centred at x with radius $r > 0$. Let $\mathbf{N} := N + \sum_{v \in R} k(v)$. Note that

$$w(B(tx, tr)) = t^{\mathbf{N}} w(B(x, r)), \text{ for } x \in \mathbb{R}^N, t, r > 0.$$

Furthermore,

$$(1) \quad w(B(x, r)) \sim r^{\mathbf{N}} \prod_{v \in R} (|\langle x, v \rangle| + r)^{k(v)},$$

where the implicit constant only depends on N, R and k . It follows from (1) that there exists a constant $C \geq 1$ such that for every $x \in \mathbb{R}^N$,

$$(2) \quad C^{-1} \left(\frac{r_2}{r_1} \right)^{\mathbf{N}} \leq \frac{w(B(x, r_2))}{w(B(x, r_1))} \leq C \left(\frac{r_2}{r_1} \right)^{\mathbf{N}}, \text{ for } 0 < r_1 < r_2,$$

so the numbers $\mathbf{N}$ and N are called the upper and lower homogeneous dimensions, respectively.

[Add the definition of A_∞ . Zhijie will do it.]

Lemma 1.1. *Let*

$$dw(x) = \varpi(x) dx, \quad \varpi(x) = \prod_{v \in R_+} |\langle x, v \rangle|^{2k(v)}.$$

Then $\varpi \in A_\infty(\mathbb{R}^N, dx)$.

Proof. The assertion is immediate if $k \equiv 0$. Otherwise, choose $p > 1$ sufficiently large so that

$$\frac{2|R_+|k(v)}{p-1} < 1, \quad v \in R_+.$$

For every $0 < \beta < 1$, $v \in R_+$, and Euclidean ball $B(x_0, r)$, we apply an orthogonal change of variables to deduce that

$$\begin{aligned} \frac{1}{|B(x_0, r)|} \int_{B(x_0, r)} |\langle x, v \rangle|^{-\beta} dx &\lesssim \frac{1}{r|v|} \int_{\langle x_0, v \rangle - |v|r}^{\langle x_0, v \rangle + |v|r} |s|^{-\beta} ds \\ &= (|v|r)^{-\beta} \int_{\frac{\langle x_0, v \rangle}{|v|r} - 1}^{\frac{\langle x_0, v \rangle}{|v|r} + 1} |s|^{-\beta} ds \\ &\lesssim (|v|r)^{-\beta} \left(1 + \frac{|\langle x_0, v \rangle|}{|v|r} \right)^{-\beta} \\ &\lesssim (|\langle x_0, v \rangle| + r)^{-\beta}. \end{aligned}$$

Hence, by Hölder's inequality,

$$\begin{aligned} \frac{1}{|B(x_0, r)|} \int_{B(x_0, r)} \varpi^{-\frac{1}{p-1}} dx &= \frac{1}{|B(x_0, r)|} \int_{B(x_0, r)} \prod_{v \in R_+} |\langle x, v \rangle|^{-\frac{2k(v)}{p-1}} dx \\ &\leq \prod_{v \in R_+} \left(\frac{1}{|B(x_0, r)|} \int_{B(x_0, r)} |\langle x, v \rangle|^{-\frac{2|R_+|k(v)}{p-1}} dx \right)^{\frac{1}{|R_+|}} \\ &\lesssim \prod_{v \in R_+} (|\langle x_0, v \rangle| + r)^{-\frac{2k(v)}{p-1}}. \end{aligned}$$

On the other hand, by (1) and the G -invariance of k ,

$$\frac{1}{|B(x_0, r)|} \int_{B(x_0, r)} \varpi dx = \frac{w(B(x_0, r))}{|B(x_0, r)|} \lesssim \prod_{v \in R_+} (|\langle x_0, v \rangle| + r)^{2k(v)}.$$

Therefore,

$$\left(\frac{1}{|B|} \int_B \varpi dx \right) \left(\frac{1}{|B|} \int_B \varpi^{-\frac{1}{p-1}} dx \right)^{p-1} \lesssim 1$$

uniformly over all Euclidean balls B . Thus $\varpi \in A_p(\mathbb{R}^N, dx)$, and hence $\varpi \in A_\infty(\mathbb{R}^N, dx)$. $\square$

The Dunkl operators T_ξ are defined by

$$T_\xi f(x) := \partial_\xi f(x) + \sum_{v \in R} \frac{k(v)}{2} \langle v, \xi \rangle \frac{f(x) - f(\sigma_v(x))}{\langle v, x \rangle},$$

where $\partial_\xi f$ denotes the directional derivative of f in the direction ξ . Furthermore, the Dunkl Laplacian is defined by

$$\Delta_{\text{Dunkl}} f(x) := \sum_{j=1}^N T_{e_j}^2 f(x) = \Delta_{\mathbb{R}^N} f(x) + \sum_{v \in R} k(v) \delta_v f(x),$$

where $\Delta_{\mathbb{R}^N}$ is the Laplacian on $\mathbb{R}^N$ and δ_v is defined by

$$\delta_v f(x) := \frac{\partial_v f(x)}{\langle v, x \rangle} - \frac{f(x) - f(\sigma_v(x))}{\langle v, x \rangle^2}.$$

In this article, we consider the Dunkl-Riesz transforms defined by $R_{\text{Dunkl}, j} := T_{e_j}(-\Delta_{\text{Dunkl}})^{-1/2}$, $j = 1, 2, \dots, N$.

Definition 1.2. The Weyl chambers are defined as the closures of connected components of the set

$$(3) \quad \{x \in \mathbb{R}^N : \langle x, v \rangle \neq 0 \text{ for all } v \in R\}.$$

Define $d(x, y) := \min_{\sigma \in G} \|x - \sigma(y)\|$ as the distance between two G -orbits $\mathcal{O}(x)$ and $\mathcal{O}(y)$. Recall from [7, Chapter VII, proof of Theorem 2.12] that

$$(4) \quad d(x, y) = \|x - y\| \text{ if and only if } x, y \in \Gamma$$

for some closed Weyl chamber Γ .

It is clear that $\mathcal{O}(B(x, r)) = \{y \in \mathbb{R}^N : d(x, y) < r\}$. Moreover,

$$w(B(x, r)) \leq w(\mathcal{O}(B(x, r))) \leq |G|w(B(x, r))$$

for all $x \in \mathbb{R}^N$ and $r > 0$. We also denote $B^d(x, r) := \{y \in \mathbb{R}^N : d(x, y) < r\}$, i.e., $B^d(x, r) = \mathcal{O}(B(x, r))$.

Denote by $K_j(x, y)$ the integral kernel of $R_{\text{Dunkl}, j}$. For every $x, y \in \mathbb{R}^N$ and $r > 0$, let

$$V(x, y, r) := \max\{w(B(x, r)), w(B(y, r))\}.$$

The following lemma establishes derivative estimates for the Dunkl–Riesz transform kernel.

Lemma 1.3. *Assume that $N \geq 1$. Then for every pair of multi-indices α, β , every $j \in \{1, \dots, N\}$, and every $x, y \in \mathbb{R}^N$ with $d(x, y) > 0$,*

$$(5) \quad |\partial_x^\alpha \partial_y^\beta K_{\text{Dunkl}, j}(x, y)| \lesssim_{\alpha, \beta, R, k} \frac{d(x, y)}{\|x - y\|} \frac{d(x, y)^{-(|\alpha| + |\beta|)}}{V(x, y, d(x, y))}.$$

Proof. Case 1. Suppose that $|\alpha| + |\beta| = 0$. By [5, Theorem 1.1],

$$|K_{\text{Dunkl}, j}(x, y)| \lesssim \frac{d(x, y)}{\|x - y\|} \frac{1}{w(B(x, d(x, y)))}.$$

By the definition of d , the G -invariance of dw , and (2),

$$V(x, y, d(x, y)) \lesssim w(B(x, d(x, y))),$$

and hence (5) follows.

Case 2. Suppose that $|\alpha| + |\beta| \geq 1$. By [?, Theorem 3.1],

$$|\partial_x^\alpha \partial_y^\beta h_t(x, y)| \lesssim_{\alpha, \beta, R, k} t^{-\frac{|\alpha| + |\beta|}{2}} \left(1 + \frac{\|x - y\|}{\sqrt{t}}\right)^{-2} V(x, y, \sqrt{t})^{-1} e^{-cd(x, y)^2/t}.$$

Recall from the proof of Lemma ?? that

$$K_{\text{Dunkl}, j}(x, y) = -\frac{C}{2} \int_0^\infty (y_j - x_j) h_t(x, y) \frac{dt}{t^{3/2}}.$$

Since

$$\partial_x^\alpha \partial_y^\beta ((y_j - x_j) h_t(x, y)) = (y_j - x_j) \partial_x^\alpha \partial_y^\beta h_t(x, y) - \alpha_j \partial_x^{\alpha - e_j} \partial_y^\beta h_t(x, y) + \beta_j \partial_x^\alpha \partial_y^{\beta - e_j} h_t(x, y),$$

where the corresponding term is omitted if $\alpha_j = 0$ or $\beta_j = 0$, we obtain

$$\begin{aligned} |\partial_x^\alpha \partial_y^\beta K_{\text{Dunkl}, j}(x, y)| &\lesssim \frac{1}{\|x - y\|} \int_0^\infty t^{-\frac{|\alpha| + |\beta| + 1}{2}} V(x, y, \sqrt{t})^{-1} e^{-cd(x, y)^2/t} dt \\ &\quad + \frac{1}{\|x - y\|^2} \int_0^\infty t^{-\frac{|\alpha| + |\beta|}{2}} V(x, y, \sqrt{t})^{-1} e^{-cd(x, y)^2/t} dt. \end{aligned}$$

By (2),

$$V(x, y, \sqrt{t})^{-1} \lesssim V(x, y, d(x, y))^{-1} \begin{cases} \left(\frac{d(x, y)}{\sqrt{t}}\right)^N, & 0 < t \leq d(x, y)^2, \\ \left(\frac{d(x, y)}{\sqrt{t}}\right)^N, & t \geq d(x, y)^2. \end{cases}$$

Hence,

$$\begin{aligned} &\int_0^\infty t^{-\frac{|\alpha| + |\beta| + 1}{2}} V(x, y, \sqrt{t})^{-1} e^{-cd(x, y)^2/t} dt \\ &\lesssim \frac{1}{V(x, y, d(x, y))} \left(d(x, y)^N \int_0^{d(x, y)^2} t^{-\frac{N + |\alpha| + |\beta| + 1}{2}} e^{-cd(x, y)^2/t} dt + d(x, y)^N \int_{d(x, y)^2}^\infty t^{-\frac{N + |\alpha| + |\beta| + 1}{2}} dt \right) \\ &\lesssim \frac{d(x, y)^{1 - |\alpha| - |\beta|}}{V(x, y, d(x, y))}. \end{aligned}$$

The first integral in the parentheses is estimated by the change of variables $s = d(x, y)^2/t$, while the second one converges since $N + |\alpha| + |\beta| > 1$.

Case 2.1. Suppose that $N + |\alpha| + |\beta| > 2$. We apply the same argument to see that

$$\begin{aligned} & \int_0^\infty t^{-\frac{|\alpha|+|\beta|}{2}} V(x, y, \sqrt{t})^{-1} e^{-cd(x, y)^2/t} dt \\ & \lesssim \frac{1}{V(x, y, d(x, y))} \left(d(x, y)^N \int_0^{d(x, y)^2} t^{-\frac{N+|\alpha|+|\beta|}{2}} e^{-cd(x, y)^2/t} dt + d(x, y)^N \int_{d(x, y)^2}^\infty t^{-\frac{N+|\alpha|+|\beta|}{2}} dt \right) \\ & \lesssim \frac{d(x, y)^{2-|\alpha|-|\beta|}}{V(x, y, d(x, y))}. \end{aligned}$$

Since $d(x, y) \leq \|x - y\|$, it follows that

$$|\partial_x^\alpha \partial_y^\beta K_{\text{Dunkl}, j}(x, y)| \lesssim_{\alpha, \beta, R, k} \frac{d(x, y)}{\|x - y\|} \frac{d(x, y)^{-(|\alpha|+|\beta|)}}{V(x, y, d(x, y))}.$$

Case 2.2. Suppose that $N = 1$ and $|\alpha| + |\beta| = 1$. For the terms in which a derivative falls on $y_j - x_j$, retaining the factor $(1 + \|x - y\|/\sqrt{t})^{-2}$ and splitting the integral at $d(x, y)^2$ and $\|x - y\|^2$, we obtain from (2)

$$\begin{aligned} & \int_0^\infty t^{-3/2} \left(1 + \frac{\|x - y\|}{\sqrt{t}} \right)^{-2} V(x, y, \sqrt{t})^{-1} e^{-cd(x, y)^2/t} dt \\ & \lesssim \frac{d(x, y)^N}{\|x - y\|^2 V(x, y, d(x, y))} \int_0^{d(x, y)^2} t^{-\frac{N+1}{2}} e^{-cd(x, y)^2/t} dt \\ & \quad + \frac{d(x, y)}{\|x - y\|^2 V(x, y, d(x, y))} \int_{d(x, y)^2}^{\|x - y\|^2} \frac{dt}{t} + \frac{d(x, y)}{V(x, y, d(x, y))} \int_{\|x - y\|^2}^\infty \frac{dt}{t^2} \\ & \lesssim \frac{1}{\|x - y\| V(x, y, d(x, y))}, \end{aligned}$$

where we used

$$\frac{d(x, y)}{\|x - y\|} \log \frac{\|x - y\|}{d(x, y)} \lesssim 1.$$

Combining this estimate with the preceding estimate for the first term, we obtain

$$|\partial_x^\alpha \partial_y^\beta K_{\text{Dunkl}, j}(x, y)| \lesssim_{\alpha, \beta, R, k} \frac{1}{\|x - y\| V(x, y, d(x, y))} = \frac{d(x, y)}{\|x - y\|} \frac{d(x, y)^{-1}}{V(x, y, d(x, y))}.$$

This completes the proof of Lemma 1.3. $\square$

1.2. Systems of dyadic cubes.

Notation 1.4. For every dyadic cube $Q \in \mathbb{D}$, we denote by c_Q and $\ell(Q)$ its center and side-length, respectively.

Notation 1.5. Let

$$\mathbb{D}^0 := \bigcup_{k \in \mathbb{Z}} \mathbb{D}_k^0$$

be the standard system of dyadic cubes over $\mathbb{R}^N$. Here each dyadic cube is a product of half-open intervals

$$[m_1 2^{-k}, (m_1 + 1) 2^{-k}) \times \cdots \times [m_N 2^{-k}, (m_N + 1) 2^{-k}),$$

with $(m_1, \dots, m_N) \in \mathbb{Z}^N$. Moreover, we let

$$\mathbb{D}_\Gamma := \bigcup_{k \in \mathbb{Z}} \mathbb{D}_{\Gamma, k}$$

be the system of dyadic cubes over a fixed chamber Γ , whose construction will be given in Corollary 2.7.

1.3. Nearly weakly orthonormal sequences. Rochberg and Semmes [12] introduced the notion of *nearly weakly orthonormal* (NWO) sequences of functions. This notion has since proved highly effective in characterizing Schatten class membership of singular integral commutators in a variety of contexts (see [3, 8, 11, 1, 2, 9, 12, 14, 13]). The properties of NWO sequences were originally investigated in the Euclidean setting (equipped with Lebesgue measure), but the extension to spaces of homogeneous type is straightforward once a system of dyadic cubes is available in this general framework (see [8, Section 13] for a complete proof). For our purposes, we adapt the extension to the Dunkl setting $(\mathbb{R}^N, \|\cdot\|, w)$ rather than recalling the general definition of NWO sequences and the general extension to spaces of homogeneous type.

Definition 1.6. Let Γ be a chamber. Let Φ be as in Corollary 2.6 and let $\mathbb{D}_\Gamma$ be as in Corollary 2.7. Let $c > 0$ and $2 < r \leq \infty$. A sequence of functions $\{e_I\}_{I \in \mathbb{D}_\Gamma}$ is called (r, c) -NWO if

- (1) $\text{supp}(e_I) \subseteq cI$ for every $I \in \mathbb{D}_\Gamma$;
- (2) $\|e_I\|_{L_r(\mathbb{R}^N, w(x)dx)} \leq w(I)^{\frac{1}{r} - \frac{1}{2}}$ for every $I \in \mathbb{D}_\Gamma$;

Lemma 1.7. Let $0 < p < \infty$. Let Γ be a chamber. Let Φ be as in Corollary 2.6 and let $\mathbb{D}_\Gamma$ be as in Corollary 2.7. Suppose that A is an integral operator on $L_2(\Gamma, w(x)dx)$ with integral kernel

$$(x, y) \rightarrow \sum_{I \in \mathbb{D}_\Gamma} \lambda_I e_I(x) f_I(y), \quad x, y \in \Gamma,$$

where $\{e_I\}_{I \in \mathbb{D}_\Gamma}$ and $\{f_I\}_{I \in \mathbb{D}_\Gamma}$ are (r_1, c_1) -NWO and (r_2, c_2) -NWO sequences, respectively. We have

$$\begin{aligned} \|A\|_{\mathcal{L}_{p, \infty}(L_2(\Gamma, w(x)dx))} &\leq C_{\Phi, \kappa, r_1, c_1, r_2, c_2, p} \|\{\lambda_I\}_{I \in \mathbb{D}_\Gamma}\|_{p, \infty}, \\ \text{dist}_{\mathcal{L}_{p, \infty}(L_2(\Gamma, w(x)dx))} \left(A, (\mathcal{L}_{p, \infty})_0(L_2(\Gamma, w(x)dx)) \right) &\leq \\ &\leq C_{\Phi, \kappa, r_1, c_1, r_2, c_2, p} \text{dist}_{l_{p, \infty}(\mathbb{D}_\Gamma)} \left(\{\lambda_I\}_{I \in \mathbb{D}_\Gamma}, (l_{p, \infty})_0(\mathbb{D}_\Gamma) \right). \end{aligned}$$

Proof. state a similar lemma with Γ replaced by $\mathbb{R}_+^{n_1} \times \mathbb{R}^{n_2}$ and more-or-less-generic weight instead of w . Provide reference to the LATTER assertion. this assertion would trivially follow from the to-be-stated one. This property can be found in [12, (1.9)] (see also [8, Section 7]). $\square$

2. ANALYTIC TOOLS FOR WEYL CHAMBER GEOMETRY

Convention. Throughout this paper, we will use the following notation:

- Let $R \subset \mathbb{R}^N$ be a finite normalized reduced root system;
- Let $W := \text{span}(R)$ with $N_0 := \dim W$;
- Let $\Omega := \mathbb{R}_+^{N_0} \times \mathbb{R}^{N-N_0}$;

- Let Γ_+ be a fixed open Weyl chamber (see Definition 2.1) and Γ be its closure.

Definition 2.1. [6, Definition 2.7] *A connected component of*

$$(1) \quad \{x \in \mathbb{R}^N : \langle x, v \rangle \neq 0 \text{ for all } v \in R\}$$

is called an open Weyl chamber. Its closure is called a Weyl chamber. Define the corresponding sets of positive and negative roots by

$$R_+ := \{\alpha \in R : \langle x, \alpha \rangle > 0 \text{ for all } x \in \Gamma_+\}, \quad R_- := -R_+.$$

Definition 2.2. [6, Definition 2.9] *A root $\alpha \in R_+$ is called simple in R_+ if it cannot be written as*

$$\alpha = \lambda_1 \alpha_1 + \lambda_2 \alpha_2 \quad \text{with } \lambda_1, \lambda_2 \geq 1 \text{ and } \alpha_1, \alpha_2 \in R_+.$$

The set $\Delta \subset R_+$ of all simple roots is called the simple system.

Lemma 2.3. *With the notation given in Definition 2.2, Δ is a basis of W . If we denote by $\Delta = \{\alpha_1, \alpha_2, \dots, \alpha_{N_0}\}$, then the open and closed chambers can be written as*

$$(2) \quad \Gamma_+ = \left\{x \in \mathbb{R}^N : \langle x, \alpha_i \rangle > 0, i = 1, \dots, N_0\right\},$$

$$(3) \quad \Gamma = \left\{x \in \mathbb{R}^N : \langle x, \alpha_i \rangle \geq 0, i = 1, \dots, N_0\right\},$$

and every root $\alpha \in R$ admits a unique expansion

$$\alpha = \sum_{i=1}^{N_0} c_i \alpha_i$$

such that either $c_i \geq 0$ for all i (when $\alpha \in R_+$), or $c_i \leq 0$ for all i (when $\alpha \in R_-$). In particular, the simple system Δ is uniquely determined by Γ_+ (equivalently, by Γ).

Proof. Let $\Delta = \{\alpha_1, \dots, \alpha_n\} \subset R_+$ be the set of simple roots in R_+ . By [6, Proposition 2.10], every $\alpha \in R_+$ is a nonnegative linear combination of $\alpha_1, \dots, \alpha_n$. Since $R = R_+ \cup R_-$ (this follows from the fact that for each $\alpha \in R$, the continuous function $x \mapsto \langle x, \alpha \rangle$ does not vanish on the connected set Γ_+) and $R_- = -R_+$, it follows that $R \subset \text{span}(\Delta)$ and hence,

$$W = \text{span}(R) \subset \text{span}(\Delta).$$

The reverse inclusion $\text{span}(\Delta) \subset W$ is immediate from $\Delta \subset R \subset W$, so $\text{span}(\Delta) = W$. On the other hand, it follows from [6, Theorem 2.16] that the simple roots are linearly independent. Consequently, Δ is a basis of W , and therefore $n = \dim W = N_0$. Moreover, every $\alpha \in R$ has a unique expansion $\alpha = \sum_{i=1}^{N_0} c_i \alpha_i$. If $\alpha \in R_+$, the coefficients can be chosen nonnegative by [6, Proposition 2.10]. If $\alpha \in R_- = -R_+$, then $-\alpha \in R_+$ has an expansion with all coefficients nonnegative, and multiplying by -1 gives an expansion of α with all coefficients nonpositive. The descriptions (2) and (3) follow from [6, Corollary 2.11].

To see that Δ is uniquely determined by Γ_+ , note that Γ_+ determines R_+ by Definition 2.1, and the set of simple roots in R_+ is characterized intrinsically by Definition 2.2. Finally, since Γ_+ is the (unique) interior component of Γ , specifying Γ determines Γ_+ and vice versa. This completes the proof of Lemma 2.3. $\square$

Notation 2.4. With the notation given in Lemma 2.3, denote by

$$\Delta = \{\alpha_1, \dots, \alpha_{N_0}\}$$

the simple system associated with Γ .

Lemma 2.5. Each Weyl chamber Γ admits the decomposition

$$\Gamma = K(\Gamma) \times W^\perp, \quad K(\Gamma) := \{w \in W : \langle w, \alpha \rangle \geq 0 \text{ for all } \alpha \in \Delta\}.$$

Proof. Write $x = w + z$ with $w \in W$ and $z \in W^\perp$. Since $\alpha_i \in W$, we have $\langle x, \alpha_i \rangle = \langle w, \alpha_i \rangle$ for all $i = 1, \dots, N_0$. This, together with (3), implies that $x \in \Gamma$ if and only if $w \in K(\Gamma)$. This implies that $\Gamma = K(\Gamma) \times W^\perp$, and thus completes the proof. $\square$

Corollary 2.6. Let $\Phi_{W^\perp}$ be any linear isomorphism from $W^\perp$ to $\mathbb{R}^{N-N_0}$, and let

$$\Phi_W : W \longrightarrow \mathbb{R}^{N_0}, \quad \Phi_W(w) := (\langle w, \alpha_1 \rangle, \dots, \langle w, \alpha_{N_0} \rangle),$$

$$\Phi : \mathbb{R}^N = W \oplus W^\perp \longrightarrow \mathbb{R}^{N_0} \oplus \mathbb{R}^{N-N_0} \cong \mathbb{R}^N, \quad \Phi(w + z) := (\Phi_W(w), \Phi_{W^\perp}(z)).$$

Then Φ is a linear isomorphism and

$$\Phi(\Gamma) = \mathbb{R}_+^{N_0} \times \mathbb{R}^{N-N_0}.$$

Proof. By Lemma 2.5 we have $\Gamma = K(\Gamma) \times W^\perp$, where

$$K(\Gamma) = \{w \in W : \langle w, \alpha_i \rangle \geq 0, i = 1, \dots, N_0\}.$$

Since Δ is a basis of W , the Gram matrix $G = (\langle \alpha_i, \alpha_j \rangle)_{i,j=1}^{N_0}$ is nonsingular. Hence, $\Phi_W : W \longrightarrow \mathbb{R}^{N_0}$ is a linear isomorphism. Moreover, for $w \in W$ we have $w \in K(\Gamma)$ if and only if $\Phi_W(w) \in \mathbb{R}_+^{N_0}$, so

$$\Phi_W(K(\Gamma)) = \mathbb{R}_+^{N_0}.$$

By construction, Φ is a linear isomorphism (as a direct sum of isomorphisms), and therefore

$$\Phi(\Gamma) = \Phi(K(\Gamma) \times W^\perp) = \Phi_W(K(\Gamma)) \times \Phi_{W^\perp}(W^\perp) = \mathbb{R}_+^{N_0} \times \mathbb{R}^{N-N_0},$$

which proves the desired conclusion. $\square$

Corollary 2.7. Let Γ be an arbitrary Weyl chamber. There exists a system of dyadic cubes over Γ .

Proof. Let $\Phi : \mathbb{R}^N \rightarrow \mathbb{R}^N$ be the linear isomorphism from Corollary 2.6 so that

$$\Phi(\Gamma) = \Omega := \mathbb{R}_+^{N_0} \times \mathbb{R}^{N-N_0}.$$

For each $k \in \mathbb{Z}$, define $\mathbb{D}_{\Omega,k}^0$ to be the family of half-open cubes

$$Q = \prod_{i=1}^N I_i, \quad I_i = [m_i 2^{-k}, (m_i + 1) 2^{-k}),$$

where $m_i \in \mathbb{Z}_+$ for $1 \leq i \leq N_0$ and $m_i \in \mathbb{Z}$ for $N_0 < i \leq N$. Set

$$\mathbb{D}_\Omega^0 := \bigcup_{k \in \mathbb{Z}} \mathbb{D}_{\Omega,k}^0.$$

It is standard that $\mathbb{D}_\Omega^0$ is a system of dyadic cubes over Ω .

Next, for each $k \in \mathbb{Z}$, define

$$\mathbb{D}_{\Gamma,k} := \{\Phi^{-1}(Q) : Q \in \mathbb{D}_{\Omega,k}^0\}, \quad \mathbb{D}_{\Gamma} := \bigcup_{k \in \mathbb{Z}} \mathbb{D}_{\Gamma,k}.$$

Since $\Phi : \mathbb{R}^N \rightarrow \mathbb{R}^N$ is a linear isomorphism, we see that $\mathbb{D}_{\Gamma}$ is a system of dyadic cubes over Γ . $\square$

Notation 2.8. Let Φ be the linear isomorphism given in Corollary 2.6. For every Borel set $E \subset \Omega$, define

$$(4) \quad \mu(E) := w(\Phi^{-1}(E)).$$

Lemma 2.9. We have

$$(5) \quad d\mu(u, z) = \phi(u) du dz, \quad \phi(u) = |\det(\Phi)|^{-1} \prod_{v \in R_+} \left(\sum_{i=1}^{N_0} a_{v,i} u_i \right)^{2k(v)}.$$

where for each $v \in R_+$, the coefficients $a_{v,i} \geq 0$ are uniquely determined by the simple-root expansion $v = \sum_{i=1}^{N_0} a_{v,i} \alpha_i$ and satisfy $(a_{v,1}, \dots, a_{v,N_0}) \neq \mathbf{0}$.

Proof. Since $\Phi(\Gamma) = \Omega$, we have $\Phi^{-1}(\Omega) = \Gamma$. Hence, for every nonnegative Borel function f on Ω ,

$$\int_{\Omega} f(u, z) d\mu(u, z) = \int_{\Gamma} f(\Phi(x)) dw(x).$$

We make the linear change of variables $(u, z) = \Phi(x)$, so that $x = \Phi^{-1}(u, z)$. Since Φ is linear and invertible, the Jacobian is constant and

$$dx = |\det(\Phi)|^{-1} du dz.$$

It follows from the definition of Φ that $u_i = \langle x, \alpha_i \rangle$ for $i = 1, \dots, N_0$. This implies that

$$(6) \quad \langle x, v \rangle = \left\langle x, \sum_{i=1}^{N_0} a_{v,i} \alpha_i \right\rangle = \sum_{i=1}^{N_0} a_{v,i} \langle x, \alpha_i \rangle = \sum_{i=1}^{N_0} a_{v,i} u_i.$$

Substituting these identities into the previous integral, we obtain

$$\begin{aligned} \int_{\Omega} f(u, z) d\mu(u, z) &= \int_{\Gamma} f(\Phi(x)) \left(\prod_{v \in R_+} |\langle x, v \rangle|^{2k(v)} \right) dx \\ &= \int_{\Omega} f(u, z) |\det(\Phi)|^{-1} \prod_{v \in R_+} \left(\sum_{i=1}^{N_0} a_{v,i} u_i \right)^{2k(v)} du dz. \end{aligned}$$

Therefore, (5) holds. This completes the proof of Lemma 2.9. $\square$

Lemma 2.10. Let Γ_1 and Γ_2 be two Weyl chambers. Then there exist an integer $M \geq 0$ and $\alpha_1, \dots, \alpha_M \in R$ such that, if we set $\Gamma^{(0)} := \Gamma_2$ and

$$\Gamma^{(k)} := \sigma_{\alpha_k}(\Gamma^{(k-1)}), \quad k = 1, \dots, M,$$

then the following hold:

- (1) $\alpha_k \in \Delta(\Gamma^{(k-1)})$ for each $k = 1, \dots, M$;
- (2) $\Gamma^{(M)} = \Gamma_1$.

Proof. If $\Gamma_1 = \Gamma_2$, we take $M = 0$ and there is nothing to prove. Assume $\Gamma_1 \neq \Gamma_2$. Since the Weyl group G acts transitively on the set of Weyl chambers (see [6, Theorem 2.12 (2)]), there exists $g \in G$ such that $\Gamma_1 = g(\Gamma_2)$.

Let $\Delta(\Gamma_2) = \{\beta_1, \beta_2, \dots, \beta_{N_0}\}$ be the simple system associated with Γ_2 . By [6, Theorem 2.12(4)], the Weyl group G is generated by the simple reflections $\sigma_{\beta_1}, \sigma_{\beta_2}, \dots, \sigma_{\beta_{N_0}}$. Hence, there exist indices $i_1, \dots, i_M \in \{1, \dots, N_0\}$ such that

$$g = \sigma_{\beta_{i_1}} \sigma_{\beta_{i_2}} \cdots \sigma_{\beta_{i_M}}.$$

Define

$$w_0 := \text{Id}, \quad w_k := \sigma_{\beta_{i_1}} \cdots \sigma_{\beta_{i_k}} \quad (k = 1, \dots, M),$$

and set

$$\Gamma^{(k)} := w_k(\Gamma_2) \quad (k = 0, 1, \dots, M).$$

It follows that $\Gamma^{(0)} = \Gamma_2$ and $\Gamma^{(M)} = w_M(\Gamma_2) = g(\Gamma_2) = \Gamma_1$, which implies the statement (2).

For each $k = 1, \dots, M$, define

$$\alpha_k := w_{k-1}(\beta_{i_k}) \in R.$$

The rest of the proof is divided into two steps.

• **Step 1.** Verify that $\Gamma^{(k)} = \sigma_{\alpha_k}(\Gamma^{(k-1)})$.

Since each $w \in G$ is orthogonal, for any root $\beta \in R$ one has the conjugation identity

$$w \sigma_{\beta} w^{-1} = \sigma_{w(\beta)}, \quad w \in G.$$

Applying this with $w = w_{k-1}$ and $\beta = \beta_{i_k}$ yields

$$\sigma_{\alpha_k} = \sigma_{w_{k-1}(\beta_{i_k})} = w_{k-1} \sigma_{\beta_{i_k}} w_{k-1}^{-1}.$$

Therefore,

$$\sigma_{\alpha_k}(\Gamma^{(k-1)}) = w_{k-1} \sigma_{\beta_{i_k}} w_{k-1}^{-1}(w_{k-1}(\Gamma_2)) = w_{k-1} \sigma_{\beta_{i_k}}(\Gamma_2) = w_k(\Gamma_2) = \Gamma^{(k)}.$$

• **Step 2.** Verify that $\alpha_k \in \Delta(\Gamma^{(k-1)})$.

Recall that if Γ is a Weyl chamber associated with simple system $\Delta(\Gamma)$, then

$$\Gamma = \{x \in \mathbb{R}^N : \langle x, \gamma \rangle \geq 0 \text{ for all } \gamma \in \Delta(\Gamma)\}.$$

Applying $w_{k-1} \in G$ to the above description for Γ_2 and using $\langle w_{k-1}x, \beta \rangle = \langle x, w_{k-1}^{-1}\beta \rangle$, we obtain

$$\Gamma^{(k-1)} = w_{k-1}(\Gamma_2) = \left\{ x \in \mathbb{R}^N : \langle x, w_{k-1}(\beta_j) \rangle \geq 0 \text{ for } j = 1, \dots, N_0 \right\}.$$

By the uniqueness of the simple system associated with a Weyl chamber (see Lemma 2.3), we conclude that

$$\Delta(\Gamma^{(k-1)}) = \Delta(w_{k-1}(\Gamma_2)) = w_{k-1}(\Delta(\Gamma_2)) = \{w_{k-1}(\beta_1), \dots, w_{k-1}(\beta_{N_0})\}.$$

In particular, $\alpha_k = w_{k-1}(\beta_{i_k}) \in \Delta(\Gamma^{(k-1)})$, proving the first statement (1).

This completes the proof of Lemma 2.10. $\square$

Definition 2.11. We say that $R' \subset \Omega$ is an axis-parallel rectangle if

$$R' := I_1 \times \cdots \times I_N,$$

with $I_i = [I_i^-, I_i^+] \subset [0, \infty)$ for $1 \leq i \leq N_0$ and $I_i = [I_i^-, I_i^+] \subset \mathbb{R}$ for $i > N_0$.

Denote by

$$B_\Omega(x_0, r) := \{x \in \Omega : \|x - x_0\| < r\}.$$

The following lemma demonstrates that $(\Omega, \|\cdot\|, d\mu)$ is a doubling metric measure space.

Lemma 2.12. *There exists a constant $C \geq 1$ such that for every $x_0 = (u_0, z_0) \in \Omega$ and every $r > 0$,*

$$(7) \quad C^{-1} r^N \prod_{v \in R_+} (L_{v,0} + r)^{2k(v)} \leq \mu(B_\Omega(x_0, r)) \leq C r^N \prod_{v \in R_+} (L_{v,0} + r)^{2k(v)},$$

where

$$L_v(u) := \sum_{i=1}^{N_0} a_{v,i} u_i, \quad L_{v,0} := L_v(u_0) = \sum_{i=1}^{N_0} a_{v,i} (u_0)_i.$$

In particular, there exists a constant $C_{\text{dbl}, \Omega} \geq 1$ such that for every $x \in \Omega$ and every $r > 0$,

$$\mu(B_\Omega(x, 2r)) \leq C_{\text{dbl}, \Omega} \mu(B_\Omega(x, r)).$$

Proof. Fix $x_0 = (u_0, z_0) \in \Omega$ and $r > 0$. For each $v \in R_+$ define the linear form

$$L_v(u) := a_v \cdot u, \quad a_v := (a_{v,1}, \dots, a_{v,N_0}) \in \mathbb{R}_+^{N_0} \setminus \{0\}.$$

Since R_+ is finite and each $a_v \neq 0$, we have

$$L_* := \max_{v \in R_+} \|a_v\| < \infty, \quad m_* := \min_{v \in R_+} \sum_{i=1}^{N_0} a_{v,i} > 0.$$

Upper bound. Let $x = (u, z) \in B_\Omega(x_0, r)$. Then

$$\|u - u_0\| \leq \|x - x_0\| \leq r.$$

Hence, for every $v \in R_+$,

$$|L_v(u) - L_{v,0}| = |a_v \cdot (u - u_0)| \leq \|a_v\| \cdot \|u - u_0\| \leq L_* r.$$

Since $L_v(u) \geq 0$ on Ω (all coefficients and coordinates are nonnegative), we obtain

$$L_v(u) \leq L_{v,0} + L_* r \leq C_1 (L_{v,0} + r), \quad C_1 := \max\{1, L_*\}.$$

Therefore,

$$\phi(u) = \prod_{v \in R_+} L_v(u)^{2k(v)} \leq C_1^{2 \sum_{v \in R_+} k(v)} \prod_{v \in R_+} (L_{v,0} + r)^{2k(v)} =: C_2 \prod_{v \in R_+} (L_{v,0} + r)^{2k(v)}.$$

Integrating over $B_\Omega(x_0, r)$ and using $|B_\Omega(x_0, r)| \leq |B(x_0, r)| = \omega_N r^N$, where ω_N is the volume of the unit ball in $\mathbb{R}^N$, gives

$$\mu(B_\Omega(x_0, r)) = \int_{B_\Omega(x_0, r)} \phi(u) dx \leq C_2 |B(x_0, r)| \prod_{v \in R_+} (L_{v,0} + r)^{2k(v)} \leq C_3 r^N \prod_{v \in R_+} (L_{v,0} + r)^{2k(v)},$$

with $C_3 := C_2 \omega_N$.

Lower bound. We construct a smaller Euclidean ball B_c contained in $B_\Omega(x_0, r)$ on which ϕ admits a uniform positive lower bound in terms of $\prod_{v \in R_+} (L_{v,0} + r)^{2k(v)}$.

To this end, we first choose

$$\lambda := 1 + \frac{L_*}{m_*} > 1, \quad \eta := \frac{1}{4(\sqrt{N_0} \lambda + 1)}, \quad \theta := \lambda \eta.$$

These parameters depend only on L_* and m_* , and satisfy

$$\theta > \eta > 0, \quad \sqrt{N_0} \theta + \eta = \eta(\sqrt{N_0} \lambda + 1) = \frac{1}{4} < 1.$$

Let $\mathbf{1}_{N_0} := (1, \dots, 1) \in \mathbb{R}^{N_0}$ and define the center

$$x_c := (u_c, z_c) := (u_0 + \theta r \mathbf{1}_{N_0}, z_0), \quad B_c := B(x_c, \eta r).$$

We *claim* that $B_c \subset B_\Omega(x_0, r)$. To see this inclusion, we first apply the triangle inequality to see that for any $x \in B_c$,

$$\|x - x_0\| \leq \|x - x_c\| + \|x_c - x_0\| \leq \eta r + \theta r \sqrt{N_0} \leq r,$$

since $\sqrt{N_0} \theta + \eta < 1$. Hence $B_c \subset B(x_0, r)$. Moreover, for any $x = (u, z) \in B_c$, we have $\|u - u_c\| \leq \|x - x_c\| \leq \eta r$. This implies that for each $1 \leq i \leq N_0$,

$$u_i \geq (u_c)_i - \eta r = (u_0)_i + (\theta - \eta)r \geq 0,$$

since $u_0 \in \mathbb{R}_+^{N_0}$ and $\theta \geq \eta$. Thus $u \in \mathbb{R}_+^{N_0}$ and $x \in \Omega$. Therefore, $B_c \subset B_\Omega(x_0, r)$. This finishes the proof of the claim.

Next we obtain a uniform lower bound for $L_v(u)$ on B_c . Write any $x = (u, z) \in B_c$ as

$$u = u_0 + \theta r \mathbf{1}_{N_0} + \Delta_u, \quad z = z_0 + \Delta_z, \quad \|\Delta_u\|^2 + \|\Delta_z\|^2 \leq (\eta r)^2,$$

so in particular $\|\Delta_u\| \leq \eta r$. Then for all $x = (u, z) \in B_c$ and all $v \in R_+$,

$$\begin{aligned} L_v(u) &= a_v \cdot u = a_v \cdot u_0 + \theta r a_v \cdot \mathbf{1}_{N_0} + a_v \cdot \Delta_u \\ &\geq L_{v,0} + \theta r \sum_{i=1}^{N_0} a_{v,i} - \|a_v\| \|\Delta_u\| \geq L_{v,0} + r K_0 \geq c_0 (L_{v,0} + r). \end{aligned}$$

with

$$K_0 := \theta m_* - \eta L_* = \eta(\lambda m_* - L_*) = \eta m_* > 0, \quad c_0 := \min\{1, K_0\} \in (0, 1].$$

Consequently, for all $u \in B_c$,

$$\phi(u) = \prod_{v \in R_+} L_v(u)^{2k(v)} \geq c_0^{2 \sum_{v \in R_+} k(v)} \prod_{v \in R_+} (L_{v,0} + r)^{2k(v)} =: C_4 \prod_{v \in R_+} (L_{v,0} + r)^{2k(v)}.$$

This, in combination with the fact that $B_c \subset B_\Omega(x_0, r)$, yields

$$\mu(B_\Omega(x_0, r)) \geq \mu(B_c) = \int_{B_c} \phi(u) dx \geq C_4 |B_c| \prod_{v \in R_+} (L_{v,0} + r)^{2k(v)} = C_5 r^N \prod_{v \in R_+} (L_{v,0} + r)^{2k(v)},$$

where $C_5 := C_4 \omega_N \eta^N > 0$.

The second statement follows immediately from the first one. Indeed, by the first statement,

$$\mu(B_\Omega(x_0, 2r)) \leq C_3 (2r)^N \prod_{v \in R_+} (L_{v,0} + 2r)^{2k(v)} \leq C_3 2^{N+\mathcal{K}} r^N \prod_{v \in R_+} (L_{v,0} + r)^{2k(v)} \leq C_3 C_5^{-1} 2^{N+\mathcal{K}} \mu(B_\Omega(x_0, r)),$$

where we set $\mathcal{K} := 2 \sum_{v \in R_+} k(v)$. Therefore, $(\Omega, \|\cdot\|, \mu)$ is doubling with the doubling constant $C_{\text{dbl}, \Omega} := C_3 C_5^{-1} 2^{N+\mathcal{K}}$. This finishes the proof of Lemma 2.12. $\square$

Let $\Phi : \mathbb{R}^N \longrightarrow \mathbb{R}^N$ be the linear isomorphism given in Corollary 2.6 such that $\Phi(\Gamma) = \Omega$. Since Φ is linear and invertible, there exist positive constants c_Φ and C_Φ such that for any $x, y \in \mathbb{R}^N$,

$$(8) \quad c_\Phi \|x - y\| \leq \|\Phi(x) - \Phi(y)\| \leq C_\Phi \|x - y\|.$$

Denote by $B_\Gamma(x, r) := \{y \in \Gamma : \|y - x\| < r\}$. It follows from (8) that for any $x_0 \in \Gamma$ and $r > 0$,

$$(9) \quad B_\Omega(\Phi(x_0), c_\Phi r) \subset \Phi(B_\Gamma(x_0, r)) \subset B_\Omega(\Phi(x_0), C_\Phi r).$$

This, together with Lemma 2.12, yields

$$(10) \quad \mu(\Phi(B_\Gamma(x_0, r))) \simeq \mu(B_\Omega(\Phi(x_0), C_\Phi r)).$$

Lemma 2.13. *For all $x_0 \in \Gamma$ and $r > 0$, one can find an axis-parallel rectangle R' on Ω such that*

$$(11) \quad B_\Omega(\Phi(x_0), C_\Phi r) \subset R' \subset B_\Omega(\Phi(x_0), \sqrt{N} C_\Phi r).$$

Furthermore, we have

$$(12) \quad \mu(R') \simeq \mu(B_\Omega(\Phi(x_0), C_\Phi r)).$$

Proof. We write $(u_0, z_0) := \Phi(x_0) \in \Omega$. and define the axis-parallel rectangle

$$(13) \quad R' := \left(\prod_{i=1}^{N_0} [\max\{0, (u_0)_i - C_\Phi r\}, (u_0)_i + C_\Phi r] \right) \times \left(\prod_{j=1}^{N-N_0} [(z_0)_j - C_\Phi r, (z_0)_j + C_\Phi r] \right).$$

On the one hand, it follows immediately that $B_\Omega(\Phi(x_0), C_\Phi r) \subset R'$. On the other hand, since R' is a rectangle with half-side lengths bounded by $C_\Phi r$ in each coordinate, we have

$$R' \subset B_\Omega(\Phi(x_0), \sqrt{N} C_\Phi r).$$

The second statement follows from the first one together with Lemma 2.12. The proof of Lemma 2.13 is finished. $\square$

The following theorem demonstrates that $(\Gamma, \|\cdot\|, dw)$ is a doubling metric measure space.

Theorem 2.14. *There exists a constant $C \geq 1$ such that for every $x \in \Gamma$ and every $r > 0$,*

$$(14) \quad C^{-1} r^N \prod_{v \in R_+} (|\langle x, v \rangle| + r)^{2k(v)} \leq w(B_\Gamma(x, r)) \leq C r^N \prod_{v \in R_+} (|\langle x, v \rangle| + r)^{2k(v)}.$$

In particular, there exists a constant $C_{\text{dbl}, \Gamma} \geq 1$ such that for every $x \in \Gamma$ and every $r > 0$,

$$w(B_\Gamma(x, 2r)) \leq C_{\text{dbl}, \Gamma} w(B_\Gamma(x, r)).$$

Moreover, for all $x \in \Gamma$ and $r > 0$, we have

$$(15) \quad w(B_\Gamma(x, r)) \sim w(B(x, r)).$$

Proof. The main strategy of the proof is to transfer the ball estimates from $(\Omega, \|\cdot\|, \mu)$ back to $(\Gamma, \|\cdot\|, dw)$. Fix $x \in \Gamma$ and $r > 0$. It follows from the equality (4) that

$$(16) \quad w(B_\Gamma(x, r)) = \mu(\Phi(B_\Gamma(x, r))).$$

Applying μ to the inclusions (9) and using (16) yields

$$(17) \quad \mu(B_\Omega(\Phi(x), c_\Phi r)) \leq w(B_\Gamma(x, r)) \leq \mu(B_\Omega(\Phi(x), C_\Phi r)).$$

Write $\Phi(x) = (u, z) \in \Omega$. For each $v \in R_+$ recall the linear form $L_{v,0} := \sum_{i=1}^{N_0} a_{v,i} u_i$. Combining the inequality (17) with Lemma 2.12 and the equality (6), we obtain that

$$(18) \quad w(B_\Gamma(x, r)) \simeq r^N \prod_{v \in R_+} (L_{v,0} + r)^{2k(v)} = r^N \prod_{v \in R_+} (|\langle x, v \rangle| + r)^{2k(v)}.$$

This proves (14). The second statement follows immediately from (14). The third statement follows from the inequalities (14) and (1). This completes the proof of Theorem 2.14. $\square$

3. SPECTRAL ASYMPTOTIC FORMULA

Definition 3.1. For $g \in G$ and $j = 1, \dots, N$, set

$$d_{g,j} := g^{-1}e_j, \quad f_g(x) := f(gx).$$

For $\alpha \in R_+$, define $S_\alpha : \mathbb{C}^G \rightarrow \mathbb{C}^G$ by

$$(S_\alpha z)_g := z_{g\sigma_\alpha}, \quad g \in G.$$

For $f \in \dot{W}^{1,N}(\mathbb{R}^N)$, $x \in \Gamma_+$, and $\theta \in \mathbb{S}^{N-1}$, define

$$\begin{aligned} \mathfrak{c}_{f,j}(x, \theta) &:= \text{diag}_{g \in G} \left(\langle d_{g,j}, \nabla f_g(x) \rangle - \langle d_{g,j}, \theta \rangle \langle \theta, \nabla f_g(x) \rangle \right) \\ &\quad + \sum_{\alpha \in R_+} k(\alpha) \text{diag}_{g \in G} \left(\frac{\langle \alpha, d_{g,j} \rangle}{\langle \alpha, x \rangle} (f_g(x) - f_{g\sigma_\alpha}(x)) \right) S_\alpha. \end{aligned}$$

For smooth f , the quotients in the second term are understood by their smooth extensions across the corresponding reflection hyperplanes. For $f \in \dot{W}^{1,N}(\mathbb{R}^N)$, $\mathfrak{c}_{f,j}$ is understood almost everywhere on $\Gamma \times \mathbb{S}^{N-1}$.

Lemma 3.2. Let $f \in W_{\text{loc}}^{1,1}(\mathbb{R}^N)$. For every $u \in C_c^\infty(\mathbb{R}^N)$,

$$(1) \quad [T_{e_j}, M_f]u(x) = \partial_j f(x)u(x) + \sum_{\alpha \in R_+} k(\alpha) \langle \alpha, e_j \rangle \frac{f(x) - f(\sigma_\alpha x)}{\langle \alpha, x \rangle} u(\sigma_\alpha x)$$

for almost every $x \in \mathbb{R}^N$. If $N > 1$ and $f \in \dot{W}^{1,N}(\mathbb{R}^N)$, then, for every $\alpha \in R_+$,

$$(2) \quad \left\| \frac{f - f \circ \sigma_\alpha}{\langle \alpha, \cdot \rangle} \right\|_{L^N(\mathbb{R}^N, dx)} \lesssim_{N,G} \|\nabla f\|_{L^N(\mathbb{R}^N, dx)}.$$

Proof. Pairing α and $-\alpha$ in the definition of T_{e_j} , and using $k(-\alpha) = k(\alpha)$ and $\sigma_{-\alpha} = \sigma_\alpha$, we have

$$T_{e_j}u(x) = \partial_j u(x) + \sum_{\alpha \in R_+} k(\alpha) \langle \alpha, e_j \rangle \frac{u(x) - u(\sigma_\alpha x)}{\langle \alpha, x \rangle}.$$

Therefore,

$$[T_{e_j}, M_f]u(x) = \partial_j f(x)u(x) + \sum_{\alpha \in R_+} k(\alpha) \langle \alpha, e_j \rangle \frac{f(x) - f(\sigma_\alpha x)}{\langle \alpha, x \rangle} u(\sigma_\alpha x),$$

which proves (1).

Since R is normalized,

$$x - \sigma_\alpha x = \langle \alpha, x \rangle \alpha, \quad \|x - \sigma_\alpha x\| = \sqrt{2} |\langle \alpha, x \rangle|.$$

Hence, for almost every $x \notin \alpha^\perp$,

$$\frac{|f(x) - f(\sigma_\alpha x)|}{|\langle \alpha, x \rangle|} = \sqrt{2} \mathcal{G}_{I, \sigma_\alpha} f(x).$$

By Lemma ?? (??),

$$\left\| \frac{f - f \circ \sigma_\alpha}{\langle \alpha, \cdot \rangle} \right\|_{L^N(\mathbb{R}^N, dx)} \lesssim_{N, G} \|\nabla f\|_{L^N(\mathbb{R}^N, dx)}.$$

This completes the proof of Lemma 3.2. $\square$

Lemma 3.3. *Let Γ_+ be an open Weyl chamber, let Γ be its closure, and let R_+ be the corresponding positive subsystem. Set*

$$\rho(x) := \varpi(x)^{1/2},$$

and define

$$\begin{aligned} U : L^2(\mathbb{R}^N, dw) &\longrightarrow L^2(\Gamma, dw; \mathbb{C}^G), & (Uu)_g(x) &:= u(gx), \\ V : L^2(\Gamma, dw; \mathbb{C}^G) &\longrightarrow L^2(\Gamma, dx; \mathbb{C}^G), & (Vh)(x) &:= \rho(x)h(x). \end{aligned}$$

Then U and V are unitary. If

$$\tilde{T}_j := VUT_{e_j}U^{-1}V^{-1}, \quad \tilde{P} := VU(-\Delta_{\text{Dunkl}})U^{-1}V^{-1},$$

then, on $C_c^\infty(\Gamma_+; \mathbb{C}^G)$,

$$(3) \quad (\tilde{T}_j h)_g = \partial_{d_{g,j}} h_g - \sum_{\alpha \in R_+} k(\alpha) \frac{\langle \alpha, d_{g,j} \rangle}{\langle \alpha, x \rangle} h_{g\sigma_\alpha},$$

$$(4) \quad \tilde{P} = -\Delta_{\mathbb{C}^G} + |\nabla \log \rho|^2 I_{\mathbb{C}^G} - 2 \sum_{\alpha \in R_+} k(\alpha) \frac{S_\alpha}{\langle \alpha, x \rangle^2}.$$

Assume in addition that $N > 1$, and let

$$\tilde{Q} := VU(-\Delta_{\text{Dunkl}})^{-1/2}U^{-1}V^{-1},$$

where $(-\Delta_{\text{Dunkl}})^{-1/2}$ is defined by the spectral calculus. Then, for every $\chi \in C_c^\infty(\Gamma_+)$, the operator $M_\chi \tilde{Q} M_\chi$, initially defined on its natural domain, admits a bounded extension to $L^2(\Gamma, dx; \mathbb{C}^G)$, and this extension is a classical matrix-valued pseudodifferential operator of order -1 . Its principal symbol is

$$\sigma_{-1}(M_\chi \tilde{Q} M_\chi)(x, \xi) = \chi(x)^2 |\xi|^{-1} I_{\mathbb{C}^G}.$$

Consequently, if $x \in \Gamma_+$, $\xi \neq 0$, and $\chi \in C_c^\infty(\Gamma_+)$ is identically 1 in a neighborhood of x , then

$$(5) \quad \sigma_{-1}(M_\chi \tilde{Q} M_\chi)(x, \xi) = |\xi|^{-1} I_{\mathbb{C}^G}.$$

Proof. The Weyl chambers $g\Gamma_+$, $g \in G$, are pairwise disjoint and cover $\mathbb{R}^N$ up to the reflection hyperplanes; see [6, Theorem 2.12]. Since dw is G -invariant,

$$\begin{aligned}\|Uu\|_{L^2(\Gamma, dw; \mathbb{C}^G)}^2 &= \sum_{g \in G} \int_{\Gamma} |u(gx)|^2 dw(x) \\ &= \int_{\mathbb{R}^N} |u(x)|^2 dw(x).\end{aligned}$$

Conversely, for $h = (h_g)_{g \in G} \in L^2(\Gamma, dw; \mathbb{C}^G)$, define $u(gx) := h_g(x)$ for $x \in \Gamma_+$. This determines u almost everywhere on $\mathbb{R}^N$ and gives $Uu = h$. Hence U is unitary. Since $dw = \rho^2 dx$ and $\rho > 0$ on Γ_+ , V is unitary as well.

Pairing the roots α and $-\alpha$ in the definition of T_{e_j} , changing variables in the root system by g , and using the G -invariance of k , we obtain, for $h \in C_c^\infty(\Gamma_+; \mathbb{C}^G)$,

$$(UT_{e_j}U^{-1}h)_g = \partial_{d_{g,j}} h_g + \sum_{\alpha \in R_+} k(\alpha) \frac{\langle \alpha, d_{g,j} \rangle}{\langle \alpha, x \rangle} (h_g - h_{g\sigma_\alpha}).$$

On Γ_+ ,

$$\nabla \log \rho(x) = \sum_{\alpha \in R_+} k(\alpha) \frac{\alpha}{\langle \alpha, x \rangle},$$

and therefore

$$\rho \partial_{d_{g,j}} \rho^{-1} = \partial_{d_{g,j}} - \sum_{\alpha \in R_+} k(\alpha) \frac{\langle \alpha, d_{g,j} \rangle}{\langle \alpha, x \rangle}.$$

Conjugation by V cancels the diagonal terms and gives (3).

Similarly, pairing α and $-\alpha$ in the formula for the Dunkl Laplacian and using $\|\alpha\|^2 = 2$, we have

$$\begin{aligned}(U(-\Delta_{\text{Dunkl}})U^{-1}h)_g &= -\Delta h_g - 2 \sum_{\alpha \in R_+} k(\alpha) \frac{\partial_\alpha h_g}{\langle \alpha, x \rangle} \\ &\quad + 2 \sum_{\alpha \in R_+} k(\alpha) \frac{h_g - h_{g\sigma_\alpha}}{\langle \alpha, x \rangle^2}.\end{aligned}$$

Writing $\phi = \log \rho$,

$$\rho(-\Delta - 2\nabla \phi \cdot \nabla) \rho^{-1} = -\Delta + |\nabla \phi|^2 + \Delta \phi,$$

while

$$\Delta \phi = - \sum_{\alpha \in R_+} k(\alpha) \frac{\|\alpha\|^2}{\langle \alpha, x \rangle^2} = -2 \sum_{\alpha \in R_+} \frac{k(\alpha)}{\langle \alpha, x \rangle^2}.$$

The latter term cancels the diagonal part of

$$2 \sum_{\alpha \in R_+} k(\alpha) \frac{I_{\mathbb{C}^G} - S_\alpha}{\langle \alpha, x \rangle^2},$$

and hence

$$\tilde{P} = -\Delta I_{\mathbb{C}^G} + |\nabla \log \rho|^2 I_{\mathbb{C}^G} - 2 \sum_{\alpha \in R_+} k(\alpha) \frac{S_\alpha}{\langle \alpha, x \rangle^2},$$

which proves (4).

Assume now that $N > 1$, fix $\chi \in C_c^\infty(\Gamma_+)$, and set

$$K := \text{supp } \chi.$$

On Γ_+ , write

$$\tilde{P} = -\Delta_{\mathbb{C}^G} + W(x), \quad W(x) = |\nabla \log \rho(x)|^2 I_{\mathbb{C}^G} - 2 \sum_{\alpha \in R_+} k(\alpha) \frac{S_\alpha}{\langle \alpha, x \rangle^2}.$$

Thus W is a smooth Hermitian matrix-valued function on Γ_+ .

Choose open sets

$$K \Subset \Omega_0 \Subset \Omega_1 \Subset \Gamma_+$$

and $\eta \in C_c^\infty(\Gamma_+)$ such that

$$0 \leq \eta \leq 1, \quad \eta = 1 \quad \text{on a neighborhood of } \overline{\Omega_1}.$$

Choose a flat torus M containing $\text{supp } \eta$ in one Euclidean coordinate chart. Since η is compactly supported in that chart, the differential operator $M_\eta \tilde{P} M_\eta$ extends smoothly by zero to M . Define on $L^2(M; \mathbb{C}^G)$

$$(6) \quad P_0 := M_\eta \tilde{P} M_\eta + M_{1-\eta}(-\Delta_M + 1)M_{1-\eta}.$$

It is a nonnegative self-adjoint elliptic differential operator of order 2. Indeed,

$$\langle P_0 u, u \rangle = \langle \tilde{P}(\eta u), \eta u \rangle + \|\nabla((1-\eta)u)\|_2^2 + \|(1-\eta)u\|_2^2 \geq 0,$$

and its principal symbol is

$$p_{0,2}(x, \xi) = (\eta(x)^2 + (1-\eta(x))^2)|\xi|^2 I_{\mathbb{C}^G},$$

which is elliptic since

$$\eta^2 + (1-\eta)^2 \geq \frac{1}{2}.$$

Moreover,

$$(7) \quad P_0 = \tilde{P} \quad \text{on } \Omega_1.$$

Let

$$H_t := e^{-t\tilde{P}}, \quad H_t^0 := e^{-tP_0}.$$

We first consider

$$B_0 := \frac{1}{\sqrt{\pi}} \int_0^1 t^{-1/2} H_t^0 dt.$$

Let Π_0 be the orthogonal projection onto $\ker P_0$. Since P_0 is elliptic on the closed manifold M , Π_0 is smoothing. Applying the complex powers calculus [?, Theorem 2 and p. 290] to $P_0 + \Pi_0$, we obtain

$$(P_0 + \Pi_0)^{-1/2} \in \Psi_{\text{cl}}^{-1}(M; \mathbb{C}^G)$$

with principal symbol $p_{0,2}^{-1/2}$.

Set

$$F(\lambda) := \frac{1}{\sqrt{\pi}} \int_0^1 t^{-1/2} e^{-t\lambda} dt, \quad \lambda \geq 0.$$

Then $B_0 = F(P_0)$, and for $\lambda > 0$,

$$F(\lambda) - \lambda^{-1/2} = -\frac{1}{\sqrt{\pi}} \int_1^\infty t^{-1/2} e^{-t\lambda} dt.$$

Hence

$$F(\lambda) - \lambda^{-1/2} = O(\lambda^{-m}) \quad (\lambda \rightarrow \infty)$$

for every $m > 0$. On $\ker P_0$ the difference between B_0 and $(P_0 + \Pi_0)^{-1/2}$ is finite rank. It follows from elliptic spectral regularity that

$$B_0 - (P_0 + \Pi_0)^{-1/2} \in \Psi^{-\infty}(M; \mathbb{C}^G).$$

Consequently,

$$(8) \quad B_0 \in \Psi_{\text{cl}}^{-1}(M; \mathbb{C}^G), \quad \sigma_{-1}(B_0)(x, \xi) = p_{0,2}(x, \xi)^{-1/2}.$$

In particular, by (7),

$$(9) \quad \sigma_{-1}(B_0)(x, \xi) = |\xi|^{-1} I_{\mathbb{C}^G}, \quad x \in \Omega_1.$$

We next compare H_t and H_t^0 in the interior. Choose $\zeta \in C_c^\infty(\Omega_1)$ such that

$$\zeta = 1 \quad \text{on a neighborhood of } K.$$

Since $P_0 = \tilde{P}$ on Ω_1 , Duhamel's formula gives

$$(10) \quad M_\chi(H_t - H_t^0)M_\chi = - \int_0^t M_\chi H_{t-s}[\tilde{P}, M_\zeta]H_s^0 M_\chi ds.$$

Indeed, for data supported in K ,

$$v(t) := M_\zeta H_t^0 M_\chi u$$

satisfies

$$(\partial_t + \tilde{P})v = [\tilde{P}, M_\zeta]H_t^0 M_\chi u, \quad v(0) = M_\chi u,$$

which yields (10).

Let $h_t(x, y)$ be the Dunkl heat kernel. The integral kernel of H_t , with respect to Lebesgue measure on Γ , is the matrix

$$(11) \quad \tilde{h}_t(x, y)_{g, \ell} = \rho(x)\rho(y)h_t(gx, \ell y), \quad g, \ell \in G.$$

For $x, y \in \Gamma$,

$$d(gx, \ell y) = d(x, y) = |x - y|.$$

Together with the mixed derivative estimate for the Dunkl heat kernel [?, Theorem 3.1], this implies that, whenever $E, F \Subset \Gamma_+$ and $\text{dist}(E, F) > 0$, for every pair of multi-indices α, β and every $m > 0$,

$$(12) \quad \sup_{\substack{x \in E \\ y \in F}} \left\| \partial_x^\alpha \partial_y^\beta \tilde{h}_t(x, y) \right\| = O(t^m), \quad t \downarrow 0.$$

The heat kernel of the smooth elliptic operator P_0 satisfies the same off-diagonal estimate on M .

Since

$$\text{supp}[\tilde{P}, M_\zeta] \cap K = \emptyset,$$

the two heat kernels occurring in (10) are evaluated between compact sets separated by a positive distance. The standard Gaussian bounds underlying (12) therefore give, for every multi-indices α, β and every $m > 0$,

$$(13) \quad \sup_{x, y} \left\| \partial_x^\alpha \partial_y^\beta K_{M_\chi(H_t - H_t^0)M_\chi}(x, y) \right\| = O(t^m), \quad t \downarrow 0.$$

Hence

$$(14) \quad S_0 := \frac{1}{\sqrt{\pi}} \int_0^1 t^{-1/2} M_\chi (H_t - H_t^0) M_\chi dt \in \Psi^{-\infty}.$$

It remains to treat the large-time part. Let $K_1 \Subset \Gamma_+$ be a compact set containing K . By (11), the heat-kernel derivative estimate cited above, and (2), for $x, y \in K_1$ and $t \geq 1$,

$$V(gx, \ell y, \sqrt{t})^{-1} \lesssim_{K_1} t^{-N/2}.$$

Since ρ and all its derivatives are bounded on K_1 , it follows that, for every pair of multi-indices α, β ,

$$(15) \quad \sup_{x, y \in K_1} \left\| \partial_x^\alpha \partial_y^\beta \tilde{h}_t(x, y) \right\| \lesssim_{\alpha, \beta, K_1} t^{-N/2}, \quad t \geq 1.$$

Since $N > 1$,

$$\int_1^\infty t^{-1/2} t^{-N/2} dt < \infty.$$

Thus all derivatives of

$$K_{S_\infty}(x, y) := \frac{\chi(x)\chi(y)}{\sqrt{\pi}} \int_1^\infty t^{-1/2} \tilde{h}_t(x, y) dt$$

converge locally uniformly, and therefore

$$(16) \quad S_\infty := \frac{1}{\sqrt{\pi}} \int_1^\infty t^{-1/2} M_\chi H_t M_\chi dt \in \Psi^{-\infty}.$$

For completeness, the preceding integrals give the canonical bounded localization of $\tilde{P}^{-1/2}$. Indeed, for $\varepsilon > 0$,

$$(\tilde{P} + \varepsilon)^{-1/2} = \frac{1}{\sqrt{\pi}} \int_0^\infty t^{-1/2} e^{-\varepsilon t} H_t dt$$

by the spectral theorem. The estimates (13) and (15) imply that

$$M_\chi (\tilde{P} + \varepsilon)^{-1/2} M_\chi$$

converges in operator norm, as $\varepsilon \downarrow 0$, to

$$\frac{1}{\sqrt{\pi}} \int_0^\infty t^{-1/2} M_\chi H_t M_\chi dt.$$

On the natural domain of $\tilde{P}^{-1/2}$, this limit agrees with $M_\chi \tilde{P}^{-1/2} M_\chi$; hence it is its bounded extension.

Combining (14), (16), and the definition of B_0 , we obtain

$$M_\chi \tilde{Q} M_\chi = M_\chi B_0 M_\chi + S_0 + S_\infty.$$

By (8), $M_\chi B_0 M_\chi$ is a classical matrix-valued pseudodifferential operator of order -1 , while $S_0 + S_\infty$ is smoothing. Therefore

$$M_\chi \tilde{Q} M_\chi \in \Psi_{\text{cl}}^{-1},$$

and, by (9),

$$\sigma_{-1}(M_\chi \tilde{Q} M_\chi)(x, \xi) = \chi(x)^2 |\xi|^{-1} I_{\mathbb{C}^G}.$$

If $\chi = 1$ in a neighborhood of x , this reduces to

$$\sigma_{-1}(M_\chi \tilde{Q} M_\chi)(x, \xi) = |\xi|^{-1} I_{\mathbb{C}^G},$$

which proves (5). $\square$

Lemma 3.4. *Let $j \in \{1, \dots, N\}$ and $f \in C_c^\infty(\mathbb{R}^N)$, and set*

$$A_{f,j} := VU[R_{\text{Dunkl},j}, M_f]U^{-1}V^{-1}.$$

Then, for every $\chi \in C_c^\infty(\Gamma_+)$,

$$M_\chi A_{f,j} M_\chi$$

is a compactly supported classical matrix valued pseudodifferential operator of order -1 , with principal symbol

$$(17) \quad \sigma_{-1}(M_\chi A_{f,j} M_\chi)(x, \xi) = \frac{\chi(x)^2}{\|\xi\|} \mathbf{c}_{f,j} \left(x, \frac{\xi}{\|\xi\|} \right), \quad x \in \Gamma_+, \quad \xi \neq 0.$$

Proof. Set

$$F(x) := \text{diag}_{g \in G} (f_g(x)).$$

Since $VUM_fU^{-1}V^{-1} = M_F$ and $R_{\text{Dunkl},j} = T_{e_j}(-\Delta_{\text{Dunkl}})^{-1/2}$,

$$(18) \quad A_{f,j} = [\tilde{T}_j, M_F] \tilde{Q} + \tilde{T}_j [\tilde{Q}, M_F].$$

By (3),

$$(19) \quad \begin{aligned} [\tilde{T}_j, M_F] &= \text{diag}_{g \in G} (\langle d_{g,j}, \nabla f_g \rangle) \\ &+ \sum_{\alpha \in R_+} k(\alpha) \text{diag}_{g \in G} \left(\frac{\langle \alpha, d_{g,j} \rangle}{\langle \alpha, x \rangle} (f_g - f_{g\sigma_\alpha}) \right) S_\alpha. \end{aligned}$$

Fix $\chi \in C_c^\infty(\Gamma_+)$. Choose $\eta, \psi \in C_c^\infty(\Gamma_+)$ such that $\eta = 1$ on a neighborhood of $\text{supp } \chi$ and $\psi = 1$ on a neighborhood of $\text{supp } \eta$, and set

$$Q_\psi := M_\psi \tilde{Q} M_\psi.$$

By Lemma 3.3, Q_ψ is a classical matrix valued pseudodifferential operator of order -1 , and on a neighborhood of $\text{supp } \eta$,

$$(20) \quad q_{-1}(x, \xi) = \|\xi\|^{-1} I_{\mathbb{C}^G}.$$

We use the Kohn–Nirenberg convention

$$\text{Op}(a)u(x) = (2\pi)^{-N} \int_{\mathbb{R}^{2N}} e^{i(x-y) \cdot \xi} a(x, \xi) u(y) dy d\xi.$$

If a and b are classical symbols, the composition formula [?, Theorem 18.1.8, p. 71] gives

$$(21) \quad a \# b \sim \sum_{\beta \in \mathbb{N}_0^N} \frac{1}{\beta!} \partial_\xi^\beta a D_x^\beta b, \quad D_x := \frac{1}{i} \partial_x.$$

Write, on a neighborhood of $\text{supp } \eta$,

$$\sigma(Q_\psi)(x, \xi) \sim q_{-1}(x, \xi) + q_{-2}(x, \xi) + \dots$$

By (4),

$$\sigma(\tilde{P})(x, \xi) \sim p_2(x, \xi) + p_1(x, \xi) + p_0(x, \xi),$$

where

$$p_2(x, \xi) = \|\xi\|^2 I_{\mathbb{C}^G}, \quad p_1(x, \xi) = 0.$$

Since $\tilde{Q} = \tilde{P}^{-1/2}$, $\tilde{Q}\tilde{P}\tilde{Q} = I$. Applying (21) on the region where $\psi = 1$ and comparing the homogeneous terms of degree -1 , the terms containing x -derivatives of p_2 and q_{-1} vanish, and hence

$$q_{-2}p_2q_{-1} + q_{-1}p_2q_{-2} = 0.$$

Together with (20), this gives

$$(22) \quad q_{-2}(x, \xi) = 0$$

on a neighborhood of $\text{supp } \eta$.

Applying (21) to the commutator $[Q_\psi, M_F]$, and using (20)–(22), we obtain, on the same neighborhood,

$$(23) \quad \begin{aligned} \sigma_{-2}([Q_\psi, M_F])(x, \xi) &= \frac{1}{i} \sum_{\ell=1}^N \partial_{\xi_\ell} (\|\xi\|^{-1}) \partial_{x_\ell} F(x) \\ &= i \|\xi\|^{-3} \text{diag}_{g \in G} (\langle \xi, \nabla f_g(x) \rangle). \end{aligned}$$

Since $\psi = 1$ on a neighborhood of $\text{supp } \eta$ and $\eta = 1$ on a neighborhood of $\text{supp } \chi$,

$$\begin{aligned} M_\chi[\tilde{T}_j, M_F]\tilde{Q}M_\chi &= M_\chi[\tilde{T}_j, M_F]Q_\psi M_\chi, \\ M_\chi\tilde{T}_j[\tilde{Q}, M_F]M_\chi &= M_\chi\tilde{T}_jM_\eta[Q_\psi, M_F]M_\chi. \end{aligned}$$

Both operators on the right are classical matrix valued pseudodifferential operators of order -1 .

By (19), (20), and the order-zero part of the composition formula,

$$(24) \quad \begin{aligned} \sigma_{-1}(M_\chi[\tilde{T}_j, M_F]\tilde{Q}M_\chi)(x, \xi) &= \frac{\chi(x)^2}{\|\xi\|} \text{diag}_{g \in G} (\langle d_{g,j}, \nabla f_g(x) \rangle) \\ &+ \frac{\chi(x)^2}{\|\xi\|} \sum_{\alpha \in R_+} k(\alpha) \text{diag}_{g \in G} \left(\frac{\langle \alpha, d_{g,j} \rangle}{\langle \alpha, x \rangle} (f_g(x) - f_{g\sigma_\alpha}(x)) \right) S_\alpha. \end{aligned}$$

Under the above Kohn–Nirenberg convention, (3) gives

$$\sigma_1(\tilde{T}_j)(x, \xi) = i \text{diag}_{g \in G} (\langle d_{g,j}, \xi \rangle).$$

Therefore, by (23),

$$(25) \quad \begin{aligned} \sigma_{-1}(M_\chi\tilde{T}_j[\tilde{Q}, M_F]M_\chi)(x, \xi) &= -\chi(x)^2 \|\xi\|^{-3} \text{diag}_{g \in G} (\langle d_{g,j}, \xi \rangle \langle \xi, \nabla f_g(x) \rangle). \end{aligned}$$

Combining (24) and (25), and setting

$$\theta := \frac{\xi}{\|\xi\|},$$

we obtain

$$\begin{aligned} \sigma_{-1}(M_\chi A_{f,j} M_\chi)(x, \xi) &= \frac{\chi(x)^2}{\|\xi\|} \text{diag}_{g \in G} \left(\langle d_{g,j}, \nabla f_g(x) \rangle - \langle d_{g,j}, \theta \rangle \langle \theta, \nabla f_g(x) \rangle \right) \\ &+ \frac{\chi(x)^2}{\|\xi\|} \sum_{\alpha \in R_+} k(\alpha) \text{diag}_{g \in G} \left(\frac{\langle \alpha, d_{g,j} \rangle}{\langle \alpha, x \rangle} (f_g(x) - f_{g\sigma_\alpha}(x)) \right) S_\alpha \\ &= \frac{\chi(x)^2}{\|\xi\|} \mathbf{c}_{f,j}(x, \theta). \end{aligned}$$

This proves (17). $\square$

Lemma 3.5. *Assume that $N > 1$, $j \in \{1, \dots, N\}$, and $f \in \dot{W}^{1,N}(\mathbb{R}^N)$. Then*

$$\left(\int_{\Gamma} \int_{\mathbb{S}^{N-1}} \operatorname{tr}_{\mathbb{C}^G} |\mathbf{c}_{f,j}(x, \theta)|^N d\theta dx \right)^{1/N} \simeq_{N,R,k} \|\nabla f\|_{L^N(\mathbb{R}^N, dx)}.$$

Proof. By the definition of $\mathbf{c}_{f,j}$, the unitary invariance of the Schatten norm, and $\|d_{g,j}\| = 1$,

$$\begin{aligned} \|\mathbf{c}_{f,j}(x, \theta)\|_{S^N} &\lesssim_{R,k} \left(\sum_{g \in G} |\nabla f_g(x)|^N \right)^{1/N} \\ &\quad + \sum_{\alpha \in R_+} \left(\sum_{g \in G} \left| \frac{f_g(x) - f_{g\sigma_\alpha}(x)}{\langle \alpha, x \rangle} \right|^N \right)^{1/N}. \end{aligned}$$

The chamber decomposition used in Lemma 3.3 and the orthogonality of $g \in G$ imply

$$\sum_{g \in G} \int_{\Gamma} |\nabla f_g(x)|^N dx = \|\nabla f\|_{L^N(\mathbb{R}^N, dx)}^N.$$

Moreover, since $\|\alpha\|^2 = 2$,

$$|x - \sigma_\alpha x| = \sqrt{2} |\langle \alpha, x \rangle|,$$

and hence

$$\left| \frac{f_g(x) - f_{g\sigma_\alpha}(x)}{\langle \alpha, x \rangle} \right| = \sqrt{2} \mathcal{G}_{g, g\sigma_\alpha} f(x)$$

for almost every $x \in \mathbb{R}^N$. By Lemma ??,

$$\int_{\Gamma} \left| \frac{f_g(x) - f_{g\sigma_\alpha}(x)}{\langle \alpha, x \rangle} \right|^N dx \lesssim_{N,G} \|\nabla f\|_{L^N(\mathbb{R}^N, dx)}^N$$

uniformly in $g \in G$ and $\alpha \in R_+$. Since G and R_+ are finite,

$$(26) \quad \left(\int_{\Gamma} \int_{\mathbb{S}^{N-1}} \operatorname{tr}_{\mathbb{C}^G} |\mathbf{c}_{f,j}(x, \theta)|^N d\theta dx \right)^{1/N} \lesssim_{N,R,k} \|\nabla f\|_{L^N(\mathbb{R}^N, dx)}.$$

For the converse estimate, let P_g denote the orthogonal projection of $\mathbb{C}^G$ onto the g -th coordinate and define

$$\mathcal{E}(A) := \sum_{g \in G} P_g A P_g, \quad A \in \mathcal{L}(\mathbb{C}^G).$$

Thus, $\mathcal{E}(A)$ is the diagonal part of A . The pinching inequality for unitarily invariant norms [?, p. 97, (IV.51)–(IV.52)] gives

$$(27) \quad \|\mathcal{E}(A)\|_{S^N} \leq \|A\|_{S^N}.$$

For every $\alpha \in R_+$, S_α has zero diagonal, since $g\sigma_\alpha \neq g$ for all $g \in G$. Hence, by the definition of $\mathbf{c}_{f,j}$,

$$\mathcal{E}(\mathbf{c}_{f,j}(x, \theta)) = \operatorname{diag}_{g \in G} \left(\langle d_{g,j}, \nabla f_g(x) \rangle - \langle d_{g,j}, \theta \rangle \langle \theta, \nabla f_g(x) \rangle \right).$$

Therefore, by (27),

$$(28) \quad \|\mathbf{c}_{f,j}(x, \theta)\|_{S^N}^N \geq \sum_{g \in G} |\langle d_{g,j}, \nabla f_g(x) \rangle - \langle d_{g,j}, \theta \rangle \langle \theta, \nabla f_g(x) \rangle|^N.$$

For each $g \in G$, set

$$y := gx, \quad \eta := g\theta.$$

Since

$$\nabla f_g(x) = g^{-1} \nabla f(y), \quad d_{g,j} = g^{-1} e_j,$$

and g is orthogonal,

$$\begin{aligned} \langle d_{g,j}, \nabla f_g(x) \rangle &= \partial_j f(y), \\ \langle d_{g,j}, \theta \rangle &= \eta_j, \\ \langle \theta, \nabla f_g(x) \rangle &= \langle \eta, \nabla f(y) \rangle. \end{aligned}$$

Using the chamber decomposition and the orthogonal invariance of surface measure, (28) therefore gives

$$(29) \quad \begin{aligned} & \int_{\Gamma} \int_{\mathbb{S}^{N-1}} \|\mathbf{c}_{f,j}(x, \theta)\|_{S^N}^N d\theta dx \\ & \geq \int_{\mathbb{R}^N} \int_{\mathbb{S}^{N-1}} |\partial_j f(y) - \eta_j \langle \eta, \nabla f(y) \rangle|^N d\eta dy. \end{aligned}$$

For $v \in \mathbb{C}^N$, set

$$\Phi_j(v) := \left(\int_{\mathbb{S}^{N-1}} |v_j - \eta_j \langle \eta, v \rangle|^N d\eta \right)^{1/N}.$$

Then

$$(30) \quad \Phi_j(v) \simeq_N \|v\|.$$

Indeed, Φ_j is a seminorm on $\mathbb{C}^N$. If $\Phi_j(v) = 0$, continuity gives

$$v_j - \eta_j \langle \eta, v \rangle = 0, \quad \eta \in \mathbb{S}^{N-1}.$$

Since $N > 1$, there exists $\eta \in \mathbb{S}^{N-1}$ with $\eta_j = 0$, and therefore $v_j = 0$. It follows that

$$\langle \eta, v \rangle = 0$$

whenever $\eta_j \neq 0$. Since such η form a dense subset of $\mathbb{S}^{N-1}$, continuity yields $\langle \eta, v \rangle = 0$ for every $\eta \in \mathbb{S}^{N-1}$, and hence $v = 0$. Thus Φ_j is a norm, and (30) follows from the equivalence of norms on the finite dimensional space $\mathbb{C}^N$.

Applying (30) with $v = \nabla f(y)$ in (29), we obtain

$$\|\nabla f\|_{L^N(\mathbb{R}^N, dx)} \lesssim_N \left(\int_{\Gamma} \int_{\mathbb{S}^{N-1}} \text{tr}_{\mathbb{C}^G} |\mathbf{c}_{f,j}(x, \theta)|^N d\theta dx \right)^{1/N}.$$

Combining this estimate with (26) proves the assertion. $\square$

Lemma 3.6. *Assume that $N > 1$, $j \in \{1, \dots, N\}$, and $f \in C_c^\infty(\mathbb{R}^N)$. Define*

$$A_f := VU[R_{\text{Dunkl},j}, M_f]U^{-1}V^{-1}.$$

Then there exist $\chi_m \in C_c^\infty(\Gamma_+)$, $0 \leq \chi_m \leq 1$, such that $\chi_m(x) \rightarrow 1$ for every $x \in \Gamma_+$ and

$$(31) \quad \lim_{m \rightarrow \infty} \|A_f - M_{\chi_m} A_f M_{\chi_m}\|_{S^{N,\infty}} = 0.$$

Proof. Fix $s > 2$, choose $2 < s_0 < s$, and set

$$\gamma := \frac{1}{s_0} - \frac{1}{s} > 0.$$

Let $r_s > s$ be as in Lemma ?? and set $\beta := N/r_s$. By Lemma ?? (??), $f \in OSC^{N,\infty}(\mathbb{R}^N)$, so Lemma ?? applies with $p = N$ and $q = \infty$.

Set

$$B_f := U[R_{\text{Dunkl},j}, M_f]U^{-1}.$$

Since $V = M_\rho$ commutes with scalar multiplication operators on Γ ,

$$(32) \quad \|A_f - M_\chi A_f M_\chi\|_{S^{N,\infty}} = \|B_f - M_\chi B_f M_\chi\|_{S^{N,\infty}}$$

for every bounded scalar multiplier χ .

For fixed $\sigma, \tau \in G$, $u, v \in \mathbb{Z}^N$, and $\varepsilon \in \{0, 1, 2, 3\}$, define

$$T_{u,v,\varepsilon}^{\sigma,\tau} := \sum_{I \in \mathbb{D}_\Gamma} A_{\sigma,\tau,I,\varepsilon}(f) \left\langle \cdot, \overline{H_{u,v,I,\varepsilon}^{\sigma,\tau}(\tau \cdot)} \right\rangle_{L^2(\Gamma, dw)} E_{u,v,I,\varepsilon}^{\sigma,\tau}(\sigma \cdot).$$

By the operator series identification in the proof of Theorem ??, the (σ, τ) matrix entry of B_f is

$$\sum_{\varepsilon=0}^3 \sum_{u,v \in \mathbb{Z}^N} c_{u,v} T_{u,v,\varepsilon}^{\sigma,\tau},$$

with convergence in $S^{N,\infty}$. Moreover, by Lemma ?? (??)–(??),

$$(33) \quad \text{supp } E_{u,v,I,\varepsilon}^{\sigma,\tau}(\sigma \cdot) \subset I, \quad \text{supp } H_{u,v,I,\varepsilon}^{\sigma,\tau}(\tau \cdot) \subset I^*,$$

(34)

$$\|E_{u,v,I,\varepsilon}^{\sigma,\tau}(\sigma \cdot)\|_{L^s(dw)} + \|H_{u,v,I,\varepsilon}^{\sigma,\tau}(\tau \cdot)\|_{L^s(dw)} \lesssim w(I)^{1/s-1/2}.$$

The definitions of $A_{\sigma,\tau,I,\varepsilon}(f)$ in the proof of Lemma ?? imply

$$(35) \quad |A_{\sigma,\tau,I,\varepsilon}(f)| \lesssim_f \begin{cases} \ell(I), & \ell(I) \leq 1, \\ \ell(I)^{-\beta}, & \ell(I) \geq 1. \end{cases}$$

Indeed, for $\varepsilon = 0, 1, 2$, the corresponding coefficients are bounded by an $L^{r_s}(dx)$ oscillation of a smooth compactly supported function on a set of diameter comparable with $\ell(I)$. Hence the first estimate follows from the Lipschitz bound, while for $\ell(I) \geq 1$,

$$\text{osc}_{r_s}(g; E) \lesssim |E|^{-1/r_s} \|g\|_{L^{r_s}} \lesssim_f \ell(I)^{-N/r_s}.$$

For $\varepsilon = 3$, writing $D := |\sigma z_I - \tau z_I|$, the definitions of $\theta_{\sigma,\tau,I}$ and $a_{\sigma,\tau,I}(f)$ imply, for $\ell(I) \leq 1$,

$$|A_{\sigma,\tau,I,3}(f)| \lesssim_f \frac{\ell(I)}{\ell(I) + D} (\ell(I) + D) \lesssim_f \ell(I),$$

whereas, for $\ell(I) \geq 1$,

$$|A_{\sigma,\tau,I,3}(f)| \leq |I|^{-1} \|H_{\sigma,\tau}(f)\|_{L^1} \lesssim_f \ell(I)^{-N} \lesssim \ell(I)^{-\beta}.$$

Let $\Delta = \{\alpha_1, \dots, \alpha_{N_0}\}$. For $1 \leq i \leq N_0$ and $0 < \delta \leq 1$, set

$$W_{i,\delta} := \{x \in \Gamma : 0 \leq \langle x, \alpha_i \rangle \leq 2\delta\}.$$

We have

$$(36) \quad \frac{w(W_{i,\delta} \cap I)}{w(I)} + \frac{w(W_{i,\delta} \cap I^*)}{w(I)} \lesssim \min\left(1, \frac{\delta}{\ell(I)}\right).$$

To prove this, write $I = \Phi^{-1}(Q)$ and $\ell := \ell(I)$. The assertion is immediate when $\delta > \ell/8$, since $w(I^*) \lesssim w(I)$. Suppose that $\delta \leq \ell/8$. If I or I^* meets $W_{i,\delta}$, then the i -th coordinate interval of Q is $[0, \ell)$. By Lemma 2.9 (5), the density of $\Phi_{\#}dw$ is

$$\phi(u) = |\det(\Phi)|^{-1} \prod_{v \in R_+} \left(\sum_{h=1}^{N_0} a_{v,h} u_h \right)^{2k(v)},$$

which is nondecreasing in every u_i . If g is nonnegative and nondecreasing on an interval of length comparable with ℓ , then

$$\int_0^{2\delta} g(t) dt \lesssim \frac{\delta}{\ell} \int g(t) dt.$$

Applying this estimate fiberwise to ϕ on Q , respectively on $(2Q) \cap \Omega$, and using $w(I^*) \lesssim w(I)$, we obtain (36).

Define

$$\omega_{I,\delta} := \min\left(1, \left(\frac{\delta}{\ell(I)}\right)^\gamma\right).$$

If $0 \leq a \leq 1$ and $\text{supp } a \subset W_{i,\delta}$, then by Hölder's inequality, (34), and (36),

$$\|aE_{u,v,I,\varepsilon}^{\sigma,\tau}(\sigma \cdot)\|_{L^{s_0}(dw)} + \|aH_{u,v,I,\varepsilon}^{\sigma,\tau}(\tau \cdot)\|_{L^{s_0}(dw)} \lesssim \omega_{I,\delta} w(I)^{1/s_0-1/2}.$$

After normalization by $\omega_{I,\delta}$, the localized factors are NWO families with exponent s_0 . Hence, by Lemma ??,

$$(37) \quad \|M_a T_{u,v,\varepsilon}^{\sigma,\tau}\|_{S^{N,\infty}} + \|T_{u,v,\varepsilon}^{\sigma,\tau} M_a\|_{S^{N,\infty}} \lesssim \left\| \left\{ A_{\sigma,\tau,I,\varepsilon}(f) \omega_{I,\delta} \right\}_{I \in \mathbb{D}_\Gamma} \right\|_{\ell^{N,\infty}}.$$

If $\ell(I) \leq 1$ and $A_{\sigma,\tau,I,\varepsilon}(f) \neq 0$, then z_I belongs to a fixed bounded set depending only on f . Consequently, at every dyadic length $0 < \ell \leq 1$,

$$(38) \quad \#\left\{ I \in \mathbb{D}_\Gamma : \ell(I) = \ell, A_{\sigma,\tau,I,\varepsilon}(f) \neq 0, (I \cup I^*) \cap W_{i,\delta} \neq \emptyset \right\} \lesssim_f \left(1 + \frac{\delta}{\ell}\right) \ell^{-(N-1)}.$$

Indeed, under Φ , there are $O(1 + \delta/\ell)$ admissible intervals in the u_i coordinate and $O(\ell^{-1})$ admissible intervals in each of the remaining $N - 1$ coordinates.

For $\ell(I) \leq \delta$, (35) and (38) imply, for every $\lambda > 0$,

$$\begin{aligned} & \# \left\{ I : \ell(I) \leq \delta, |A_{\sigma,\tau,I,\varepsilon}(f)\omega_{I,\delta}| > \lambda \right\} \\ & \lesssim_f \delta \sum_{\substack{\ell \leq \delta \\ \lambda \lesssim_f \ell}} \ell^{-N} \lesssim_f \delta \lambda^{-N}. \end{aligned}$$

Therefore,

$$(39) \quad \left\| \left\{ A_{\sigma,\tau,I,\varepsilon}(f)\omega_{I,\delta} \right\}_{\ell(I) \leq \delta} \right\|_{\ell^{N,\infty}} \lesssim_f \delta^{1/N}.$$

For $\delta < \ell(I) \leq 1$,

$$|A_{\sigma,\tau,I,\varepsilon}(f)\omega_{I,\delta}| \lesssim_f \delta^\gamma \ell(I)^{1-\gamma},$$

and (38) reduces to $O_f(\ell^{-(N-1)})$. Given $\lambda > 0$, set

$$L := \max \left(\delta, c_f \left(\frac{\lambda}{\delta^\gamma} \right)^{1/(1-\gamma)} \right).$$

Since $N > 1$,

$$\# \left\{ I : \delta < \ell(I) \leq 1, |A_{\sigma,\tau,I,\varepsilon}(f)\omega_{I,\delta}| > \lambda \right\} \lesssim_f L^{-(N-1)}.$$

If $L = \delta$, then $\lambda \lesssim_f \delta$, and hence

$$\lambda L^{-(N-1)/N} \lesssim_f \delta^{1/N}.$$

Otherwise, $\lambda \lesssim_f \delta^\gamma L^{1-\gamma}$, so

$$\lambda L^{-(N-1)/N} \lesssim_f \delta^\gamma L^{1/N-\gamma} \lesssim_f \delta^{\min(\gamma, 1/N)}.$$

Thus,

$$(40) \quad \left\| \left\{ A_{\sigma,\tau,I,\varepsilon}(f)\omega_{I,\delta} \right\}_{\delta < \ell(I) \leq 1} \right\|_{\ell^{N,\infty}} \lesssim_f \delta^{\min(\gamma, 1/N)}.$$

For $\ell(I) \geq 1$, only $O_f(1)$ cubes at each dyadic scale have nonzero coefficients.

By (35),

$$\sum_{\ell(I) \geq 1} |A_{\sigma,\tau,I,\varepsilon}(f)\omega_{I,\delta}|^N \lesssim_f \delta^{\gamma N} \sum_{\substack{\ell \geq 1 \\ \text{dyadic}}} \ell^{-N(\beta+\gamma)} \lesssim_f \delta^{\gamma N}.$$

Combining this with (39) and (40), we obtain

$$(41) \quad \left\| \left\{ A_{\sigma,\tau,I,\varepsilon}(f)\omega_{I,\delta} \right\}_{I \in \mathbb{D}_\Gamma} \right\|_{\ell^{N,\infty}} \lesssim_f \delta^\kappa, \quad \kappa := \min \left(\gamma, \frac{1}{N} \right).$$

By (37),

$$(42) \quad \|M_a T_{u,v,\varepsilon}^{\sigma,\tau}\|_{S^{N,\infty}} + \|T_{u,v,\varepsilon}^{\sigma,\tau} M_a\|_{S^{N,\infty}} \lesssim_f \delta^\kappa$$

whenever $0 \leq a \leq 1$ and $\text{supp } a \subset W_{i,\delta}$.

We also need localization at infinity. Let $0 \leq b_R \leq 1$ with $\text{supp } b_R \subset \{x \in \Gamma : |x| \geq R\}$. If $A_{\sigma,\tau,I,\varepsilon}(f) \neq 0$, then

$$(43) \quad |z_I| \lesssim_f 1 + \ell(I).$$

For $\varepsilon = 0, 1$, the ball $B_\Gamma(z_I, C_1 \ell(I))$ must meet the support of $f \circ \tau$; for $\varepsilon = 2, 3$, I must meet the support of $H_{\sigma,\tau}(f)$. Since $|x - z_I| \lesssim \ell(I)$ for $x \in I^*$, (43) shows that, for sufficiently large R ,

$$A_{\sigma,\tau,I,\varepsilon}(f) \neq 0, \quad (I \cup I^*) \cap \{x : |x| \geq R\} \neq \emptyset$$

can occur only when $\ell(I) \gtrsim_f R$. At each such dyadic scale there are $O_f(1)$ cubes with nonzero coefficients. Hence, by (35) and Lemma ??,

$$(44) \quad \|M_{b_R} T_{u,v,\varepsilon}^{\sigma,\tau}\|_{S^{N,\infty}} + \|T_{u,v,\varepsilon}^{\sigma,\tau} M_{b_R}\|_{S^{N,\infty}} \lesssim_f R^{-\beta}.$$

Choose $\eta \in C^\infty(\mathbb{R})$, $0 \leq \eta \leq 1$, such that

$$\eta(t) = 0 \quad \text{for } t \leq 1, \quad \eta(t) = 1 \quad \text{for } t \geq 2,$$

and choose $\zeta \in C_c^\infty(\mathbb{R}^N)$, $0 \leq \zeta \leq 1$, such that

$$\zeta = 1 \quad \text{on } B(0, 1), \quad \text{supp } \zeta \subset B(0, 2).$$

For $0 < \delta \leq 1$ and $R \geq 1$, define

$$\chi_{\delta,R}(x) := \zeta(x/R) \prod_{i=1}^{N_0} \eta\left(\frac{\langle x, \alpha_i \rangle}{\delta}\right), \quad x \in \Gamma_+.$$

Then $\chi_{\delta,R} \in C_c^\infty(\Gamma_+)$ and $0 \leq \chi_{\delta,R} \leq 1$. Moreover,

$$\begin{aligned} 1 - \chi_{\delta,R} &= 1 - \zeta(\cdot/R) \\ &+ \sum_{i=1}^{N_0} \zeta(\cdot/R) \prod_{h < i} \eta\left(\frac{\langle \cdot, \alpha_h \rangle}{\delta}\right) \left(1 - \eta\left(\frac{\langle \cdot, \alpha_i \rangle}{\delta}\right)\right). \end{aligned}$$

The first term is supported in $\{x : |x| \geq R\}$, while the i -th term in the sum is supported in $W_{i,\delta}$. Since G is finite, $S^{N,\infty}$ is normable for $N > 1$, and Lemma ?? (??) implies

$$\sum_{u,v \in \mathbb{Z}^N} |c_{u,v}| < \infty,$$

we may sum (42) and (44) over the operator series for B_f . Therefore,

$$\|(I - M_{\chi_{\delta,R}})B_f\|_{S^{N,\infty}} + \|B_f(I - M_{\chi_{\delta,R}})\|_{S^{N,\infty}} \lesssim_f \delta^\kappa + R^{-\beta}.$$

Since

$$B_f - M_{\chi_{\delta,R}} B_f M_{\chi_{\delta,R}} = (I - M_{\chi_{\delta,R}})B_f + M_{\chi_{\delta,R}} B_f (I - M_{\chi_{\delta,R}}),$$

the ideal property of $S^{N,\infty}$ and (32) imply

$$\|A_f - M_{\chi_{\delta,R}} A_f M_{\chi_{\delta,R}}\|_{S^{N,\infty}} \lesssim_f \delta^\kappa + R^{-\beta}.$$

Define

$$\chi_m := \chi_{2^{-m}, 2^m}.$$

Then $\chi_m(x) \rightarrow 1$ for every $x \in \Gamma_+$, while

$$\|A_f - M_{\chi_m} A_f M_{\chi_m}\|_{S^{N,\infty}} \lesssim_f 2^{-m\kappa} + 2^{-m\beta} \rightarrow 0.$$

This completes the proof of Lemma 3.6. $\square$

Lemma 3.7. *Assume that $N > 1$, $j \in \{1, \dots, N\}$, and $f \in C_c^\infty(\mathbb{R}^N)$. Then*

$$(45) \quad \lim_{t \rightarrow \infty} t^{1/N} \mu(t, [R_{\text{Dunkl},j}, M_f]) = \left(\frac{1}{N(2\pi)^N} \int_{\Gamma} \int_{\mathbb{S}^{N-1}} \text{tr}_{\mathbb{C}^G} |\mathbf{c}_{f,j}(x, \theta)|^N d\theta dx \right)^{1/N}.$$

Proof. Define

$$A_f := VU[R_{\text{Dunkl},j}, M_f]U^{-1}V^{-1}.$$

By Lemma 3.6 (31), there exist $\chi_m \in C_c^\infty(\Gamma_+)$, $0 \leq \chi_m \leq 1$, such that $\chi_m(x) \rightarrow 1$ for every $x \in \Gamma_+$ and

$$(46) \quad \|A_f - M_{\chi_m} A_f M_{\chi_m}\|_{S^{N,\infty}} \rightarrow 0.$$

Set

$$A_{f,m} := M_{\chi_m} A_f M_{\chi_m}, \quad K_m := \text{supp } \chi_m.$$

By Lemma 3.4 (17), $A_{f,m}$ is a compactly supported classical matrix valued pseudodifferential operator of order -1 on Γ_+ , with

$$(47) \quad \sigma_{-1}(A_{f,m})(x, \xi) = \frac{\chi_m(x)^2}{\|\xi\|} \mathbf{c}_{f,j}\left(x, \frac{\xi}{\|\xi\|}\right), \quad x \in \Gamma_+, \quad \xi \neq 0.$$

Choose a bounded open set U_m with $K_m \Subset U_m \Subset \Gamma_+$, and choose $L_m > 0$ such that

$$L_m > \sup_{x,y \in U_m} \|x - y\|_\infty.$$

For

$$\mathbb{T}_m^N := \mathbb{R}^N / L_m \mathbb{Z}^N,$$

the quotient map restricts to a diffeomorphism

$$\kappa_m : U_m \rightarrow \Omega_m := \kappa_m(U_m),$$

which is a local isometry. Since the Schwartz kernel of $A_{f,m}$ is supported in $K_m \times K_m$, with respect to

$$L^2(\Gamma; \mathbb{C}^G) = L^2(U_m; \mathbb{C}^G) \oplus L^2(\Gamma \setminus U_m; \mathbb{C}^G)$$

we have

$$A_{f,m} = B_{f,m} \oplus 0$$

for some $B_{f,m}$ on $L^2(U_m; \mathbb{C}^G)$.

Let

$$W_m : L^2(U_m; \mathbb{C}^G) \rightarrow L^2(\Omega_m; \mathbb{C}^G), \quad W_m u := u \circ \kappa_m^{-1}.$$

Then W_m is unitary. By the invariance of classical pseudodifferential operators under smooth coordinate changes [?, Theorem 18.1.17],

$$T_{f,m} := W_m B_{f,m} W_m^{-1} \in \Psi_{\text{cl}}^{-1}(\Omega_m; \mathbb{C}^G).$$

Its Schwartz kernel is supported in $\kappa_m(K_m) \times \kappa_m(K_m) \Subset \Omega_m \times \Omega_m$. Choose $\vartheta_m \in C_c^\infty(\Omega_m)$ equal to 1 on a neighborhood of $\kappa_m(K_m)$. Then

$$T_{f,m} = M_{\vartheta_m} T_{f,m} M_{\vartheta_m}.$$

Hence the zero extension of the Schwartz kernel of $T_{f,m}$ from $\Omega_m \times \Omega_m$ to $\mathbb{T}_m^N \times \mathbb{T}_m^N$ is a classical pseudodifferential kernel of order -1 : locally in $\Omega_m \times \Omega_m$ this follows

from $T_{f,m} \in \Psi_{\text{cl}}^{-1}$, while near the boundary of $\Omega_m \times \Omega_m$ the kernel vanishes. Denote the corresponding operator on $\mathbb{T}_m^N$ by $\widehat{A}_{f,m}$. Thus,

$$(48) \quad \widehat{A}_{f,m} \in \Psi_{\text{cl}}^{-1}(\mathbb{T}_m^N; \mathbb{C}^G).$$

Since κ_m is a local isometry, by (47) and the transformation law for principal symbols,

$$(49) \quad \sigma_{-1}(\widehat{A}_{f,m})(\kappa_m(x), \xi) = \frac{\chi_m(x)^2}{\|\xi\|} \mathbf{c}_{f,j} \left(x, \frac{\xi}{\|\xi\|} \right), \quad x \in U_m, \quad \xi \neq 0,$$

and the principal symbol vanishes outside Ω_m .

With respect to

$$L^2(\mathbb{T}_m^N; \mathbb{C}^G) = L^2(\Omega_m; \mathbb{C}^G) \oplus L^2(\mathbb{T}_m^N \setminus \Omega_m; \mathbb{C}^G),$$

we have

$$\widehat{A}_{f,m} = T_{f,m} \oplus 0.$$

Since $T_{f,m}$ and $B_{f,m}$ are unitarily equivalent,

$$(50) \quad \mu(t, \widehat{A}_{f,m}) = \mu(t, A_{f,m}), \quad t > 0.$$

By (48) and the Weyl law [?, Theorem 6.1],

$$\lim_{t \rightarrow \infty} t^{1/N} \mu(t, \widehat{A}_{f,m}) = \left(\frac{1}{N(2\pi)^N} \int_{S^* \mathbb{T}_m^N} \text{tr}_{\mathbb{C}^G} |\sigma_{-1}(\widehat{A}_{f,m})(z, \theta)|^N dz d\theta \right)^{1/N}.$$

Since the metric on $\mathbb{T}_m^N$ is flat, combining (49) with (50), we obtain

$$(51) \quad \lim_{t \rightarrow \infty} t^{1/N} \mu(t, A_{f,m}) = \left(\frac{1}{N(2\pi)^N} \int_{\Gamma} \int_{\mathbb{S}^{N-1}} \chi_m(x)^{2N} \text{tr}_{\mathbb{C}^G} |\mathbf{c}_{f,j}(x, \theta)|^N d\theta dx \right)^{1/N},$$

where χ_m is identified with its zero extension to Γ .

By Lemma 3.5,

$$(x, \theta) \mapsto \text{tr}_{\mathbb{C}^G} |\mathbf{c}_{f,j}(x, \theta)|^N$$

belongs to $L^1(\Gamma \times \mathbb{S}^{N-1})$. Since $0 \leq \chi_m \leq 1$, $\chi_m \rightarrow 1$ on Γ_+ , and $\Gamma \setminus \Gamma_+$ is contained in finitely many hyperplanes, the dominated convergence theorem implies

$$(52) \quad \begin{aligned} & \lim_{m \rightarrow \infty} \left(\frac{1}{N(2\pi)^N} \int_{\Gamma} \int_{\mathbb{S}^{N-1}} \chi_m(x)^{2N} \text{tr}_{\mathbb{C}^G} |\mathbf{c}_{f,j}(x, \theta)|^N d\theta dx \right)^{1/N} \\ &= \left(\frac{1}{N(2\pi)^N} \int_{\Gamma} \int_{\mathbb{S}^{N-1}} \text{tr}_{\mathbb{C}^G} |\mathbf{c}_{f,j}(x, \theta)|^N d\theta dx \right)^{1/N}. \end{aligned}$$

By (46), $A_{f,m} \rightarrow A_f$ in $S^{N,\infty}$. Combining (51), (52), and Lemma 3.8, we obtain

$$\lim_{t \rightarrow \infty} t^{1/N} \mu(t, A_f) = \left(\frac{1}{N(2\pi)^N} \int_{\Gamma} \int_{\mathbb{S}^{N-1}} \text{tr}_{\mathbb{C}^G} |\mathbf{c}_{f,j}(x, \theta)|^N d\theta dx \right)^{1/N}.$$

Since U and V are unitary,

$$\mu(t, A_f) = \mu(t, [R_{\text{Dunkl},j}, M_f]), \quad t > 0.$$

This completes the proof of Lemma 3.7. $\square$

Lemma 3.8. *Let $0 < p < \infty$, let $\mathcal{H}$ be a Hilbert space, and let $A, A_m \in S^{p,\infty}(\mathcal{H})$, $m \geq 1$, satisfy*

$$\|A_m - A\|_{S^{p,\infty}} \longrightarrow 0.$$

Suppose that, for every $m \geq 1$, the limit

$$c_m := \lim_{t \rightarrow \infty} t^{1/p} \mu(t, A_m)$$

exists. Then both limits

$$\lim_{m \rightarrow \infty} c_m \quad \text{and} \quad \lim_{t \rightarrow \infty} t^{1/p} \mu(t, A)$$

exist and

$$\lim_{t \rightarrow \infty} t^{1/p} \mu(t, A) = \lim_{m \rightarrow \infty} c_m.$$

Proof. The assertion is the Birman–Solomyak approximation lemma; see [?, Section 11.6]. $\square$

Lemma 3.9. *Assume that $N > 1$, $j \in \{1, \dots, N\}$, and $f \in \dot{W}^{1,N}(\mathbb{R}^N)$. Then there exists $f_m \in C_c^\infty(\mathbb{R}^N)$ such that*

$$\begin{aligned} \|\nabla(f_m - f)\|_{L^N(\mathbb{R}^N, dx)} &\longrightarrow 0, \\ \|[R_{\text{Dunkl},j}, M_{f_m}] - [R_{\text{Dunkl},j}, M_f]\|_{S^{N,\infty}} &\longrightarrow 0, \\ \|\mathbf{c}_{f_m,j} - \mathbf{c}_{f,j}\|_{L^N(\Gamma \times \mathbb{S}^{N-1}; S^N(\mathbb{C}^G))} &\longrightarrow 0. \end{aligned}$$

Proof. By the density of $C_c^\infty(\mathbb{R}^N)$ in $\dot{W}^{1,N}(\mathbb{R}^N)$, choose $f_m \in C_c^\infty(\mathbb{R}^N)$ such that

$$\|\nabla(f_m - f)\|_{L^N(\mathbb{R}^N, dx)} \longrightarrow 0.$$

By Theorem ?? and Lemma ??,

$$\begin{aligned} \|[R_{\text{Dunkl},j}, M_{f_m}] - [R_{\text{Dunkl},j}, M_f]\|_{S^{N,\infty}} &= \|[R_{\text{Dunkl},j}, M_{f_m-f}]\|_{S^{N,\infty}} \\ &\lesssim \|f_m - f\|_{\text{OSC}^{N,\infty}(\mathbb{R}^N)} \lesssim \|\nabla(f_m - f)\|_{L^N(\mathbb{R}^N, dx)} \longrightarrow 0. \end{aligned}$$

By the linearity of $f \mapsto \mathbf{c}_{f,j}$ and Lemma 3.5,

$$\begin{aligned} \|\mathbf{c}_{f_m,j} - \mathbf{c}_{f,j}\|_{L^N(\Gamma \times \mathbb{S}^{N-1}; S^N(\mathbb{C}^G))} &= \|\mathbf{c}_{f_m-f,j}\|_{L^N(\Gamma \times \mathbb{S}^{N-1}; S^N(\mathbb{C}^G))} \\ &\lesssim_{N,R,k} \|\nabla(f_m - f)\|_{L^N(\mathbb{R}^N, dx)} \longrightarrow 0. \end{aligned}$$

This completes the proof of Lemma 3.9. $\square$

Theorem 3.10. *Assume that $N > 1$ and $j \in \{1, \dots, N\}$. Then, for every $f \in \dot{W}^{1,N}(\mathbb{R}^N)$,*

(53)

$$\lim_{t \rightarrow \infty} t^{1/N} \mu(t, [R_{\text{Dunkl},j}, M_f]) = \left(\frac{1}{N(2\pi)^N} \int_{\Gamma} \int_{\mathbb{S}^{N-1}} \text{tr}_{\mathbb{C}^G} |\mathbf{c}_{f,j}(x, \theta)|^N d\theta dx \right)^{1/N}.$$

Moreover,

$$(54) \quad \left(\int_{\Gamma} \int_{\mathbb{S}^{N-1}} \text{tr}_{\mathbb{C}^G} |\mathbf{c}_{f,j}(x, \theta)|^N d\theta dx \right)^{1/N} \simeq_{N,R,k} \|\nabla f\|_{L^N(\mathbb{R}^N, dx)}.$$

Proof. By Lemma 3.9, choose $f_m \in C_c^\infty(\mathbb{R}^N)$ such that

$$(55) \quad \|[R_{\text{Dunkl},j}, M_{f_m}] - [R_{\text{Dunkl},j}, M_f]\|_{S^{N,\infty}} \longrightarrow 0,$$

$$(56) \quad \|\mathfrak{c}_{f_m,j} - \mathfrak{c}_{f,j}\|_{L^N(\Gamma \times \mathbb{S}^{N-1}; S^N(\mathbb{C}^G))} \longrightarrow 0.$$

By Lemma 3.7, for every m ,

$$\lim_{t \rightarrow \infty} t^{1/N} \mu(t, [R_{\text{Dunkl},j}, M_{f_m}]) = \frac{1}{(N(2\pi)^N)^{1/N}} \|\mathfrak{c}_{f_m,j}\|_{L^N(\Gamma \times \mathbb{S}^{N-1}; S^N(\mathbb{C}^G))}.$$

By (56),

$$\|\mathfrak{c}_{f_m,j}\|_{L^N(\Gamma \times \mathbb{S}^{N-1}; S^N(\mathbb{C}^G))} \longrightarrow \|\mathfrak{c}_{f,j}\|_{L^N(\Gamma \times \mathbb{S}^{N-1}; S^N(\mathbb{C}^G))}.$$

Combining this convergence with (55), we apply Lemma 3.8 with $p = N$ and obtain (53). Finally, (54) follows from Lemma 3.5. This completes the proof of Theorem 3.10. $\square$

4. FIRST ROCHBERG–SEMMES BASED ESTIMATE

it seems that we need $N > 2$, not $N > 1$. indeed, in Lemma 4.9 we use Lemma 6.1 — and this requires $N > r$. I mentioned this while in Guangzhou, but at the time we just ignored this issue. this is THE ONLY place where $N > 2$ is used.

Theorem 4.1. *Let $N > 1$. Fix a closed Weyl chamber Γ and $1_G \neq g \in G$. Suppose that $K : \Gamma \times \Gamma \rightarrow \mathbb{C}$ satisfies*

$$|\partial_x^\alpha \partial_y^\beta K|(x, y) \leq c_{K,\alpha,\beta} \frac{|x - y|^{1-|\alpha|-|\beta|_1}}{|x - gy| \cdot V(x, |x - y|)}, \quad x, y \in \Gamma, \quad \alpha, \beta \in \mathbb{Z}_+^N.$$

Let V be an integral operator with integral kernel K . For every $f \in \dot{W}^{1,N}(\Gamma)$, we have

$$\|M_{f-f \circ g} V\|_{\mathcal{L}_{N,\infty}(L_2(\Gamma, w(x)dx))} \leq c_{K,\kappa} (\|f\|_{\dot{W}^{1,N}(\Gamma)} + \|f\|_{\dot{W}^{1,N}(g\Gamma)}).$$

Theorem 4.2. *In the setting of Theorem 4.1, let $f \in C_c^\infty$. If $0 \leq \phi_m \leq 1$ for $m \geq 1$ are in $C_c^\infty(\mathbb{R}^N)$ and $\text{supp}(1 - \phi_m)$ increases to $\mathbb{R}^N$, then*

$$\text{dist}_{\mathcal{L}_{N,\infty}(L_2(\Gamma, w(x)dx))} \left(M_{\phi_m} \cdot M_f V \cdot M_{\phi_m} - M_f V, (\mathcal{L}_{N,\infty})_0(L_2(\Gamma, w(x)dx)) \right) \rightarrow 0$$

as $m \rightarrow \infty$.

Notation 4.3. *Fix a closed Weyl chamber Γ and let $\Phi : \mathbb{R}^N \rightarrow \mathbb{R}^N$ be the linear isomorphism appearing in Corollary 2.6.*

Let $\mathbb{D}_\Gamma$ be the dyadic system over Γ constructed in Corollary 2.7. If $I \in \mathbb{D}_\Gamma$ and if $Q = \Phi(I)$, then we denote $d_I = \ell(Q)$.

4.1. Proof of Theorem 4.1.

Lemma 4.4. *There exists a smooth function ψ supported on $[-1, 1]^N \setminus (-\frac{1}{4}, \frac{1}{4})^N$ such that*

$$1 = \sum_{I \in \mathbb{D}_\Gamma} \chi_I(x) \chi_{3I}(y) \psi\left(\frac{\Phi(x) - \Phi(y)}{d_I}\right).$$

Proof. Choose $\rho \in C_c^\infty(\mathbb{R}^N)$ such that $\rho = 1$ on $(-\frac{1}{2}, \frac{1}{2})^N$ and $\rho = 0$ outside $(-1, 1)^N$. Set $\psi(z) = \rho(z) - \rho(2z)$. Clearly,

$$(57) \quad \text{supp}(\psi) \subset \left\{ z \in \mathbb{R}^N : \frac{1}{4} \leq |z|_\infty \leq 1 \right\}.$$

For every $z \neq 0$, we have

$$(58) \quad \sum_{j \in \mathbb{Z}} \psi(2^j z) = 1.$$

For every $x \in \Gamma$, there exists a unique $I \in \mathbb{D}_{\Gamma,j}$ containing x . For this I , we have $d_I = 2^{-j}$. We have

$$\begin{aligned} 1 &= \sum_{j \in \mathbb{Z}} \psi(2^j(\Phi(x) - \Phi(y))) = \sum_{j \in \mathbb{Z}} \left(\sum_{I \in \mathbb{D}_{\Gamma,j}} \chi_I(x) \right) \psi(2^j(\Phi(x) - \Phi(y))) = \\ &= \sum_{j \in \mathbb{Z}} \sum_{I \in \mathbb{D}_{\Gamma,j}} \chi_I(x) \psi\left(\frac{\Phi(x) - \Phi(y)}{d_I}\right) = \sum_{I \in \mathbb{D}_{\Gamma}} \chi_I(x) \psi\left(\frac{\Phi(x) - \Phi(y)}{d_I}\right). \end{aligned}$$

If $x \in I$ and $\psi\left(\frac{\Phi(x) - \Phi(y)}{d_I}\right) \neq 0$, then

$$\Phi(x) \in \Phi(I) = Q, \quad |\Phi(x) - \Phi(y)| \leq d_I = \ell(Q).$$

We obviously have $|\Phi(x) - c_Q|_{\infty} \leq \frac{1}{2}\ell(Q)$ and $|\Phi(x) - \Phi(y)|_{\infty} \leq \ell(Q)$. Thus, $|\Phi(y) - c_Q| \leq \frac{3}{2}\ell(Q)$. In other words, $\Phi(y) \in 3Q$ or, equivalently, $y \in 3I$. Thus,

$$\chi_I(x) \psi\left(\frac{\Phi(x) - \Phi(y)}{d_I}\right) = \chi_I(x) \chi_{3I}(y) \psi\left(\frac{\Phi(x) - \Phi(y)}{d_I}\right).$$

This completes the proof. $\square$

Lemma 4.5. Assume that $N > 1$ and $2 < r < \infty$. Let $f \in L_r^{\text{loc}}(\mathbb{R}^N)$. Fix a closed Weyl chamber Γ and $1_G \neq g \in G$. Let K be as in Theorem 4.1. We have

$$f(x)K(x, y) = \sum_{k, l \in \mathbb{Z}^N} (1 + |k|^2 + |l|^2)^{-N-1} \sum_{I \in \mathbb{D}_{\Gamma}} e_{k, l, I}(x) f_{k, l, I}(y),$$

where

- (1) $e_{k, l, I}$ is supported in I and $f_{k, l, I}$ is supported in $3I$;
- (2) for every $k, l \in \mathbb{Z}^N$ and for every $I \in \mathbb{D}_{\Gamma}$, we have

$$\|e_{k, l, I}\|_{L_r(\mathbb{R}^N, w(x)dx)} = \|f\|_{L_r(I, w(x)dx)};$$

- (3) for every $k, l \in \mathbb{Z}^N$ and for every $I \in \mathbb{D}_{\Gamma}$, we have

$$\|f_{k, l, I}\|_{L_{\infty}(\mathbb{R}^N)} \leq c_{K, \kappa, \Phi} \frac{d_I}{d_I + |c_I - gc_I|};$$

Proof. We have

$$\begin{aligned} K(x, y) &= \sum_{I \in \mathbb{D}_{\Gamma}} \chi_I(x) \chi_{3I}(y) K(x, y) \psi\left(\frac{\Phi(x) - \Phi(y)}{d_I}\right) = \\ &= \sum_{\substack{I \in \mathbb{D}_{\Gamma} \\ 3I \subset \Gamma}} \chi_I(x) \chi_{3I}(y) K(x, y) \psi\left(\frac{\Phi(x) - \Phi(y)}{d_I}\right) + \sum_{\substack{I \in \mathbb{D}_{\Gamma} \\ 3I \not\subset \Gamma}} \chi_I(x) \chi_{3I}(y) K(x, y) \psi\left(\frac{\Phi(x) - \Phi(y)}{d_I}\right). \end{aligned}$$

We will only consider the first sum. The second some can be treated similarly, but the notations become heavier. **feel free to write the expression for the second sum if you think it is needed. the only difference is that the domain of the map K_I becomes more complicated.**

For every $I \in \mathbb{D}_{\Gamma}$ with $3I \subset \Gamma$, define the map

$$K_I : \left[-\frac{1}{2}, \frac{1}{2}\right]^N \times \left[-\frac{3}{2}, \frac{3}{2}\right]^N \rightarrow \mathbb{C}$$

by setting

$$K_I(a, b) = K(c_I + d_I \Phi^{-1}(a), c_I + d_I \Phi^{-1}(b)) \psi(a - b), \quad a \in [-\frac{1}{2}, \frac{1}{2}]^N, \quad b \in [-\frac{3}{2}, \frac{3}{2}]^N.$$

It follows from the assumption on K that **feel free to prove this if you think this requires a proof. however, do not prove it here — make a separate lemma instead.**

$$\|K_I\|_{C^{2N+2}([-\frac{1}{2}, \frac{1}{2}]^N \times [-\frac{3}{2}, \frac{3}{2}]^N)} \leq \frac{c'_{K, \kappa, \Phi}}{w(I)} \cdot \frac{d_I}{d_I + |c_I - gc_I|}.$$

Hence, there exists an extension $K_I : \mathbb{T}^N \times \mathbb{T}^N \rightarrow \mathbb{C}$ such that

$$\|K_I\|_{C^{2N+2}(\mathbb{T}^N \times \mathbb{T}^N)} \leq \frac{c''_{K, \kappa, \Phi}}{w(I)} \cdot \frac{d_I}{d_I + |c_I - gc_I|}.$$

In particular,

$$\|K_I\|_{W^{2N+2, 2}(\mathbb{T}^N \times \mathbb{T}^N)} \leq \frac{c_{K, \kappa, \Phi}}{w(I)} \cdot \frac{d_I}{d_I + |c_I - gc_I|}.$$

We may, therefore, write

$$K_I(a, b) = \sum_{k, l \in \mathbb{Z}^N} \text{coef}_{k, l, I} e^{i\langle k, a \rangle} e^{i\langle l, b \rangle}, \quad a, b \in \mathbb{T}^N,$$

where

$$\sum_{k, l \in \mathbb{Z}^N} (1 + |k|^2 + |l|^2)^{2N+2} |\text{coef}_{k, l, I}|^2 \leq \left(\frac{c_{K, \kappa, \Phi}}{w(I)} \cdot \frac{d_I}{d_I + |c_I - gc_I|} \right)^2.$$

In particular,

$$(1 + |k|^2 + |l|^2)^{-N-1} \cdot |\text{coef}_{k, l, I}| \leq \frac{c_{K, \kappa, \Phi}}{w(I)} \cdot \frac{d_I}{d_I + |c_I - gc_I|}, \quad k, l \in \mathbb{Z}^N, \quad I \in \mathbb{D}_\Gamma.$$

We now set

$$e_{k, l, I}(x) = f(x) \chi_I(x) e^{i\langle k, \frac{\Phi^{-1}(x - c_I)}{d_I} \rangle},$$

$$f_{k, l, I}(y) = (1 + |k|^2 + |l|^2)^{N+1} \text{coef}_{k, l, I} \cdot \chi_{3I}(y) e^{i\langle l, \frac{\Phi^{-1}(y - c_I)}{d_I} \rangle}.$$

This allows us to write

$$f(x) K(x, y) = \sum_{k, l \in \mathbb{Z}^N} (1 + |k|^2 + |l|^2)^{-N-1} \sum_{I \in \mathbb{D}_\Gamma} e_{k, l, I}(x) f_{k, l, I}(y).$$

Obviously,

$$\text{supp}(e_{k, l, I}) \subset I, \quad \text{supp}(f_{k, l, I}) \subset 3I,$$

$$\|e_{k, l, I}\|_{L_r(\mathbb{R}^N, w(x) dx)} = \|f\|_{L_r(I, w(x) dx)},$$

$$\|f_{k, l, I}\|_{L_\infty(\mathbb{R}^N, w(x) dx)} = (1 + |k|^2 + |l|^2)^{N+1} |\text{coef}_{I, k, l}| \leq \frac{c_{K, \kappa, \Phi}}{w(I)} \cdot \frac{d_I}{d_I + |c_I - gc_I|}.$$

□

Lemma 4.6. *For every $I \in \mathbb{D}_\Gamma$, we have*

$$\sup_{x \in I} w(x) \leq c_{\kappa, \Phi} \frac{w(I)}{m(I)}.$$

Proof. We have

$$w(I) \approx_{\Phi, \kappa} w(B(c_I, d_I)) \approx_{\kappa} d_I^N \prod_{v \in R} (|\langle c_I, v \rangle| + d_I)^{k(v)}.$$

On the other hand, if $x \in I$, then $|x - c_I| \leq c_{\Phi} d_I$ and, therefore,

$$|\langle x, v \rangle| \leq |\langle x - c_I, v \rangle| + |\langle c_I, v \rangle| \leq |x - c_I| + |\langle c_I, v \rangle| \leq c_{\Phi} d_I + |\langle c_I, v \rangle|.$$

Thus,

$$w(x) \leq \prod_{v \in R} (c_{\Phi} d_I + |\langle c_I, v \rangle|)^{k(v)} \leq c_{\Phi, \kappa} \prod_{v \in R} (|\langle c_I, v \rangle| + d_I)^{k(v)}.$$

This completes the proof. $\square$

Lemma 4.7. Assume that $N > 1$ and $2 < r < \infty$. Let $f \in L_r^{\text{loc}}(\mathbb{R}^N)$. We have

$$\|M_f V\|_{\mathcal{L}_{N, \infty}(L_2(\Gamma, w(x)dx))} \leq c_{K, \kappa, \Phi, r} \left\| \left\{ m(I)^{-\frac{1}{r}} \|f\|_{L_r(I)} \cdot \frac{d_I}{d_I + |c_I - gc_I|} \right\}_{I \in \mathbb{D}_{\Gamma}} \right\|_{N, \infty}.$$

Proof. Let $A_{k, l}$ be the integral operator with integral kernel given by the formula

$$(x, y) \rightarrow \sum_{I \in \mathbb{D}_{\Gamma}} e_{k, l, I}(x) f_{k, l, I}(y), \quad x, y \in \Gamma, \quad k, l \in \mathbb{Z}^N.$$

By Lemma 4.5, we have

$$M_f V = \sum_{k, l \in \mathbb{Z}^N} (1 + |k|^2 + |l|^2)^{-N-1} A_{k, l}.$$

By the triangle inequality,

$$\begin{aligned} & \|M_f V\|_{\mathcal{L}_{N, \infty}(L_2(\Gamma, w(x)dx))} \leq \\ & \leq \frac{N}{N-1} \sum_{k, l \in \mathbb{Z}^N} (1 + |k|^2 + |l|^2)^{-N-1} \|A_{k, l}\|_{\mathcal{L}_{N, \infty}(L_2(\Gamma, w(x)dx))}. \end{aligned}$$

By Lemma 4.5, the sequences

$$\left\{ w(I)^{\frac{1}{r} - \frac{1}{2}} \frac{e_{k, l, I}}{\|f\|_{L_r(I, w(x)dx)}} \right\}_{I \in \mathbb{D}_{\Gamma}}, \quad \left\{ c_{K, \kappa, \Phi}^{-1} w(I)^{\frac{1}{2}} \cdot \frac{d_I + |c_I - gc_I|}{d_I} f_{k, l, I} \right\}_{I \in \mathbb{D}_{\Gamma}}$$

are $(r, 1)$ -NWO and, respectively, $(\infty, 3)$ -NWO in the sense of Definition 1.6. Using Lemma 1.7, we conclude that

$$\begin{aligned} & \|A_{k, l}\|_{\mathcal{L}_{N, \infty}(L_2(\Gamma, w(x)dx))} \leq \\ & \leq c_{\Phi, \kappa, r, 1, \infty, 3, N} c_{K, \kappa} \left\| \left\{ w(I)^{-\frac{1}{r}} \|f\|_{L_r(I, w(x)dx)} \cdot \frac{d_I}{d_I + |c_I - gc_I|} \right\}_{I \in \mathbb{D}_{\Gamma}} \right\|_{N, \infty}. \end{aligned}$$

Therefore,

$$\begin{aligned} & \|M_f V\|_{\mathcal{L}_{N, \infty}(L_2(\Gamma, w(x)dx))} \leq \\ & \leq \frac{N}{N-1} c_{\Phi, \kappa, r, 1, \infty, 3, N} c_{K, \kappa, \Phi} \sum_{k, l \in \mathbb{Z}^N} (1 + |k|^2 + |l|^2)^{-N-1} \times \\ & \times \left\| \left\{ w(I)^{-\frac{1}{r}} \|f\|_{L_r(I, w(x)dx)} \cdot \frac{d_I}{d_I + |c_I - gc_I|} \right\}_{I \in \mathbb{D}_{\Gamma}} \right\|_{N, \infty}. \end{aligned}$$

By Lemma 4.6,

$$w(I)^{-\frac{1}{r}} \|f\|_{L_r(I, w(x)dx)} \leq c_{\kappa, \Phi}^{\frac{1}{r}} m(I)^{-\frac{1}{r}} \|f\|_{L_r(I)}, \quad I \in \mathbb{D}_{\Gamma}.$$

Combining the last two estimates, we complete the proof. $\square$

Lemma 4.8.

$$\left\| \left\{ m(I)^{-1} \|f - f \circ g\|_{L_1(I)} \cdot \frac{d_I}{d_I + |c_I - gc_I|} \right\}_{I \in \mathbb{D}_\Gamma} \right\|_{N, \infty} \leq c_\Phi (\|f\|_{\dot{W}^{1,N}(\Gamma)} + \|f\|_{\dot{W}^{1,N}(g\Gamma)}).$$

Proof. Let $\mathcal{M}$ be the Hardy–Littlewood maximal operator on $\mathbb{R}^N$. By [?, Theorem 3], for almost every $x \in \mathbb{R}^N$,

$$|f(x) - f(gx)| \leq c_N |x - gx| \cdot ((\mathcal{M}(|\nabla f|))(x) + (\mathcal{M}(|\nabla f|))(gx)).$$

We have

$$|x - gx| \leq 2|x - c_I| + |c_I - gc_I|, \quad x \in I, \quad y \in gI.$$

Obviously,

$$|x - c_I| \leq \frac{d_I^{\frac{1}{2}}}{2} \|\Phi\|_\infty d_I, \quad x \in I.$$

Hence,

$$|x - gx| \leq \max\{1, d_I^{\frac{1}{2}} \|\Phi\|_\infty\} \cdot (d_I + |c_I - gc_I|), \quad x \in I, \quad y \in gI.$$

Thus,

$$|f(x) - f(gx)| \leq c'_\Phi \cdot (d_I + |c_I - gc_I|) \cdot ((\mathcal{M}(|\nabla f|))(x) + (\mathcal{M}(|\nabla f|))(gx)).$$

This immediately implies

$$\frac{1}{d_I + |c_I - gc_I|} \|f - f \circ g\|_{L_1(I)} \leq c''_\Phi \cdot (\|\mathcal{M}(|\nabla f|)\|_{L_1(I)} + \|\mathcal{M}(|\nabla f|)\|_{L_1(gI)})$$

for every $I \in \mathbb{D}_\Gamma$. Hence,

$$\begin{aligned} m(I)^{-1} \|f - f \circ g\|_{L_1(I)} \cdot \frac{d_I}{d_I + |c_I - gc_I|} &\leq \\ &\leq c'''_\Phi \cdot m(I)^{\frac{1}{N}-1} (\|\mathcal{M}(|\nabla f|)\|_{L_1(I)} + \|\mathcal{M}(|\nabla f|)\|_{L_1(gI)}) \end{aligned}$$

for every $I \in \mathbb{D}_\Gamma$. Thus,

$$\begin{aligned} &\left\| \left\{ m(I)^{-1} \|f - f \circ g\|_{L_1(I)} \cdot \frac{d_I}{d_I + |c_I - gc_I|} \right\}_{I \in \mathbb{D}_\Gamma} \right\|_{N, \infty} \leq \\ &\leq c'''_\Phi \frac{N}{N-1} \left(\left\| \left\{ m(I)^{\frac{1}{N}-1} \|\mathcal{M}(|\nabla f|)\|_{L_1(I)} \right\}_{I \in \mathbb{D}_\Gamma} \right\|_{N, \infty} + \right. \\ &\quad \left. + \left\| \left\{ m(I)^{\frac{1}{N}-1} \|\mathcal{M}(|\nabla f|)\|_{L_1(gI)} \right\}_{I \in \mathbb{D}_\Gamma} \right\|_{N, \infty} \right) \leq \\ &\stackrel{L.6.1}{\leq} c'''_\Phi \frac{N}{N-1} c_N \left(\|\mathcal{M}(|\nabla f|)\|_{L_N(\Gamma)} + \|\mathcal{M}(|\nabla f|)\|_{L_N(g\Gamma)} \right) \leq \\ &\leq c'''_\Phi \frac{N}{N-1} c_N \|\mathcal{M}\|_{L_N(\mathbb{R}^N) \odot} \left(\|\nabla f\|_{L_N(\Gamma)} + \|\nabla f\|_{L_N(g\Gamma)} \right). \end{aligned}$$

□

Lemma 4.9. Let $2 < r < N$ and let $f \in \dot{W}^{1,N}(\Gamma) \cap \dot{W}^{1,N}(g\Gamma)$. We have

$$\left\| \left\{ m(I)^{-\frac{1}{r}} \|f - f \circ g\|_{L_r(I)} \cdot \frac{d_I}{d_I + |c_I - gc_I|} \right\}_{I \in \mathbb{D}_\Gamma} \right\|_{N, \infty} \leq c_\Phi (\|f\|_{\dot{W}^{1,N}(\Gamma)} + \|f\|_{\dot{W}^{1,N}(g\Gamma)}).$$

Proof. We obviously have

$$\begin{aligned} \|f - f \circ g\|_{L_r(I)} &\leq \left\| f - \frac{1}{m(I)} \int_I f \right\|_{L_r(I)} + \\ &+ \left\| f \circ g - \frac{1}{m(I)} \int_I (f \circ g) \right\|_{L_r(I)} + m(I)^{\frac{1}{r}-1} \|f - f \circ g\|_{L_1(I)}. \end{aligned}$$

By Poincare inequality,

$$\begin{aligned} &\|f - f \circ g\|_{L_r(I)} \leq \\ &\leq c_{\Phi,r} \left(m(I)^{\frac{1}{N}} \|\nabla f\|_{L_r(I)} + m(I)^{\frac{1}{N}} \|\nabla(f \circ g)\|_{L_r(I)} + m(I)^{\frac{1}{r}-1} \|f - f \circ g\|_{L_1(I)} \right). \end{aligned}$$

By the triangle inequality,

$$\begin{aligned} &\left\| \left\{ m(I)^{-\frac{1}{r}} \|f - f \circ g\|_{L_r(I)} \cdot \frac{d_I}{d_I + |c_I - gc_I|} \right\}_{I \in \mathbb{D}_\Gamma} \right\|_{N,\infty} \leq \\ &\leq \frac{N}{N-1} c_{\Phi,r} \left\| \left\{ m(I)^{\frac{1}{N}-\frac{1}{r}} \|\nabla f\|_{L_r(I)} \right\}_{I \in \mathbb{D}_\Gamma} \right\|_{N,\infty} + \\ &+ \frac{N}{N-1} c_{\Phi,r} \left\| \left\{ m(I)^{\frac{1}{N}-\frac{1}{r}} \|\nabla f\|_{L_r(gI)} \right\}_{I \in \mathbb{D}_\Gamma} \right\|_{N,\infty} + \\ &+ \frac{N}{N-1} c_{\Phi,r} \left\| \left\{ m(I)^{-1} \|f - f \circ g\|_{L_1(I)} \cdot \frac{d_I}{d_I + |c_I - gc_I|} \right\}_{I \in \mathbb{D}_\Gamma} \right\|_{N,\infty}. \end{aligned}$$

The assertion follows now from Lemma 6.1 (for the first two summands) and Lemma 4.8 (for the last summand). $\square$

Proof of Theorem 4.1. The assertion follows by combining Lemmas 4.7 (taken with $f - f \circ g$ instead of f) and 4.9. $\square$

4.2. Proof of Theorem 4.2.

Lemma 4.10. *Let $f, \phi \in C_c^\infty(\mathbb{R}^N)$. If $0 \leq \phi \leq 1$, then*

$$\begin{aligned} &\text{dist}_{\mathcal{L}_{N,\infty}(L_2(\Gamma, w(x)dx))} \left(M_f V - M_{f\phi} V, (\mathcal{L}_{N,\infty})_0(L_2(\Gamma, w(x)dx)) \right) \leq c_{K,\kappa,\phi,r} \times \\ &\times \text{dist}_{l_{N,\infty}(\mathbb{D}_\Gamma)} \left(\left\{ \frac{d_I m(I)^{-\frac{1}{r}}}{d_I + |c_I - gc_I|} \|f\|_{L_r(I)} \right\}_{\substack{I \in \mathbb{D}_\Gamma \\ I \cap \text{supp}(f) \neq \emptyset \\ I \cap \text{supp}(1-\phi) \neq \emptyset}}, (l_{N,\infty})_0(\mathbb{D}_\Gamma) \right), \\ &\text{dist}_{\mathcal{L}_{N,\infty}(L_2(\Gamma, w(x)dx))} \left(M_f V - M_f V M_\phi, (\mathcal{L}_{N,\infty})_0(L_2(\Gamma, w(x)dx)) \right) \leq c_{K,\kappa,\phi,r} \times \\ &\times \text{dist}_{l_{N,\infty}(\mathbb{D}_\Gamma)} \left(\left\{ \frac{d_I m(I)^{-\frac{1}{r}}}{d_I + |c_I - gc_I|} \|f\|_{L_r(I)} \right\}_{\substack{I \in \mathbb{D}_\Gamma \\ I \cap \text{supp}(f) \neq \emptyset \\ 3I \cap \text{supp}(1-\phi) \neq \emptyset}}, (l_{N,\infty})_0(\mathbb{D}_\Gamma) \right). \end{aligned}$$

Proof. We only prove the second inequality. The proof of the first one is similar, but simpler.

Let $A_{k,l}$ be as in the proof of Lemma 4.7. We have

$$M_f V - M_f V M_\phi = \sum_{k,l \in \mathbb{Z}^N} (1 + |k|^2 + |l|^2)^{-N-1} A_{k,l} M_{1-\phi}.$$

By the triangle inequality,

$$\text{dist}_{\mathcal{L}_{N,\infty}(L_2(\Gamma, w(x)dx))} \left(M_f V - M_f V M_\phi, (\mathcal{L}_{N,\infty})_0(L_2(\Gamma, w(x)dx)) \right) \leq$$

$$\leq \frac{N}{N-1} \sum_{k,l \in \mathbb{Z}^N} \left((1 + |k|^2 + |l|^2)^{-N-1} \times \right. \\ \left. \times \text{dist}_{\mathcal{L}_{N,\infty}(L_2(\Gamma, w(x)dx))} \left(A_{k,l} M_{1-\phi}, (\mathcal{L}_{N,\infty})_0(L_2(\Gamma, w(x)dx)) \right) \right).$$

By Lemma 4.5, the sequences

$$\left\{ w(I)^{\frac{1}{r}-\frac{1}{2}} \frac{e_{k,l,I}}{\|f\|_{L_r(I, w(x)dx)}} \right\}_{I \in \mathbb{D}_\Gamma}, \quad \left\{ c_{K,\kappa,\Phi}^{-1} w(I)^{\frac{1}{2}} \cdot \frac{d_I + |c_I - gc_I|}{d_I} f_{k,l,I} \cdot (1-\phi) \right\}_{I \in \mathbb{D}_\Gamma}$$

are $(r, 1)$ -NWO and, respectively, $(\infty, 3)$ -NWO in the sense of Definition 1.6. Obviously, $e_{k,l,I} \neq 0$ only if $I \cap \text{supp}(f) \neq \emptyset$ and $f_{k,l,I} \cdot (1-\phi) \neq 0$ only if $I \cap \text{supp}(f) \neq \emptyset$ and $3I \cap \text{supp}(1-\phi) \neq \emptyset$. Using Lemma 1.7, we conclude that

$$\text{dist}_{\mathcal{L}_{N,\infty}(L_2(\Gamma, w(x)dx))} \left(A_{k,l}(1-\phi), (\mathcal{L}_{N,\infty})_0(L_2(\Gamma, w(x)dx)) \right) \leq \\ \leq c_{\Phi,\kappa,r,1,\infty,3,N} c_{K,\kappa,\Phi} \times \\ \times \text{dist}_{l_{N,\infty}(\mathbb{D}_\Gamma)} \left(\left\{ \frac{d_I w(I)^{-\frac{1}{r}}}{d_I + |c_I - gc_I|} \|f\|_{L_r(I, w(x)dx)} \right\}_{\substack{I \in \mathbb{D}_\Gamma \\ I \cap \text{supp}(f) \neq \emptyset \\ 3I \cap \text{supp}(1-\phi) \neq \emptyset}}, (l_{N,\infty})_0(\mathbb{D}_\Gamma) \right).$$

By Lemma 4.6,

$$w(I)^{-\frac{1}{r}} \|f\|_{L_r(I, w(x)dx)} \leq c_{\kappa,\Phi}^{\frac{1}{r}} m(I)^{-\frac{1}{r}} \|f\|_{L_r(I)}, \quad I \in \mathbb{D}_\Gamma.$$

Thus,

$$\text{dist}_{\mathcal{L}_{N,\infty}(L_2(\Gamma, w(x)dx))} \left(A_{k,l}(1-\phi), (\mathcal{L}_{N,\infty})_0(L_2(\Gamma, w(x)dx)) \right) \leq \\ \leq c_{\Phi,\kappa,r,1,\infty,3,N} c_{K,\kappa,\Phi} c_{\kappa,\Phi}^{\frac{1}{r}} \times \\ \times \text{dist}_{l_{N,\infty}(\mathbb{D}_\Gamma)} \left(\left\{ \frac{d_I m(I)^{-\frac{1}{r}}}{d_I + |c_I - gc_I|} \|f\|_{L_r(I)} \right\}_{\substack{I \in \mathbb{D}_\Gamma \\ I \cap \text{supp}(f) \neq \emptyset \\ 3I \cap \text{supp}(1-\phi) \neq \emptyset}}, (l_{N,\infty})_0(\mathbb{D}_\Gamma) \right).$$

Combining this estimate with the one in the first paragraph **assign label and refer to it** yields the second inequality. $\square$

Lemma 4.11. *For a bounded Borel $A \subset \mathbb{R}^N$, we have*

$$\lim_{n \rightarrow \infty} \left\| \left\{ m(Q)^{\frac{1}{N}} \right\}_{\substack{Q \in \mathbb{D} \\ Q \subset A \\ \ell(Q) \leq 2^{-n}}} \right\|_{l_{N,\infty}(\mathbb{D})} \leq c_N m(A)^{\frac{1}{N}}.$$

Proof. For a given $\epsilon > 0$, choose j (depending on ϵ) and finite set $B \subset \mathbb{D}_j$ such that $A \subset \cup_{Q' \in B} Q'$ and such that $m((\cup_{Q' \in B} Q') \setminus A) < \epsilon$. For $n \geq m$, we have (here, we use the notation $Q_0 = [0, 1)^N$)

$$\left\| \left\{ m(Q)^{\frac{1}{N}} \right\}_{\substack{Q \in \mathbb{D} \\ Q \subset A \\ \ell(Q) \leq 2^{-n}}} \right\|_{l_{N,\infty}(\mathbb{D})} \leq \left\| \left\{ m(Q)^{\frac{1}{N}} \right\}_{\substack{Q \in \mathbb{D} \\ Q \subset Q' \text{ for some } Q' \in B}} \right\|_{l_{N,\infty}(\mathbb{D})} = \\ = \left(\frac{|B|}{2^j} \right)^{\frac{1}{N}} \left\| \left\{ m(Q)^{\frac{1}{N}} \right\}_{\substack{Q \in \mathbb{D} \\ Q \subset Q_0}} \right\|_{l_{N,\infty}(\mathbb{D})} = c_N \cdot m^{\frac{1}{N}}(\cup_{Q' \in B} Q') \leq c_N \cdot (m(A) + \epsilon)^{\frac{1}{N}}.$$

Passing $n \rightarrow \infty$ and, after that, passing $\epsilon \rightarrow 0$, we complete the proof. $\square$

Lemma 4.12. *We have*

$$\lim_{n \rightarrow \infty} \left\| \left\{ m(I)^{\frac{1}{N}} \right\}_{\substack{I \in \mathbb{D}_\Gamma \\ I \cap \text{supp}(f) \neq \emptyset \\ 3I \cap \text{supp}(1-\phi) \neq \emptyset \\ d_I \leq 2^{-n}}} \right\|_{l_{N,\infty}(\mathbb{D}_\Gamma)} \leq c_N m^{\frac{1}{N}}(\text{supp}(f) \cap \text{supp}(1-\phi)).$$

Proof. Denote for brevity $A = \Phi^{-1}(\text{supp}(f))$ and $B = \Phi^{-1}(\text{supp}(1-\phi))$. For $\epsilon > 0$, denote by A_ϵ (respectively, B_ϵ) the ϵ -neighborhood of A (respectively, of B) with respect to the norm $|\cdot|_\infty$.

If $2^{-n} < \epsilon$, then the condition $I \cap \text{supp}(\phi) \neq \emptyset$ yields, for $Q = \Phi(I) \in \mathbb{D}$, the condition $Q \subset A_\epsilon$. If $2^{-n} < \frac{\epsilon}{3}$, then the condition $I \cap \text{supp}(1-\phi) \neq \emptyset$ yields, for $Q = \Phi(I) \in \mathbb{D}$, the condition $Q \subset B_\epsilon$. Hence, for $2^{-n} < \frac{\epsilon}{3}$, we have

$$\left\| \left\{ m(I)^{\frac{1}{N}} \right\}_{\substack{I \in \mathbb{D}_\Gamma \\ I \cap \text{supp}(f) \neq \emptyset \\ 3I \cap \text{supp}(1-\phi) \neq \emptyset \\ d_I \leq 2^{-n}}} \right\|_{l_{N,\infty}(\mathbb{D}_\Gamma)} \leq \det(\Phi)^{-\frac{1}{N}} \left\| \left\{ m(Q)^{\frac{1}{N}} \right\}_{\substack{Q \in \mathbb{D} \\ Q \subset A_\epsilon \cap B_\epsilon \\ \ell(Q) \leq 2^{-n}}} \right\|_{l_{N,\infty}(\mathbb{D})}.$$

Using Lemma 4.11 and passing $n \rightarrow \infty$, we obtain

$$\lim_{n \rightarrow \infty} \left\| \left\{ m(I)^{\frac{1}{N}} \right\}_{\substack{I \in \mathbb{D}_\Gamma \\ I \cap \text{supp}(f) \neq \emptyset \\ 3I \cap \text{supp}(1-\phi) \neq \emptyset \\ d_I \leq 2^{-n}}} \right\|_{l_{N,\infty}(\mathbb{D}_\Gamma)} \leq c_N \det(\Phi)^{-\frac{1}{N}} m^{\frac{1}{N}}(A_\epsilon \cap B_\epsilon).$$

Since A and B are both closed, it follows that

$$\lim_{\epsilon \rightarrow 0} m(A_\epsilon \cap B_\epsilon) = m(A \cap B).$$

Passing $\epsilon \rightarrow 0$ in the second last display, we complete the proof. $\square$

Lemma 4.13. *Let $r > 2$. For every $f \in C_c^\infty(\mathbb{R}^N)$ and for every $n \in \mathbb{N}$, we have*

$$\begin{aligned} \text{dist}_{l_{N,\infty}(\mathbb{D}_\Gamma)} \left(\left\{ m(I)^{\frac{1}{N} - \frac{1}{r}} \|\nabla f\|_{L_r(I)} \right\}_{\substack{I \in \mathbb{D}_\Gamma \\ I \cap \text{supp}(f) \neq \emptyset \\ 3I \cap \text{supp}(1-\phi) \neq \emptyset}}, (l_{N,\infty})_0(\mathbb{D}_\Gamma) \right) &\leq \\ &\leq c_N \|\nabla f\|_{L_\infty(\mathbb{R}^N)} \times m^{\frac{1}{N}}(\text{supp}(f) \cap \text{supp}(1-\phi)). \end{aligned}$$

Proof. For a given $n \in \mathbb{Z}$, we have

$$\left\{ m(I)^{\frac{1}{N} - \frac{1}{r}} \|\nabla f\|_{L_r(I)} \right\}_{\substack{I \in \mathbb{D}_\Gamma \\ I \cap \text{supp}(f) \neq \emptyset \\ 3I \cap \text{supp}(1-\phi) \neq \emptyset \\ d_I > 2^{-n}}} \leq \|\nabla f\|_{L_r(\mathbb{R}^N)} \left\{ m(I)^{\frac{1}{N} - \frac{1}{r}} \right\}_{\substack{I \in \mathbb{D}_\Gamma \\ I \cap \text{supp}(f) \neq \emptyset \\ 3I \cap \text{supp}(1-\phi) \neq \emptyset \\ d_I > 2^{-n}}}.$$

The right hand side obviously belongs to $(l_{N,\infty})_0(\mathbb{D}_\Gamma)$ and, hence, so does the left hand side. Thus,

$$\begin{aligned} \text{dist}_{l_{N,\infty}(\mathbb{D}_\Gamma)} \left(\left\{ m(I)^{\frac{1}{N} - \frac{1}{r}} \|\nabla f\|_{L_r(I)} \right\}_{\substack{I \in \mathbb{D}_\Gamma \\ I \cap \text{supp}(f) \neq \emptyset \\ 3I \cap \text{supp}(1-\phi) \neq \emptyset}}, (l_{N,\infty})_0(\mathbb{D}_\Gamma) \right) &\leq \\ &\leq \left\| \left\{ m(I)^{\frac{1}{N} - \frac{1}{r}} \|\nabla f\|_{L_r(I)} \right\}_{\substack{I \in \mathbb{D}_\Gamma \\ I \cap \text{supp}(f) \neq \emptyset \\ 3I \cap \text{supp}(1-\phi) \neq \emptyset \\ d_I \leq 2^{-n}}} \right\|_{l_{N,\infty}(\mathbb{D}_\Gamma)} \leq \\ &\leq \|\nabla f\|_{L_\infty(\mathbb{R}^N)} \times \left\| \left\{ m(I)^{\frac{1}{N}} \right\}_{\substack{I \in \mathbb{D}_\Gamma \\ I \cap \text{supp}(f) \neq \emptyset \\ 3I \cap \text{supp}(1-\phi) \neq \emptyset \\ d_I \leq 2^{-n}}} \right\|_{l_{N,\infty}(\mathbb{D}_\Gamma)}. \end{aligned}$$

Passing $n \rightarrow \infty$ and using Lemma 4.12, we complete the proof. $\square$

Lemma 4.14. *For every $f \in C_c^\infty(\mathbb{R}^N)$ and for every $n \in \mathbb{N}$, we have*

$$\begin{aligned} \text{dist}_{l_{N,\infty}(\mathbb{D}_\Gamma)} \left(\left\{ \frac{d_I m(I)^{-1}}{d_I + |c_I - gc_I|} \|f - f \circ g\|_{L_1(I)} \right\}_{\substack{I \in \mathbb{D}_\Gamma \\ I \cap \text{supp}(f) \neq \emptyset \\ 3I \cap \text{supp}(1-\phi) \neq \emptyset}}, (l_{N,\infty})_0(\mathbb{D}_\Gamma) \right) \leq \\ \leq c_\Phi \|\nabla f\|_{L_\infty(\mathbb{R}^N)} \times m^{\frac{1}{N}}(\text{supp}(f) \cap \text{supp}(1-\phi)). \end{aligned}$$

Proof. For a given $n \in \mathbb{Z}$, we have

$$\left\{ \frac{d_I m(I)^{-1}}{d_I + |c_I - gc_I|} \|f - f \circ g\|_{L_1(I)} \right\}_{\substack{I \in \mathbb{D}_\Gamma \\ I \cap \text{supp}(f) \neq \emptyset \\ 3I \cap \text{supp}(1-\phi) \neq \emptyset \\ d_I > 2^{-n}}} \leq 2 \|f\|_{L_1(\mathbb{R}^N)} \left\{ m(I)^{-1} \right\}_{\substack{I \in \mathbb{D}_\Gamma \\ I \cap \text{supp}(f) \neq \emptyset \\ 3I \cap \text{supp}(1-\phi) \neq \emptyset}}.$$

The right hand side obviously belongs to $(l_{N,\infty})_0(\mathbb{D}_\Gamma)$ and, hence, so does the left hand side. Thus,

$$\begin{aligned} \text{dist}_{l_{N,\infty}(\mathbb{D}_\Gamma)} \left(\left\{ \frac{d_I m(I)^{-1}}{d_I + |c_I - gc_I|} \|f - f \circ g\|_{L_1(I)} \right\}_{\substack{I \in \mathbb{D}_\Gamma \\ I \cap \text{supp}(f) \neq \emptyset \\ 3I \cap \text{supp}(1-\phi) \neq \emptyset}}, (l_{N,\infty})_0(\mathbb{D}_\Gamma) \right) \leq \\ \leq \left\| \left\{ \frac{d_I m(I)^{-1}}{d_I + |c_I - gc_I|} \|f - f \circ g\|_{L_1(I)} \right\}_{\substack{I \in \mathbb{D}_\Gamma \\ I \cap \text{supp}(f) \neq \emptyset \\ 3I \cap \text{supp}(1-\phi) \neq \emptyset \\ d_I \leq 2^{-n}}} \right\|_{l_{N,\infty}(\mathbb{D}_\Gamma)}. \end{aligned}$$

Obviously,

$$|f(x) - f(y)| \leq c_N |x - y| \cdot \|\nabla f\|_{L_\infty(\mathbb{R}^N)}.$$

Hence,

$$\begin{aligned} \|f - f \circ g\|_{L_1(I)} &\leq c_N \|\nabla f\|_{L_\infty(\mathbb{R}^N)} \times \int_I |x - gx| dx \leq \\ &\leq c'_\Phi m(I) \times (d_I + |c_I - gc_I|) \times \|\nabla f\|_{L_\infty(\mathbb{R}^N)}. \end{aligned}$$

Thus,

$$\frac{d_I m(I)^{-1}}{d_I + |c_I - gc_I|} \|f - f \circ g\|_{L_1(I)} \leq c_\Phi m(I)^{\frac{1}{N}} \|\nabla f\|_{L_\infty(\mathbb{R}^N)}$$

Finally,

$$\begin{aligned} \text{dist}_{l_{N,\infty}(\mathbb{D}_\Gamma)} \left(\left\{ \frac{d_I m(I)^{-1}}{d_I + |c_I - gc_I|} \|f - f \circ g\|_{L_1(I)} \right\}_{\substack{I \in \mathbb{D}_\Gamma \\ I \cap \text{supp}(f) \neq \emptyset \\ 3I \cap \text{supp}(1-\phi) \neq \emptyset}}, (l_{N,\infty})_0(\mathbb{D}_\Gamma) \right) \leq \\ \leq c_\Phi \|\nabla f\|_{L_\infty(\mathbb{R}^N)} \times \left\| \left\{ m(I)^{\frac{1}{N}} \right\}_{\substack{I \in \mathbb{D}_\Gamma \\ I \cap \text{supp}(f) \neq \emptyset \\ 3I \cap \text{supp}(1-\phi) \neq \emptyset \\ d_I \leq 2^{-n}}} \right\|_{l_{N,\infty}(\mathbb{D}_\Gamma)}. \end{aligned}$$

Passing $n \rightarrow \infty$ and using Lemma 4.12, we complete the proof. $\square$

Lemma 4.15. *Let $r > 2$. For every $f \in C_c^\infty(\mathbb{R}^N)$ and for every $n \in \mathbb{N}$, we have*

$$\begin{aligned} \text{dist}_{l_{N,\infty}(\mathbb{D}_\Gamma)} \left(\left\{ \frac{d_I m(I)^{-\frac{1}{r}}}{d_I + |c_I - gc_I|} \|f - f \circ g\|_{L_r(I)} \right\}_{\substack{I \in \mathbb{D}_\Gamma \\ I \cap \text{supp}(f) \neq \emptyset \\ 3I \cap \text{supp}(1-\phi) \neq \emptyset}}, (l_{N,\infty})_0(\mathbb{D}_\Gamma) \right) \leq \\ \leq c_\Phi \|\nabla f\|_{L_\infty(\mathbb{R}^N)} \times m^{\frac{1}{N}}(\text{supp}(f) \cap \text{supp}(1-\phi)). \end{aligned}$$

Proof. We obviously have

$$\begin{aligned} \|f - f \circ g\|_{L_r(I)} &\leq \left\| f - \frac{1}{m(I)} \int_I f \right\|_{L_r(I)} + \\ &+ \left\| f \circ g - \frac{1}{m(I)} \int_I (f \circ g) \right\|_{L_r(I)} + m(I)^{\frac{1}{r}-1} \|f - f \circ g\|_{L_1(I)}. \end{aligned}$$

By Poincare inequality,

$$\begin{aligned} &\|f - f \circ g\|_{L_r(I)} \leq \\ &\leq c_{\Phi,r} \left(m(I)^{\frac{1}{N}} \|\nabla f\|_{L_r(I)} + m(I)^{\frac{1}{N}} \|\nabla(f \circ g)\|_{L_r(I)} + m(I)^{\frac{1}{r}-1} \|f - f \circ g\|_{L_1(I)} \right). \end{aligned}$$

By the triangle inequality,

$$\begin{aligned} &\text{dist}_{l_{N,\infty}(\mathbb{D}_\Gamma)} \left(\left\{ \frac{d_I m(I)^{-\frac{1}{r}}}{d_I + |c_I - g c_I|} \|f - f \circ g\|_{L_r(I)} \right\}_{\substack{I \in \mathbb{D}_\Gamma \\ I \cap \text{supp}(f) \neq \emptyset \\ 3I \cap \text{supp}(1-\phi) \neq \emptyset}}, (l_{N,\infty})_0(\mathbb{D}_\Gamma) \right) \leq \\ &\leq \frac{N}{N-1} c_{\Phi,r} \text{dist}_{l_{N,\infty}(\mathbb{D}_\Gamma)} \left(\left\{ m(I)^{\frac{1}{N}-\frac{1}{r}} \|\nabla f\|_{L_r(I)} \right\}_{\substack{I \in \mathbb{D}_\Gamma \\ I \cap \text{supp}(f) \neq \emptyset \\ 3I \cap \text{supp}(1-\phi) \neq \emptyset}}, (l_{N,\infty})_0(\mathbb{D}_\Gamma) \right) + \\ &+ \frac{N}{N-1} c_{\Phi,r} \text{dist}_{l_{N,\infty}(\mathbb{D}_\Gamma)} \left(\left\{ m(I)^{\frac{1}{N}-\frac{1}{r}} \|\nabla f\|_{L_r(gI)} \right\}_{\substack{I \in \mathbb{D}_\Gamma \\ I \cap \text{supp}(f) \neq \emptyset \\ 3I \cap \text{supp}(1-\phi) \neq \emptyset}}, (l_{N,\infty})_0(\mathbb{D}_\Gamma) \right) + \\ &+ \text{dist}_{l_{N,\infty}(\mathbb{D}_\Gamma)} \left(\left\{ \frac{d_I m(I)^{-1}}{d_I + |c_I - g c_I|} \|f - f \circ g\|_{L_1(I)} \right\}_{\substack{I \in \mathbb{D}_\Gamma \\ I \cap \text{supp}(f) \neq \emptyset \\ 3I \cap \text{supp}(1-\phi) \neq \emptyset}}, (l_{N,\infty})_0(\mathbb{D}_\Gamma) \right). \end{aligned}$$

The assertion follows from Lemmas 4.13 and 4.14. $\square$

Proof of Theorem 4.2. The assertion follows by combining Lemmas 4.10 (applied with $f - f \circ g$ instead of f) and 4.15. $\square$

5. SECOND ROCHBERG-SEMMES BASED ESTIMATE

Theorem 5.1. *Let $N > 1$. Fix a closed Weyl chamber Γ . Suppose that $K : \Gamma \times \Gamma \rightarrow \mathbb{C}$ satisfies*

$$|\partial_x^\alpha \partial_y^\beta K|(x, y) \leq c_{K,\alpha,\beta} \frac{|x - y|^{-|\alpha|_1 - |\beta|_1}}{V(x, |x - y|)}, \quad x, y \in \Gamma, \quad \alpha, \beta \in \mathbb{Z}_+^N.$$

Let V be an integral operator with integral kernel K . For every $f \in \dot{W}^{1,N}(\Gamma)$, we have

$$\|[M_f, V]\|_{\mathcal{L}_{N,\infty}(L_2(\Gamma, w(x)dx))} \leq c_{K,\kappa} (\|f\|_{\dot{W}^{1,N}(\Gamma)} + \|f\|_{\dot{W}^{1,N}(g\Gamma)}).$$

Theorem 5.2. *In the setting of Theorem 4.1, let $f \in C_c^\infty$. If $0 \leq \phi_m \leq 1$ for $m \geq 1$ are in $C_c^\infty(\mathbb{R}^N)$ and $\text{supp}(1 - \phi_m)$ increases to $\mathbb{R}^N$, then*

$$\begin{aligned} &\text{dist}_{\mathcal{L}_{N,\infty}(L_2(\Gamma, w(x)dx))} \left(M_{\phi_m} \cdot [M_f, V] \cdot M_{\phi_m} - [M_f, V], (\mathcal{L}_{N,\infty})_0(L_2(\Gamma, w(x)dx)) \right) \rightarrow 0 \\ &\text{as } m \rightarrow \infty. \end{aligned}$$

Lemma 5.3. *Assume that $N > 1$ and $2 < r < \infty$. Let $f \in L_r^{\text{loc}}(\mathbb{R}^N)$. Fix a closed Weyl chamber Γ . Let K be as in Theorem 4.1. We have*

$$\begin{aligned} (f(x) - f(y)) \cdot K(x, y) &= \sum_{k, l \in \mathbb{Z}^N} (1 + |k|^2 + |l|^2)^{-N-1} \sum_{I \in \mathbb{D}_\Gamma} e_{k, l, I}^{(1)}(x) f_{k, l, I}^{(1)}(y) - \\ &\quad - \sum_{k, l \in \mathbb{Z}^N} (1 + |k|^2 + |l|^2)^{-N-1} \sum_{I \in \mathbb{D}_\Gamma} e_{k, l, I}^{(2)}(x) f_{k, l, I}^{(2)}(y) \end{aligned}$$

where

- (1) $e_{k, l, I}^{(1)}$ and $e_{k, l, I}^{(2)}$ are supported in I and $f_{k, l, I}^{(1)}$ and $f_{k, l, I}^{(2)}$ are supported in $3I$;
- (2) for every $k, l \in \mathbb{Z}^N$ and for every $I \in \mathbb{D}_\Gamma$, we have

$$\|e_{k, l, I}^{(1)}\|_{L_r(\mathbb{R}^N, w(x)dx)} \leq \left\| f - \frac{1}{m(3I)} \int_{3I} f \right\|_{L_r(3I, w(x)dx)};$$

$$\|f_{k, l, I}^{(2)}\|_{L_r(\mathbb{R}^N, w(x)dx)} = \left\| f - \frac{1}{m(3I)} \int_{3I} f \right\|_{L_r(3I, w(x)dx)};$$

- (3) for every $k, l \in \mathbb{Z}^N$ and for every $I \in \mathbb{D}_\Gamma$, we have

$$\|f_{k, l, I}^{(1)}\|_{L_\infty(\mathbb{R}^N)}, \|e_{k, l, I}^{(2)}\|_{L_\infty(\mathbb{R}^N)} \leq c_{K, \kappa, \Phi};$$

Proof. Repeating the argument in Lemma 4.5 *mutatis mutandi*, we obtain

$$(1 + |k|^2 + |l|^2)^{-N-1} \cdot |\text{coef}_{k, l, I}| \leq \frac{c_{K, \kappa, \Phi}}{w(I)}, \quad k, l \in \mathbb{Z}^N, \quad I \in \mathbb{D}_\Gamma.$$

Set

$$\begin{aligned} e_{k, l, I}^{(1)}(x) &= \left(f(x) - \frac{1}{m(3I)} \int_{3I} f \right) \cdot \chi_I(x) e^{i \langle k, \frac{\Phi^{-1}(x - c_I)}{d_I} \rangle}, \\ f_{k, l, I}^{(1)}(y) &= (1 + |k|^2 + |l|^2)^{N+1} \text{coef}_{k, l, I} \cdot \chi_{3I}(y) e^{i \langle l, \frac{\Phi^{-1}(y - c_I)}{d_I} \rangle}, \\ e_{k, l, I}^{(2)}(x) &= (1 + |k|^2 + |l|^2)^{N+1} \text{coef}_{k, l, I} \cdot \chi_I(x) e^{i \langle k, \frac{\Phi^{-1}(x - c_I)}{d_I} \rangle}, \\ f_{k, l, I}^{(2)}(y) &= \left(f(y) - \frac{1}{m(3I)} \int_{3I} f \right) \cdot \chi_{3I}(y) e^{i \langle l, \frac{\Phi^{-1}(y - c_I)}{d_I} \rangle}. \end{aligned}$$

The assertion follows immediately. $\square$

6. APPENDIX

Lemma 6.1. *Let $1 < s < \infty$, and let $\mathbb{D}^0$ be the standard dyadic grid in $\mathbb{R}^N$. Suppose that $F \in L^s(\mathbb{R}^N)$. Then there exists a constant $C > 0$ such that*

$$\left\| \left\{ |Q|^{-1/s'} \int_Q |F(x)| dx \right\}_{Q \in \mathbb{D}^0} \right\|_{\ell^{s, \infty}} \leq C \|F\|_{L^s(\mathbb{R}^N)}.$$

Proof. Let $\mathcal{E} \subset \mathbb{D}^0$ be finite and set

$$G_{\mathcal{E}}(x) := \sum_{Q \in \mathcal{E}} |Q|^{-1/s'} \chi_Q(x).$$

Since the dyadic cubes containing any fixed x are totally ordered by inclusion, for each $x \in \bigcup_{Q \in \mathcal{E}} Q$ we may choose a cube $Q_x \in \mathcal{E}$ of minimal volume such that

$x \in Q_x$. Every $Q \in \mathcal{E}$ containing x is then a dyadic ancestor of Q_x . Since the volume increases by the factor 2^N at each generation,

$$G_{\mathcal{E}}(x) = \sum_{\substack{Q \in \mathcal{E} \\ x \in Q}} |Q|^{-1/s'} \leq |Q_x|^{-1/s'} \sum_{k=0}^{\infty} 2^{-kN/s'} \lesssim_{N,s} \sup_{Q \in \mathcal{E}} |Q|^{-1/s'} \chi_Q(x).$$

Consequently,

$$\|G_{\mathcal{E}}\|_{L^{s'}(\mathbb{R}^N)}^{s'} \lesssim_{N,s} \int_{\mathbb{R}^N} \sup_{Q \in \mathcal{E}} |Q|^{-1} \chi_Q(x) dx \leq \int_{\mathbb{R}^N} \sum_{Q \in \mathcal{E}} |Q|^{-1} \chi_Q(x) dx = \#\mathcal{E}.$$

and hence,

$$\|G_{\mathcal{E}}\|_{L^{s'}(\mathbb{R}^N)} \lesssim_{N,s} (\#\mathcal{E})^{1/s'}.$$

By Hölder's inequality,

$$\sum_{Q \in \mathcal{E}} |Q|^{-1/s'} \int_Q |F(x)| dx = \int_{\mathbb{R}^N} |F(x)| G_{\mathcal{E}}(x) dx \lesssim_{N,s} \|F\|_{L^s(\mathbb{R}^N)} (\#\mathcal{E})^{1/s'}.$$

Fix $t > 0$ and let $\mathcal{E}$ be any finite subcollection of the dyadic cubes satisfying

$$|Q|^{-1/s'} \int_Q |F(x)| dx > t.$$

Then

$$t \#\mathcal{E} < \sum_{Q \in \mathcal{E}} |Q|^{-1/s'} \int_Q |F(x)| dx \lesssim_{N,s} \|F\|_{L^s(\mathbb{R}^N)} (\#\mathcal{E})^{1/s'},$$

so

$$\#\mathcal{E} \lesssim_{N,s} t^{-s} \|F\|_{L^s(\mathbb{R}^N)}^s.$$

Since $\mathcal{E}$ is arbitrary,

$$\#\left\{Q \in \mathbb{D}^0 : |Q|^{-1/s'} \int_Q |F(x)| dx > t\right\} \lesssim_{N,s} t^{-s} \|F\|_{L^s(\mathbb{R}^N)}^s, \quad t > 0.$$

The asserted weak ℓ^s estimate follows. This completes the proof of Lemma 6.1. $\square$

REFERENCES

- [1] Z. Fan, M. Lacey, and J. Li. Schatten classes and commutators of Riesz transform on Heisenberg group and applications. *J. Fourier Anal. Appl.*, 29(2):Paper No. 17, 28, 2023. [6](#)
- [2] Z. Fan, M. Lacey, J. Li, M.N. Vempati, and B.D. Wick. Besov space, Schatten classes, and commutators of Riesz transforms associated with the Neumann Laplacian. *Michigan Math. J.*, 75(1):199–224, 2025. [6](#)
- [3] Z. Fan, M.T. Lacey, J. Li, and X. Xiong. Schatten–Lorentz characterization of Riesz transform commutator associated with Bessel operators. Preprint available at <https://arxiv.org/abs/2403.08249>, 2024. [6](#)
- [4] L. Grafakos. *Classical Fourier analysis*, volume 249 of *Graduate Texts in Mathematics*. Springer, New York, third edition, 2014.
- [5] Y. Han, M. Lee, J. Li, and B.D. Wick. Riesz transforms and commutators in the Dunkl setting. *Anal. Math. Phys.*, 14(3):Paper No. 46, 32, 2024. [4](#)
- [6] G. Heckman. Coxeter groups. Lecture notes available at <https://www.math.ru.nl/~heckman/CoxeterGroups.pdf>. Radboud University Nijmegen, 2018. [7](#), [10](#), [16](#)
- [7] S. Helgason. *Differential geometry, Lie groups, and symmetric spaces*, volume 80 of *Pure and Applied Mathematics*. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York-London, 1978. [3](#)
- [8] T. Hytonen. Schatten properties of commutators on metric spaces. Preprint available at <https://arxiv.org/abs/2411.02613>, 2024. [6](#)

- [9] M.T. Lacey, J. Li, and B.D. Wick. Schatten classes and commutator in the two weight setting, I. Hilbert transform. *Potential Anal.*, 60(2):875–894, 2024. [6](#)
- [10] J. Li, J. Pipher, and L.A. Ward. Dyadic structure theorems for multiparameter function spaces. *Rev. Mat. Iberoam.*, 31(3):767–797, 2015.
- [11] J. Li, X. Xiong, and F. Yang. Schatten properties of Calderón–Zygmund singular integral commutator on stratified Lie groups. *J. Math. Pures Appl. (9)*, 188:73–113, 2024. [6](#)
- [12] R. Rochberg and S. Semmes. Nearly weakly orthonormal sequences, singular value estimates, and Calderon-Zygmund operators. *J. Funct. Anal.*, 86(2):237–306, 1989. [6](#)
- [13] Z. Wei and H. Zhang. Complex median method and Schatten class membership of commutators. Preprint available at <https://arxiv.org/abs/2411.05810>, 2024. [6](#)
- [14] Z. Wei and H. Zhang. On the boundedness and Schatten class property of noncommutative martingale paraproducts and operator-valued commutators. Preprint available at <https://arxiv.org/abs/2407.14988>, 2024. [6](#)

ZHIJIE FAN, SCHOOL OF MATHEMATICAL SCIENCES, SOUTH CHINA NORMAL UNIVERSITY, GUANGZHOU 510631, CHINA

Email address: fanzhj3@mail2.sysu.edu.cn

Ji LI, DEPARTMENT OF MATHEMATICS, MACQUARIE UNIVERSITY, NSW2109, AUSTRALIA

Email address: ji.li@mq.edu.au

SCHOOL OF MATHEMATICS AND STATISTICS, HNP-LAMA, CENTRAL SOUTH UNIVERSITY, CHANGSHA 410075, CHINA

Email address: d.zanin@csu.edu.cn