

WEYL–CHAMBER GEOMETRY, POINCARÉ INEQUALITY MEETS SCHATTEN CLASS PROPERTIES OF DUNKL–RIESZ TRANSFORM COMMUTATORS

ZHIJIE FAN, JI LI, AND DMITRIY ZANIN

Abstract.

Contents

1. Introduction	1
2. Preliminaries	2
3. Analytic tools for Weyl chamber geometry	9
4. Proof of upper bound	17
5. Proof of lower bound	19
6. Spectral asymptotic formula	24
7. Alternative proof by using Dima's approach	41
8. Appendix	46
9. Answer to Dima's question	47
References	56

1. Introduction

Definition 1.1. Let $1 \leq r < \infty$. For a measurable set $E \subset \mathbb{R}^N$ of finite positive measure and $f \in L^r(E)$, define

$$\text{osc}_r(f; E) := \left(\frac{1}{|E|} \int_E |f - (f)_E|^r dx \right)^{1/r}.$$

Definition 1.2. Let $0 < p < \infty$, $0 < q \leq \infty$, and $1 \leq r < \infty$. Define

$$OSC^{p,q}(\mathbb{R}^N) := \left\{ f \in L^r_{\text{loc}}(\mathbb{R}^N) : \|f\|_{OSC^{p,q}(\mathbb{R}^N)} < \infty \right\},$$

where

$$\|f\|_{OSC^{p,q}(\mathbb{R}^N)} := \sum_{v=1}^{2^N} \left\| \{\text{osc}_r(f; Q)\}_{Q \in \mathbb{D}^v} \right\|_{\ell^{p,q}}.$$

By [11, Proposition 4.1], the above definition is independent of r .

Date: September 11, 2026.

2010 Mathematics Subject Classification. 47B10, 42B20, 43A85.

Key words and phrases. Schatten class, Riesz transform commutators, Dunkl operator, Besov space, Nearly weakly orthogonal sequence.

1.1. Notation. Conventionally, we let $\mathbb{N}$ denote the set of positive integers and set $\mathbb{Z}_+ := \{0\} \cup \mathbb{N}$. Throughout the paper, the symbols C and c denote positive constants (possibly varying from line to line) that may depend only on N , p , and the structural data (R, k) of the Dunkl setting, but are independent of the main parameters (e.g., the symbol b and auxiliary decay factors). We write $A \lesssim B$ to mean that $A \leq CB$ for some constant $C > 0$, and $A \sim B$ to mean that both $A \lesssim B$ and $B \lesssim A$ hold. For $\lambda > 0$ and a cube Q with sidelength $\ell(Q)$, write λQ for the cube concentric with Q and sidelength $\lambda \ell(Q)$. Similarly, for every ball $B = B(x, r)$, set $\lambda B := B(x, \lambda r)$. For any $1 \leq p \leq \infty$, we denote by p' the conjugate of p , which satisfies $\frac{1}{p} + \frac{1}{p'} = 1$. We denote the average of a function f over a measurable set E by

$$(f)_{E,dx} := \frac{1}{|E|} \int_E f(x) dx.$$

For a subset $E \subset \mathbb{R}^N$, we denote by χ_E its characteristic function.

Denote by $\{e_j\}_{j=1}^N$ the standard orthonormal basis on $\mathbb{R}^N$.

2. Preliminaries

2.1. Harmonic analysis in the Dunkl setting. Let $\mathbb{R}^N$ be the Euclidean space equipped with the usual scalar product $\langle \cdot, \cdot \rangle$ and the induced norm $\|\cdot\|$. For a nonzero vector $v \in \mathbb{R}^N$, the reflection σ_v with respect to the hyperplane $v^\perp$ orthogonal to v is defined by

$$\sigma_v x := x - 2 \frac{\langle x, v \rangle}{\|v\|^2} v.$$

A finite set $R \subset \mathbb{R}^N \setminus \{0\}$ is called a root system if $\sigma_v(R) = R$ for every $v \in R$. A root system R is called a normalized reduced root system if $\|v\|^2 = 2$ for every $v \in R$. The finite group G generated by the reflections $\{\sigma_v\}_{v \in R}$ is called the Weyl group (reflection group) of the root system R . Let $k : R \rightarrow [0, \infty)$ be a multiplicity function, which is a G -invariant function fixed throughout the paper. We will denote by $\mathcal{O}(x) = \{\sigma(x) : \sigma \in G\}$ the G -orbit of a point $x \in \mathbb{R}^N$, and $\mathcal{O}(E) = \bigcup_{x \in E} \mathcal{O}(x)$ the G -orbit of $E \subset \mathbb{R}^N$.

Define the Dunkl measure dw on $\mathbb{R}^N$ by

$$dw(x) := \prod_{v \in R} |\langle x, v \rangle|^{k(v)} dx,$$

where dx stands for the Lebesgue measure on $\mathbb{R}^N$. Let $B(x, r) := \{y \in \mathbb{R}^N : \|x - y\| < r\}$ be the Euclidean ball centred at x with radius $r > 0$. Let $\mathbf{N} := N + \sum_{v \in R} k(v)$. Note that

$$w(B(tx, tr)) = t^{\mathbf{N}} w(B(x, r)), \text{ for } x \in \mathbb{R}^N, t, r > 0.$$

Furthermore,

$$(2.1) \quad w(B(x, r)) \sim r^{\mathbf{N}} \prod_{v \in R} (|\langle x, v \rangle| + r)^{k(v)},$$

where the implicit constant only depends on N, R and k . It follows from (2.1) that there exists a constant $C \geq 1$ such that for every $x \in \mathbb{R}^N$,

$$(2.2) \quad C^{-1} \left(\frac{r_2}{r_1} \right)^N \leq \frac{w(B(x, r_2))}{w(B(x, r_1))} \leq C \left(\frac{r_2}{r_1} \right)^N, \text{ for } 0 < r_1 < r_2,$$

so the numbers $\mathbf{N}$ and N are called the upper and lower homogeneous dimensions, respectively.

[Add the definition of A_∞ . Zhijie will do it.]

Lemma 2.1. Let

$$dw(x) = \varpi(x) dx, \quad \varpi(x) = \prod_{v \in R_+} |\langle x, v \rangle|^{2k(v)}.$$

Then $\varpi \in A_\infty(\mathbb{R}^N, dx)$.

Proof. The assertion is immediate if $k \equiv 0$. Otherwise, choose $p > 1$ sufficiently large so that

$$\frac{2|R_+|k(v)}{p-1} < 1, \quad v \in R_+.$$

For every $0 < \beta < 1$, $v \in R_+$, and Euclidean ball $B(x_0, r)$, we apply an orthogonal change of variables to deduce that

$$\begin{aligned} \int_{B(x_0, r)} |\langle x, v \rangle|^{-\beta} dx &\lesssim \frac{1}{r|v|} \int_{\langle x_0, v \rangle - |v|r}^{\langle x_0, v \rangle + |v|r} |s|^{-\beta} ds \\ &= (|v|r)^{-\beta} \int_{\frac{\langle x_0, v \rangle}{|v|r} - 1}^{\frac{\langle x_0, v \rangle}{|v|r} + 1} |s|^{-\beta} ds \\ &\lesssim (|v|r)^{-\beta} \left(1 + \frac{|\langle x_0, v \rangle|}{|v|r} \right)^{-\beta} \\ &\lesssim (|\langle x_0, v \rangle| + r)^{-\beta}. \end{aligned}$$

Hence, by Hölder's inequality,

$$\begin{aligned} \int_{B(x_0, r)} \varpi^{-\frac{1}{p-1}} dx &= \int_{B(x_0, r)} \prod_{v \in R_+} |\langle x, v \rangle|^{-\frac{2k(v)}{p-1}} dx \\ &\leq \prod_{v \in R_+} \left(\int_{B(x_0, r)} |\langle x, v \rangle|^{-\frac{2|R_+|k(v)}{p-1}} dx \right)^{\frac{1}{|R_+|}} \\ &\lesssim \prod_{v \in R_+} (|\langle x_0, v \rangle| + r)^{-\frac{2k(v)}{p-1}}. \end{aligned}$$

On the other hand, by (2.1) and the G -invariance of k ,

$$\int_{B(x_0, r)} \varpi dx = \frac{w(B(x_0, r))}{|B(x_0, r)|} \lesssim \prod_{v \in R_+} (|\langle x_0, v \rangle| + r)^{2k(v)}.$$

Therefore,

$$\left(\int_B \varpi dx \right) \left(\int_B \varpi^{-\frac{1}{p-1}} dx \right)^{p-1} \lesssim 1$$

uniformly over all Euclidean balls B . Thus $\varpi \in A_p(\mathbb{R}^N, dx)$, and hence $\varpi \in A_\infty(\mathbb{R}^N, dx)$. $\square$

The Dunkl operators T_ξ are defined by

$$T_\xi f(x) := \partial_\xi f(x) + \sum_{v \in R} \frac{k(v)}{2} \langle v, \xi \rangle \frac{f(x) - f(\sigma_v(x))}{\langle v, x \rangle},$$

where $\partial_\xi f$ denotes the directional derivative of f in the direction ξ . Furthermore, the Dunkl Laplacian is defined by

$$\Delta_{\text{Dunkl}} f(x) := \sum_{j=1}^N T_{e_j}^2 f(x) = \Delta_{\mathbb{R}^N} f(x) + \sum_{v \in R} k(v) \delta_v f(x),$$

where $\Delta_{\mathbb{R}^N}$ is the Laplacian on $\mathbb{R}^N$ and δ_v is defined by

$$\delta_v f(x) := \frac{\partial_v f(x)}{\langle v, x \rangle} - \frac{f(x) - f(\sigma_v(x))}{\langle v, x \rangle^2}.$$

In this article, we consider the Dunkl-Riesz transforms defined by $R_{\text{Dunkl},j} := T_{e_j}(-\Delta_{\text{Dunkl}})^{-1/2}$, $j = 1, 2, \dots, N$.

Definition 2.2. The Weyl chambers are defined as the closures of connected components of the set

$$(2.3) \quad \{x \in \mathbb{R}^N : \langle x, v \rangle \neq 0 \text{ for all } v \in R\}.$$

Define $d(x, y) := \min_{\sigma \in G} \|x - \sigma(y)\|$ as the distance between two G -orbits $O(x)$ and $O(y)$. Recall from [6, Chapter VII, proof of Theorem 2.12] that

$$(2.4) \quad d(x, y) = \|x - y\| \text{ if and only if } x, y \in \Gamma$$

for some closed Weyl chamber Γ .

It is clear that $O(B(x, r)) = \{y \in \mathbb{R}^N : d(x, y) < r\}$. Moreover,

$$w(B(x, r)) \leq w(O(B(x, r))) \leq |G|w(B(x, r))$$

for all $x \in \mathbb{R}^N$ and $r > 0$. We also denote $B^d(x, r) := \{y \in \mathbb{R}^N : d(x, y) < r\}$, i.e., $B^d(x, r) = O(B(x, r))$.

Denote by $K_{\text{Dunkl},j}(x, y)$ the integral kernel of $R_{\text{Dunkl},j}$. For every $x, y \in \mathbb{R}^N$ and $r > 0$, let

$$V(x, y, r) := \max\{w(B(x, r)), w(B(y, r))\}.$$

The following lemma establishes derivative estimates for the Dunkl-Riesz transform kernel.

Lemma 2.3. Assume that $N \geq 1$. Then for every pair of multi-indices α, β , every $j \in \{1, \dots, N\}$, and every $x, y \in \mathbb{R}^N$ with $d(x, y) > 0$,

$$(2.5) \quad \left| \partial_x^\alpha \partial_y^\beta K_{\text{Dunkl},j}(x, y) \right| \lesssim_{\alpha, \beta, R, k} \frac{d(x, y)}{\|x - y\|} \frac{d(x, y)^{-(|\alpha| + |\beta|)}}{V(x, y, d(x, y))}.$$

Proof. Case 1. Suppose that $|\alpha| + |\beta| = 0$. By [4, Theorem 1.1],

$$|K_{\text{Dunkl},j}(x, y)| \lesssim \frac{d(x, y)}{\|x - y\|} \frac{1}{w(B(x, d(x, y)))}.$$

By the definition of d , the G -invariance of dw , and (2.2),

$$V(x, y, d(x, y)) \lesssim w(B(x, d(x, y))),$$

and hence (2.5) follows.

Case 2. Suppose that $|\alpha| + |\beta| \geq 1$. By [?, Theorem 3.1],

$$|\partial_x^\alpha \partial_y^\beta h_t(x, y)| \lesssim_{\alpha, \beta, R, k} t^{-\frac{|\alpha|+|\beta|}{2}} \left(1 + \frac{\|x - y\|}{\sqrt{t}}\right)^{-2} V(x, y, \sqrt{t})^{-1} e^{-cd(x, y)^2/t}.$$

Recall from the proof of Lemma 2.4 that

$$K_{\text{Dunkl}, j}(x, y) = -\frac{C}{2} \int_0^\infty (y_j - x_j) h_t(x, y) \frac{dt}{t^{3/2}}.$$

Since

$$\partial_x^\alpha \partial_y^\beta ((y_j - x_j) h_t(x, y)) = (y_j - x_j) \partial_x^\alpha \partial_y^\beta h_t(x, y) - \alpha_j \partial_x^{\alpha - e_j} \partial_y^\beta h_t(x, y) + \beta_j \partial_x^\alpha \partial_y^{\beta - e_j} h_t(x, y),$$

where the corresponding term is omitted if $\alpha_j = 0$ or $\beta_j = 0$, we obtain

$$\begin{aligned} |\partial_x^\alpha \partial_y^\beta K_{\text{Dunkl}, j}(x, y)| &\lesssim \frac{1}{\|x - y\|} \int_0^\infty t^{-\frac{|\alpha|+|\beta|+1}{2}} V(x, y, \sqrt{t})^{-1} e^{-cd(x, y)^2/t} dt \\ &\quad + \frac{1}{\|x - y\|^2} \int_0^\infty t^{-\frac{|\alpha|+|\beta|}{2}} V(x, y, \sqrt{t})^{-1} e^{-cd(x, y)^2/t} dt. \end{aligned}$$

By (2.2),

$$V(x, y, \sqrt{t})^{-1} \lesssim V(x, y, d(x, y))^{-1} \begin{cases} \left(\frac{d(x, y)}{\sqrt{t}}\right)^N, & 0 < t \leq d(x, y)^2, \\ \left(\frac{d(x, y)}{\sqrt{t}}\right)^N, & t \geq d(x, y)^2. \end{cases}$$

Hence,

$$\begin{aligned} &\int_0^\infty t^{-\frac{|\alpha|+|\beta|+1}{2}} V(x, y, \sqrt{t})^{-1} e^{-cd(x, y)^2/t} dt \\ &\lesssim \frac{1}{V(x, y, d(x, y))} \left(d(x, y)^N \int_0^{d(x, y)^2} t^{-\frac{N+|\alpha|+|\beta|+1}{2}} e^{-cd(x, y)^2/t} dt + d(x, y)^N \int_{d(x, y)^2}^\infty t^{-\frac{N+|\alpha|+|\beta|+1}{2}} dt \right) \\ &\lesssim \frac{d(x, y)^{1-|\alpha|-|\beta|}}{V(x, y, d(x, y))}. \end{aligned}$$

The first integral in the parentheses is estimated by the change of variables $s = d(x, y)^2/t$, while the second one converges since $N + |\alpha| + |\beta| > 1$.

Case 2.1. Suppose that $N + |\alpha| + |\beta| > 2$. We apply the same argument to see that

$$\begin{aligned} &\int_0^\infty t^{-\frac{|\alpha|+|\beta|}{2}} V(x, y, \sqrt{t})^{-1} e^{-cd(x, y)^2/t} dt \\ &\lesssim \frac{1}{V(x, y, d(x, y))} \left(d(x, y)^N \int_0^{d(x, y)^2} t^{-\frac{N+|\alpha|+|\beta|}{2}} e^{-cd(x, y)^2/t} dt + d(x, y)^N \int_{d(x, y)^2}^\infty t^{-\frac{N+|\alpha|+|\beta|}{2}} dt \right) \\ &\lesssim \frac{d(x, y)^{2-|\alpha|-|\beta|}}{V(x, y, d(x, y))}. \end{aligned}$$

Since $d(x, y) \leq \|x - y\|$, it follows that

$$|\partial_x^\alpha \partial_y^\beta K_{\text{Dunkl}, j}(x, y)| \lesssim_{\alpha, \beta, R, k} \frac{d(x, y)}{\|x - y\|} \frac{d(x, y)^{-(|\alpha|+|\beta|)}}{V(x, y, d(x, y))}.$$

Case 2.2. Suppose that $N = 1$ and $|\alpha| + |\beta| = 1$. For the terms in which a derivative falls on $y_j - x_j$, retaining the factor $(1 + \|x - y\|/\sqrt{t})^{-2}$ and splitting the integral at $d(x, y)^2$ and $\|x - y\|^2$, we obtain from (2.2)

$$\begin{aligned} & \int_0^\infty t^{-3/2} \left(1 + \frac{\|x - y\|}{\sqrt{t}}\right)^{-2} V(x, y, \sqrt{t})^{-1} e^{-cd(x, y)^2/t} dt \\ & \lesssim \frac{d(x, y)^N}{\|x - y\|^2 V(x, y, d(x, y))} \int_0^{d(x, y)^2} t^{-\frac{N+1}{2}} e^{-cd(x, y)^2/t} dt \\ & \quad + \frac{d(x, y)}{\|x - y\|^2 V(x, y, d(x, y))} \int_{d(x, y)^2}^{\|x - y\|^2} \frac{dt}{t} + \frac{d(x, y)}{V(x, y, d(x, y))} \int_{\|x - y\|^2}^\infty \frac{dt}{t^2} \\ & \lesssim \frac{1}{\|x - y\| V(x, y, d(x, y))}, \end{aligned}$$

where we used

$$\frac{d(x, y)}{\|x - y\|} \log \frac{\|x - y\|}{d(x, y)} \lesssim 1.$$

Combining this estimate with the preceding estimate for the first term, we obtain

$$|\partial_x^\alpha \partial_y^\beta K_{\text{Dunkl}, j}(x, y)| \lesssim_{\alpha, \beta, R, k} \frac{1}{\|x - y\| V(x, y, d(x, y))} = \frac{d(x, y)}{\|x - y\|} \frac{d(x, y)^{-1}}{V(x, y, d(x, y))}.$$

This completes the proof of Lemma 2.3. $\square$

Lemma 2.4. For $j \in \{1, 2, \dots, N\}$, and for every ball $B = B(x_0, r) \subset \mathbb{R}^N$, there is another ball $\widehat{B} = B(y_0, r)$ with $y_0 = x_0 + 5re_j$ such that for every $(x, y) \in B \times \widehat{B}$, $K_{\text{Dunkl}, j}(x, y)$ does not change sign and satisfies

$$|K_{\text{Dunkl}, j}(x, y)| \geq \frac{C}{w(B(x_0, r))}.$$

Proof. The lower bound is given in [4, Theorem 1.2], while the fact that $K_{\text{Dunkl}, j}(x, y)$ does not change sign can be seen in the proof of [4, Theorem 1.2]. Indeed, with the choice $y_0 := x_0 + 5re_j$, we have $y_j - x_j \geq 3r$ for all $(x, y) \in B \times \widehat{B}$. Together with the fact that the heat kernel $h_t(x, y)$ associated with the Dunkl Laplacian is positive, the representation

$$K_{\text{Dunkl}, j}(x, y) = -\frac{C}{2} \int_0^\infty \frac{y_j - x_j}{t} h_t(x, y) \frac{dt}{\sqrt{t}},$$

implies that $K_{\text{Dunkl}, j}(x, y) < 0$ (and thus, does not change sign) on $B \times \widehat{B}$. $\square$

2.2. Systems of dyadic cubes.

Definition 2.5. Let $X \subseteq \mathbb{R}^N$ be a nonempty open set. A countable family $\mathcal{D} := \bigcup_{k \in \mathbb{Z}} \mathcal{D}_k$, $\mathcal{D}_k := \{Q_\alpha^k : \alpha \in \mathcal{A}_k\}$, of Borel sets $Q_\alpha^k \subseteq X$ is called a system of dyadic cubes over X if it has the following properties:

(i) For each $k \in \mathbb{Z}$, we have

$$X = \bigcup_{\alpha \in \mathcal{A}_k} Q_\alpha^k;$$

- (ii) If $\ell \geq k$, then either $Q_\beta^\ell \subseteq Q_\alpha^k$ or $Q_\alpha^k \cap Q_\beta^\ell = \emptyset$;
- (iii) For each (k, α) and each $\ell \leq k$, there exists a unique β such that $Q_\alpha^k \subseteq Q_\beta^\ell$;
- (iv) For each (k, α) there exist 2^N indices β such that

$$Q_\beta^{k+1} \subseteq Q_\alpha^k, \text{ and } Q_\alpha^k = \bigcup_{\substack{Q \in \mathcal{D}_{k+1} \\ Q \subseteq Q_\alpha^k}} Q;$$

- (v) For each (k, α) , one has

$$B_X(x_\alpha^k, A_1 2^{-k}) \subseteq Q_\alpha^k \subseteq B_X(x_\alpha^k, A_2 2^{-k}) =: B(Q_\alpha^k)$$

for some positive constants A_1, A_2 depending only on the geometry of X , where we set $B_X(x, r) := B(x, r) \cap X$ for all $x \in X$ and $r > 0$.

- (vi) If $\ell \geq k$ and $Q_\beta^\ell \subseteq Q_\alpha^k$, then

$$B(Q_\beta^\ell) \subseteq B(Q_\alpha^k).$$

Each Q_α^k is called a dyadic cube of generation k , with center $x_\alpha^k \in Q_\alpha^k$ and side-length $\ell(Q_\alpha^k) = 2^{-k}$.

Notation 2.6. For every dyadic cube $Q \in \mathcal{D}$, we denote by c_Q and $\ell(Q)$ its center and side-length, respectively.

Notation 2.7. Let

$$\mathbb{D}^0 := \bigcup_{k \in \mathbb{Z}} \mathbb{D}_k^0$$

be the standard system of dyadic cubes over $\mathbb{R}^N$. Here each dyadic cube is a product of half-open intervals

$$[m_1 2^{-k}, (m_1 + 1) 2^{-k}) \times \cdots \times [m_N 2^{-k}, (m_N + 1) 2^{-k}),$$

with $(m_1, \dots, m_N) \in \mathbb{Z}^N$. Moreover, we let

$$\mathbb{D}_\Gamma := \bigcup_{k \in \mathbb{Z}} \mathbb{D}_{\Gamma, k}$$

be the system of dyadic cubes over a fixed chamber Γ , whose construction will be given in Corollary 3.7.

Lemma 2.8. There exist 2^N systems of dyadic cubes $\mathbb{D}^1, \mathbb{D}^2, \dots, \mathbb{D}^{2^N}$ such that for any ball $B = B(x_B, r_B) \subset \mathbb{R}^N$ with $\frac{2^{-k-2}}{3} \leq r_B < \frac{2^{-k-1}}{3}$ for some $k \in \mathbb{Z}$, there exists a cube $Q \in \bigcup_{v=1}^{2^N} \mathbb{D}_k^v$ such that

$$B \subset Q \subset 12 \sqrt{N} B.$$

Proof. • Step 1. Recall the one-dimensional setting given in [9, Section 2].

Let $\mathbb{D}^{(0)}(\mathbb{R})$ be the standard system of dyadic intervals over $\mathbb{R}$ (see Notation 2.7). Let $\delta = 1/3$ and define

$$\mathbb{D}^{(\delta)}(\mathbb{R}) = \bigcup_{k \in \mathbb{Z}} \mathbb{D}_k^{(\delta)}(\mathbb{R}),$$

where for $k \geq 0$,

$$\mathbb{D}_k^{(\delta)}(\mathbb{R}) = \{I + \delta \mid I \in \mathbb{D}_k^{(0)}(\mathbb{R})\},$$

while for $k < 0$ and k even,

$$\mathbb{D}_k^{(\delta)}(\mathbb{R}) = \left\{ \left[\frac{m}{2^k} + \delta + \sum_{j=(k/2)+1}^0 2^{-2j}, \frac{m+1}{2^k} + \delta + \sum_{j=(k/2)+1}^0 2^{-2j} \right) \mid m \in \mathbb{Z} \right\}.$$

These choices together with the nestedness property completely determine the collections $\mathbb{D}_k^{(\delta)}(\mathbb{R})$ for $k < 0$, k odd.

Fix an interval $J \subset \mathbb{R}$ with $2^{-k-1}/3 \leq |J| < 2^{-k}/3$ for some $k \in \mathbb{Z}$. Now we set $A_k = \{m \cdot 2^{-k} \mid m \in \mathbb{Z}\}$ for every fixed k and

- (1) $A_k^{(\delta)} = \{\delta + m \cdot 2^{-k} \mid m \in \mathbb{Z}\}$ for $k \geq 0$,
- (2) $A_k^{(\delta)} = \{\delta + \sum_{j=(k/2)+1}^0 2^{-2j} + m \cdot 2^{-k} \mid m \in \mathbb{Z}\}$ for $k < 0$, k even, and
- (3) $A_k^{(\delta)} = \{\delta + \sum_{j=(k-1)/2+1}^0 2^{-2j} + m \cdot 2^{-k} \mid m \in \mathbb{Z}\}$ for $k < 0$, k odd.

Note that for any two different points $a, b \in A_k \cup A_k^{(\delta)}$, we have $|a - b| \geq 2^{-k}/3 > |J|$. Thus, there is at most one element of $A_k \cup A_k^{(\delta)}$ contained in J . Hence, $J \cap A_k = \emptyset$ or $J \cap A_k^{(\delta)} = \emptyset$. Therefore, J must be contained in some dyadic interval $I \in \mathbb{D}_k^{(0)}$ or $I \in \mathbb{D}_k^{(\delta)}$ and $|I| = 2^{-k} \leq 6|J|$.

• Step 2. We extend the construction to $\mathbb{R}^N$ ($N \geq 1$).

For $\vartheta = (\vartheta_1, \vartheta_2, \dots, \vartheta_N) \in \{0, \delta\}^N$, define

$$\mathbb{D}^{(\vartheta)} := \bigcup_{k \in \mathbb{Z}} \mathbb{D}_k^{(\vartheta)}, \quad \mathbb{D}_k^{(\vartheta)} := \left\{ \prod_{j=1}^N I_j : I_j \in \mathbb{D}_k^{(\vartheta_j)} \right\}.$$

Applying Step 1 to $\pi_i(B)$, where $\pi_i(B)$ denotes the i -th coordinate projection, we obtain $I_i \in \mathbb{D}_k^{(0)}$ or $I_i \in \mathbb{D}_k^{(\delta)}$ with $\pi_i(B) \subset I_i$ and $|I_i| = 2^{-k} \leq 12r_B$. Let $\vartheta_i \in \{0, \delta\}$ be the corresponding shift so that $I_i \in \mathbb{D}_k^{(\vartheta_i)}$, set $\vartheta := (\vartheta_1, \dots, \vartheta_N)$, and let $Q = \prod_{j=1}^N I_j \in \mathbb{D}_k^{(\vartheta)}$. It follows that $B \subset Q$ and $\ell(Q) = 2^{-k} \leq 12r_B$. Hence, for every $y \in Q$,

$$\|y - x_B\| \leq \sqrt{N} \ell(Q) \leq 12 \sqrt{N} r_B.$$

This implies that $Q \subset B(x_B, 12 \sqrt{N} r_B)$. Enumerating $\{\mathbb{D}^{(\vartheta)} : \vartheta \in \{0, \delta\}^N\}$ as $\mathbb{D}^1, \dots, \mathbb{D}^{2^N}$, we complete the proof of Lemma 2.8. $\square$

2.3. Nearly weakly orthonormal sequences. Rochberg and Semmes [11] introduced the notion of nearly weakly orthonormal (NWO) sequences of functions. This notion has since proved highly effective in characterizing Schatten class membership of singular integral commutators in a variety of contexts (see [1–3, 7, 8, 10–13]). The properties of NWO sequences were originally investigated in the Euclidean setting (equipped with Lebesgue measure), but the extension to spaces of homogeneous type is straightforward once a system of dyadic cubes is available in this general framework (see [7, Section 13] for a complete proof). For our purposes, we adapt the extension to the Dunkl setting $(\mathbb{R}^N, \|\cdot\|, w)$ rather than recalling the general definition of NWO sequences and the general extension to spaces of homogeneous type.

Let $\mathcal{D}$ be a system of dyadic cubes. A sequence of functions $\{e_Q\}_{Q \in \mathcal{D}}$ is an NWO sequence if

- (1) $\text{supp } e_Q \subseteq cQ$ for some positive constant c independent of Q
- (2) $\|e_Q\|_{L^s(\mathbb{R}^N, dw)} \leq w(Q)^{1/s-1/2}$ for some $s > 2$.

[We need to replace the following Lemma by Hytonen's version on space of homogeneous type. Zhijie Will do it later]

Lemma 2.9. Let $1 < p < \infty$, $1 \leq q \leq \infty$, and let $\mathcal{D}$ be a system of dyadic cubes over $\mathbb{R}^N$. If

$$T \in S^{p,q}(L^2(\mathbb{R}^N, dw)),$$

then

$$\left\| \left\{ \langle T e_Q, f_Q \rangle_{L^2(\mathbb{R}^N, dw)} \right\}_{Q \in \mathcal{D}} \right\|_{\ell^{p,q}} \leq C_{N,p,q,w} \|T\|_{S^{p,q}(L^2(\mathbb{R}^N, dw))},$$

where $\{e_Q\}_{Q \in \mathcal{D}}$ and $\{f_Q\}_{Q \in \mathcal{D}}$ are NWO sequences.

Proof. This property can be found in [11, (1.10)] (see also [7, Section 7]). $\square$

Lemma 2.10. Let $0 < p < \infty$, $0 < q \leq \infty$, and let $\mathcal{D}$ be a system of dyadic cubes over $\mathbb{R}^N$. Assume that

$$A = \sum_{Q \in \mathcal{D}} \lambda_Q \langle \cdot, e_Q \rangle_{L^2(\mathbb{R}^N, dw)} f_Q,$$

where $\{e_Q\}_{Q \in \mathcal{D}}$ and $\{f_Q\}_{Q \in \mathcal{D}}$ are NWO sequences and $\{\lambda_Q\}_{Q \in \mathcal{D}}$ is a sequence of scalars. We have

$$(2.6) \quad \|A\|_{S^{p,q}} \leq C_{N,p,q,w} \|\{\lambda_Q\}_Q\|_{\ell^{p,q}}.$$

Proof. This property can be found in [11, (1.9)] (see also [7, Section 7]). $\square$

3. Analytic tools for Weyl chamber geometry

Convention. Throughout this paper, we will use the following notation:

- Let $R \subset \mathbb{R}^N$ be a finite normalized reduced root system;
- Let $W := \text{span}(R)$ with $N_0 := \dim W$;
- Let $\Omega := \mathbb{R}_+^{N_0} \times \mathbb{R}^{N-N_0}$;
- Let Γ_+ be a fixed open Weyl chamber (see Definition 3.1) and Γ be its closure.

Definition 3.1. [5, Definition 2.7] A connected component of

$$(3.1) \quad \{x \in \mathbb{R}^N : \langle x, v \rangle \neq 0 \text{ for all } v \in R\}$$

is called an open Weyl chamber. Its closure is called a Weyl chamber. Define the corresponding sets of positive and negative roots by

$$R_+ := \{\alpha \in R : \langle x, \alpha \rangle > 0 \text{ for all } x \in \Gamma_+\}, \quad R_- := -R_+.$$

Definition 3.2. [5, Definition 2.9] A root $\alpha \in R_+$ is called simple in R_+ if it cannot be written as

$$\alpha = \lambda_1 \alpha_1 + \lambda_2 \alpha_2 \quad \text{with } \lambda_1, \lambda_2 \geq 1 \text{ and } \alpha_1, \alpha_2 \in R_+.$$

The set $\Delta \subset R_+$ of all simple roots is called the simple system.

Lemma 3.3. With the notation given in Definition 3.2, Δ is a basis of W . If we denote by $\Delta = \{\alpha_1, \alpha_2, \dots, \alpha_{N_0}\}$, then the open and closed chambers can be written as

$$(3.2) \quad \Gamma_+ = \{x \in \mathbb{R}^N : \langle x, \alpha_i \rangle > 0, i = 1, \dots, N_0\},$$

$$(3.3) \quad \Gamma = \{x \in \mathbb{R}^N : \langle x, \alpha_i \rangle \geq 0, i = 1, \dots, N_0\},$$

and every root $\alpha \in R$ admits a unique expansion

$$\alpha = \sum_{i=1}^{N_0} c_i \alpha_i$$

such that either $c_i \geq 0$ for all i (when $\alpha \in R_+$), or $c_i \leq 0$ for all i (when $\alpha \in R_-$). In particular, the simple system Δ is uniquely determined by Γ_+ (equivalently, by Γ).

Proof. Let $\Delta = \{\alpha_1, \dots, \alpha_n\} \subset R_+$ be the set of simple roots in R_+ . By [5, Proposition 2.10], every $\alpha \in R_+$ is a nonnegative linear combination of $\alpha_1, \dots, \alpha_n$. Since $R = R_+ \cup R_-$ (this follows from the fact that for each $\alpha \in R$, the continuous function $x \mapsto \langle x, \alpha \rangle$ does not vanish on the connected set Γ_+) and $R_- = -R_+$, it follows that $R \subset \text{span}(\Delta)$ and hence,

$$W = \text{span}(R) \subset \text{span}(\Delta).$$

The reverse inclusion $\text{span}(\Delta) \subset W$ is immediate from $\Delta \subset R \subset W$, so $\text{span}(\Delta) = W$. On the other hand, it follows from [5, Theorem 2.16] that the simple roots are linearly independent. Consequently, Δ is a basis of W , and therefore $n = \dim W = N_0$. Moreover, every $\alpha \in R$ has a unique expansion $\alpha = \sum_{i=1}^{N_0} c_i \alpha_i$. If $\alpha \in R_+$, the coefficients can be chosen nonnegative by [5, Proposition 2.10]. If $\alpha \in R_- = -R_+$, then $-\alpha \in R_+$ has an expansion with all coefficients nonnegative, and multiplying by -1 gives an expansion of α with all coefficients nonpositive. The descriptions (3.2) and (3.3) follow from [5, Corollary 2.11].

To see that Δ is uniquely determined by Γ_+ , note that Γ_+ determines R_+ by Definition 3.1, and the set of simple roots in R_+ is characterized intrinsically by Definition 3.2. Finally, since Γ_+ is the (unique) interior component of Γ , specifying Γ determines Γ_+ and vice versa. This completes the proof of Lemma 3.3. $\square$

Notation 3.4. With the notation given in Lemma 3.3, denote by

$$\Delta = \{\alpha_1, \dots, \alpha_{N_0}\}$$

the simple system associated with Γ .

Lemma 3.5. Each Weyl chamber Γ admits the decomposition

$$\Gamma = K(\Gamma) \times W^\perp, \quad K(\Gamma) := \{w \in W : \langle w, \alpha \rangle \geq 0 \text{ for all } \alpha \in \Delta\}.$$

Proof. Write $x = w + z$ with $w \in W$ and $z \in W^\perp$. Since $\alpha_i \in W$, we have $\langle x, \alpha_i \rangle = \langle w, \alpha_i \rangle$ for all $i = 1, \dots, N_0$. This, together with (3.3), implies that $x \in \Gamma$ if and only if $w \in K(\Gamma)$. This implies that $\Gamma = K(\Gamma) \times W^\perp$, and thus completes the proof. $\square$

Corollary 3.6. Let $\Phi_{W^\perp}$ be any linear isomorphism from $W^\perp$ to $\mathbb{R}^{N-N_0}$, and let

$$\begin{aligned}\Phi_W : W &\longrightarrow \mathbb{R}^{N_0}, & \Phi_W(w) &:= (\langle w, \alpha_1 \rangle, \dots, \langle w, \alpha_{N_0} \rangle), \\ \Phi : \mathbb{R}^N = W \oplus W^\perp &\longrightarrow \mathbb{R}^{N_0} \oplus \mathbb{R}^{N-N_0} \cong \mathbb{R}^N, & \Phi(w+z) &:= (\Phi_W(w), \Phi_{W^\perp}(z)).\end{aligned}$$

Then Φ is a linear isomorphism and

$$\Phi(\Gamma) = \mathbb{R}_+^{N_0} \times \mathbb{R}^{N-N_0}.$$

Proof. By Lemma 3.5 we have $\Gamma = K(\Gamma) \times W^\perp$, where

$$K(\Gamma) = \{w \in W : \langle w, \alpha_i \rangle \geq 0, i = 1, \dots, N_0\}.$$

Since Δ is a basis of W , the Gram matrix $G = (\langle \alpha_i, \alpha_j \rangle)_{i,j=1}^{N_0}$ is nonsingular. Hence, $\Phi_W : W \longrightarrow \mathbb{R}^{N_0}$ is a linear isomorphism. Moreover, for $w \in W$ we have $w \in K(\Gamma)$ if and only if $\Phi_W(w) \in \mathbb{R}_+^{N_0}$, so

$$\Phi_W(K(\Gamma)) = \mathbb{R}_+^{N_0}.$$

By construction, Φ is a linear isomorphism (as a direct sum of isomorphisms), and therefore

$$\Phi(\Gamma) = \Phi(K(\Gamma) \times W^\perp) = \Phi_W(K(\Gamma)) \times \Phi_{W^\perp}(W^\perp) = \mathbb{R}_+^{N_0} \times \mathbb{R}^{N-N_0},$$

which proves the desired conclusion. $\square$

Corollary 3.7. Let Γ be an arbitrary Weyl chamber. There exists a system of dyadic cubes over Γ .

Proof. Let $\Phi : \mathbb{R}^N \rightarrow \mathbb{R}^N$ be the linear isomorphism from Corollary 3.6 so that

$$\Phi(\Gamma) = \Omega := \mathbb{R}_+^{N_0} \times \mathbb{R}^{N-N_0}.$$

For each $k \in \mathbb{Z}$, define $\mathbb{D}_{\Omega,k}^0$ to be the family of half-open cubes

$$\mathcal{Q} = \prod_{i=1}^N I_i, \quad I_i = [m_i 2^{-k}, (m_i + 1) 2^{-k}),$$

where $m_i \in \mathbb{Z}_+$ for $1 \leq i \leq N_0$ and $m_i \in \mathbb{Z}$ for $N_0 < i \leq N$. Set

$$\mathbb{D}_\Omega^0 := \bigcup_{k \in \mathbb{Z}} \mathbb{D}_{\Omega,k}^0.$$

It is standard that $\mathbb{D}_\Omega^0$ is a system of dyadic cubes over Ω .

Next, for each $k \in \mathbb{Z}$, define

$$\mathbb{D}_{\Gamma,k} := \{\Phi^{-1}(\mathcal{Q}) : \mathcal{Q} \in \mathbb{D}_{\Omega,k}^0\}, \quad \mathbb{D}_\Gamma := \bigcup_{k \in \mathbb{Z}} \mathbb{D}_{\Gamma,k}.$$

Since $\Phi : \mathbb{R}^N \rightarrow \mathbb{R}^N$ is a linear isomorphism, we see that $\mathbb{D}_\Gamma$ is a system of dyadic cubes over Γ . $\square$

Notation 3.8. Let Φ be the linear isomorphism given in Corollary 3.6. For every Borel set $E \subset \Omega$, define

$$(3.4) \quad \mu(E) := w(\Phi^{-1}(E)).$$

Lemma 3.9. We have

$$(3.5) \quad d\mu(u, z) = \phi(u) du dz, \quad \phi(u) = |\det(\Phi)|^{-1} \prod_{v \in R_+} \left(\sum_{i=1}^{N_0} a_{v,i} u_i \right)^{2k(v)}.$$

where for each $v \in R_+$, the coefficients $a_{v,i} \geq 0$ are uniquely determined by the simple-root expansion $v = \sum_{i=1}^{N_0} a_{v,i} \alpha_i$ and satisfy $(a_{v,1}, \dots, a_{v,N_0}) \neq \mathbf{0}$.

Proof. Since $\Phi(\Gamma) = \Omega$, we have $\Phi^{-1}(\Omega) = \Gamma$. Hence, for every nonnegative Borel function f on Ω ,

$$\int_{\Omega} f(u, z) d\mu(u, z) = \int_{\Gamma} f(\Phi(x)) dw(x).$$

We make the linear change of variables $(u, z) = \Phi(x)$, so that $x = \Phi^{-1}(u, z)$. Since Φ is linear and invertible, the Jacobian is constant and

$$dx = |\det(\Phi)|^{-1} du dz.$$

It follows from the definition of Φ that $u_i = \langle x, \alpha_i \rangle$ for $i = 1, \dots, N_0$. This implies that

$$(3.6) \quad \langle x, v \rangle = \left\langle x, \sum_{i=1}^{N_0} a_{v,i} \alpha_i \right\rangle = \sum_{i=1}^{N_0} a_{v,i} \langle x, \alpha_i \rangle = \sum_{i=1}^{N_0} a_{v,i} u_i.$$

Substituting these identities into the previous integral, we obtain

$$\begin{aligned} \int_{\Omega} f(u, z) d\mu(u, z) &= \int_{\Gamma} f(\Phi(x)) \left(\prod_{v \in R_+} |\langle x, v \rangle|^{2k(v)} \right) dx \\ &= \int_{\Omega} f(u, z) |\det(\Phi)|^{-1} \prod_{v \in R_+} \left(\sum_{i=1}^{N_0} a_{v,i} u_i \right)^{2k(v)} du dz. \end{aligned}$$

Therefore, (3.5) holds. This completes the proof of Lemma 3.9. $\square$

Lemma 3.10. Let Γ_1 and Γ_2 be two Weyl chambers. Then there exist an integer $M \geq 0$ and $\alpha_1, \dots, \alpha_M \in R$ such that, if we set $\Gamma^{(0)} := \Gamma_2$ and

$$\Gamma^{(k)} := \sigma_{\alpha_k}(\Gamma^{(k-1)}), \quad k = 1, \dots, M,$$

then the following hold:

- (1) $\alpha_k \in \Delta(\Gamma^{(k-1)})$ for each $k = 1, \dots, M$;
- (2) $\Gamma^{(M)} = \Gamma_1$.

Proof. If $\Gamma_1 = \Gamma_2$, we take $M = 0$ and there is nothing to prove. Assume $\Gamma_1 \neq \Gamma_2$. Since the Weyl group G acts transitively on the set of Weyl chambers (see [5, Theorem 2.12 (2)]), there exists $g \in G$ such that $\Gamma_1 = g(\Gamma_2)$.

Let $\Delta(\Gamma_2) = \{\beta_1, \beta_2, \dots, \beta_{N_0}\}$ be the simple system associated with Γ_2 . By [5, Theorem 2.12(4)], the Weyl group G is generated by the simple reflections $\sigma_{\beta_1}, \sigma_{\beta_2}, \dots, \sigma_{\beta_{N_0}}$. Hence, there exist indices $i_1, \dots, i_M \in \{1, \dots, N_0\}$ such that

$$g = \sigma_{\beta_{i_1}} \sigma_{\beta_{i_2}} \cdots \sigma_{\beta_{i_M}}.$$

Define

$$w_0 := \text{Id}, \quad w_k := \sigma_{\beta_{i_1}} \cdots \sigma_{\beta_{i_k}} \quad (k = 1, \dots, M),$$

and set

$$\Gamma^{(k)} := w_k(\Gamma_2) \quad (k = 0, 1, \dots, M).$$

It follows that $\Gamma^{(0)} = \Gamma_2$ and $\Gamma^{(M)} = w_M(\Gamma_2) = g(\Gamma_2) = \Gamma_1$, which implies the statement (2).

For each $k = 1, \dots, M$, define

$$\alpha_k := w_{k-1}(\beta_{i_k}) \in R.$$

The rest of the proof is divided into two steps.

- Step 1. Verify that $\Gamma^{(k)} = \sigma_{\alpha_k}(\Gamma^{(k-1)})$.

Since each $w \in G$ is orthogonal, for any root $\beta \in R$ one has the conjugation identity

$$w \sigma_{\beta} w^{-1} = \sigma_{w(\beta)}, \quad w \in G.$$

Applying this with $w = w_{k-1}$ and $\beta = \beta_{i_k}$ yields

$$\sigma_{\alpha_k} = \sigma_{w_{k-1}(\beta_{i_k})} = w_{k-1} \sigma_{\beta_{i_k}} w_{k-1}^{-1}.$$

Therefore,

$$\sigma_{\alpha_k}(\Gamma^{(k-1)}) = w_{k-1} \sigma_{\beta_{i_k}} w_{k-1}^{-1}(w_{k-1}(\Gamma_2)) = w_{k-1} \sigma_{\beta_{i_k}}(\Gamma_2) = w_k(\Gamma_2) = \Gamma^{(k)}.$$

- Step 2. Verify that $\alpha_k \in \Delta(\Gamma^{(k-1)})$.

Recall that if Γ is a Weyl chamber associated with simple system $\Delta(\Gamma)$, then

$$\Gamma = \{x \in \mathbb{R}^N : \langle x, \gamma \rangle \geq 0 \text{ for all } \gamma \in \Delta(\Gamma)\}.$$

Applying $w_{k-1} \in G$ to the above description for Γ_2 and using $\langle w_{k-1}x, \beta \rangle = \langle x, w_{k-1}^{-1}\beta \rangle$, we obtain

$$\Gamma^{(k-1)} = w_{k-1}(\Gamma_2) = \{x \in \mathbb{R}^N : \langle x, w_{k-1}(\beta_j) \rangle \geq 0 \text{ for } j = 1, \dots, N_0\}.$$

By the uniqueness of the simple system associated with a Weyl chamber (see Lemma 3.3), we conclude that

$$\Delta(\Gamma^{(k-1)}) = \Delta(w_{k-1}(\Gamma_2)) = w_{k-1}(\Delta(\Gamma_2)) = \{w_{k-1}(\beta_1), \dots, w_{k-1}(\beta_{N_0})\}.$$

In particular, $\alpha_k = w_{k-1}(\beta_{i_k}) \in \Delta(\Gamma^{(k-1)})$, proving the first statement (1).

This completes the proof of Lemma 3.10. □

Definition 3.11. We say that $R' \subset \Omega$ is an axis-parallel rectangle if

$$R' := I_1 \times \cdots \times I_N,$$

with $I_i = [I_i^-, I_i^+] \subset [0, \infty)$ for $1 \leq i \leq N_0$ and $I_i = [I_i^-, I_i^+] \subset \mathbb{R}$ for $i > N_0$.

Denote by

$$B_{\Omega}(x_0, r) := \{x \in \Omega : \|x - x_0\| < r\}.$$

The following lemma demonstrates that $(\Omega, \|\cdot\|, d\mu)$ is a doubling metric measure space.

Lemma 3.12. There exists a constant $C \geq 1$ such that for every $x_0 = (u_0, z_0) \in \Omega$ and every $r > 0$,

$$(3.7) \quad C^{-1} r^N \prod_{v \in R_+} (L_{v,0} + r)^{2k(v)} \leq \mu(B_\Omega(x_0, r)) \leq C r^N \prod_{v \in R_+} (L_{v,0} + r)^{2k(v)},$$

where

$$L_v(u) := \sum_{i=1}^{N_0} a_{v,i} u_i, \quad L_{v,0} := L_v(u_0) = \sum_{i=1}^{N_0} a_{v,i} (u_0)_i.$$

In particular, there exists a constant $C_{\text{dbl}, \Omega} \geq 1$ such that for every $x \in \Omega$ and every $r > 0$,

$$\mu(B_\Omega(x, 2r)) \leq C_{\text{dbl}, \Omega} \mu(B_\Omega(x, r)).$$

Proof. Fix $x_0 = (u_0, z_0) \in \Omega$ and $r > 0$. For each $v \in R_+$ define the linear form

$$L_v(u) := a_v \cdot u, \quad a_v := (a_{v,1}, \dots, a_{v,N_0}) \in \mathbb{R}_+^{N_0} \setminus \{0\}.$$

Since R_+ is finite and each $a_v \neq 0$, we have

$$L_* := \max_{v \in R_+} \|a_v\| < \infty, \quad m_* := \min_{v \in R_+} \sum_{i=1}^{N_0} a_{v,i} > 0.$$

Upper bound. Let $x = (u, z) \in B_\Omega(x_0, r)$. Then

$$\|u - u_0\| \leq \|x - x_0\| \leq r.$$

Hence, for every $v \in R_+$,

$$|L_v(u) - L_{v,0}| = |a_v \cdot (u - u_0)| \leq \|a_v\| \cdot \|u - u_0\| \leq L_* r.$$

Since $L_v(u) \geq 0$ on Ω (all coefficients and coordinates are nonnegative), we obtain

$$L_v(u) \leq L_{v,0} + L_* r \leq C_1 (L_{v,0} + r), \quad C_1 := \max\{1, L_*\}.$$

Therefore,

$$\phi(u) = \prod_{v \in R_+} L_v(u)^{2k(v)} \leq C_1^{2 \sum_{v \in R_+} k(v)} \prod_{v \in R_+} (L_{v,0} + r)^{2k(v)} =: C_2 \prod_{v \in R_+} (L_{v,0} + r)^{2k(v)}.$$

Integrating over $B_\Omega(x_0, r)$ and using $|B_\Omega(x_0, r)| \leq |B(x_0, r)| = \omega_N r^N$, where ω_N is the volume of the unit ball in $\mathbb{R}^N$, gives

$$\mu(B_\Omega(x_0, r)) = \int_{B_\Omega(x_0, r)} \phi(u) dx \leq C_2 |B(x_0, r)| \prod_{v \in R_+} (L_{v,0} + r)^{2k(v)} \leq C_3 r^N \prod_{v \in R_+} (L_{v,0} + r)^{2k(v)},$$

with $C_3 := C_2 \omega_N$.

Lower bound. We construct a smaller Euclidean ball B_c contained in $B_\Omega(x_0, r)$ on which ϕ admits a uniform positive lower bound in terms of $\prod_{v \in R_+} (L_{v,0} + r)^{2k(v)}$. To this end, we first choose

$$\lambda := 1 + \frac{L_*}{m_*} > 1, \quad \eta := \frac{1}{4(\sqrt{N_0} \lambda + 1)}, \quad \theta := \lambda \eta.$$

These parameters depend only on L_* and m_* , and satisfy

$$\theta > \eta > 0, \quad \sqrt{N_0} \theta + \eta = \eta(\sqrt{N_0} \lambda + 1) = \frac{1}{4} < 1.$$

Let $\mathbf{1}_{N_0} := (1, \dots, 1) \in \mathbb{R}^{N_0}$ and define the center

$$x_c := (u_c, z_c) := (u_0 + \theta r \mathbf{1}_{N_0}, z_0), \quad B_c := B(x_c, \eta r).$$

We claim that $B_c \subset B_\Omega(x_0, r)$. To see this inclusion, we first apply the triangle inequality to see that for any $x \in B_c$,

$$\|x - x_0\| \leq \|x - x_c\| + \|x_c - x_0\| \leq \eta r + \theta r \sqrt{N_0} \leq r,$$

since $\sqrt{N_0}\theta + \eta < 1$. Hence $B_c \subset B(x_0, r)$. Moreover, for any $x = (u, z) \in B_c$, we have $\|u - u_c\| \leq \|x - x_c\| \leq \eta r$. This implies that for each $1 \leq i \leq N_0$,

$$u_i \geq (u_c)_i - \eta r = (u_0)_i + (\theta - \eta)r \geq 0,$$

since $u_0 \in \mathbb{R}_+^{N_0}$ and $\theta \geq \eta$. Thus $u \in \mathbb{R}_+^{N_0}$ and $x \in \Omega$. Therefore, $B_c \subset B_\Omega(x_0, r)$. This finishes the proof of the claim.

Next we obtain a uniform lower bound for $L_v(u)$ on B_c . Write any $x = (u, z) \in B_c$ as

$$u = u_0 + \theta r \mathbf{1}_{N_0} + \Delta_u, \quad z = z_0 + \Delta_z, \quad \|\Delta_u\|^2 + \|\Delta_z\|^2 \leq (\eta r)^2,$$

so in particular $\|\Delta_u\| \leq \eta r$. Then for all $x = (u, z) \in B_c$ and all $v \in R_+$,

$$\begin{aligned} L_v(u) &= a_v \cdot u = a_v \cdot u_0 + \theta r a_v \cdot \mathbf{1}_{N_0} + a_v \cdot \Delta_u \\ &\geq L_{v,0} + \theta r \sum_{i=1}^{N_0} a_{v,i} - \|a_v\| \|\Delta_u\| \geq L_{v,0} + r K_0 \geq c_0(L_{v,0} + r). \end{aligned}$$

with

$$K_0 := \theta m_* - \eta L_* = \eta(\lambda m_* - L_*) = \eta m_* > 0, \quad c_0 := \min\{1, K_0\} \in (0, 1].$$

Consequently, for all $u \in B_c$,

$$\phi(u) = \prod_{v \in R_+} L_v(u)^{2k(v)} \geq c_0^{2 \sum_{v \in R_+} k(v)} \prod_{v \in R_+} (L_{v,0} + r)^{2k(v)} =: C_4 \prod_{v \in R_+} (L_{v,0} + r)^{2k(v)}.$$

This, in combination with the fact that $B_c \subset B_\Omega(x_0, r)$, yields

$$\mu(B_\Omega(x_0, r)) \geq \mu(B_c) = \int_{B_c} \phi(u) dx \geq C_4 |B_c| \prod_{v \in R_+} (L_{v,0} + r)^{2k(v)} = C_5 r^N \prod_{v \in R_+} (L_{v,0} + r)^{2k(v)},$$

where $C_5 := C_4 \omega_N \eta^N > 0$.

The second statement follows immediately from the first one. Indeed, by the first statement,

$$\mu(B_\Omega(x_0, 2r)) \leq C_3 (2r)^N \prod_{v \in R_+} (L_{v,0} + 2r)^{2k(v)} \leq C_3 2^{N+\mathcal{K}} r^N \prod_{v \in R_+} (L_{v,0} + r)^{2k(v)} \leq C_3 C_5^{-1} 2^{N+\mathcal{K}} \mu(B_\Omega(x_0, r)),$$

where we set $\mathcal{K} := 2 \sum_{v \in R_+} k(v)$. Therefore, $(\Omega, \|\cdot\|, \mu)$ is doubling with the doubling constant $C_{\text{dbl}, \Omega} := C_3 C_5^{-1} 2^{N+\mathcal{K}}$. This finishes the proof of Lemma 3.12. $\square$

Let $\Phi : \mathbb{R}^N \rightarrow \mathbb{R}^N$ be the linear isomorphism given in Corollary 3.6 such that $\Phi(\Gamma) = \Omega$. Since Φ is linear and invertible, there exist positive constants c_Φ and C_Φ such that for any $x, y \in \mathbb{R}^N$,

$$(3.8) \quad c_\Phi \|x - y\| \leq \|\Phi(x) - \Phi(y)\| \leq C_\Phi \|x - y\|.$$

Denote by $B_\Gamma(x, r) := \{y \in \Gamma : \|y - x\| < r\}$. It follows from (3.8) that for any $x_0 \in \Gamma$ and $r > 0$,

$$(3.9) \quad B_\Omega(\Phi(x_0), c_\Phi r) \subset \Phi(B_\Gamma(x_0, r)) \subset B_\Omega(\Phi(x_0), C_\Phi r).$$

This, together with Lemma 3.12, yields

$$(3.10) \quad \mu(\Phi(B_\Gamma(x_0, r))) \simeq \mu(B_\Omega(\Phi(x_0), C_\Phi r)).$$

Lemma 3.13. For all $x_0 \in \Gamma$ and $r > 0$, one can find an axis-parallel rectangle R' on Ω such that

$$(3.11) \quad B_\Omega(\Phi(x_0), C_\Phi r) \subset R' \subset B_\Omega(\Phi(x_0), \sqrt{N}C_\Phi r).$$

Furthermore, we have

$$(3.12) \quad \mu(R') \simeq \mu(B_\Omega(\Phi(x_0), C_\Phi r)).$$

Proof. We write $(u_0, z_0) := \Phi(x_0) \in \Omega$. and define the axis-parallel rectangle

$$(3.13) \quad R' := \left(\prod_{i=1}^{N_0} [\max\{0, (u_0)_i - C_\Phi r\}, (u_0)_i + C_\Phi r] \right) \times \left(\prod_{j=1}^{N-N_0} [(z_0)_j - C_\Phi r, (z_0)_j + C_\Phi r] \right).$$

On the one hand, it follows immediately that $B_\Omega(\Phi(x_0), C_\Phi r) \subset R'$. On the other hand, since R' is a rectangle with half-side lengths bounded by $C_\Phi r$ in each coordinate, we have

$$R' \subset B_\Omega(\Phi(x_0), \sqrt{N}C_\Phi r).$$

The second statement follows from the first one together with Lemma 3.12. The proof of Lemma 3.13 is finished. $\square$

The following theorem demonstrates that $(\Gamma, \|\cdot\|, dw)$ is a doubling metric measure space.

Theorem 3.14. There exists a constant $C \geq 1$ such that for every $x \in \Gamma$ and every $r > 0$,

$$(3.14) \quad C^{-1} r^N \prod_{v \in R_+} (|\langle x, v \rangle| + r)^{2k(v)} \leq w(B_\Gamma(x, r)) \leq C r^N \prod_{v \in R_+} (|\langle x, v \rangle| + r)^{2k(v)}.$$

In particular, there exists a constant $C_{\text{dbl}, \Gamma} \geq 1$ such that for every $x \in \Gamma$ and every $r > 0$,

$$w(B_\Gamma(x, 2r)) \leq C_{\text{dbl}, \Gamma} w(B_\Gamma(x, r)).$$

Moreover, for all $x \in \Gamma$ and $r > 0$, we have

$$(3.15) \quad w(B_\Gamma(x, r)) \sim w(B(x, r)).$$

Proof. The main strategy of the proof is to transfer the ball estimates from $(\Omega, \|\cdot\|, \mu)$ back to $(\Gamma, \|\cdot\|, dw)$. Fix $x \in \Gamma$ and $r > 0$. It follows from the equality (3.4) that

$$(3.16) \quad w(B_\Gamma(x, r)) = \mu(\Phi(B_\Gamma(x, r))).$$

Applying μ to the inclusions (3.9) and using (3.16) yields

$$(3.17) \quad \mu(B_\Omega(\Phi(x), c_\Phi r)) \leq w(B_\Gamma(x, r)) \leq \mu(B_\Omega(\Phi(x), C_\Phi r)).$$

Write $\Phi(x) = (u, z) \in \Omega$. For each $v \in R_+$ recall the linear form $L_{v,0} := \sum_{i=1}^{N_0} a_{v,i} u_i$. Combining the inequality (3.17) with Lemma 3.12 and the equality (3.6), we obtain that

$$(3.18) \quad w(B_\Gamma(x, r)) \simeq r^N \prod_{v \in R_+} (L_{v,0} + r)^{2k(v)} = r^N \prod_{v \in R_+} (|\langle x, v \rangle| + r)^{2k(v)}.$$

This proves (3.14). The second statement follows immediately from (3.14). The third statement follows from the inequalities (3.14) and (2.1). This completes the proof of Theorem 3.14. $\square$

4. Proof of upper bound

Lemma 4.1. Assume that $N > 1$. For $\sigma, \tau \in G$, define

$$(4.1) \quad \mathcal{G}_{\sigma,\tau} f(z) := \begin{cases} \frac{|f(\tau z) - f(\sigma z)|}{|\tau z - \sigma z|}, & \tau z \neq \sigma z, \\ 0, & \tau z = \sigma z. \end{cases}$$

Then for every $f \in \dot{W}^{1,N}(\mathbb{R}^N)$,

$$(4.2) \quad \|\mathcal{G}_{\sigma,\tau} f\|_{L^N(\mathbb{R}^N, dx)} \lesssim_{N,G} \|\nabla f\|_{L^N(\mathbb{R}^N, dx)}.$$

The implicit constant is independent of $\sigma, \tau \in G$.

Proof. Let $\mathcal{M}$ be the Hardy–Littlewood maximal operator on $\mathbb{R}^N$. By [?, Theorem 3], after choosing the precise representative of f , there exists a null set $E \subset \mathbb{R}^N$ such that

$$|f(x) - f(y)| \lesssim_N |x - y|(\mathcal{M}(|\nabla f|)(x) + \mathcal{M}(|\nabla f|)(y)), \quad x, y \in \mathbb{R}^N \setminus E.$$

Since σ and τ are orthogonal, they preserve Lebesgue measure. Hence, for almost every $z \in \mathbb{R}^N$ with $\tau z \neq \sigma z$,

$$\mathcal{G}_{\sigma,\tau} f(z) \lesssim_N \mathcal{M}(|\nabla f|)(\tau z) + \mathcal{M}(|\nabla f|)(\sigma z).$$

The same estimate holds when $\tau z = \sigma z$ by the definition of $\mathcal{G}_{\sigma,\tau} f$. Therefore, using again the measure-preserving property of σ and τ and the L^N -boundedness of $\mathcal{M}$, which is valid since $N > 1$,

$$\begin{aligned} \|\mathcal{G}_{\sigma,\tau} f\|_{L^N(\mathbb{R}^N)} &\lesssim_N \|\mathcal{M}(|\nabla f|) \circ \tau\|_{L^N} + \|\mathcal{M}(|\nabla f|) \circ \sigma\|_{L^N} \\ &= 2\|\mathcal{M}(|\nabla f|)\|_{L^N} \lesssim_N \|\nabla f\|_{L^N(\mathbb{R}^N)}. \end{aligned}$$

The implicit constant is independent of $\sigma, \tau \in G$. $\square$

Lemma 4.2. Assume that $N \geq 1$, $1 < p < \infty$, and $1 \leq q \leq \infty$. For $\sigma, \tau \in G$ and $I \in \mathbb{D}_\Gamma$, let $\theta_{\sigma,\tau,I}$ be as in (??). For $f \in L^1_{\text{loc}}(\mathbb{R}^N)$, define

$$(4.3) \quad a_{\sigma,\tau,I}(f) := \frac{1}{|I|} \int_I (f(\tau z) - f(\sigma z)) dz.$$

We have

$$(4.4) \quad \left\| \{\theta_{\sigma,\tau,I} a_{\sigma,\tau,I}(f)\}_{I \in \mathbb{D}_\Gamma} \right\|_{\ell^{N,\infty}} \lesssim \|f\|_{W^{1,N}(\Gamma)}.$$

Proof. By part 1 of Lemma 8.2,

$$\|\nabla f\|_{L^N(\mathbb{R}^N)} \simeq \|f\|_{OSC^{N,\infty}(\mathbb{R}^N)}.$$

For $I = \Phi^{-1}(Q) \in \mathbb{D}_\Gamma$ and $z \in I$, we use the bi-Lipschitz property (3.8) to see that

$$|z - z_I| \lesssim |\Phi(z) - \Phi(z_I)| \leq \frac{\sqrt{N}}{2} \ell(Q) \lesssim \ell(I).$$

Since σ and τ are orthogonal,

$$|\tau z - \sigma z| \leq 2|z - z_I| + |\tau z_I - \sigma z_I| \lesssim \ell(I) + |\tau z_I - \sigma z_I|.$$

Hence, by the definition of $\mathcal{G}_{\sigma,\tau}f$ (see (4.1)) and (??),

$$\begin{aligned} \theta_{\sigma,\tau,I} |a_{\sigma,\tau,I}(f)| &\leq \frac{\theta_{\sigma,\tau,I}}{|I|} \int_I |f(\tau z) - f(\sigma z)| dz \\ &\lesssim \frac{\ell(I)}{\ell(I) + |\sigma z_I - \tau z_I|} (\ell(I) + |\tau z_I - \sigma z_I|) \frac{1}{|I|} \int_I \mathcal{G}_{\sigma,\tau}f(z) dz \\ &\lesssim \ell(I) \frac{1}{|I|} \int_I \mathcal{G}_{\sigma,\tau}f(z) dz. \end{aligned}$$

Set

$$F_{\sigma,\tau}(u) := \mathcal{G}_{\sigma,\tau}f(\Phi^{-1}u) \chi_\Omega(u).$$

By the change of variables $u = \Phi(z)$, together with Lemma 4.1,

$$\|F_{\sigma,\tau}\|_{L^N(\mathbb{R}^N)} \lesssim \|\mathcal{G}_{\sigma,\tau}f\|_{L^N(\mathbb{R}^N)} \lesssim \|\nabla f\|_{L^N(\mathbb{R}^N)},$$

uniformly in $\sigma, \tau \in G$. Moreover, for $I = \Phi^{-1}(Q)$,

$$\ell(I) \frac{1}{|I|} \int_I \mathcal{G}_{\sigma,\tau}f(z) dz = \ell(Q) \frac{1}{|Q|} \int_Q F_{\sigma,\tau}(u) du = |Q|^{\frac{1}{N}-1} \int_Q F_{\sigma,\tau}(u) du.$$

By the construction of $\mathbb{D}_\Gamma$, every $I \in \mathbb{D}_\Gamma$ is of the form $I = \Phi^{-1}(Q)$ for a unique $Q \in \mathbb{D}_\Omega^0 \subset \mathbb{D}^0$. Applying Lemma 8.1 with $s = N$, we deduce that

$$\left\| \left\{ \ell(I) \frac{1}{|I|} \int_I \mathcal{G}_{\sigma,\tau}f(z) dz \right\}_{I \in \mathbb{D}_\Gamma} \right\|_{\ell^{N,\infty}} \leq \left\| \left\{ |Q|^{\frac{1}{N}-1} \int_Q F_{\sigma,\tau}(u) du \right\}_{Q \in \mathbb{D}^0} \right\|_{\ell^{N,\infty}} \lesssim \|F_{\sigma,\tau}\|_{L^N(\mathbb{R}^N)}.$$

Combining the preceding estimates, we deduce that

$$\left\| \{\theta_{\sigma,\tau,I} a_{\sigma,\tau,I}(f)\}_{I \in \mathbb{D}_\Gamma} \right\|_{\ell^{N,\infty}} \lesssim \|\nabla f\|_{L^N(\mathbb{R}^N)} \simeq \|f\|_{OSC^{N,\infty}(\mathbb{R}^N)},$$

where the final equivalence follows from Lemma 8.2, and where the implicit constant is uniform in $\sigma, \tau \in G$. $\square$

Lemma 4.3. Assume that $N \geq 1$, $1 < p < \infty$, and $1 \leq q \leq \infty$. Fix $s > 2$ and a closed Weyl chamber Γ . Let r_s be the exponent in Lemma ???. For $f \in L^1_{\text{loc}}(\mathbb{R}^N)$ and $\sigma, \tau \in G$, define

$$(4.5) \quad H_{\sigma,\tau}(f)(z) := f(\tau z) - f(\sigma z), \quad z \in \mathbb{R}^N.$$

Then for any $f \in OSC^{p,q}(\mathbb{R}^N)$,

$$\left\| \{\text{osc}_{r_s}(H_{\sigma,\tau}(f); I)\}_{I \in \mathbb{D}_\Gamma} \right\|_{\ell^{p,q}} \lesssim \|f\|_{OSC^{p,q}(\mathbb{R}^N)},$$

The implicit constant is independent of $\sigma, \tau \in G$.

Proof. Since σ and τ are orthogonal,

$$(H_{\sigma,\tau}(f))_{I,dx} = (f)_{\tau I,dx} - (f)_{\sigma I,dx},$$

and hence, by Minkowski's inequality,

$$\text{osc}_{r_s}(H_{\sigma,\tau}(f); I) \leq \text{osc}_{r_s}(f; \tau I) + \text{osc}_{r_s}(f; \sigma I).$$

Fix $\gamma \in \{\sigma, \tau\}$. By the construction of $\mathbb{D}_\Gamma$ and (3.8),

$$\gamma I \subset B(\gamma z_I, C\ell(I)), \quad |\gamma I| = |I| \simeq \ell(I)^N.$$

By Lemma 2.8, for every $I \in \mathbb{D}_\Gamma$ there exist $\nu_\gamma(I) \in \{1, \dots, 2^N\}$ and $J_I^\gamma \in \mathbb{D}^{\nu_\gamma(I)}$ such that

$$\gamma I \subset B(\gamma z_I, C\ell(I)) \subset J_I^\gamma \subset C_N B(\gamma z_I, C\ell(I)).$$

Thus, $\ell(J_I^\gamma) \simeq \ell(I)$, and $|J_I^\gamma| \simeq |\gamma I|$. This implies that

$$\text{osc}_{r_s}(f; \gamma I) \lesssim \text{osc}_{r_s}(f; J_I^\gamma).$$

Note that for each $\nu \in \{1, \dots, 2^N\}$, the map

$$I \mapsto J_I^\gamma, \quad \{I \in \mathbb{D}_\Gamma : \nu_\gamma(I) = \nu\} \longrightarrow \mathbb{D}^\nu,$$

has uniformly bounded multiplicity. That is,

$$\sup_{J \in \mathbb{D}^\nu} \# \{I \in \mathbb{D}_\Gamma : \nu_\gamma(I) = \nu, J_I^\gamma = J\} \lesssim 1.$$

Therefore, by Lemma ??,

$$\begin{aligned} \left\| \{\text{osc}_{r_s}(f; \gamma I)\}_{I \in \mathbb{D}_\Gamma} \right\|_{\ell^{p,q}} &\lesssim \sum_{\nu=1}^{2^N} \left\| \{\text{osc}_{r_s}(f; J_I^\gamma)\}_{I \in \mathbb{D}_\Gamma, \nu_\gamma(I)=\nu} \right\|_{\ell^{p,q}} \\ &\lesssim \sum_{\nu=1}^{2^N} \left\| \{\text{osc}_{r_s}(f; J)\}_{J \in \mathbb{D}^\nu} \right\|_{\ell^{p,q}} = \|f\|_{\text{osc}^{p,q}(\mathbb{R}^N)}. \end{aligned}$$

Applying this estimate with $\gamma = \sigma$ and $\gamma = \tau$, we obtain

$$\left\| \{\text{osc}_{r_s}(H_{\sigma,\tau}(f); I)\}_{I \in \mathbb{D}_\Gamma} \right\|_{\ell^{p,q}} \lesssim \|f\|_{\text{osc}^{p,q}(\mathbb{R}^N)}.$$

This completes the proof of Lemma 4.3. $\square$

5. Proof of lower bound

Definition 5.1. Let $1 \leq r < \infty$. For a measurable set $E \subset \mathbb{R}^N$ with $0 < w(E) < \infty$, set

$$(f)_{E,dw} := \frac{1}{w(E)} \int_E f dw$$

and

$$\text{osc}_{r,dw}(f; E) := \left(\frac{1}{w(E)} \int_E |f - (f)_{E,dw}|^r dw \right)^{1/r}.$$

For $0 < p < \infty$ and $0 < q \leq \infty$, define

$$\|f\|_{\text{osc}_{dw}^{p,q}} := \sum_{\nu=1}^{2^N} \left\| \{\text{osc}_{1,dw}(f; Q)\}_{Q \in \mathbb{D}^\nu} \right\|_{\ell^{p,q}}.$$

Lemma 5.2. Let $0 < p < \infty$, $0 < q \leq \infty$, and $1 \leq r < \infty$. For $f \in L_{\text{loc}}^r(\mathbb{R}^N, dx)$, set

$$\text{osc}_{r,dw}(f; Q) := \left(\frac{1}{w(Q)} \int_Q |f - (f)_{Q,dw}|^r dw \right)^{1/r}$$

and

$$\|f\|_{\text{osc}_{dw}^{p,q}} := \sum_{\nu=1}^{2^N} \left\| \{\text{osc}_{r,dw}(f; Q)\}_{Q \in \mathbb{D}^\nu} \right\|_{\ell^{p,q}}.$$

Then

$$\|f\|_{\text{osc}^{p,q}(\mathbb{R}^N)} \sim \|f\|_{\text{osc}_{dw}^{p,q}},$$

where the implicit constants are independent of f .

Proof. By Lemma 2.1,

$$dw = \varpi dx, \quad \varpi \in A_\infty(\mathbb{R}^N, dx).$$

Moreover, both dx and dw are doubling measures. Hence dx and dw satisfy the A_∞ -relation in the sense of [?, Proposition 1.2]. Therefore,

$$\|f\|_{\text{osc}^{p,q}(dx)} \sim \|f\|_{\text{osc}^{p,q}(dw)}$$

by [?, Proposition 1.2].

By [?, (3.1) and Lemma 3.4], together with Lemma 2.8, these two oscillatory norms are equivalent, respectively, to the dyadic norms

$$\|f\|_{\text{osc}^{p,q}(\mathbb{R}^N)} \quad \text{and} \quad \|f\|_{\text{osc}_{dw}^{p,q}}.$$

The assertion follows. □

Lemma 5.3. Fix $j \in \{1, \dots, N\}$. For every $Q \in \bigcup_{\nu=1}^{2^N} \mathbb{D}^\nu$, let

$$r_Q := \sqrt{N} \ell(Q), \quad B_Q := B(c_Q, r_Q),$$

and define

$$\widehat{B}_Q := B(c_Q + 5r_Q e_j, r_Q).$$

Then

$$(5.1) \quad w(Q) \sim w(B_Q) \sim w(\widehat{B}_Q),$$

and there is a constant C_N , independent of Q , such that

$$Q \cup \widehat{B}_Q \subset C_N Q.$$

Moreover, for every $(x, y) \in Q \times \widehat{B}_Q$, the kernel $K_{\text{Dunkl},j}(x, y)$ does not change sign and

$$(5.2) \quad |K_{\text{Dunkl},j}(x, y)| \gtrsim \frac{1}{w(Q)}.$$

Proof. Since Q is a Euclidean cube,

$$B\left(c_Q, \frac{\ell(Q)}{2}\right) \subset Q \subset B_Q.$$

Hence the doubling property (2.2) gives

$$w(B_Q) \lesssim w\left(B\left(c_Q, \frac{\ell(Q)}{2}\right)\right) \leq w(Q) \leq w(B_Q),$$

and therefore

$$w(Q) \sim w(B_Q).$$

Since the distance between the centers of B_Q and $\widehat{B}_Q$ is $5r_Q$,

$$\widehat{B}_Q \subset 6B_Q.$$

Similarly,

$$B_Q \subset B(c_Q + 5r_Q e_j, 6r_Q) = 6\widehat{B}_Q.$$

The doubling property therefore yields

$$w(B_Q) \sim w(\widehat{B}_Q),$$

which proves (5.1).

For $y \in \widehat{B}_Q$,

$$\|y - c_Q\| \leq 6r_Q = 6\sqrt{N}\ell(Q).$$

Thus $Q \cup \widehat{B}_Q$ is contained in a fixed concentric enlargement $C_N Q$.

Finally, $Q \subset B_Q$, so Lemma 2.4, applied to B_Q , shows that $K_{\text{Dunkl},j}$ has a fixed sign on $Q \times \widehat{B}_Q$ and

$$|K_{\text{Dunkl},j}(x, y)| \gtrsim \frac{1}{w(B_Q)} \sim \frac{1}{w(Q)}.$$

□

Lemma 5.4. Let $f \in L^1_{\text{loc}}(\mathbb{R}^N, dw)$ be real-valued, and assume that $[R_{\text{Dunkl},j}, M_f]$ is a compact operator on $L^2(\mathbb{R}^N, dw)$. For every $\nu \in \{1, \dots, 2^N\}$, there exist NWO sequences

$$\{e_{Q,m}\}_{Q \in \mathbb{D}^\nu}, \quad \{h_{Q,m}\}_{Q \in \mathbb{D}^\nu}, \quad m = 1, 2,$$

such that, for every $Q \in \mathbb{D}^\nu$,

$$\text{osc}_{1,dw}(f; Q) \lesssim \sum_{m=1}^2 \left| \langle [R_{\text{Dunkl},j}, M_f] e_{Q,m}, h_{Q,m} \rangle_{L^2(\mathbb{R}^N, dw)} \right|.$$

Proof. Fix $\nu \in \{1, \dots, 2^N\}$ and $Q \in \mathbb{D}^\nu$. Set

$$r_Q := \sqrt{N}\ell(Q), \quad B_Q := B(c_Q, r_Q).$$

Since

$$B(c_Q, \ell(Q)/2) \subset Q \subset B_Q,$$

by (2.2),

$$w(B_Q) \simeq w(Q).$$

By Lemma 2.4, for

$$\widehat{B}_Q := B(c_Q + 5r_Q e_j, r_Q),$$

the kernel $K_{\text{Dunkl},j}$ has a fixed sign on $B_Q \times \widehat{B}_Q$ and

$$|K_{\text{Dunkl},j}(x, y)| \gtrsim \frac{1}{w(Q)}, \quad (x, y) \in Q \times \widehat{B}_Q.$$

Moreover,

$$\widehat{B}_Q \subset 6B_Q, \quad B_Q \subset 6\widehat{B}_Q,$$

so (2.2) implies

$$w(\widehat{B}_Q) \simeq w(Q).$$

Since $\widehat{B}_Q \subset B(c_Q, 6r_Q)$, there is a fixed $C_N \geq 1$ such that

$$\widehat{B}_Q \subset C_N Q, \quad w(C_N Q) \simeq w(Q).$$

Let α_Q be a dw -median of f on $\widehat{B}_Q$, and define

$$\begin{aligned} E_{Q,1} &:= \{x \in Q : f(x) < \alpha_Q\}, & F_{Q,1} &:= \{y \in \widehat{B}_Q : f(y) \geq \alpha_Q\}, \\ E_{Q,2} &:= \{x \in Q : f(x) \geq \alpha_Q\}, & F_{Q,2} &:= \{y \in \widehat{B}_Q : f(y) \leq \alpha_Q\}. \end{aligned}$$

Then

$$w(F_{Q,m}) \geq \frac{1}{2}w(\widehat{B}_Q) \simeq w(Q), \quad m = 1, 2.$$

For each $m = 1, 2$, $f(y) - f(x)$ has a fixed sign on $E_{Q,m} \times F_{Q,m}$, and

$$|f(x) - \alpha_Q| \leq |f(y) - f(x)|, \quad (x, y) \in E_{Q,m} \times F_{Q,m}.$$

Since $K_{\text{Dunkl},j}$ also has a fixed sign on $Q \times \widehat{B}_Q$, we obtain

$$\begin{aligned} \frac{1}{w(Q)} \int_{E_{Q,m}} |f - \alpha_Q| dw &\leq \frac{1}{w(Q)w(F_{Q,m})} \int_{E_{Q,m}} \int_{F_{Q,m}} |f(y) - f(x)| dw(y) dw(x) \\ &\lesssim \frac{1}{w(Q)} \left| \int_{E_{Q,m}} \int_{F_{Q,m}} (f(y) - f(x)) K_{\text{Dunkl},j}(x, y) dw(y) dw(x) \right|. \end{aligned}$$

Define

$$e_{Q,m} := w(C_N Q)^{-1/2} \chi_{F_{Q,m}}, \quad h_{Q,m} := w(Q)^{-1/2} \chi_{E_{Q,m}}.$$

By the off-diagonal kernel identity

$$K_{[R_{\text{Dunkl},j}, M_f]}(x, y) = (f(y) - f(x)) K_{\text{Dunkl},j}(x, y),$$

and $w(C_N Q) \simeq w(Q)$,

$$\frac{1}{w(Q)} \int_{E_{Q,m}} |f - \alpha_Q| dw \lesssim \left| \langle [R_{\text{Dunkl},j}, M_f] e_{Q,m}, h_{Q,m} \rangle_{L^2(\mathbb{R}^N, dw)} \right|.$$

Furthermore,

$$\text{supp } h_{Q,m} \subset Q, \quad \text{supp } e_{Q,m} \subset C_N Q.$$

For any fixed $s > 2$,

$$\begin{aligned} \|h_{Q,m}\|_{L^s(\mathbb{R}^N, dw)} &\leq w(Q)^{1/s-1/2}, \\ \|e_{Q,m}\|_{L^s(\mathbb{R}^N, dw)} &\leq w(C_N Q)^{1/s-1/2} \leq w(Q)^{1/s-1/2}. \end{aligned}$$

Hence, $\{e_{Q,m}\}_{Q \in \mathbb{D}^N}$ and $\{h_{Q,m}\}_{Q \in \mathbb{D}^N}$ are NWO sequences for $m = 1, 2$.

Finally,

$$|(f)_{Q,dw} - \alpha_Q| \leq \frac{1}{w(Q)} \int_Q |f - \alpha_Q| dw,$$

and therefore

$$\begin{aligned} \text{osc}_{1,dw}(f; Q) &\leq \frac{2}{w(Q)} \int_Q |f - \alpha_Q| dw \\ &= 2 \sum_{m=1}^2 \frac{1}{w(Q)} \int_{E_{Q,m}} |f - \alpha_Q| dw \\ &\lesssim \sum_{m=1}^2 \left| \langle [R_{\text{Dunkl},j}, M_f] e_{Q,m}, h_{Q,m} \rangle_{L^2(\mathbb{R}^N, dw)} \right|. \end{aligned}$$

This completes the proof of Lemma 5.4. $\square$

Theorem 5.5. Assume that $N \geq 1$, $1 < p < \infty$, and $1 \leq q \leq \infty$. Fix $j \in \{1, \dots, N\}$. Let

$$f \in L^1_{\text{loc}}(\mathbb{R}^N, dx).$$

If

$$[R_{\text{Dunkl},j}, M_f] \in S^{p,q}(L^2(\mathbb{R}^N, dw)),$$

then $f \in OSC^{p,q}(\mathbb{R}^N)$ and

$$(5.3) \quad \|f\|_{OSC^{p,q}(\mathbb{R}^N)} \lesssim \|[R_{\text{Dunkl},j}, M_f]\|_{S^{p,q}(L^2(\mathbb{R}^N, dw))}.$$

Proof. Assume first that f is real valued. Fix $\nu \in \{1, \dots, 2^N\}$. By Lemma 5.4, for $m = 1, 2$ there are NWO sequences

$$\{e_{Q,m}\}_{Q \in \mathbb{D}^\nu}, \quad \{h_{Q,m}\}_{Q \in \mathbb{D}^\nu},$$

such that

$$\text{osc}_{1,dw}(f; Q) \lesssim \sum_{m=1}^2 \left| \langle [R_{\text{Dunkl},j}, M_f] e_{Q,m}, h_{Q,m} \rangle_{L^2(\mathbb{R}^N, dw)} \right|.$$

By the quasi triangle inequality in $\ell^{p,q}$ and Lemma 2.9,

$$\begin{aligned} \left\| \left\{ \text{osc}_{1,dw}(f; Q) \right\}_{Q \in \mathbb{D}^\nu} \right\|_{\ell^{p,q}} &\lesssim \sum_{m=1}^2 \left\| \left\{ \langle [R_{\text{Dunkl},j}, M_f] e_{Q,m}, h_{Q,m} \rangle_{L^2(\mathbb{R}^N, dw)} \right\}_{Q \in \mathbb{D}^\nu} \right\|_{\ell^{p,q}} \\ &\lesssim \|[R_{\text{Dunkl},j}, M_f]\|_{S^{p,q}(L^2(\mathbb{R}^N, dw))}. \end{aligned}$$

Summing over ν and using Definition 5.1, we obtain

$$\|f\|_{OSC^{p,q}_{dw}} \lesssim \|[R_{\text{Dunkl},j}, M_f]\|_{S^{p,q}(L^2(\mathbb{R}^N, dw))}.$$

By Lemma 5.2 with $r = 1$,

$$(5.4) \quad \|f\|_{OSC^{p,q}(\mathbb{R}^N)} \lesssim \|[R_{\text{Dunkl},j}, M_f]\|_{S^{p,q}(L^2(\mathbb{R}^N, dw))}.$$

For general f , let C denote complex conjugation on $L^2(\mathbb{R}^N, dw)$. Since $K_{\text{Dunkl},j}$ is real valued by its heat kernel representation,

$$CR_{\text{Dunkl},j}C = R_{\text{Dunkl},j}, \quad C[R_{\text{Dunkl},j}, M_f]C = [R_{\text{Dunkl},j}, M_{\bar{f}}].$$

Hence,

$$\begin{aligned} [R_{\text{Dunkl},j}, M_{\text{Re } f}] &= \frac{1}{2}([R_{\text{Dunkl},j}, M_f] + C[R_{\text{Dunkl},j}, M_f]C), \\ [R_{\text{Dunkl},j}, M_{\text{Im } f}] &= \frac{1}{2i}([R_{\text{Dunkl},j}, M_f] - C[R_{\text{Dunkl},j}, M_f]C). \end{aligned}$$

Since conjugation preserves singular values,

$$\|[R_{\text{Dunkl},j}, M_{\text{Re } f}]\|_{S^{p,q}} + \|[R_{\text{Dunkl},j}, M_{\text{Im } f}]\|_{S^{p,q}} \lesssim \|[R_{\text{Dunkl},j}, M_f]\|_{S^{p,q}}.$$

By (5.4) applied to $\text{Re } f$ and $\text{Im } f$,

$$\|\text{Re } f\|_{OS^{p,q}(\mathbb{R}^N)} + \|\text{Im } f\|_{OS^{p,q}(\mathbb{R}^N)} \lesssim \|[R_{\text{Dunkl},j}, M_f]\|_{S^{p,q}(L^2(\mathbb{R}^N, dw))}.$$

Since

$$\text{osc}_1(f; Q) \leq \text{osc}_1(\text{Re } f; Q) + \text{osc}_1(\text{Im } f; Q),$$

the quasi triangle inequality in $\ell^{p,q}$ proves (5.3). This completes the proof of Theorem 5.5. $\square$

6. Spectral asymptotic formula

Definition 6.1. For $g \in G$ and $j = 1, \dots, N$, set

$$d_{g,j} := g^{-1}e_j, \quad f_g(x) := f(gx).$$

For $\alpha \in R_+$, define $S_\alpha : \mathbb{C}^G \rightarrow \mathbb{C}^G$ by

$$(S_\alpha z)_g := z_{g\sigma_\alpha}, \quad g \in G.$$

For $f \in \dot{W}^{1,N}(\mathbb{R}^N)$, $x \in \Gamma_+$, and $\theta \in \mathbb{S}^{N-1}$, define

$$\begin{aligned} \mathfrak{c}_{f,j}(x, \theta) &:= \text{diag}_{g \in G} \left(\langle d_{g,j}, \nabla f_g(x) \rangle - \langle d_{g,j}, \theta \rangle \langle \theta, \nabla f_g(x) \rangle \right) \\ &\quad + \sum_{\alpha \in R_+} k(\alpha) \text{diag}_{g \in G} \left(\frac{\langle \alpha, d_{g,j} \rangle}{\langle \alpha, x \rangle} (f_g(x) - f_{g\sigma_\alpha}(x)) \right) S_\alpha. \end{aligned}$$

For smooth f , the quotients in the second term are understood by their smooth extensions across the corresponding reflection hyperplanes. For $f \in \dot{W}^{1,N}(\mathbb{R}^N)$, $\mathfrak{c}_{f,j}$ is understood almost everywhere on $\Gamma \times \mathbb{S}^{N-1}$.

Lemma 6.2. Let $f \in W_{\text{loc}}^{1,1}(\mathbb{R}^N)$. For every $u \in C_c^\infty(\mathbb{R}^N)$,

$$(6.1) \quad [T_{e_j}, M_f]u(x) = \partial_j f(x)u(x) + \sum_{\alpha \in R_+} k(\alpha) \langle \alpha, e_j \rangle \frac{f(x) - f(\sigma_\alpha x)}{\langle \alpha, x \rangle} u(\sigma_\alpha x)$$

for almost every $x \in \mathbb{R}^N$. If $N > 1$ and $f \in \dot{W}^{1,N}(\mathbb{R}^N)$, then, for every $\alpha \in R_+$,

$$(6.2) \quad \left\| \frac{f - f \circ \sigma_\alpha}{\langle \alpha, \cdot \rangle} \right\|_{L^N(\mathbb{R}^N, dx)} \lesssim_{N,G} \|\nabla f\|_{L^N(\mathbb{R}^N, dx)}.$$

Proof. Pairing α and $-\alpha$ in the definition of T_{e_j} , and using $k(-\alpha) = k(\alpha)$ and $\sigma_{-\alpha} = \sigma_\alpha$, we have

$$T_{e_j}u(x) = \partial_j u(x) + \sum_{\alpha \in R_+} k(\alpha) \langle \alpha, e_j \rangle \frac{u(x) - u(\sigma_\alpha x)}{\langle \alpha, x \rangle}.$$

Therefore,

$$[T_{e_j}, M_f]u(x) = \partial_j f(x)u(x) + \sum_{\alpha \in R_+} k(\alpha) \langle \alpha, e_j \rangle \frac{f(x) - f(\sigma_\alpha x)}{\langle \alpha, x \rangle} u(\sigma_\alpha x),$$

which proves (6.1).

Since R is normalized,

$$x - \sigma_\alpha x = \langle \alpha, x \rangle \alpha, \quad \|x - \sigma_\alpha x\| = \sqrt{2} |\langle \alpha, x \rangle|.$$

Hence, for almost every $x \notin \alpha^\perp$,

$$\frac{|f(x) - f(\sigma_\alpha x)|}{|\langle \alpha, x \rangle|} = \sqrt{2} \mathcal{G}_{I, \sigma_\alpha} f(x).$$

By Lemma 4.1 (4.2),

$$\left\| \frac{f - f \circ \sigma_\alpha}{\langle \alpha, \cdot \rangle} \right\|_{L^N(\mathbb{R}^N, dx)} \lesssim_{N,G} \|\nabla f\|_{L^N(\mathbb{R}^N, dx)}.$$

This completes the proof of Lemma 6.2. $\square$

Lemma 6.3. Let Γ_+ be an open Weyl chamber, let Γ be its closure, and let R_+ be the corresponding positive subsystem. Set

$$\rho(x) := \varpi(x)^{1/2},$$

and define

$$\begin{aligned} U : L^2(\mathbb{R}^N, dw) &\longrightarrow L^2(\Gamma, dw; \mathbb{C}^G), & (Uu)_g(x) &:= u(gx), \\ V : L^2(\Gamma, dw; \mathbb{C}^G) &\longrightarrow L^2(\Gamma, dx; \mathbb{C}^G), & (Vh)(x) &:= \rho(x)h(x). \end{aligned}$$

Then U and V are unitary. If

$$\widetilde{T}_j := VUT_{e_j}U^{-1}V^{-1}, \quad \widetilde{P} := VU(-\Delta_{\text{Dunkl}})U^{-1}V^{-1},$$

then, on $C_c^\infty(\Gamma_+; \mathbb{C}^G)$,

$$(6.3) \quad (\widetilde{T}_j h)_g = \partial_{d_{g,j}} h_g - \sum_{\alpha \in R_+} k(\alpha) \frac{\langle \alpha, d_{g,j} \rangle}{\langle \alpha, x \rangle} h_{g\sigma_\alpha},$$

$$(6.4) \quad \widetilde{P} = -\Delta_{\mathbb{C}^G} + |\nabla \log \rho|^2 I_{\mathbb{C}^G} - 2 \sum_{\alpha \in R_+} k(\alpha) \frac{S_\alpha}{\langle \alpha, x \rangle^2}.$$

Assume in addition that $N > 1$, and let

$$\widetilde{Q} := VU(-\Delta_{\text{Dunkl}})^{-1/2}U^{-1}V^{-1},$$

where $(-\Delta_{\text{Dunkl}})^{-1/2}$ is defined by the spectral calculus. Then, for every $\chi \in C_c^\infty(\Gamma_+)$, the operator $M_\chi \widetilde{Q} M_\chi$, initially defined on its natural domain, admits a bounded extension to $L^2(\Gamma, dx; \mathbb{C}^G)$, and this extension is a classical matrix-valued pseudodifferential operator of order -1 . Its principal symbol is

$$\sigma_{-1}(M_\chi \widetilde{Q} M_\chi)(x, \xi) = \chi(x)^2 |\xi|^{-1} I_{\mathbb{C}^G}.$$

Consequently, if $x \in \Gamma_+$, $\xi \neq 0$, and $\chi \in C_c^\infty(\Gamma_+)$ is identically 1 in a neighborhood of x , then

$$(6.5) \quad \sigma_{-1}(M_\chi \widetilde{Q} M_\chi)(x, \xi) = |\xi|^{-1} I_{\mathbb{C}^G}.$$

Proof. The Weyl chambers $g\Gamma_+$, $g \in G$, are pairwise disjoint and cover $\mathbb{R}^N$ up to the reflection hyperplanes; see [5, Theorem 2.12]. Since dw is G -invariant,

$$\begin{aligned} \|Uu\|_{L^2(\Gamma, dw; \mathbb{C}^G)}^2 &= \sum_{g \in G} \int_{\Gamma} |u(gx)|^2 dw(x) \\ &= \int_{\mathbb{R}^N} |u(x)|^2 dw(x). \end{aligned}$$

Conversely, for $h = (h_g)_{g \in G} \in L^2(\Gamma, dw; \mathbb{C}^G)$, define $u(gx) := h_g(x)$ for $x \in \Gamma_+$. This determines u almost everywhere on $\mathbb{R}^N$ and gives $Uu = h$. Hence U is unitary. Since $dw = \rho^2 dx$ and $\rho > 0$ on Γ_+ , V is unitary as well.

Pairing the roots α and $-\alpha$ in the definition of T_{e_j} , changing variables in the root system by g , and using the G -invariance of k , we obtain, for $h \in C_c^\infty(\Gamma_+; \mathbb{C}^G)$,

$$(UT_{e_j}U^{-1}h)_g = \partial_{d_{g,j}}h_g + \sum_{\alpha \in R_+} k(\alpha) \frac{\langle \alpha, d_{g,j} \rangle}{\langle \alpha, x \rangle} (h_g - h_{g\sigma_\alpha}).$$

On Γ_+ ,

$$\nabla \log \rho(x) = \sum_{\alpha \in R_+} k(\alpha) \frac{\alpha}{\langle \alpha, x \rangle},$$

and therefore

$$\rho \partial_{d_{g,j}} \rho^{-1} = \partial_{d_{g,j}} - \sum_{\alpha \in R_+} k(\alpha) \frac{\langle \alpha, d_{g,j} \rangle}{\langle \alpha, x \rangle}.$$

Conjugation by V cancels the diagonal terms and gives (6.3).

Similarly, pairing α and $-\alpha$ in the formula for the Dunkl Laplacian and using $\|\alpha\|^2 = 2$, we have

$$\begin{aligned} (U(-\Delta_{\text{Dunkl}})U^{-1}h)_g &= -\Delta h_g - 2 \sum_{\alpha \in R_+} k(\alpha) \frac{\partial_\alpha h_g}{\langle \alpha, x \rangle} \\ &\quad + 2 \sum_{\alpha \in R_+} k(\alpha) \frac{h_g - h_{g\sigma_\alpha}}{\langle \alpha, x \rangle^2}. \end{aligned}$$

Writing $\phi = \log \rho$,

$$\rho(-\Delta - 2\nabla \phi \cdot \nabla) \rho^{-1} = -\Delta + |\nabla \phi|^2 + \Delta \phi,$$

while

$$\Delta \phi = - \sum_{\alpha \in R_+} k(\alpha) \frac{\|\alpha\|^2}{\langle \alpha, x \rangle^2} = -2 \sum_{\alpha \in R_+} \frac{k(\alpha)}{\langle \alpha, x \rangle^2}.$$

The latter term cancels the diagonal part of

$$2 \sum_{\alpha \in R_+} k(\alpha) \frac{I_{\mathbb{C}^G} - S_\alpha}{\langle \alpha, x \rangle^2},$$

and hence

$$\widetilde{P} = -\Delta_{\mathbb{C}^G} + |\nabla \log \rho|^2 I_{\mathbb{C}^G} - 2 \sum_{\alpha \in R_+} k(\alpha) \frac{S_\alpha}{\langle \alpha, x \rangle^2},$$

which proves (6.4).

Assume now that $N > 1$, fix $\chi \in C_c^\infty(\Gamma_+)$, and set

$$K := \text{supp } \chi.$$

On Γ_+ , write

$$\widetilde{P} = -\Delta_{\mathbb{C}^G} + W(x), \quad W(x) = |\nabla \log \rho(x)|^2 I_{\mathbb{C}^G} - 2 \sum_{\alpha \in R_+} k(\alpha) \frac{S_\alpha}{\langle \alpha, x \rangle^2}.$$

Thus W is a smooth Hermitian matrix-valued function on Γ_+ .

Choose open sets

$$K \Subset \Omega_0 \Subset \Omega_1 \Subset \Gamma_+$$

and $\eta \in C_c^\infty(\Gamma_+)$ such that

$$0 \leq \eta \leq 1, \quad \eta = 1 \quad \text{on a neighborhood of } \overline{\Omega_1}.$$

Choose a flat torus M containing $\text{supp } \eta$ in one Euclidean coordinate chart. Since η is compactly supported in that chart, the differential operator $M_\eta \widetilde{P} M_\eta$ extends smoothly by zero to M . Define on $L^2(M; \mathbb{C}^G)$

$$(6.6) \quad P_0 := M_\eta \widetilde{P} M_\eta + M_{1-\eta}(-\Delta_M + 1)M_{1-\eta}.$$

It is a nonnegative self-adjoint elliptic differential operator of order 2. Indeed,

$$\langle P_0 u, u \rangle = \langle \widetilde{P}(\eta u), \eta u \rangle + \|\nabla((1-\eta)u)\|_2^2 + \|(1-\eta)u\|_2^2 \geq 0,$$

and its principal symbol is

$$p_{0,2}(x, \xi) = (\eta(x)^2 + (1-\eta(x))^2)|\xi|^2 I_{\mathbb{C}^G},$$

which is elliptic since

$$\eta^2 + (1-\eta)^2 \geq \frac{1}{2}.$$

Moreover,

$$(6.7) \quad P_0 = \widetilde{P} \quad \text{on } \Omega_1.$$

Let

$$H_t := e^{-t\widetilde{P}}, \quad H_t^0 := e^{-tP_0}.$$

We first consider

$$B_0 := \frac{1}{\sqrt{\pi}} \int_0^1 t^{-1/2} H_t^0 dt.$$

Let Π_0 be the orthogonal projection onto $\ker P_0$. Since P_0 is elliptic on the closed manifold M , Π_0 is smoothing. Applying the complex powers calculus [?, Theorem 2 and p. 290] to $P_0 + \Pi_0$, we obtain

$$(P_0 + \Pi_0)^{-1/2} \in \Psi_{\text{cl}}^{-1}(M; \mathbb{C}^G)$$

with principal symbol $p_{0,2}^{-1/2}$.

Set

$$F(\lambda) := \frac{1}{\sqrt{\pi}} \int_0^1 t^{-1/2} e^{-t\lambda} dt, \quad \lambda \geq 0.$$

Then $B_0 = F(P_0)$, and for $\lambda > 0$,

$$F(\lambda) - \lambda^{-1/2} = -\frac{1}{\sqrt{\pi}} \int_1^\infty t^{-1/2} e^{-t\lambda} dt.$$

Hence

$$F(\lambda) - \lambda^{-1/2} = O(\lambda^{-m}) \quad (\lambda \rightarrow \infty)$$

for every $m > 0$. On $\ker P_0$ the difference between B_0 and $(P_0 + \Pi_0)^{-1/2}$ is finite rank. It follows from elliptic spectral regularity that

$$B_0 - (P_0 + \Pi_0)^{-1/2} \in \Psi^{-\infty}(M; \mathbb{C}^G).$$

Consequently,

$$(6.8) \quad B_0 \in \Psi_{\text{cl}}^{-1}(M; \mathbb{C}^G), \quad \sigma_{-1}(B_0)(x, \xi) = p_{0,2}(x, \xi)^{-1/2}.$$

In particular, by (6.7),

$$(6.9) \quad \sigma_{-1}(B_0)(x, \xi) = |\xi|^{-1} I_{\mathbb{C}^G}, \quad x \in \Omega_1.$$

We next compare H_t and H_t^0 in the interior. Choose $\zeta \in C_c^\infty(\Omega_1)$ such that

$$\zeta = 1 \quad \text{on a neighborhood of } K.$$

Since $P_0 = \widetilde{P}$ on Ω_1 , Duhamel's formula gives

$$(6.10) \quad M_\chi(H_t - H_t^0)M_\chi = - \int_0^t M_\chi H_{t-s} [\widetilde{P}, M_\zeta] H_s^0 M_\chi ds.$$

Indeed, for data supported in K ,

$$v(t) := M_\zeta H_t^0 M_\chi u$$

satisfies

$$(\partial_t + \widetilde{P})v = [\widetilde{P}, M_\zeta] H_t^0 M_\chi u, \quad v(0) = M_\chi u,$$

which yields (6.10).

Let $h_t(x, y)$ be the Dunkl heat kernel. The integral kernel of H_t , with respect to Lebesgue measure on Γ , is the matrix

$$(6.11) \quad \widetilde{h}_t(x, y)_{g,\ell} = \rho(x)\rho(y) h_t(gx, \ell y), \quad g, \ell \in G.$$

For $x, y \in \Gamma$,

$$d(gx, \ell y) = d(x, y) = |x - y|.$$

Together with the mixed derivative estimate for the Dunkl heat kernel [?, Theorem 3.1], this implies that, whenever $E, F \Subset \Gamma_+$ and $\text{dist}(E, F) > 0$, for every pair of multi-indices α, β and every $m > 0$,

$$(6.12) \quad \sup_{\substack{x \in E \\ y \in F}} \left\| \partial_x^\alpha \partial_y^\beta \widetilde{h}_t(x, y) \right\| = O(t^m), \quad t \downarrow 0.$$

The heat kernel of the smooth elliptic operator P_0 satisfies the same off-diagonal estimate on M .

Since

$$\text{supp}[\widetilde{P}, M_\varepsilon] \cap K = \emptyset,$$

the two heat kernels occurring in (6.10) are evaluated between compact sets separated by a positive distance. The standard Gaussian bounds underlying (6.12) therefore give, for every multi-indices α, β and every $m > 0$,

$$(6.13) \quad \sup_{x,y} \left\| \partial_x^\alpha \partial_y^\beta K_{M_\chi(H_t - H_t^0)M_\chi}(x, y) \right\| = O(t^m), \quad t \downarrow 0.$$

Hence

$$(6.14) \quad S_0 := \frac{1}{\sqrt{\pi}} \int_0^1 t^{-1/2} M_\chi(H_t - H_t^0) M_\chi dt \in \Psi^{-\infty}.$$

It remains to treat the large-time part. Let $K_1 \Subset \Gamma_+$ be a compact set containing K . By (6.11), the heat-kernel derivative estimate cited above, and (2.2), for $x, y \in K_1$ and $t \geq 1$,

$$V(gx, \ell y, \sqrt{t})^{-1} \lesssim_{K_1} t^{-N/2}.$$

Since ρ and all its derivatives are bounded on K_1 , it follows that, for every pair of multi-indices α, β ,

$$(6.15) \quad \sup_{x,y \in K_1} \left\| \partial_x^\alpha \partial_y^\beta \widetilde{h}_t(x, y) \right\| \lesssim_{\alpha, \beta, K_1} t^{-N/2}, \quad t \geq 1.$$

Since $N > 1$,

$$\int_1^\infty t^{-1/2} t^{-N/2} dt < \infty.$$

Thus all derivatives of

$$K_{S_\infty}(x, y) := \frac{\chi(x)\chi(y)}{\sqrt{\pi}} \int_1^\infty t^{-1/2} \widetilde{h}_t(x, y) dt$$

converge locally uniformly, and therefore

$$(6.16) \quad S_\infty := \frac{1}{\sqrt{\pi}} \int_1^\infty t^{-1/2} M_\chi H_t M_\chi dt \in \Psi^{-\infty}.$$

For completeness, the preceding integrals give the canonical bounded localization of $\widetilde{P}^{-1/2}$. Indeed, for $\varepsilon > 0$,

$$(\widetilde{P} + \varepsilon)^{-1/2} = \frac{1}{\sqrt{\pi}} \int_0^\infty t^{-1/2} e^{-\varepsilon t} H_t dt$$

by the spectral theorem. The estimates (6.13) and (6.15) imply that

$$M_\chi (\widetilde{P} + \varepsilon)^{-1/2} M_\chi$$

converges in operator norm, as $\varepsilon \downarrow 0$, to

$$\frac{1}{\sqrt{\pi}} \int_0^\infty t^{-1/2} M_\chi H_t M_\chi dt.$$

On the natural domain of $\widetilde{P}^{-1/2}$, this limit agrees with $M_\chi \widetilde{P}^{-1/2} M_\chi$; hence it is its bounded extension.

Combining (6.14), (6.16), and the definition of B_0 , we obtain

$$M_\chi \widetilde{Q} M_\chi = M_\chi B_0 M_\chi + S_0 + S_\infty.$$

By (6.8), $M_\chi B_0 M_\chi$ is a classical matrix-valued pseudodifferential operator of order -1 , while $S_0 + S_\infty$ is smoothing. Therefore

$$M_\chi \widetilde{Q} M_\chi \in \Psi_{\text{cl}}^{-1},$$

and, by (6.9),

$$\sigma_{-1}(M_\chi \widetilde{Q} M_\chi)(x, \xi) = \chi(x)^2 |\xi|^{-1} I_{\mathbb{C}G}.$$

If $\chi = 1$ in a neighborhood of x , this reduces to

$$\sigma_{-1}(M_\chi \widetilde{Q} M_\chi)(x, \xi) = |\xi|^{-1} I_{\mathbb{C}G},$$

which proves (6.5). □

Lemma 6.4. Let $j \in \{1, \dots, N\}$ and $f \in C_c^\infty(\mathbb{R}^N)$, and set

$$A_{f,j} := VU[R_{\text{Dunkl},j}, M_f]U^{-1}V^{-1}.$$

Then, for every $\chi \in C_c^\infty(\Gamma_+)$,

$$M_\chi A_{f,j} M_\chi$$

is a compactly supported classical matrix valued pseudodifferential operator of order -1 , with principal symbol

$$(6.17) \quad \sigma_{-1}(M_\chi A_{f,j} M_\chi)(x, \xi) = \frac{\chi(x)^2}{\|\xi\|} c_{f,j}\left(x, \frac{\xi}{\|\xi\|}\right), \quad x \in \Gamma_+, \quad \xi \neq 0.$$

Proof. Set

$$F(x) := \text{diag}_{g \in G}(f_g(x)).$$

Since $VUM_f U^{-1}V^{-1} = M_F$ and $R_{\text{Dunkl},j} = T_{e_j}(-\Delta_{\text{Dunkl}})^{-1/2}$,

$$(6.18) \quad A_{f,j} = [\widetilde{T}_j, M_F] \widetilde{Q} + \widetilde{T}_j [\widetilde{Q}, M_F].$$

By (6.3),

$$(6.19) \quad \begin{aligned} [\widetilde{T}_j, M_F] &= \text{diag}_{g \in G}(\langle d_{g,j}, \nabla f_g \rangle) \\ &+ \sum_{\alpha \in R_+} k(\alpha) \text{diag}_{g \in G} \left(\frac{\langle \alpha, d_{g,j} \rangle}{\langle \alpha, x \rangle} (f_g - f_{g\sigma_\alpha}) \right) S_\alpha. \end{aligned}$$

Fix $\chi \in C_c^\infty(\Gamma_+)$. Choose $\eta, \psi \in C_c^\infty(\Gamma_+)$ such that $\eta = 1$ on a neighborhood of $\text{supp } \chi$ and $\psi = 1$ on a neighborhood of $\text{supp } \eta$, and set

$$Q_\psi := M_\psi \widetilde{Q} M_\psi.$$

By Lemma 6.3, Q_ψ is a classical matrix valued pseudodifferential operator of order -1 , and on a neighborhood of $\text{supp } \eta$,

$$(6.20) \quad q_{-1}(x, \xi) = \|\xi\|^{-1} I_{\mathbb{C}G}.$$

We use the Kohn–Nirenberg convention

$$\text{Op}(a)u(x) = (2\pi)^{-N} \int_{\mathbb{R}^{2N}} e^{i(x-y)\cdot\xi} a(x, \xi) u(y) dy d\xi.$$

If a and b are classical symbols, the composition formula [?, Theorem 18.1.8, p. 71] gives

$$(6.21) \quad a \# b \sim \sum_{\beta \in \mathbb{N}_0^N} \frac{1}{\beta!} \partial_\xi^\beta a D_x^\beta b, \quad D_x := \frac{1}{i} \partial_x.$$

Write, on a neighborhood of $\text{supp } \eta$,

$$\sigma(Q_\psi)(x, \xi) \sim q_{-1}(x, \xi) + q_{-2}(x, \xi) + \cdots.$$

By (6.4),

$$\sigma(\tilde{P})(x, \xi) \sim p_2(x, \xi) + p_1(x, \xi) + p_0(x, \xi),$$

where

$$p_2(x, \xi) = \|\xi\|^2 I_{\mathbb{C}^G}, \quad p_1(x, \xi) = 0.$$

Since $\tilde{Q} = \tilde{P}^{-1/2}$, $\tilde{Q}\tilde{P}\tilde{Q} = I$. Applying (6.21) on the region where $\psi = 1$ and comparing the homogeneous terms of degree -1 , the terms containing x -derivatives of p_2 and q_{-1} vanish, and hence

$$q_{-2}p_2q_{-1} + q_{-1}p_2q_{-2} = 0.$$

Together with (6.20), this gives

$$(6.22) \quad q_{-2}(x, \xi) = 0$$

on a neighborhood of $\text{supp } \eta$.

Applying (6.21) to the commutator $[Q_\psi, M_F]$, and using (6.20)–(6.22), we obtain, on the same neighborhood,

$$(6.23) \quad \begin{aligned} \sigma_{-2}([Q_\psi, M_F])(x, \xi) &= \frac{1}{i} \sum_{\ell=1}^N \partial_{\xi_\ell} (\|\xi\|^{-1}) \partial_{x_\ell} F(x) \\ &= i \|\xi\|^{-3} \text{diag}_{g \in G} (\langle \xi, \nabla f_g(x) \rangle). \end{aligned}$$

Since $\psi = 1$ on a neighborhood of $\text{supp } \eta$ and $\eta = 1$ on a neighborhood of $\text{supp } \chi$,

$$\begin{aligned} M_\chi[\tilde{T}_j, M_F] \tilde{Q} M_\chi &= M_\chi[\tilde{T}_j, M_F] Q_\psi M_\chi, \\ M_\chi \tilde{T}_j [\tilde{Q}, M_F] M_\chi &= M_\chi \tilde{T}_j M_\eta [Q_\psi, M_F] M_\chi. \end{aligned}$$

Both operators on the right are classical matrix valued pseudodifferential operators of order -1 .

By (6.19), (6.20), and the order-zero part of the composition formula,

$$\begin{aligned} \sigma_{-1}(M_\chi[\tilde{T}_j, M_F] \tilde{Q} M_\chi)(x, \xi) \\ = \frac{\chi(x)^2}{\|\xi\|} \text{diag}_{g \in G} (\langle d_{g,j}, \nabla f_g(x) \rangle) \end{aligned}$$

$$(6.24) \quad + \frac{\chi(x)^2}{\|\xi\|} \sum_{\alpha \in R_+} k(\alpha) \operatorname{diag}_{g \in G} \left(\frac{\langle \alpha, d_{g,j} \rangle}{\langle \alpha, x \rangle} (f_g(x) - f_{g\sigma_\alpha}(x)) \right) S_\alpha.$$

Under the above Kohn–Nirenberg convention, (6.3) gives

$$\sigma_1(\widetilde{T}_j)(x, \xi) = i \operatorname{diag}_{g \in G} (\langle d_{g,j}, \xi \rangle).$$

Therefore, by (6.23),

$$(6.25) \quad \begin{aligned} & \sigma_{-1}(M_\chi \widetilde{T}_j [\widetilde{Q}, M_F] M_\chi)(x, \xi) \\ &= -\chi(x)^2 \|\xi\|^{-3} \operatorname{diag}_{g \in G} (\langle d_{g,j}, \xi \rangle \langle \xi, \nabla f_g(x) \rangle). \end{aligned}$$

Combining (6.24) and (6.25), and setting

$$\theta := \frac{\xi}{\|\xi\|},$$

we obtain

$$\begin{aligned} \sigma_{-1}(M_\chi A_{f,j} M_\chi)(x, \xi) &= \frac{\chi(x)^2}{\|\xi\|} \operatorname{diag}_{g \in G} \left(\langle d_{g,j}, \nabla f_g(x) \rangle - \langle d_{g,j}, \theta \rangle \langle \theta, \nabla f_g(x) \rangle \right) \\ &\quad + \frac{\chi(x)^2}{\|\xi\|} \sum_{\alpha \in R_+} k(\alpha) \operatorname{diag}_{g \in G} \left(\frac{\langle \alpha, d_{g,j} \rangle}{\langle \alpha, x \rangle} (f_g(x) - f_{g\sigma_\alpha}(x)) \right) S_\alpha \\ &= \frac{\chi(x)^2}{\|\xi\|} c_{f,j}(x, \theta). \end{aligned}$$

This proves (6.17). □

Lemma 6.5. Assume that $N > 1$, $j \in \{1, \dots, N\}$, and $f \in \dot{W}^{1,N}(\mathbb{R}^N)$. Then

$$\left(\int_\Gamma \int_{\mathbb{S}^{N-1}} \operatorname{tr}_{\mathbb{C}^G} |c_{f,j}(x, \theta)|^N d\theta dx \right)^{1/N} \simeq_{N,R,k} \|\nabla f\|_{L^N(\mathbb{R}^N, dx)}.$$

Proof. By the definition of $c_{f,j}$, the unitary invariance of the Schatten norm, and $\|d_{g,j}\| = 1$,

$$\begin{aligned} \|c_{f,j}(x, \theta)\|_{S^N} &\lesssim_{R,k} \left(\sum_{g \in G} |\nabla f_g(x)|^N \right)^{1/N} \\ &\quad + \sum_{\alpha \in R_+} \left(\sum_{g \in G} \left| \frac{f_g(x) - f_{g\sigma_\alpha}(x)}{\langle \alpha, x \rangle} \right|^N \right)^{1/N}. \end{aligned}$$

The chamber decomposition used in Lemma 6.3 and the orthogonality of $g \in G$ imply

$$\sum_{g \in G} \int_\Gamma |\nabla f_g(x)|^N dx = \|\nabla f\|_{L^N(\mathbb{R}^N, dx)}^N.$$

Moreover, since $\|\alpha\|^2 = 2$,

$$|x - \sigma_\alpha x| = \sqrt{2} |\langle \alpha, x \rangle|,$$

and hence

$$\left| \frac{f_g(x) - f_{g\sigma_\alpha}(x)}{\langle \alpha, x \rangle} \right| = \sqrt{2} \mathcal{G}_{g, g\sigma_\alpha} f(x)$$

for almost every $x \in \mathbb{R}^N$. By Lemma 4.1,

$$\int_{\Gamma} \left| \frac{f_g(x) - f_{g\sigma_\alpha}(x)}{\langle \alpha, x \rangle} \right|^N dx \lesssim_{N,G} \|\nabla f\|_{L^N(\mathbb{R}^N, dx)}^N$$

uniformly in $g \in G$ and $\alpha \in R_+$. Since G and R_+ are finite,

$$(6.26) \quad \left(\int_{\Gamma} \int_{\mathbb{S}^{N-1}} \text{tr}_{\mathbb{C}^G} |\mathfrak{c}_{f,j}(x, \theta)|^N d\theta dx \right)^{1/N} \lesssim_{N,R,k} \|\nabla f\|_{L^N(\mathbb{R}^N, dx)}.$$

For the converse estimate, let P_g denote the orthogonal projection of $\mathbb{C}^G$ onto the g -th coordinate and define

$$\mathcal{E}(A) := \sum_{g \in G} P_g A P_g, \quad A \in \mathcal{L}(\mathbb{C}^G).$$

Thus, $\mathcal{E}(A)$ is the diagonal part of A . The pinching inequality for unitarily invariant norms [?, p. 97, (IV.51)–(IV.52)] gives

$$(6.27) \quad \|\mathcal{E}(A)\|_{S^N} \leq \|A\|_{S^N}.$$

For every $\alpha \in R_+$, S_α has zero diagonal, since $g\sigma_\alpha \neq g$ for all $g \in G$. Hence, by the definition of $\mathfrak{c}_{f,j}$,

$$\mathcal{E}(\mathfrak{c}_{f,j}(x, \theta)) = \text{diag}_{g \in G} \left(\langle d_{g,j}, \nabla f_g(x) \rangle - \langle d_{g,j}, \theta \rangle \langle \theta, \nabla f_g(x) \rangle \right).$$

Therefore, by (6.27),

$$(6.28) \quad \|\mathfrak{c}_{f,j}(x, \theta)\|_{S^N}^N \geq \sum_{g \in G} \left| \langle d_{g,j}, \nabla f_g(x) \rangle - \langle d_{g,j}, \theta \rangle \langle \theta, \nabla f_g(x) \rangle \right|^N.$$

For each $g \in G$, set

$$y := gx, \quad \eta := g\theta.$$

Since

$$\nabla f_g(x) = g^{-1} \nabla f(y), \quad d_{g,j} = g^{-1} e_j,$$

and g is orthogonal,

$$\begin{aligned} \langle d_{g,j}, \nabla f_g(x) \rangle &= \partial_j f(y), \\ \langle d_{g,j}, \theta \rangle &= \eta_j, \\ \langle \theta, \nabla f_g(x) \rangle &= \langle \eta, \nabla f(y) \rangle. \end{aligned}$$

Using the chamber decomposition and the orthogonal invariance of surface measure, (6.28) therefore gives

$$(6.29) \quad \begin{aligned} & \int_{\Gamma} \int_{\mathbb{S}^{N-1}} \|\mathfrak{c}_{f,j}(x, \theta)\|_{S^N}^N d\theta dx \\ & \geq \int_{\mathbb{R}^N} \int_{\mathbb{S}^{N-1}} \left| \partial_j f(y) - \eta_j \langle \eta, \nabla f(y) \rangle \right|^N d\eta dy. \end{aligned}$$

For $v \in \mathbb{C}^N$, set

$$\Phi_j(v) := \left(\int_{\mathbb{S}^{N-1}} |v_j - \eta_j \langle \eta, v \rangle|^N d\eta \right)^{1/N}.$$

Then

$$(6.30) \quad \Phi_j(v) \simeq_N \|v\|.$$

Indeed, Φ_j is a seminorm on $\mathbb{C}^N$. If $\Phi_j(v) = 0$, continuity gives

$$v_j - \eta_j \langle \eta, v \rangle = 0, \quad \eta \in \mathbb{S}^{N-1}.$$

Since $N > 1$, there exists $\eta \in \mathbb{S}^{N-1}$ with $\eta_j = 0$, and therefore $v_j = 0$. It follows that

$$\langle \eta, v \rangle = 0$$

whenever $\eta_j \neq 0$. Since such η form a dense subset of $\mathbb{S}^{N-1}$, continuity yields $\langle \eta, v \rangle = 0$ for every $\eta \in \mathbb{S}^{N-1}$, and hence $v = 0$. Thus Φ_j is a norm, and (6.30) follows from the equivalence of norms on the finite dimensional space $\mathbb{C}^N$.

Applying (6.30) with $v = \nabla f(y)$ in (6.29), we obtain

$$\|\nabla f\|_{L^N(\mathbb{R}^N, dx)} \lesssim_N \left(\int_{\Gamma} \int_{\mathbb{S}^{N-1}} \text{tr}_{\mathbb{C}^G} |\zeta_{f,j}(x, \theta)|^N d\theta dx \right)^{1/N}.$$

Combining this estimate with (6.26) proves the assertion. $\square$

Lemma 6.6. Assume that $N > 1$, $j \in \{1, \dots, N\}$, and $f \in C_c^\infty(\mathbb{R}^N)$. Define

$$A_f := VU[R_{\text{Dunkl},j}, M_f]U^{-1}V^{-1}.$$

Then there exist $\chi_m \in C_c^\infty(\Gamma_+)$, $0 \leq \chi_m \leq 1$, such that $\chi_m(x) \rightarrow 1$ for every $x \in \Gamma_+$ and

$$(6.31) \quad \lim_{m \rightarrow \infty} \|A_f - M_{\chi_m} A_f M_{\chi_m}\|_{S^{N,\infty}} = 0.$$

Proof. Fix $s > 2$, choose $2 < s_0 < s$, and set

$$\gamma := \frac{1}{s_0} - \frac{1}{s} > 0.$$

Let $r_s > s$ be as in Lemma ?? and set $\beta := N/r_s$. By Lemma 8.2 (1), $f \in OSC^{N,\infty}(\mathbb{R}^N)$, so Lemma ?? applies with $p = N$ and $q = \infty$.

Set

$$B_f := U[R_{\text{Dunkl},j}, M_f]U^{-1}.$$

Since $V = M_\rho$ commutes with scalar multiplication operators on Γ ,

$$(6.32) \quad \|A_f - M_\chi A_f M_\chi\|_{S^{N,\infty}} = \|B_f - M_\chi B_f M_\chi\|_{S^{N,\infty}}$$

for every bounded scalar multiplier χ .

For fixed $\sigma, \tau \in G$, $u, v \in \mathbb{Z}^N$, and $\varepsilon \in \{0, 1, 2, 3\}$, define

$$T_{u,v,\varepsilon}^{\sigma,\tau} := \sum_{l \in \mathbb{D}_\Gamma} A_{\sigma,\tau,l,\varepsilon}(f) \left\langle \cdot, \overline{H_{u,v,l,\varepsilon}^{\sigma,\tau}(\tau \cdot)} \right\rangle_{L^2(\Gamma, dw)} E_{u,v,l,\varepsilon}^{\sigma,\tau}(\sigma \cdot).$$

By the operator series identification in the proof of Theorem ??, the (σ, τ) matrix entry of B_f is

$$\sum_{\varepsilon=0}^3 \sum_{u,v \in \mathbb{Z}^N} c_{u,v} T_{u,v,\varepsilon}^{\sigma,\tau},$$

with convergence in $\mathcal{S}^{N,\infty}$. Moreover, by Lemma ?? (??)–(??),

$$(6.33) \quad \text{supp } E_{u,v,I,\varepsilon}^{\sigma,\tau}(\sigma \cdot) \subset I, \quad \text{supp } H_{u,v,I,\varepsilon}^{\sigma,\tau}(\tau \cdot) \subset I^*,$$

$$(6.34) \quad \|E_{u,v,I,\varepsilon}^{\sigma,\tau}(\sigma \cdot)\|_{L^s(dw)} + \|H_{u,v,I,\varepsilon}^{\sigma,\tau}(\tau \cdot)\|_{L^s(dw)} \lesssim w(I)^{1/s-1/2}.$$

The definitions of $A_{\sigma,\tau,I,\varepsilon}(f)$ in the proof of Lemma ?? imply

$$(6.35) \quad |A_{\sigma,\tau,I,\varepsilon}(f)| \lesssim_f \begin{cases} \ell(I), & \ell(I) \leq 1, \\ \ell(I)^{-\beta}, & \ell(I) \geq 1. \end{cases}$$

Indeed, for $\varepsilon = 0, 1, 2$, the corresponding coefficients are bounded by an $L^{r_s}(dx)$ oscillation of a smooth compactly supported function on a set of diameter comparable with $\ell(I)$. Hence the first estimate follows from the Lipschitz bound, while for $\ell(I) \geq 1$,

$$\text{osc}_{r_s}(g; E) \lesssim |E|^{-1/r_s} \|g\|_{L^{r_s}} \lesssim_f \ell(I)^{-N/r_s}.$$

For $\varepsilon = 3$, writing $D := |\sigma z_I - \tau z_I|$, the definitions of $\theta_{\sigma,\tau,I}$ and $a_{\sigma,\tau,I}(f)$ imply, for $\ell(I) \leq 1$,

$$|A_{\sigma,\tau,I,3}(f)| \lesssim_f \frac{\ell(I)}{\ell(I) + D} (\ell(I) + D) \lesssim_f \ell(I),$$

whereas, for $\ell(I) \geq 1$,

$$|A_{\sigma,\tau,I,3}(f)| \leq |I|^{-1} \|H_{\sigma,\tau}(f)\|_{L^1} \lesssim_f \ell(I)^{-N} \lesssim \ell(I)^{-\beta}.$$

Let $\Delta = \{\alpha_1, \dots, \alpha_{N_0}\}$. For $1 \leq i \leq N_0$ and $0 < \delta \leq 1$, set

$$W_{i,\delta} := \{x \in \Gamma : 0 \leq \langle x, \alpha_i \rangle \leq 2\delta\}.$$

We have

$$(6.36) \quad \frac{w(W_{i,\delta} \cap I)}{w(I)} + \frac{w(W_{i,\delta} \cap I^*)}{w(I)} \lesssim \min\left(1, \frac{\delta}{\ell(I)}\right).$$

To prove this, write $I = \Phi^{-1}(Q)$ and $\ell := \ell(I)$. The assertion is immediate when $\delta > \ell/8$, since $w(I^*) \lesssim w(I)$. Suppose that $\delta \leq \ell/8$. If I or I^* meets $W_{i,\delta}$, then the i -th coordinate interval of Q is $[0, \ell)$. By Lemma 3.9 (3.5), the density of $\Phi_* dw$ is

$$\phi(u) = |\det(\Phi)|^{-1} \prod_{v \in R_+} \left(\sum_{h=1}^{N_0} a_{v,h} u_h \right)^{2k(v)},$$

which is nondecreasing in every u_i . If g is nonnegative and nondecreasing on an interval of length comparable with ℓ , then

$$\int_0^{2\delta} g(t) dt \lesssim \frac{\delta}{\ell} \int g(t) dt.$$

Applying this estimate fiberwise to ϕ on Q , respectively on $(2Q) \cap \Omega$, and using $w(I^*) \lesssim w(I)$, we obtain (6.36).

Define

$$\omega_{I,\delta} := \min\left(1, \left(\frac{\delta}{\ell(I)}\right)^\gamma\right).$$

If $0 \leq a \leq 1$ and $\text{supp } a \subset W_{i,\delta}$, then by Hölder's inequality, (6.34), and (6.36),

$$\|aE_{u,v,I,\varepsilon}^{\sigma,\tau}(\sigma \cdot)\|_{L^{s_0}(dw)} + \|aH_{u,v,I,\varepsilon}^{\sigma,\tau}(\tau \cdot)\|_{L^{s_0}(dw)} \lesssim \omega_{I,\delta} w(I)^{1/s_0-1/2}.$$

After normalization by $\omega_{I,\delta}$, the localized factors are NWO families with exponent s_0 . Hence, by Lemma 2.10,

$$(6.37) \quad \|M_a T_{u,v,\varepsilon}^{\sigma,\tau}\|_{S^{N,\infty}} + \|T_{u,v,\varepsilon}^{\sigma,\tau} M_a\|_{S^{N,\infty}} \lesssim \left\| \left\{ A_{\sigma,\tau,I,\varepsilon}(f) \omega_{I,\delta} \right\}_{I \in \mathbb{D}_\Gamma} \right\|_{\ell^{N,\infty}}.$$

If $\ell(I) \leq 1$ and $A_{\sigma,\tau,I,\varepsilon}(f) \neq 0$, then z_I belongs to a fixed bounded set depending only on f . Consequently, at every dyadic length $0 < \ell \leq 1$,

$$(6.38) \quad \#\left\{ I \in \mathbb{D}_\Gamma : \ell(I) = \ell, A_{\sigma,\tau,I,\varepsilon}(f) \neq 0, (I \cup I^*) \cap W_{i,\delta} \neq \emptyset \right\} \lesssim_f \left(1 + \frac{\delta}{\ell}\right) \ell^{-(N-1)}.$$

Indeed, under Φ , there are $O(1 + \delta/\ell)$ admissible intervals in the u_i coordinate and $O(\ell^{-1})$ admissible intervals in each of the remaining $N-1$ coordinates.

For $\ell(I) \leq \delta$, (6.35) and (6.38) imply, for every $\lambda > 0$,

$$\begin{aligned} & \#\left\{ I : \ell(I) \leq \delta, |A_{\sigma,\tau,I,\varepsilon}(f) \omega_{I,\delta}| > \lambda \right\} \\ & \lesssim_f \delta \sum_{\substack{\ell \leq \delta \\ \lambda \lesssim_f \ell}} \ell^{-N} \lesssim_f \delta \lambda^{-N}. \end{aligned}$$

Therefore,

$$(6.39) \quad \left\| \left\{ A_{\sigma,\tau,I,\varepsilon}(f) \omega_{I,\delta} \right\}_{\ell(I) \leq \delta} \right\|_{\ell^{N,\infty}} \lesssim_f \delta^{1/N}.$$

For $\delta < \ell(I) \leq 1$,

$$|A_{\sigma,\tau,I,\varepsilon}(f) \omega_{I,\delta}| \lesssim_f \delta^\gamma \ell(I)^{1-\gamma},$$

and (6.38) reduces to $O_f(\ell^{-(N-1)})$. Given $\lambda > 0$, set

$$L := \max\left(\delta, c_f \left(\frac{\lambda}{\delta^\gamma}\right)^{1/(1-\gamma)}\right).$$

Since $N > 1$,

$$\#\left\{ I : \delta < \ell(I) \leq 1, |A_{\sigma,\tau,I,\varepsilon}(f) \omega_{I,\delta}| > \lambda \right\} \lesssim_f L^{-(N-1)}.$$

If $L = \delta$, then $\lambda \lesssim_f \delta$, and hence

$$\lambda L^{-(N-1)/N} \lesssim_f \delta^{1/N}.$$

Otherwise, $\lambda \lesssim_f \delta^\gamma L^{1-\gamma}$, so

$$\lambda L^{-(N-1)/N} \lesssim_f \delta^\gamma L^{1/N-\gamma} \lesssim_f \delta^{\min(\gamma, 1/N)}.$$

Thus,

$$(6.40) \quad \left\| \left\{ A_{\sigma,\tau,I,\varepsilon}(f) \omega_{I,\delta} \right\}_{\delta < \ell(I) \leq 1} \right\|_{\ell^{N,\infty}} \lesssim_f \delta^{\min(\gamma, 1/N)}.$$

For $\ell(I) \geq 1$, only $O_f(1)$ cubes at each dyadic scale have nonzero coefficients. By (6.35),

$$\sum_{\ell(I) \geq 1} |A_{\sigma,\tau,I,\varepsilon}(f) \omega_{I,\delta}|^N \lesssim_f \delta^{\gamma N} \sum_{\substack{\ell \geq 1 \\ \ell \text{ dyadic}}} \ell^{-N(\beta+\gamma)} \lesssim_f \delta^{\gamma N}.$$

Combining this with (6.39) and (6.40), we obtain

$$(6.41) \quad \left\| \left\{ A_{\sigma,\tau,I,\varepsilon}(f) \omega_{I,\delta} \right\}_{I \in \mathbb{D}_\Gamma} \right\|_{\ell^{N,\infty}} \lesssim_f \delta^\kappa, \quad \kappa := \min\left(\gamma, \frac{1}{N}\right).$$

By (6.37),

$$(6.42) \quad \|M_a T_{u,v,\varepsilon}^{\sigma,\tau}\|_{S^{N,\infty}} + \|T_{u,v,\varepsilon}^{\sigma,\tau} M_a\|_{S^{N,\infty}} \lesssim_f \delta^\kappa$$

whenever $0 \leq a \leq 1$ and $\text{supp } a \subset W_{i,\delta}$.

We also need localization at infinity. Let $0 \leq b_R \leq 1$ with $\text{supp } b_R \subset \{x \in \Gamma : |x| \geq R\}$. If $A_{\sigma,\tau,I,\varepsilon}(f) \neq 0$, then

$$(6.43) \quad |z_I| \lesssim_f 1 + \ell(I).$$

For $\varepsilon = 0, 1$, the ball $B_\Gamma(z_I, C_1 \ell(I))$ must meet the support of $f \circ \tau$; for $\varepsilon = 2, 3$, I must meet the support of $H_{\sigma,\tau}(f)$. Since $|x - z_I| \lesssim \ell(I)$ for $x \in I^*$, (6.43) shows that, for sufficiently large R ,

$$A_{\sigma,\tau,I,\varepsilon}(f) \neq 0, \quad (I \cup I^*) \cap \{x : |x| \geq R\} \neq \emptyset$$

can occur only when $\ell(I) \gtrsim_f R$. At each such dyadic scale there are $O_f(1)$ cubes with nonzero coefficients. Hence, by (6.35) and Lemma 2.10,

$$(6.44) \quad \|M_{b_R} T_{u,v,\varepsilon}^{\sigma,\tau}\|_{S^{N,\infty}} + \|T_{u,v,\varepsilon}^{\sigma,\tau} M_{b_R}\|_{S^{N,\infty}} \lesssim_f R^{-\beta}.$$

Choose $\eta \in C^\infty(\mathbb{R})$, $0 \leq \eta \leq 1$, such that

$$\eta(t) = 0 \quad \text{for } t \leq 1, \quad \eta(t) = 1 \quad \text{for } t \geq 2,$$

and choose $\zeta \in C_c^\infty(\mathbb{R}^N)$, $0 \leq \zeta \leq 1$, such that

$$\zeta = 1 \quad \text{on } B(0, 1), \quad \text{supp } \zeta \subset B(0, 2).$$

For $0 < \delta \leq 1$ and $R \geq 1$, define

$$\chi_{\delta,R}(x) := \zeta(x/R) \prod_{i=1}^{N_0} \eta\left(\frac{\langle x, \alpha_i \rangle}{\delta}\right), \quad x \in \Gamma_+.$$

Then $\chi_{\delta,R} \in C_c^\infty(\Gamma_+)$ and $0 \leq \chi_{\delta,R} \leq 1$. Moreover,

$$\begin{aligned} 1 - \chi_{\delta,R} &= 1 - \zeta(\cdot/R) \\ &\quad + \sum_{i=1}^{N_0} \zeta(\cdot/R) \prod_{h < i} \eta\left(\frac{\langle \cdot, \alpha_h \rangle}{\delta}\right) \left(1 - \eta\left(\frac{\langle \cdot, \alpha_i \rangle}{\delta}\right)\right). \end{aligned}$$

The first term is supported in $\{x : |x| \geq R\}$, while the i -th term in the sum is supported in $W_{i,\delta}$. Since G is finite, $S^{N,\infty}$ is normable for $N > 1$, and Lemma ?? (??) implies

$$\sum_{u,v \in \mathbb{Z}^N} |c_{u,v}| < \infty,$$

we may sum (6.42) and (6.44) over the operator series for B_f . Therefore,

$$\|(I - M_{\chi_{\delta,R}})B_f\|_{S^{N,\infty}} + \|B_f(I - M_{\chi_{\delta,R}})\|_{S^{N,\infty}} \lesssim_f \delta^\kappa + R^{-\beta}.$$

Since

$$B_f - M_{\chi_{\delta,R}}B_fM_{\chi_{\delta,R}} = (I - M_{\chi_{\delta,R}})B_f + M_{\chi_{\delta,R}}B_f(I - M_{\chi_{\delta,R}}),$$

the ideal property of $S^{N,\infty}$ and (6.32) imply

$$\|A_f - M_{\chi_{\delta,R}}A_fM_{\chi_{\delta,R}}\|_{S^{N,\infty}} \lesssim_f \delta^\kappa + R^{-\beta}.$$

Define

$$\chi_m := \chi_{2^{-m}, 2^m}.$$

Then $\chi_m(x) \rightarrow 1$ for every $x \in \Gamma_+$, while

$$\|A_f - M_{\chi_m}A_fM_{\chi_m}\|_{S^{N,\infty}} \lesssim_f 2^{-m\kappa} + 2^{-m\beta} \rightarrow 0.$$

This completes the proof of Lemma 9.1. □

Lemma 6.7. Assume that $N > 1$, $j \in \{1, \dots, N\}$, and $f \in C_c^\infty(\mathbb{R}^N)$. Then

$$(6.45) \quad \lim_{t \rightarrow \infty} t^{1/N} \mu(t, [R_{\text{Dunkl},j}, M_f]) = \left(\frac{1}{N(2\pi)^N} \int_{\Gamma} \int_{\mathbb{S}^{N-1}} \text{tr}_{\mathbb{C}G} |\mathfrak{c}_{f,j}(x, \theta)|^N d\theta dx \right)^{1/N}.$$

Proof. Define

$$A_f := VU[R_{\text{Dunkl},j}, M_f]U^{-1}V^{-1}.$$

By Lemma 9.1 (9.2), there exist $\chi_m \in C_c^\infty(\Gamma_+)$, $0 \leq \chi_m \leq 1$, such that $\chi_m(x) \rightarrow 1$ for every $x \in \Gamma_+$ and

$$(6.46) \quad \|A_f - M_{\chi_m}A_fM_{\chi_m}\|_{S^{N,\infty}} \rightarrow 0.$$

Set

$$A_{f,m} := M_{\chi_m}A_fM_{\chi_m}, \quad K_m := \text{supp } \chi_m.$$

By Lemma 6.4 (6.17), $A_{f,m}$ is a compactly supported classical matrix valued pseudodifferential operator of order -1 on Γ_+ , with

$$(6.47) \quad \sigma_{-1}(A_{f,m})(x, \xi) = \frac{\chi_m(x)^2}{\|\xi\|} \mathfrak{c}_{f,j}\left(x, \frac{\xi}{\|\xi\|}\right), \quad x \in \Gamma_+, \quad \xi \neq 0.$$

Choose a bounded open set U_m with $K_m \Subset U_m \Subset \Gamma_+$, and choose $L_m > 0$ such that

$$L_m > \sup_{x,y \in U_m} \|x - y\|_\infty.$$

For

$$\mathbb{T}_m^N := \mathbb{R}^N / L_m \mathbb{Z}^N,$$

the quotient map restricts to a diffeomorphism

$$\kappa_m : U_m \longrightarrow \Omega_m := \kappa_m(U_m),$$

which is a local isometry. Since the Schwartz kernel of $A_{f,m}$ is supported in $K_m \times K_m$, with respect to

$$L^2(\Gamma; \mathbb{C}^G) = L^2(U_m; \mathbb{C}^G) \oplus L^2(\Gamma \setminus U_m; \mathbb{C}^G)$$

we have

$$A_{f,m} = B_{f,m} \oplus 0$$

for some $B_{f,m}$ on $L^2(U_m; \mathbb{C}^G)$.

Let

$$W_m : L^2(U_m; \mathbb{C}^G) \longrightarrow L^2(\Omega_m; \mathbb{C}^G), \quad W_m u := u \circ \kappa_m^{-1}.$$

Then W_m is unitary. By the invariance of classical pseudodifferential operators under smooth coordinate changes [?, Theorem 18.1.17],

$$T_{f,m} := W_m B_{f,m} W_m^{-1} \in \Psi_{\text{cl}}^{-1}(\Omega_m; \mathbb{C}^G).$$

Its Schwartz kernel is supported in $\kappa_m(K_m) \times \kappa_m(K_m) \Subset \Omega_m \times \Omega_m$. Choose $\vartheta_m \in C_c^\infty(\Omega_m)$ equal to 1 on a neighborhood of $\kappa_m(K_m)$. Then

$$T_{f,m} = M_{\vartheta_m} T_{f,m} M_{\vartheta_m}.$$

Hence the zero extension of the Schwartz kernel of $T_{f,m}$ from $\Omega_m \times \Omega_m$ to $\mathbb{T}_m^N \times \mathbb{T}_m^N$ is a classical pseudodifferential kernel of order -1 : locally in $\Omega_m \times \Omega_m$ this follows from $T_{f,m} \in \Psi_{\text{cl}}^{-1}$, while near the boundary of $\Omega_m \times \Omega_m$ the kernel vanishes. Denote the corresponding operator on $\mathbb{T}_m^N$ by $\widehat{A}_{f,m}$. Thus,

$$(6.48) \quad \widehat{A}_{f,m} \in \Psi_{\text{cl}}^{-1}(\mathbb{T}_m^N; \mathbb{C}^G).$$

Since κ_m is a local isometry, by (6.47) and the transformation law for principal symbols,

$$(6.49) \quad \sigma_{-1}(\widehat{A}_{f,m})(\kappa_m(x), \xi) = \frac{\chi_m(x)^2}{\|\xi\|} c_{f,j} \left(x, \frac{\xi}{\|\xi\|} \right), \quad x \in U_m, \quad \xi \neq 0,$$

and the principal symbol vanishes outside Ω_m .

With respect to

$$L^2(\mathbb{T}_m^N; \mathbb{C}^G) = L^2(\Omega_m; \mathbb{C}^G) \oplus L^2(\mathbb{T}_m^N \setminus \Omega_m; \mathbb{C}^G),$$

we have

$$\widehat{A}_{f,m} = T_{f,m} \oplus 0.$$

Since $T_{f,m}$ and $B_{f,m}$ are unitarily equivalent,

$$(6.50) \quad \mu(t, \widehat{A}_{f,m}) = \mu(t, A_{f,m}), \quad t > 0.$$

By (6.48) and the Weyl law [?, Theorem 6.1],

$$\lim_{t \rightarrow \infty} t^{1/N} \mu(t, \widehat{A}_{f,m}) = \left(\frac{1}{N(2\pi)^N} \int_{S^* \mathbb{T}_m^N} \text{tr}_{\mathbb{C}^G} |\sigma_{-1}(\widehat{A}_{f,m})(z, \theta)|^N dz d\theta \right)^{1/N}.$$

Since the metric on $\mathbb{T}_m^N$ is flat, combining (6.49) with (6.50), we obtain

$$(6.51) \quad \lim_{t \rightarrow \infty} t^{1/N} \mu(t, A_{f,m}) = \left(\frac{1}{N(2\pi)^N} \int_{\Gamma} \int_{\mathbb{S}^{N-1}} \chi_m(x)^{2N} \operatorname{tr}_{\mathbb{C}^G} |\mathfrak{c}_{f,j}(x, \theta)|^N d\theta dx \right)^{1/N},$$

where χ_m is identified with its zero extension to Γ .

By Lemma 6.5,

$$(x, \theta) \mapsto \operatorname{tr}_{\mathbb{C}^G} |\mathfrak{c}_{f,j}(x, \theta)|^N$$

belongs to $L^1(\Gamma \times \mathbb{S}^{N-1})$. Since $0 \leq \chi_m \leq 1$, $\chi_m \rightarrow 1$ on Γ_+ , and $\Gamma \setminus \Gamma_+$ is contained in finitely many hyperplanes, the dominated convergence theorem implies

$$(6.52) \quad \begin{aligned} \lim_{m \rightarrow \infty} \left(\frac{1}{N(2\pi)^N} \int_{\Gamma} \int_{\mathbb{S}^{N-1}} \chi_m(x)^{2N} \operatorname{tr}_{\mathbb{C}^G} |\mathfrak{c}_{f,j}(x, \theta)|^N d\theta dx \right)^{1/N} \\ = \left(\frac{1}{N(2\pi)^N} \int_{\Gamma} \int_{\mathbb{S}^{N-1}} \operatorname{tr}_{\mathbb{C}^G} |\mathfrak{c}_{f,j}(x, \theta)|^N d\theta dx \right)^{1/N}. \end{aligned}$$

By (6.46), $A_{f,m} \rightarrow A_f$ in $S^{N,\infty}$. Combining (6.51), (6.52), and Lemma 6.8, we obtain

$$\lim_{t \rightarrow \infty} t^{1/N} \mu(t, A_f) = \left(\frac{1}{N(2\pi)^N} \int_{\Gamma} \int_{\mathbb{S}^{N-1}} \operatorname{tr}_{\mathbb{C}^G} |\mathfrak{c}_{f,j}(x, \theta)|^N d\theta dx \right)^{1/N}.$$

Since U and V are unitary,

$$\mu(t, A_f) = \mu(t, [R_{\text{Dunkl},j}, M_f]), \quad t > 0.$$

This completes the proof of Lemma 6.7. $\square$

Lemma 6.8. Let $0 < p < \infty$, let $\mathcal{H}$ be a Hilbert space, and let $A, A_m \in S^{p,\infty}(\mathcal{H})$, $m \geq 1$, satisfy

$$\|A_m - A\|_{S^{p,\infty}} \longrightarrow 0.$$

Suppose that, for every $m \geq 1$, the limit

$$c_m := \lim_{t \rightarrow \infty} t^{1/p} \mu(t, A_m)$$

exists. Then both limits

$$\lim_{m \rightarrow \infty} c_m \quad \text{and} \quad \lim_{t \rightarrow \infty} t^{1/p} \mu(t, A)$$

exist and

$$\lim_{t \rightarrow \infty} t^{1/p} \mu(t, A) = \lim_{m \rightarrow \infty} c_m.$$

Proof. The assertion is the Birman–Solomyak approximation lemma; see [?, Section 11.6]. $\square$

Lemma 6.9. Assume that $N > 1$, $j \in \{1, \dots, N\}$, and $f \in \dot{W}^{1,N}(\mathbb{R}^N)$. Then there exists $f_m \in C_c^\infty(\mathbb{R}^N)$ such that

$$\begin{aligned} \|\nabla(f_m - f)\|_{L^N(\mathbb{R}^N, dx)} &\longrightarrow 0, \\ \|[R_{\text{Dunkl},j}, M_{f_m}] - [R_{\text{Dunkl},j}, M_f]\|_{S^{N,\infty}} &\longrightarrow 0, \\ \|\mathfrak{c}_{f_m,j} - \mathfrak{c}_{f,j}\|_{L^N(\Gamma \times \mathbb{S}^{N-1}; S^N(\mathbb{C}^G))} &\longrightarrow 0. \end{aligned}$$

Proof. By the density of $C_c^\infty(\mathbb{R}^N)$ in $\dot{W}^{1,N}(\mathbb{R}^N)$, choose $f_m \in C_c^\infty(\mathbb{R}^N)$ such that

$$\|\nabla(f_m - f)\|_{L^N(\mathbb{R}^N, dx)} \longrightarrow 0.$$

By Theorem ?? and Lemma 8.2,

$$\begin{aligned} \|[R_{\text{Dunkl},j}, M_{f_m}] - [R_{\text{Dunkl},j}, M_f]\|_{S^{N,\infty}} &= \|[R_{\text{Dunkl},j}, M_{f_m-f}]\|_{S^{N,\infty}} \\ &\lesssim \|f_m - f\|_{\text{OSC}^{N,\infty}(\mathbb{R}^N)} \lesssim \|\nabla(f_m - f)\|_{L^N(\mathbb{R}^N, dx)} \longrightarrow 0. \end{aligned}$$

By the linearity of $f \mapsto \mathfrak{c}_{f,j}$ and Lemma 6.5,

$$\begin{aligned} \|\mathfrak{c}_{f_m,j} - \mathfrak{c}_{f,j}\|_{L^N(\Gamma \times \mathbb{S}^{N-1}; S^N(\mathbb{C}^G))} &= \|\mathfrak{c}_{f_m-f,j}\|_{L^N(\Gamma \times \mathbb{S}^{N-1}; S^N(\mathbb{C}^G))} \\ &\lesssim_{N,R,k} \|\nabla(f_m - f)\|_{L^N(\mathbb{R}^N, dx)} \longrightarrow 0. \end{aligned}$$

This completes the proof of Lemma 6.9. $\square$

Theorem 6.10. Assume that $N > 1$ and $j \in \{1, \dots, N\}$. Then, for every $f \in \dot{W}^{1,N}(\mathbb{R}^N)$,

$$(6.53) \quad \lim_{t \rightarrow \infty} t^{1/N} \mu(t, [R_{\text{Dunkl},j}, M_f]) = \left(\frac{1}{N(2\pi)^N} \int_{\Gamma} \int_{\mathbb{S}^{N-1}} \text{tr}_{\mathbb{C}^G} |\mathfrak{c}_{f,j}(x, \theta)|^N d\theta dx \right)^{1/N}.$$

Moreover,

$$(6.54) \quad \left(\int_{\Gamma} \int_{\mathbb{S}^{N-1}} \text{tr}_{\mathbb{C}^G} |\mathfrak{c}_{f,j}(x, \theta)|^N d\theta dx \right)^{1/N} \simeq_{N,R,k} \|\nabla f\|_{L^N(\mathbb{R}^N, dx)}.$$

Proof. By Lemma 6.9, choose $f_m \in C_c^\infty(\mathbb{R}^N)$ such that

$$(6.55) \quad \|[R_{\text{Dunkl},j}, M_{f_m}] - [R_{\text{Dunkl},j}, M_f]\|_{S^{N,\infty}} \longrightarrow 0,$$

$$(6.56) \quad \|\mathfrak{c}_{f_m,j} - \mathfrak{c}_{f,j}\|_{L^N(\Gamma \times \mathbb{S}^{N-1}; S^N(\mathbb{C}^G))} \longrightarrow 0.$$

By Lemma 6.7, for every m ,

$$\lim_{t \rightarrow \infty} t^{1/N} \mu(t, [R_{\text{Dunkl},j}, M_{f_m}]) = \frac{1}{(N(2\pi)^N)^{1/N}} \|\mathfrak{c}_{f_m,j}\|_{L^N(\Gamma \times \mathbb{S}^{N-1}; S^N(\mathbb{C}^G))}.$$

By (6.56),

$$\|\mathfrak{c}_{f_m,j}\|_{L^N(\Gamma \times \mathbb{S}^{N-1}; S^N(\mathbb{C}^G))} \longrightarrow \|\mathfrak{c}_{f,j}\|_{L^N(\Gamma \times \mathbb{S}^{N-1}; S^N(\mathbb{C}^G))}.$$

Combining this convergence with (6.55), we apply Lemma 6.8 with $p = N$ and obtain (6.53).

Finally, (6.54) follows from Lemma 6.5. This completes the proof of Theorem 6.10. $\square$

7. Alternative proof by using Dima's approach

Lemma 7.1. Fix a closed Weyl chamber Γ and $\sigma \in G \setminus \{\text{Id}\}$. Suppose that K satisfies

$$|\partial_x^\alpha \partial_y^\beta K(x, y)| \lesssim_{\alpha, \beta} \frac{\|x - y\|^{1-|\alpha|-|\beta|}}{\|x - \sigma y\| V(x, y, \|x - y\|)}$$

for every pair of multi-indices α, β and every $x, y \in \Gamma^\circ$ with $x \neq y$. For every $M_0 > 2N$, there exist positive numbers $\{c_{u,v}\}_{u,v \in \mathbb{Z}^N}$ and families $\{e_{u,v,I}\}_{u,v \in \mathbb{Z}^N, I \in \mathbb{D}_\Gamma}$ and $\{h_{u,v,I}\}_{u,v \in \mathbb{Z}^N, I \in \mathbb{D}_\Gamma}$ such that

$$(7.57) \quad |c_{u,v}| \lesssim_{M_0} (1 + |u| + |v|)^{-M_0},$$

$$(7.58) \quad \text{supp } e_{u,v,I} \subset I, \quad \text{supp } h_{u,v,I} \subset I^*,$$

$$(7.59) \quad |e_{u,v,I}| + |h_{u,v,I}| \lesssim w(I)^{-1/2},$$

and, for almost every $x, y \in \Gamma$ with $x \neq y$,

$$(7.60) \quad K(x, y) = \sum_{u,v \in \mathbb{Z}^N} c_{u,v} \sum_{I \in \mathbb{D}_\Gamma} \theta_{\text{Id}, \sigma, I} e_{u,v, I}(x) h_{u,v, I}(y).$$

Proof. Choose $\psi \in C_c^\infty(\mathbb{R}^N)$ such that

$$(7.61) \quad \text{supp } \psi \subset \left\{ z : \frac{1}{8} \leq |z| \leq \frac{1}{2} \right\}, \quad \sum_{m \in \mathbb{Z}} \psi(2^m z) = 1, \quad z \neq 0.$$

For $x, y \in \Gamma^\circ$, $x \neq y$, the unique cube $I \in \mathbb{D}_{\Gamma, m}$ containing x at each scale satisfies

$$(7.62) \quad K(x, y) = \sum_{I \in \mathbb{D}_\Gamma} \chi_I(x) K(x, y) \psi\left(\frac{\Phi(x) - \Phi(y)}{\ell(I)}\right).$$

If the I -th summand is nonzero, then $|\Phi(x) - \Phi(y)| \leq \ell(I)/2$; hence $y \in I^*$.

Fix $I = \Phi^{-1}(Q)$. For

$$a \in (-1/2, 1/2)^N, \quad c_Q + \ell(I)b \in (2Q) \cap \Omega,$$

set

$$X := \Phi^{-1}(c_Q + \ell(I)a), \quad Y := \Phi^{-1}(c_Q + \ell(I)b).$$

Whenever $\psi(a - b) \neq 0$, by the G -invariance of d and dw , (2.4), and Lemma ?? applied with (Id, σ) ,

$$(7.63) \quad \begin{aligned} \|X - Y\| &\simeq \ell(I), \\ \|X - \sigma Y\| &\simeq \ell(I) + |z_I - \sigma z_I|, \\ V(X, Y, \|X - Y\|) &\simeq w(I). \end{aligned}$$

Consequently,

$$(7.64) \quad \left| \partial_x^\alpha \partial_y^\beta K(X, Y) \right| \lesssim_{\alpha, \beta} \theta_{\text{Id}, \sigma, I} \frac{\ell(I)^{-|\alpha| - |\beta|}}{w(I)}.$$

Let

$$\mathcal{R}_Q := \left\{ (a, b) : a \in (-1/2, 1/2)^N, \ c_Q + \ell(I)b \in (2Q) \cap \Omega \right\}.$$

On $\mathcal{R}_Q$, define

$$F_I(a, b) := \frac{w(I)}{\theta_{\text{Id}, \sigma, I}} \psi(a - b) K\left(\Phi^{-1}(c_Q + \ell(I)a), \Phi^{-1}(c_Q + \ell(I)b)\right).$$

By (7.64) and the chain rule, for every integer $L \geq 0$,

$$(7.65) \quad \sup_{I \in \mathbb{D}_\Gamma} \|F_I\|_{W^{L, \infty}(\mathcal{R}_Q)} \lesssim_L 1.$$

The rectangles $\mathcal{R}_Q$ range over finitely many fixed rectangular domains. Fix an integer $L > M_0$. By the standard extension theorem for Sobolev spaces on rectangles, followed by multiplication by a fixed smooth cutoff, F_I admits an extension $\widetilde{F}_I$ supported in $(-2, 2)^N \times (-2, 2)^N$ such that

$$(7.66) \quad \widetilde{F}_I = F_I \quad \text{on } \mathcal{R}_Q, \quad \sup_{I \in \mathbb{D}_\Gamma} \|\widetilde{F}_I\|_{W^{L, \infty}(\mathbb{R}^{2N})} \lesssim_L 1.$$

The 4-periodic extension of $\widetilde{F}_I$ has a Fourier expansion

$$\widetilde{F}_I(a, b) = \sum_{u, v \in \mathbb{Z}^N} b_{u, v, I} e^{\frac{\pi i}{2} u \cdot a} e^{\frac{\pi i}{2} v \cdot b}.$$

By repeated integration by parts and (7.66),

$$\sup_{I \in \mathbb{D}_\Gamma} |b_{u, v, I}| \lesssim_L (1 + |u| + |v|)^{-L}.$$

Increasing the implicit constant if necessary, define

$$c_{u, v} := C_{M_0} (1 + |u| + |v|)^{-M_0}$$

so that $|b_{u, v, I}| \leq c_{u, v}$ for every I . For $I = \Phi^{-1}(Q)$, define

$$\begin{aligned} e_{u, v, I}(x) &:= \frac{b_{u, v, I}}{c_{u, v}} w(I)^{-1/2} \chi_I(x) \exp\left(\frac{\pi i}{2\ell(I)} u \cdot (\Phi(x) - c_Q)\right), \\ h_{u, v, I}(y) &:= w(I)^{-1/2} \chi_{I^*}(y) \exp\left(\frac{\pi i}{2\ell(I)} v \cdot (\Phi(y) - c_Q)\right). \end{aligned}$$

Then (7.57)–(7.59) hold.

If $x \in I$ and $y \in I^*$, then

$$a := \frac{\Phi(x) - c_Q}{\ell(I)}, \quad b := \frac{\Phi(y) - c_Q}{\ell(I)}$$

belong to $\mathcal{R}_Q$. Hence, by (7.66) and the Fourier expansion,

$$(7.67) \quad K(x, y) \psi\left(\frac{\Phi(x) - \Phi(y)}{\ell(I)}\right) = \sum_{u, v \in \mathbb{Z}^N} c_{u, v} \theta_{\text{Id}, \sigma, I} e_{u, v, I}(x) h_{u, v, I}(y).$$

If $x \in I$ and $y \notin I^*$, both sides of (7.67) vanish. Therefore,

$$\chi_I(x) K(x, y) \psi\left(\frac{\Phi(x) - \Phi(y)}{\ell(I)}\right) = \sum_{u, v \in \mathbb{Z}^N} c_{u, v} \theta_{\text{Id}, \sigma, I} e_{u, v, I}(x) h_{u, v, I}(y)$$

for almost every $x, y \in \Gamma$. Summing over $I \in \mathbb{D}_\Gamma$ and using (7.61), we obtain (7.60).

This completes the proof of Lemma 7.1. $\square$

Lemma 7.2. Assume that $N \geq 1$, $1 < p < \infty$, and $1 \leq q \leq \infty$. Fix $s > 2$, a closed Weyl chamber Γ , and $\sigma \in G$. Let r_s be as in Lemma ???. Then, for every $f \in OSC^{p, q}(\mathbb{R}^N)$,

$$(7.68) \quad \left\| \left\{ \theta_{\sigma, \text{Id}, I} \text{osc}_{r_s}(H_{\sigma, \text{Id}}(f); I) \right\}_{I \in \mathbb{D}_\Gamma} \right\|_{\ell^{p, q}} + \left\| \left\{ \theta_{\sigma, \text{Id}, I} a_{\sigma, \text{Id}, I}(f) \right\}_{I \in \mathbb{D}_\Gamma} \right\|_{\ell^{p, q}} \lesssim \|f\|_{OSC^{p, q}(\mathbb{R}^N)}.$$

The implicit constant is independent of f and σ .

Proof. By (??), $0 < \theta_{\sigma, \text{Id}, I} \leq 1$. Hence, by Lemma 4.3,

$$\left\| \left\{ \theta_{\sigma, \text{Id}, I} \text{osc}_{r_s}(H_{\sigma, \text{Id}}(f); I) \right\}_{I \in \mathbb{D}_\Gamma} \right\|_{\ell^{p, q}} \lesssim \left\| \left\{ \text{osc}_{r_s}(H_{\sigma, \text{Id}}(f); I) \right\}_{I \in \mathbb{D}_\Gamma} \right\|_{\ell^{p, q}} \lesssim \|f\|_{OSC^{p, q}(\mathbb{R}^N)}.$$

By Lemma 4.2,

$$\left\| \left\{ \theta_{\sigma, \text{Id}, I} a_{\sigma, \text{Id}, I}(f) \right\}_{I \in \mathbb{D}_\Gamma} \right\|_{\ell^{p, q}} \lesssim \|f\|_{OSC^{p, q}(\mathbb{R}^N)}.$$

Combining the two estimates, we obtain (7.68).

This completes the proof of Lemma 7.2. $\square$

Theorem 7.3. Assume that $N \geq 1$, $1 < p < \infty$, and $1 \leq q \leq \infty$. Fix a closed Weyl chamber Γ and $\sigma \in G \setminus \{\text{Id}\}$. Let T be an integral operator on $L^2(\Gamma, dw)$ whose kernel K satisfies

$$|\partial_x^\alpha \partial_y^\beta K(x, y)| \lesssim_{\alpha, \beta} \frac{\|x - y\|^{1-|\alpha|-|\beta|}}{\|x - \sigma y\| V(x, y, \|x - y\|)}$$

for every pair of multi-indices α, β and every $x, y \in \Gamma^\circ$ with $x \neq y$. Then, for every $f \in OSC^{p,q}(\mathbb{R}^N)$,

$$(7.69) \quad M_{f-f \circ \sigma} T \in S^{p,q}(L^2(\Gamma, dw)), \quad \|M_{f-f \circ \sigma} T\|_{S^{p,q}} \lesssim \|f\|_{OSC^{p,q}(\mathbb{R}^N)}.$$

Proof. Fix $s > 2$, let r_s be as in Lemma ??, and set

$$H := H_{\sigma, \text{Id}}(f) = f - f \circ \sigma.$$

For $I \in \mathbb{D}_\Gamma$, set

$$\omega_I := \text{osc}_{r_s}(H; I), \quad a_I := a_{\sigma, \text{Id}, I}(f) = (H)_{I, dx}.$$

By Lemma 7.2 (7.68),

$$(7.70) \quad \left\| \left\{ \theta_{\sigma, \text{Id}, I} \omega_I \right\}_{I \in \mathbb{D}_\Gamma} \right\|_{\ell^{p,q}} + \left\| \left\{ \theta_{\sigma, \text{Id}, I} a_I \right\}_{I \in \mathbb{D}_\Gamma} \right\|_{\ell^{p,q}} \lesssim \|f\|_{OSC^{p,q}(\mathbb{R}^N)}.$$

In particular, $\omega_I < \infty$ for every I . By Lemma ??, $H \in L_{\text{loc}}^s(\Gamma, dw)$.

Fix $M_0 > 2N$. By Lemma 7.1, for almost every $x, y \in \Gamma$ with $x \neq y$,

$$(7.71) \quad K(x, y) = \sum_{u, v \in \mathbb{Z}^N} c_{u, v} \sum_{I \in \mathbb{D}_\Gamma} \theta_{\text{Id}, \sigma, I} e_{u, v, I}(x) h_{u, v, I}(y),$$

where

$$\sum_{u, v \in \mathbb{Z}^N} |c_{u, v}| < \infty.$$

By (??), $\theta_{\text{Id}, \sigma, I} = \theta_{\sigma, \text{Id}, I}$.

For $u, v \in \mathbb{Z}^N$ and $I \in \mathbb{D}_\Gamma$, define

$$\widetilde{e}_{u, v, I}(x) := \begin{cases} \omega_I^{-1} (H(x) - a_I) e_{u, v, I}(x), & \omega_I > 0, \\ 0, & \omega_I = 0. \end{cases}$$

By Lemma 7.1 (7.59) and Lemma ??,

$$\|\widetilde{e}_{u, v, I}\|_{L^s(dw)} \lesssim w(I)^{-1/2} \omega_I^{-1} \left(\int_I |H - a_I|^s dw \right)^{1/s} \lesssim w(I)^{\frac{1}{s} - \frac{1}{2}}$$

when $\omega_I > 0$, while the same estimate is immediate when $\omega_I = 0$. Moreover, by Lemma 7.1 (7.58)–(7.59) and $w(I^*) \lesssim w(I)$,

$$\|e_{u, v, I}\|_{L^s(dw)} + \|h_{u, v, I}\|_{L^s(dw)} \lesssim w(I)^{\frac{1}{s} - \frac{1}{2}}.$$

Thus, for fixed u, v ,

$$\{\widetilde{e}_{u, v, I}\}_{I \in \mathbb{D}_\Gamma}, \quad \{e_{u, v, I}\}_{I \in \mathbb{D}_\Gamma}, \quad \{\overline{h_{u, v, I}}\}_{I \in \mathbb{D}_\Gamma}$$

are NWO families.

If $\omega_I = 0$, then $H = a_I$ dx -a.e. on I , and hence dw -a.e. on I . Therefore, in all cases,

$$(7.72) \quad \begin{aligned} & \theta_{\sigma, \text{Id}, I} H(x) e_{u,v,I}(x) h_{u,v,I}(y) \\ &= \theta_{\sigma, \text{Id}, I} \omega_I \widetilde{e}_{u,v,I}(x) h_{u,v,I}(y) + \theta_{\sigma, \text{Id}, I} a_I e_{u,v,I}(x) h_{u,v,I}(y). \end{aligned}$$

Hence, by Lemma 2.10 and (7.70),

$$(7.73) \quad \left\| \sum_{I \in \mathbb{D}_\Gamma} \theta_{\sigma, \text{Id}, I} \omega_I \langle \cdot, \overline{h_{u,v,I}} \rangle \widetilde{e}_{u,v,I} + \sum_{I \in \mathbb{D}_\Gamma} \theta_{\sigma, \text{Id}, I} a_I \langle \cdot, \overline{h_{u,v,I}} \rangle e_{u,v,I} \right\|_{S^{p,q}} \lesssim \|f\|_{OS^{p,q}(\mathbb{R}^N)},$$

uniformly in u, v .

Since $p > 1$, using an equivalent norm on $S^{p,q}$ when necessary, the series

$$(7.74) \quad S_f := \sum_{u,v \in \mathbb{Z}^N} c_{u,v} \sum_{I \in \mathbb{D}_\Gamma} \theta_{\sigma, \text{Id}, I} \left(\omega_I \langle \cdot, \overline{h_{u,v,I}} \rangle \widetilde{e}_{u,v,I} + a_I \langle \cdot, \overline{h_{u,v,I}} \rangle e_{u,v,I} \right)$$

converges in $S^{p,q}(L^2(\Gamma, dw))$. By (7.73),

$$(7.75) \quad \|S_f\|_{S^{p,q}} \lesssim \sum_{u,v \in \mathbb{Z}^N} |c_{u,v}| \|f\|_{OS^{p,q}(\mathbb{R}^N)} \lesssim \|f\|_{OS^{p,q}(\mathbb{R}^N)}.$$

We identify S_f with $M_H T$. Fix $r > p$. By $\ell^{p,q} \subset \ell^r$ and (7.70),

$$\left\{ \theta_{\sigma, \text{Id}, I} \omega_I \right\}_{I \in \mathbb{D}_\Gamma}, \quad \left\{ \theta_{\sigma, \text{Id}, I} a_I \right\}_{I \in \mathbb{D}_\Gamma} \in \ell^r.$$

For $m \in \mathbb{N}$, let $S_f^{(m)}$ be defined by (7.74) with the sum over I restricted to $\bigcup_{|k| \leq m} \mathbb{D}_{\Gamma,k}$. By Lemma 2.10, now with exponent r ,

$$\begin{aligned} \|S_f - S_f^{(m)}\|_{S^r} &\lesssim \sum_{u,v \in \mathbb{Z}^N} |c_{u,v}| \left(\left\| \left\{ \theta_{\sigma, \text{Id}, I} \omega_I \right\}_{I \in \bigcup_{|k| > m} \mathbb{D}_{\Gamma,k}} \right\|_{\ell^r} \right. \\ &\quad \left. + \left\| \left\{ \theta_{\sigma, \text{Id}, I} a_I \right\}_{I \in \bigcup_{|k| > m} \mathbb{D}_{\Gamma,k}} \right\|_{\ell^r} \right) \rightarrow 0. \end{aligned}$$

Let $\varphi \in C_c^\infty(\Gamma^\circ)$ and

$$\eta \in C_c^\infty(\Gamma^\circ \setminus \{x : \sigma x = x\}).$$

By the construction in the proof of Lemma 7.1 and (7.72),

$$(7.76) \quad \begin{aligned} \langle S_f^{(m)} \varphi, \eta \rangle &= \int_\Gamma \int_\Gamma H(x) K(x, y) \left(\sum_{|k| \leq m} \psi(2^k(\Phi(x) - \Phi(y))) \right) \\ &\quad \times \varphi(y) \overline{\eta(x)} dw(y) dw(x). \end{aligned}$$

Set

$$\delta := \inf_{x \in \text{supp } \eta} \|x - \sigma x\| > 0.$$

For $x \in \text{supp } \eta$ and $\|x - y\| < \delta/2$,

$$\|x - \sigma y\| \geq \|x - \sigma x\| - \|x - y\| \geq \frac{\delta}{2}.$$

Moreover, since $x, y \in \Gamma$, by (2.4) and the G -invariance of d ,

$$\|x - \sigma y\| \geq d(x, \sigma y) = d(x, y) = \|x - y\|.$$

Consequently, the kernel estimate and the doubling property imply

$$\int_{\substack{y \in \Gamma \\ \|x-y\| < \delta/2}} |K(x, y)| dw(y) \lesssim \sum_{j=1}^{\infty} \frac{2^{-j}\delta}{\delta} \frac{w(B(x, 2^{-j}\delta))}{w(B(x, 2^{-j-1}\delta))} \lesssim \sum_{j=1}^{\infty} 2^{-j} \lesssim 1$$

uniformly in $x \in \text{supp } \eta$. On $\text{supp } \eta \times \text{supp } \varphi$ with $\|x - y\| \geq \delta/2$, the same annular argument involves only finitely many scales. Hence,

$$(7.77) \quad \sup_{x \in \text{supp } \eta} \int_{\text{supp } \varphi} |K(x, y)| dw(y) < \infty.$$

Since $H \in L^s_{\text{loc}}(\Gamma, dw)$ with $s > 2$, $H\eta \in L^2(\Gamma, dw) \cap L^1(\Gamma, dw)$. The partial sums in (7.76) are uniformly bounded and converge to 1 for $x \neq y$. Therefore, by (7.77) and dominated convergence,

$$\begin{aligned} \lim_{m \rightarrow \infty} \langle S_f^{(m)} \varphi, \eta \rangle &= \int_{\Gamma} \int_{\Gamma} H(x) K(x, y) \varphi(y) \overline{\eta(x)} dw(y) dw(x) \\ &= \langle T\varphi, \overline{H}\eta \rangle = \langle M_H T\varphi, \eta \rangle. \end{aligned}$$

On the other hand, the convergence in S^r implies

$$\lim_{m \rightarrow \infty} \langle S_f^{(m)} \varphi, \eta \rangle = \langle S_f \varphi, \eta \rangle.$$

Since $\sigma \neq \text{Id}$, the fixed point space $\{x : \sigma x = x\}$ is a proper linear subspace and therefore has dw -measure zero. Thus, $C_c^\infty(\Gamma^\circ \setminus \{x : \sigma x = x\})$ is dense in $L^2(\Gamma, dw)$, and S_f is the bounded extension of $M_H T = M_{f \circ \sigma} T$. Combining this identification with (7.75), we obtain (7.69).

This completes the proof of Theorem 7.3. $\square$

8. Appendix

Lemma 8.1. Let $1 < s < \infty$, and let $\mathbb{D}^0$ be the standard dyadic grid in $\mathbb{R}^N$. Suppose that $F \in L^s(\mathbb{R}^N)$. Then there exists a constant $C > 0$ such that

$$\left\| \left\{ |Q|^{-1/s'} \int_Q |F(x)| dx \right\}_{Q \in \mathbb{D}^0} \right\|_{\ell^{s, \infty}} \leq C \|F\|_{L^s(\mathbb{R}^N)}.$$

Proof. Let $\mathcal{E} \subset \mathbb{D}^0$ be finite and set

$$G_{\mathcal{E}}(x) := \sum_{Q \in \mathcal{E}} |Q|^{-1/s'} \chi_Q(x).$$

Since the dyadic cubes containing any fixed x are totally ordered by inclusion, for each $x \in \bigcup_{Q \in \mathcal{E}} Q$ we may choose a cube $Q_x \in \mathcal{E}$ of minimal volume such that $x \in Q_x$. Every $Q \in \mathcal{E}$ containing x is then a dyadic ancestor of Q_x . Since the volume increases by the factor 2^N at each generation,

$$G_{\mathcal{E}}(x) = \sum_{\substack{Q \in \mathcal{E} \\ x \in Q}} |Q|^{-1/s'} \leq |Q_x|^{-1/s'} \sum_{k=0}^{\infty} 2^{-kN/s'} \lesssim_{N,s} \sup_{Q \in \mathcal{E}} |Q|^{-1/s'} \chi_Q(x).$$

Consequently,

$$\|G_{\mathcal{E}}\|_{L^{s'}(\mathbb{R}^N)}^{s'} \lesssim_{N,s} \int_{\mathbb{R}^N} \sup_{Q \in \mathcal{E}} |Q|^{-1} \chi_Q(x) dx \leq \int_{\mathbb{R}^N} \sum_{Q \in \mathcal{E}} |Q|^{-1} \chi_Q(x) dx = \#\mathcal{E}.$$

and hence,

$$\|G_{\mathcal{E}}\|_{L^{s'}(\mathbb{R}^N)} \lesssim_{N,s} (\#\mathcal{E})^{1/s'}.$$

By Hölder's inequality,

$$\sum_{Q \in \mathcal{E}} |Q|^{-1/s'} \int_Q |F(x)| dx = \int_{\mathbb{R}^N} |F(x)| G_{\mathcal{E}}(x) dx \lesssim_{N,s} \|F\|_{L^s(\mathbb{R}^N)} (\#\mathcal{E})^{1/s'}.$$

Fix $t > 0$ and let $\mathcal{E}$ be any finite subcollection of the dyadic cubes satisfying

$$|Q|^{-1/s'} \int_Q |F(x)| dx > t.$$

Then

$$t \#\mathcal{E} < \sum_{Q \in \mathcal{E}} |Q|^{-1/s'} \int_Q |F(x)| dx \lesssim_{N,s} \|F\|_{L^s(\mathbb{R}^N)} (\#\mathcal{E})^{1/s'},$$

so

$$\#\mathcal{E} \lesssim_{N,s} t^{-s} \|F\|_{L^s(\mathbb{R}^N)}^s.$$

Since $\mathcal{E}$ is arbitrary,

$$\#\left\{Q \in \mathbb{D}^0 : |Q|^{-1/s'} \int_Q |F(x)| dx > t\right\} \lesssim_{N,s} t^{-s} \|F\|_{L^s(\mathbb{R}^N)}^s, \quad t > 0.$$

The asserted weak ℓ^s estimate follows. This completes the proof of Lemma 8.1. $\square$

Lemma 8.2. Assume that $N > 1$. Then the following statements hold.

(1) For every $f \in L_{\text{loc}}^1(\mathbb{R}^N)$, we have

$$(8.1) \quad \|\nabla f\|_{L^N(\mathbb{R}^N)} \simeq \|f\|_{OSC^{N,\infty}(\mathbb{R}^N)}.$$

(2) If $1 \leq q < \infty$ and $f \in OSC^{N,q}(\mathbb{R}^N)$, then f is constant almost everywhere.

(3) If $1 \leq p < N$, $1 \leq q \leq \infty$, and $f \in OSC^{p,q}(\mathbb{R}^N)$, then f is constant almost everywhere.

Proof. This is a well-known result due to Rochberg–Semmes. $\square$

9. Answer to Dima's question

Lemma 9.1. Assume that $N > 1$, $j \in \{1, \dots, N\}$, and $f \in C_c^\infty(\mathbb{R}^N)$. Let Γ_1 and Γ_2 be Weyl chambers, and denote their interiors by Γ_1° and Γ_2° , respectively. Then there exist

$$\varphi_m \in C_c^\infty(\Gamma_1^\circ), \quad \psi_m \in C_c^\infty(\Gamma_2^\circ), \quad 0 \leq \varphi_m, \psi_m \leq 1,$$

such that

$$\varphi_m \longrightarrow \chi_{\Gamma_1^\circ}, \quad \psi_m \longrightarrow \chi_{\Gamma_2^\circ}$$

pointwise on $\mathbb{R}^N$, and

$$(9.2) \quad \lim_{m \rightarrow \infty} \|M_{\varphi_m}[R_{\text{Dunkl},j}, M_f]M_{\psi_m} - M_{\chi_{\Gamma_1}}[R_{\text{Dunkl},j}, M_f]M_{\chi_{\Gamma_2}}\|_{S^{N,\infty}} = 0.$$

Proof. Set

$$T_f := [R_{\text{Dunkl},j}, M_f].$$

We apply Lemma ?? with the fixed closed chamber $\Gamma = \Gamma_1$. Choose $\tau_0 \in G$ such that

$$\Gamma_2 = \tau_0 \Gamma_1.$$

By Lemma 8.2, $f \in OSC^{N,\infty}(\mathbb{R}^N)$, and hence the operator-series identification in the proof of Theorem ?? gives, with convergence in $S^{N,\infty}$,

$$T_f = \sum_{\varepsilon=0}^3 \sum_{\sigma, \tau \in G} \sum_{u, v \in \mathbb{Z}^N} c_{u,v} \sum_{I \in \mathbb{D}_{\Gamma_1}} A_{\sigma, \tau, I, \varepsilon}(f) \left\langle \cdot, \overline{H_{u,v,I,\varepsilon}^{\sigma, \tau}} \right\rangle E_{u,v,I,\varepsilon}^{\sigma, \tau}.$$

By Lemma ?? (??),

$$\text{supp } E_{u,v,I,\varepsilon}^{\sigma, \tau} \subset \sigma I, \quad \text{supp } H_{u,v,I,\varepsilon}^{\sigma, \tau} \subset \tau I^*.$$

Since distinct open Weyl chambers are disjoint and their boundaries have $d\mathbf{w}$ -measure zero,

$$(9.3) \quad M_{\chi_{\Gamma_1}} T_f M_{\chi_{\Gamma_2}} = \sum_{\varepsilon=0}^3 \sum_{u, v \in \mathbb{Z}^N} c_{u,v} T_{u,v,\varepsilon},$$

where

$$T_{u,v,\varepsilon} := \sum_{I \in \mathbb{D}_{\Gamma_1}} A_{I,\varepsilon} \left\langle \cdot, \overline{H_{I,\varepsilon}} \right\rangle E_{I,\varepsilon},$$

and, for brevity,

$$A_{I,\varepsilon} := A_{\text{Id}, \tau_0, I, \varepsilon}(f), \quad E_{I,\varepsilon} := E_{u,v,I,\varepsilon}^{\text{Id}, \tau_0}, \quad H_{I,\varepsilon} := H_{u,v,I,\varepsilon}^{\text{Id}, \tau_0}.$$

Thus

$$(9.4) \quad \text{supp } E_{I,\varepsilon} \subset I, \quad \text{supp } H_{I,\varepsilon} \subset \tau_0 I^*,$$

$$(9.5) \quad \|E_{I,\varepsilon}\|_{L^s(dw)} + \|H_{I,\varepsilon}\|_{L^s(dw)} \lesssim w(I)^{1/s-1/2}$$

for every fixed $s > 2$.

Fix $2 < s_0 < s$ and set

$$\gamma := \frac{1}{s_0} - \frac{1}{s} > 0.$$

Let $r_s > s$ be as in Lemma ??, and set

$$\beta := \frac{N}{r_s} > 0.$$

Exactly as in the coefficient estimate in the proof of the present lemma in its previous formulation, the definitions of $A_{\text{Id}, \tau_0, I, \varepsilon}(f)$ give

$$(9.6) \quad |A_{I,\varepsilon}| \lesssim_f \begin{cases} \ell(I), & \ell(I) \leq 1, \\ \ell(I)^{-\beta}, & \ell(I) \geq 1. \end{cases}$$

For $\varepsilon = 0, 1, 2$, this follows from the $L^{r_s}(dx)$ -oscillation defining the corresponding coefficient: the small-scale bound follows from the Lipschitz regularity of the relevant smooth function, whereas

$$\text{osc}_{r_s}(g; E) \lesssim |E|^{-1/r_s} \|g\|_{L^{r_s}} \lesssim_f \ell(I)^{-N/r_s}$$

when $\ell(I) \geq 1$. For $\varepsilon = 3$, if $D = |z_I - \tau_0 z_I|$, then for $\ell(I) \leq 1$,

$$|A_{I,3}| \lesssim_f \frac{\ell(I)}{\ell(I) + D} (\ell(I) + D) \lesssim_f \ell(I),$$

while, for $\ell(I) \geq 1$,

$$|A_{I,3}| \lesssim_f \ell(I)^{-N} \lesssim \ell(I)^{-\beta}.$$

Let

$$\Delta_1 = \{\alpha_1, \dots, \alpha_{N_0}\}$$

be the simple system associated with Γ_1 . For $1 \leq i \leq N_0$ and $0 < \delta \leq 1$, set

$$W_{i,\delta} := \{x \in \Gamma_1 : 0 \leq \langle x, \alpha_i \rangle \leq 2\delta\}.$$

As in the wall estimate in the previous proof,

$$(9.7) \quad \frac{w(W_{i,\delta} \cap I)}{w(I)} + \frac{w(W_{i,\delta} \cap I^*)}{w(I)} \lesssim \min\left(1, \frac{\delta}{\ell(I)}\right).$$

Indeed, the assertion is immediate for $\delta > \ell(I)/8$. If $\delta \leq \ell(I)/8$, write $I = \Phi^{-1}(Q)$. Whenever I or I^* meets $W_{i,\delta}$, the relevant i -th coordinate interval begins at the boundary of Ω . By Lemma 3.9 (3.5), the density of $\Phi_{\#}dw$ is nondecreasing in this coordinate. Hence the elementary estimate

$$\int_0^{2\delta} g(t) dt \lesssim \frac{\delta}{\ell(I)} \int g(t) dt$$

for nonnegative nondecreasing g , applied fiberwise on Q and $(2Q) \cap \Omega$, yields (9.7).

The simple system associated with Γ_2 is

$$\Delta_2 = \{\tau_0 \alpha_1, \dots, \tau_0 \alpha_{N_0}\}.$$

Set

$$\widetilde{W}_{i,\delta} := \tau_0 W_{i,\delta} = \{y \in \Gamma_2 : 0 \leq \langle y, \tau_0 \alpha_i \rangle \leq 2\delta\}.$$

Since dw is G -invariant, (9.7) also gives the corresponding estimate for $\widetilde{W}_{i,\delta} \cap \tau_0 I^*$.

Set

$$\omega_{I,\delta} := \min\left(1, \left(\frac{\delta}{\ell(I)}\right)^{\gamma}\right).$$

If $0 \leq a \leq 1$ and $\text{supp } a \subset W_{i,\delta}$, then (9.5), (9.7), and Hölder's inequality give

$$\|a E_{I,\varepsilon}\|_{L^{s_0}(dw)} \lesssim \omega_{I,\delta} w(I)^{1/s_0-1/2}.$$

Likewise, if $0 \leq b \leq 1$ and $\text{supp } b \subset \widetilde{W}_{i,\delta}$, then

$$\|b H_{I,\varepsilon}\|_{L^{s_0}(dw)} \lesssim \omega_{I,\delta} w(I)^{1/s_0-1/2}.$$

Here we use Lemma 2.10 in its standard two-family form: the two NWO families may be adapted to two dyadic systems related by a fixed measure-preserving Weyl-group element. The proof of Lemma 2.10 is unchanged, since the NWO estimates for the two families are used separately. Consequently,

$$(9.8) \quad \|M_a T_{u,v,\varepsilon}\|_{S^{N,\infty}} + \|T_{u,v,\varepsilon} M_b\|_{S^{N,\infty}} \lesssim \left\| \{A_{I,\varepsilon} \omega_{I,\delta}\}_{I \in \mathcal{I}_{i,\delta}} \right\|_{\ell^{N,\infty}},$$

where $\mathcal{I}_{i,\delta}$ denotes the cubes for which the corresponding localized factor is nonzero. In particular,

$$I \in \mathcal{I}_{i,\delta} \implies (I \cup I^*) \cap W_{i,\delta} \neq \emptyset.$$

If $\ell(I) \leq 1$ and $A_{I,\varepsilon} \neq 0$, then z_I belongs to a fixed bounded set depending only on f . Therefore, for every dyadic $0 < \ell \leq 1$,

$$(9.9) \quad \#\{I : \ell(I) = \ell, A_{I,\varepsilon} \neq 0, (I \cup I^*) \cap W_{i,\delta} \neq \emptyset\} \lesssim_f \left(1 + \frac{\delta}{\ell}\right) \ell^{-(N-1)}.$$

Indeed, in the Φ -coordinates there are $O(1 + \delta/\ell)$ admissible intervals in the i -th coordinate and $O(\ell^{-1})$ admissible intervals in each of the remaining $N - 1$ coordinates.

For $\ell(I) \leq \delta$, we have $\omega_{I,\delta} = 1$. Hence, by (9.6) and (9.9),

$$\begin{aligned} & \#\{I \in \mathcal{I}_{i,\delta} : \ell(I) \leq \delta, |A_{I,\varepsilon}\omega_{I,\delta}| > \lambda\} \\ & \lesssim_f \delta \sum_{\substack{\ell \leq \delta \\ \lambda \lesssim_f \ell}} \ell^{-N} \lesssim_f \delta \lambda^{-N}, \end{aligned}$$

and therefore

$$(9.10) \quad \left\| \{A_{I,\varepsilon}\omega_{I,\delta}\}_{I \in \mathcal{I}_{i,\delta}} \right\|_{\ell^{N,\infty}} \lesssim_f \delta^{1/N}.$$

If $\delta < \ell(I) \leq 1$, then

$$|A_{I,\varepsilon}\omega_{I,\delta}| \lesssim_f \delta^\gamma \ell(I)^{1-\gamma},$$

whereas (9.9) reduces to $O_f(\ell^{-(N-1)})$. Given $\lambda > 0$, put

$$L := \max\left(\delta, c_f \left(\frac{\lambda}{\delta^\gamma}\right)^{1/(1-\gamma)}\right).$$

Then

$$\#\{I \in \mathcal{I}_{i,\delta} : \delta < \ell(I) \leq 1, |A_{I,\varepsilon}\omega_{I,\delta}| > \lambda\} \lesssim_f L^{-(N-1)}.$$

If $L = \delta$, then $\lambda \lesssim_f \delta$, and

$$\lambda L^{-(N-1)/N} \lesssim_f \delta^{1/N}.$$

Otherwise, $\lambda \lesssim_f \delta^\gamma L^{1-\gamma}$, whence

$$\lambda L^{-(N-1)/N} \lesssim_f \delta^\gamma L^{1/N-\gamma} \lesssim_f \delta^{\min(\gamma, 1/N)}.$$

Thus

$$(9.11) \quad \left\| \{A_{I,\varepsilon}\omega_{I,\delta}\}_{I \in \mathcal{I}_{i,\delta}} \right\|_{\ell^{N,\infty}} \lesssim_f \delta^{\min(\gamma, 1/N)}.$$

For $\ell(I) \geq 1$, only $O_f(1)$ cubes at each dyadic scale have nonzero coefficients. Hence

$$\sum_{\substack{I \in \mathcal{I}_{i,\delta} \\ \ell(I) \geq 1}} |A_{I,\varepsilon}\omega_{I,\delta}|^N \lesssim_f \delta^{\gamma N} \sum_{\substack{\ell \geq 1 \\ \ell \text{ dyadic}}} \ell^{-N(\beta+\gamma)} \lesssim_f \delta^{\gamma N}.$$

Combining this estimate with (9.10) and (9.11) gives

$$(9.12) \quad \left\| \{A_{I,\varepsilon}\omega_{I,\delta}\}_{I \in \mathcal{I}_{i,\delta}} \right\|_{\ell^{N,\infty}} \lesssim_f \delta^\kappa, \quad \kappa := \min\left(\gamma, \frac{1}{N}\right).$$

It follows from (9.8) that

$$(9.13) \quad \|M_a T_{u,v,\varepsilon}\|_{S^{N,\infty}} + \|T_{u,v,\varepsilon} M_b\|_{S^{N,\infty}} \lesssim_f \delta^\kappa$$

whenever

$$0 \leq a \leq 1, \quad \text{supp } a \subset W_{i,\delta},$$

or, respectively,

$$0 \leq b \leq 1, \quad \text{supp } b \subset \widetilde{W}_{i,\delta}.$$

We next treat localization at infinity. Suppose $0 \leq a_R \leq 1$ is supported in

$$\Gamma_1 \cap \{|x| \geq R\}.$$

If $A_{I,\varepsilon} \neq 0$, then, as in the previous proof,

$$|z_I| \lesssim_f 1 + \ell(I).$$

Since $E_{I,\varepsilon}$ is supported in I , for sufficiently large R ,

$$A_{I,\varepsilon} \neq 0, \quad I \cap \{|x| \geq R\} \neq \emptyset$$

can occur only if

$$\ell(I) \gtrsim_f R.$$

At every such scale there are $O_f(1)$ cubes with nonzero coefficients. Therefore

$$\|M_{a_R} T_{u,v,\varepsilon}\|_{S^{N,\infty}} \lesssim_f \left(\sum_{\ell(I) \gtrsim_f R} |A_{I,\varepsilon}|^N \right)^{1/N} \lesssim_f R^{-\beta}.$$

If $0 \leq b_R \leq 1$ is supported in

$$\Gamma_2 \cap \{|x| \geq R\},$$

then

$$\text{supp } H_{I,\varepsilon} \subset \tau_0 I^*$$

and $|\tau_0 x| = |x|$. The same argument gives

$$(9.14) \quad \|M_{a_R} T_{u,v,\varepsilon}\|_{S^{N,\infty}} + \|T_{u,v,\varepsilon} M_{b_R}\|_{S^{N,\infty}} \lesssim_f R^{-\beta}.$$

Choose $\eta \in C^\infty(\mathbb{R})$, $0 \leq \eta \leq 1$, such that

$$\eta(t) = 0 \quad \text{for } t \leq 1, \quad \eta(t) = 1 \quad \text{for } t \geq 2,$$

and choose a radial $\zeta \in C_c^\infty(\mathbb{R}^N)$, $0 \leq \zeta \leq 1$, satisfying

$$\zeta = 1 \quad \text{on } B(0, 1), \quad \text{supp } \zeta \subset B(0, 2).$$

For $0 < \delta \leq 1$ and $R \geq 1$, define on $\mathbb{R}^N$

$$\begin{aligned} \varphi_{\delta,R}(x) &:= \zeta(x/R) \prod_{i=1}^{N_0} \eta\left(\frac{\langle x, \alpha_i \rangle}{\delta}\right), \\ \psi_{\delta,R}(x) &:= \zeta(x/R) \prod_{i=1}^{N_0} \eta\left(\frac{\langle x, \tau_0 \alpha_i \rangle}{\delta}\right). \end{aligned}$$

Then

$$\varphi_{\delta,R} \in C_c^\infty(\Gamma_1^\circ), \quad \psi_{\delta,R} \in C_c^\infty(\Gamma_2^\circ), \quad 0 \leq \varphi_{\delta,R}, \psi_{\delta,R} \leq 1.$$

On Γ_1 ,

$$1 - \varphi_{\delta,R} = 1 - \zeta(\cdot/R) + \sum_{i=1}^{N_0} \zeta(\cdot/R) \prod_{h < i} \eta\left(\frac{\langle \cdot, \alpha_h \rangle}{\delta}\right) \left(1 - \eta\left(\frac{\langle \cdot, \alpha_i \rangle}{\delta}\right)\right).$$

The first term is supported in $\Gamma_1 \cap \{|x| \geq R\}$, and the i -th term in the sum is supported in $W_{i,\delta}$. The analogous decomposition for $1 - \psi_{\delta,R}$ on Γ_2 has its i -th wall term supported in $\widetilde{W}_{i,\delta}$.

Since $N > 1$, $S^{N,\infty}$ is normable. Moreover, Lemma ?? (??) gives

$$\sum_{u,v \in \mathbb{Z}^N} |c_{u,v}| < \infty.$$

Hence, summing (9.13) and (9.14) in (9.13), over ε, u, v , yields

$$(9.15) \quad \begin{aligned} & \left\| M_{\chi_{\Gamma_1} - \varphi_{\delta,R}} T_f M_{\chi_{\Gamma_2}} \right\|_{S^{N,\infty}} + \left\| M_{\chi_{\Gamma_1}} T_f M_{\chi_{\Gamma_2} - \psi_{\delta,R}} \right\|_{S^{N,\infty}} \\ & \lesssim_f \delta^\kappa + R^{-\beta}. \end{aligned}$$

Since

$$M_{\varphi_{\delta,R}} = M_{\varphi_{\delta,R}} M_{\chi_{\Gamma_1}}, \quad M_{\psi_{\delta,R}} = M_{\chi_{\Gamma_2}} M_{\psi_{\delta,R}},$$

we have

$$\begin{aligned} & M_{\chi_{\Gamma_1}} T_f M_{\chi_{\Gamma_2}} - M_{\varphi_{\delta,R}} T_f M_{\psi_{\delta,R}} \\ & = M_{\chi_{\Gamma_1} - \varphi_{\delta,R}} T_f M_{\chi_{\Gamma_2}} + M_{\varphi_{\delta,R}} T_f M_{\chi_{\Gamma_2} - \psi_{\delta,R}}. \end{aligned}$$

By the ideal property of $S^{N,\infty}$ and (9.15),

$$\left\| M_{\varphi_{\delta,R}} T_f M_{\psi_{\delta,R}} - M_{\chi_{\Gamma_1}} T_f M_{\chi_{\Gamma_2}} \right\|_{S^{N,\infty}} \lesssim_f \delta^\kappa + R^{-\beta}.$$

Finally, set

$$\varphi_m := \varphi_{2^{-m}, 2^m}, \quad \psi_m := \psi_{2^{-m}, 2^m}.$$

For every $x \in \Gamma_1^\circ$,

$$\varphi_m(x) \longrightarrow 1,$$

whereas $\varphi_m(x) = 0$ for every $x \notin \Gamma_1^\circ$. Thus

$$\varphi_m \longrightarrow \chi_{\Gamma_1^\circ}$$

pointwise. Similarly,

$$\psi_m \longrightarrow \chi_{\Gamma_2^\circ}$$

pointwise. Moreover,

$$\left\| M_{\varphi_m} T_f M_{\psi_m} - M_{\chi_{\Gamma_1}} T_f M_{\chi_{\Gamma_2}} \right\|_{S^{N,\infty}} \lesssim_f 2^{-m\kappa} + 2^{-m\beta} \longrightarrow 0.$$

This proves (9.2). □

Lemma 9.2. Let Γ be a closed Weyl chamber, $\sigma \in G \setminus \{\text{Id}\}$, and let T satisfy the assumptions of Theorem 7.3. Assume $N > 1$ and $f \in C_c^\infty(\mathbb{R}^N)$. Set

$$H := f - f \circ \sigma.$$

Let

$$0 \leq f_m \leq 1, \quad f_m \in C_c^\infty(\Gamma^\circ),$$

and suppose that

$$f_m(x) \uparrow 1, \quad x \in \Gamma^\circ.$$

Then

$$\lim_{m \rightarrow \infty} \|M_{f_m H} T - M_H T\|_{S^{N,\infty}(L^2(\Gamma, dw))} = 0.$$

Proof. Fix $s > 2$, choose $2 < s_0 < s$, and set

$$\gamma := \frac{1}{s_0} - \frac{1}{s} > 0.$$

Let $r_s > s$ be as in Lemma ??, and put

$$\beta := \frac{N}{r_s}, \quad \kappa := \min\left(\gamma, \frac{1}{N}\right).$$

We first establish a localization estimate for $M_H T$. In the proof of Theorem 7.3, for fixed $u, v \in \mathbb{Z}^N$, one writes

$$M_H T = \sum_{\varepsilon=0}^1 \sum_{u,v \in \mathbb{Z}^N} c_{u,v} \sum_{I \in \mathbb{D}_\Gamma} A_{I,\varepsilon} \langle \cdot, \overline{F_{u,v,I,\varepsilon}} \rangle E_{u,v,I,\varepsilon},$$

with convergence in $S^{N,\infty}$, where

$$A_{I,0} = \theta_{\text{Id},\sigma,I} \text{osc}_{r_s}(H; I), \quad A_{I,1} = \theta_{\text{Id},\sigma,I}(H)_{I,dx},$$

and

$$\text{supp } E_{u,v,I,\varepsilon} \subset I, \quad \|E_{u,v,I,\varepsilon}\|_{L^s(dw)} + \|F_{u,v,I,\varepsilon}\|_{L^s(dw)} \lesssim w(I)^{1/s-1/2}.$$

We claim that

$$(9.16) \quad |A_{I,\varepsilon}| \lesssim_f \begin{cases} \ell(I), & \ell(I) \leq 1, \\ \ell(I)^{-\beta}, & \ell(I) \geq 1, \end{cases} \quad \varepsilon = 0, 1.$$

For $\varepsilon = 0$, this follows from the smoothness and compact support of H :

$$\text{osc}_{r_s}(H; I) \lesssim_f \ell(I), \quad \ell(I) \leq 1,$$

whereas

$$\text{osc}_{r_s}(H; I) \lesssim |I|^{-1/r_s} \|H\|_{L^{r_s}} \lesssim_f \ell(I)^{-\beta}, \quad \ell(I) \geq 1.$$

For $\varepsilon = 1$, put

$$D_I := |z_I - \sigma z_I|.$$

Since

$$H(z) = f(z) - f(\sigma z),$$

the Lipschitz regularity of f gives

$$|H(z)| \lesssim_f |z - \sigma z| \lesssim D_I + \ell(I), \quad z \in I.$$

Hence

$$|\theta_{\text{Id}, \sigma, I}(H)_{I, dx}| \lesssim_f \frac{\ell(I)}{\ell(I) + D_I} (\ell(I) + D_I) \lesssim_f \ell(I).$$

For $\ell(I) \geq 1$,

$$|\theta_{\text{Id}, \sigma, I}(H)_{I, dx}| \leq |I|^{-1} \|H\|_{L^1} \lesssim_f \ell(I)^{-N} \lesssim \ell(I)^{-\beta}.$$

Thus (9.16) holds.

Let

$$\Delta = \{\alpha_1, \dots, \alpha_{N_0}\}$$

be the simple system determining Γ , and set

$$W_{i, \delta} := \{x \in \Gamma : 0 \leq \langle x, \alpha_i \rangle \leq 2\delta\}.$$

The wall estimate used in the proof of Lemma 9.1 gives

$$(9.17) \quad \frac{w(W_{i, \delta} \cap I)}{w(I)} \lesssim \min\left(1, \frac{\delta}{\ell(I)}\right).$$

Consequently, if

$$0 \leq a \leq 1, \quad \text{supp } a \subset W_{i, \delta},$$

then Hölder's inequality gives

$$\|a E_{u, v, I, \varepsilon}\|_{L^{s_0}(dw)} \lesssim \omega_{I, \delta} w(I)^{1/s_0 - 1/2},$$

where

$$\omega_{I, \delta} := \min\left(1, \left(\frac{\delta}{\ell(I)}\right)^\gamma\right).$$

After normalization, Lemma 2.10 therefore yields

$$\left\| M_a \sum_{I \in \mathbb{D}_\Gamma} A_{I, \varepsilon} \langle \cdot, \overline{F_{u, v, I, \varepsilon}} \rangle E_{u, v, I, \varepsilon} \right\|_{S^{N, \infty}} \lesssim \left\| \{A_{I, \varepsilon} \omega_{I, \delta}\}_I \right\|_{\ell^{N, \infty}}.$$

Using (9.16) and exactly the dyadic counting argument in the proof of Lemma 9.1, one obtains

$$\left\| \{A_{I, \varepsilon} \omega_{I, \delta}\}_I \right\|_{\ell^{N, \infty}} \lesssim_f \delta^K.$$

Since

$$\sum_{u, v \in \mathbb{Z}^N} |c_{u, v}| < \infty,$$

it follows that

$$(9.18) \quad \|M_a M_H T\|_{S^{N, \infty}} \lesssim_f \delta^K$$

whenever a is supported in one of the strips $W_{i, \delta}$.

We also need localization at infinity. If

$$0 \leq a_R \leq 1, \quad \text{supp } a_R \subset \Gamma \cap \{|x| \geq R\},$$

then $A_{I, \varepsilon} \neq 0$ implies that I meets a fixed compact set depending only on f . Since $\text{supp } E_{u, v, I, \varepsilon} \subset I$, the additional condition

$$a_R E_{u, v, I, \varepsilon} \neq 0$$

forces

$$\ell(I) \gtrsim_f R.$$

There are only $O_f(1)$ such cubes at each dyadic scale, and (9.16) gives

$$(9.19) \quad \|M_{a_R} M_H T\|_{S^{N,\infty}} \lesssim_f R^{-\beta}.$$

Choose $\eta \in C^\infty(\mathbb{R})$, $0 \leq \eta \leq 1$, such that

$$\eta(t) = 0 \quad (t \leq 1), \quad \eta(t) = 1 \quad (t \geq 2),$$

and choose a radial $\zeta \in C_c^\infty(\mathbb{R}^N)$, $0 \leq \zeta \leq 1$, satisfying

$$\zeta = 1 \quad \text{on } B(0, 1), \quad \text{supp } \zeta \subset B(0, 2).$$

For $0 < \delta \leq 1$ and $R \geq 1$, set

$$\chi_{\delta,R}(x) := \zeta(x/R) \prod_{i=1}^{N_0} \eta\left(\frac{\langle x, \alpha_i \rangle}{\delta}\right), \quad x \in \Gamma.$$

Then

$$\chi_{\delta,R} \in C_c^\infty(\Gamma^\circ), \quad 0 \leq \chi_{\delta,R} \leq 1.$$

The function $1 - \chi_{\delta,R}$ is the sum of one function supported in

$$\Gamma \cap \{|x| \geq R\}$$

and finitely many functions supported in the strips $W_{i,\delta}$. Hence (9.18) and (9.19) imply

$$(9.20) \quad \|M_{1-\chi_{\delta,R}} M_H T\|_{S^{N,\infty}} \lesssim_f \delta^\kappa + R^{-\beta}.$$

We now return to the prescribed sequence f_m . Since

$$M_{f_m H} T - M_H T = -M_{1-f_m} M_H T,$$

it suffices to prove

$$\|M_{1-f_m} M_H T\|_{S^{N,\infty}} \longrightarrow 0.$$

Fix $\varepsilon > 0$. By (9.20), choose first $0 < \delta \leq 1$ and then $R \geq 1$ so that

$$\|M_{1-\chi_{\delta,R}} M_H T\|_{S^{N,\infty}} < \varepsilon.$$

Since

$$\text{supp } \chi_{\delta,R} \Subset \Gamma^\circ$$

and $f_m \uparrow 1$ pointwise on Γ° , Dini's theorem gives

$$\sup_{\text{supp } \chi_{\delta,R}} (1 - f_m) \longrightarrow 0.$$

Moreover,

$$1 - f_m = (1 - f_m)(1 - \chi_{\delta,R}) + (1 - f_m)\chi_{\delta,R}.$$

Therefore, by the ideal property of $S^{N,\infty}$,

$$\begin{aligned} \|M_{1-f_m} M_H T\|_{S^{N,\infty}} &\leq \|M_{1-f_m} M_{1-\chi_{\delta,R}} M_H T\|_{S^{N,\infty}} \\ &\quad + \|M_{(1-f_m)\chi_{\delta,R}} M_H T\|_{S^{N,\infty}} \\ &\leq \|M_{1-\chi_{\delta,R}} M_H T\|_{S^{N,\infty}} \end{aligned}$$

$$+ \sup_{\text{supp } \chi_{\delta,R}} (1 - f_m) \|M_H T\|_{S^{N,\infty}}.$$

The first term is $< \varepsilon$, while the second tends to zero. Hence

$$\limsup_{m \rightarrow \infty} \|M_{1-f_m} M_H T\|_{S^{N,\infty}} \leq \varepsilon.$$

Since $\varepsilon > 0$ is arbitrary,

$$\|M_{1-f_m} M_H T\|_{S^{N,\infty}} \longrightarrow 0.$$

Equivalently,

$$M_{f_m(f-f \circ \sigma)} T \longrightarrow M_{f-f \circ \sigma} T$$

in $S^{N,\infty}(L^2(\Gamma, dw))$. □

References

- [1] Z. Fan, M. Lacey, and J. Li. Schatten classes and commutators of Riesz transform on Heisenberg group and applications. *J. Fourier Anal. Appl.*, 29(2):Paper No. 17, 28, 2023. [8](#)
- [2] Z. Fan, M. Lacey, J. Li, M.N. Vempati, and B.D. Wick. Besov space, Schatten classes, and commutators of Riesz transforms associated with the Neumann Laplacian. *Michigan Math. J.*, 75(1):199–224, 2025. [8](#)
- [3] Z. Fan, M.T. Lacey, J. Li, and X. Xiong. Schatten–Lorentz characterization of Riesz transform commutator associated with Bessel operators. Preprint available at <https://arxiv.org/abs/2403.08249>, 2024. [8](#)
- [4] Y. Han, M. Lee, J. Li, and B.D. Wick. Riesz transforms and commutators in the Dunkl setting. *Anal. Math. Phys.*, 14(3):Paper No. 46, 32, 2024. [4](#), [6](#)
- [5] G. Heckman. Coxeter groups. Lecture notes available at <https://www.math.ru.nl/~heckman/CoxeterGroups.pdf>. Radboud University Nijmegen, 2018. [9](#), [10](#), [12](#), [26](#)
- [6] S. Helgason. Differential geometry, Lie groups, and symmetric spaces, volume 80 of Pure and Applied Mathematics. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York-London, 1978. [4](#)
- [7] T. Hytonen. Schatten properties of commutators on metric spaces. Preprint available at <https://arxiv.org/abs/2411.02613>, 2024. [8](#), [9](#)
- [8] M.T. Lacey, J. Li, and B.D. Wick. Schatten classes and commutator in the two weight setting, I. Hilbert transform. *Potential Anal.*, 60(2):875–894, 2024. [8](#)
- [9] J. Li, J. Pipher, and L.A. Ward. Dyadic structure theorems for multiparameter function spaces. *Rev. Mat. Iberoam.*, 31(3):767–797, 2015. [7](#)
- [10] J. Li, X. Xiong, and F. Yang. Schatten properties of Calderón–Zygmund singular integral commutator on stratified Lie groups. *J. Math. Pures Appl.* (9), 188:73–113, 2024. [8](#)
- [11] R. Rochberg and S. Semmes. Nearly weakly orthonormal sequences, singular value estimates, and Calderon–Zygmund operators. *J. Funct. Anal.*, 86(2):237–306, 1989. [1](#), [8](#), [9](#)
- [12] Z. Wei and H. Zhang. Complex median method and Schatten class membership of commutators. Preprint available at <https://arxiv.org/abs/2411.05810>, 2024. [8](#)
- [13] Z. Wei and H. Zhang. On the boundedness and Schatten class property of noncommutative martingale paraproducts and operator-valued commutators. Preprint available at <https://arxiv.org/abs/2407.14988>, 2024. [8](#)

Zhijie Fan, School of Mathematical Sciences, South China Normal University, Guangzhou 510631, China
Email address: fanzhj3@mail2.sysu.edu.cn

Ji Li, Department of Mathematics, Macquarie University, NSW2109, Australia
Email address: ji.li@mq.edu.au

Dmitriy Zanin, University of New South Wales, Kensington, NSW 2052, Australia
Email address: d.zanin@unsw.edu.au