

# SCHATTEN CLASS PROPERTIES OF DUNKL–RIESZ TRANSFORM COMMUTATORS

ZHIJIE FAN, JI LI, AND DMITRIY ZANIN

ABSTRACT.

## CONTENTS

|                                                    |    |
|----------------------------------------------------|----|
| 1. Introduction                                    | 1  |
| 1.1. Background and motivation                     | 1  |
| 1.2. Strategy of the proof                         | 2  |
| 1.3. Notation                                      | 2  |
| 2. Preliminaries                                   | 3  |
| 2.1. Harmonic analysis in the Dunkl setting        | 3  |
| 2.2. Schatten class                                | 5  |
| 2.3. Nearly weakly orthonormal sequences           | 5  |
| 3. Comparison of Sobolev norms                     | 5  |
| 4. Proof of Theorem 1.1 (i)                        | 10 |
| 5. Proof of restricted version of Theorem 1.1 (ii) | 12 |
| 6. Useful part of Zhijie argument                  | 15 |
| 7. Proof of Theorem 1.1 (ii)                       | 20 |
| 8. Constant characterization                       | 23 |
| 9. Global result from Hytonen                      | 24 |
| References                                         | 25 |

## 1. INTRODUCTION

### 1.1. Background and motivation.

**Theorem 1.1.** *Let  $R$  be a normalised root system in  $\mathbb{R}^N$  and let  $G$  be the corresponding finite reflection group. Let  $k : R \rightarrow \mathbb{R}_+$  be  $G$ -invariant multiplicity function. Let  $R_{\text{Dunkl},j}$  be the  $j$ -th Dunkl-Riesz transform corresponding to  $R$  and  $k$ . Let  $w$  be the Dunkl weight corresponding to  $R$  and  $k$ . Let  $f \in L_1^{\text{loc}}(\mathbb{R}^N)$ .*

- (i) *If  $f \in \dot{W}^{\frac{N}{p},p}(\mathbb{R}^N)$ ,  $p > N$ , then  $[R_{\text{Dunkl},j}, M_f] \in \mathcal{L}_p(L_2(\mathbb{R}^N, w(x)dx))$ . Furthermore,*

$$\|[R_{\text{Dunkl},j}, M_f]\|_{\mathcal{L}_p(L_2(\mathbb{R}^N, w(x)dx))} \leq c_{p,R,k} \|f\|_{\dot{W}^{\frac{N}{p},p}(\mathbb{R}^N)};$$

---

*Date:* May 8, 2026.

*2010 Mathematics Subject Classification.* 47B10, 42B20, 43A85.

*Key words and phrases.* Schatten class, Riesz transform commutators, Dunkl operator, Besov space, Nearly weakly orthogonal sequence.

(ii) If  $[R_{\text{Dunkl},j}, M_f] \in \mathcal{L}_p(L_2(\mathbb{R}^N, w(x)dx))$ ,  $p > N$ , then  $f \in \dot{W}^{\frac{N}{p},p}(\mathbb{R}^N)$ . Furthermore,

$$\|[R_{\text{Dunkl},j}, M_f]\|_{\mathcal{L}_p(L_2(\mathbb{R}^N, w(x)dx))} \leq c_{p,R,k}^{-1} \|f\|_{\dot{W}^{\frac{N}{p},p}(\mathbb{R}^N)};$$

(iii) If  $f \in \dot{W}^{1,N}(\mathbb{R}^N)$ , then  $[R_{\text{Dunkl},j}, M_f] \in \mathcal{L}_{N,\infty}(L_2(\mathbb{R}^N, w(x)dx))$ . Furthermore,

$$\|[R_{\text{Dunkl},j}, M_f]\|_{\mathcal{L}_{N,\infty}(L_2(\mathbb{R}^N, w(x)dx))} \leq c_{R,k} \|f\|_{\dot{W}^{1,N}(\mathbb{R}^N)};$$

(iv) If  $[R_{\text{Dunkl},j}, M_f] \in \mathcal{L}_{N,\infty}(L_2(\mathbb{R}^N, w(x)dx))$ , then  $f \in \dot{W}^{1,N}(\mathbb{R}^N)$ . Furthermore,

$$\|[R_{\text{Dunkl},j}, M_f]\|_{\mathcal{L}_{N,\infty}(L_2(\mathbb{R}^N, w(x)dx))} \geq c_{R,k}^{-1} \|f\|_{\dot{W}^{1,N}(\mathbb{R}^N)};$$

**1.2. Strategy of the proof.** In Section 3, we introduce our *first ad hoc* Sobolev space and demonstrate that its semi-norm is controlled by the usual Sobolev semi-norm. In Section 4, we use Janson-Wolff method to control the  $\|\cdot\|_{\mathcal{L}_p(L_2(\mathbb{R}^N, w(x)dx))}$ -norm of the commutator  $[R_{\text{Dunkl},j}, M_f]$  by the *first ad hoc* Sobolev semi-norm of  $f$ . Thus, we control the  $\|\cdot\|_{\mathcal{L}_p(L_2(\mathbb{R}^N, w(x)dx))}$ -norm of the commutator  $[R_{\text{Dunkl},j}, M_f]$  by the usual Sobolev semi-norm (i.e., we prove Theorem 1.1 (i)).

In Section 5, we control, in every chamber  $\Gamma$ , the Sobolev semi-norm of  $f$  via  $\|\cdot\|_{\mathcal{L}_p(L_2(\mathbb{R}^N, w(x)dx))}$ -norm of the commutator  $[R_{\text{Dunkl},j}, M_f]$ . This is done by verifying the assumptions in Theorem 1.4 in [6] and applying the latter theorem.

In Section 6, we introduce our *second ad hoc* Sobolev space and use the Rochberg-Semmes method to demonstrate that the semi-norm of the latter space is controlled by  $\|\cdot\|_{\mathcal{L}_p(L_2(\mathbb{R}^N, w(x)dx))}$ -norm of the commutator  $[R_{\text{Dunkl},j}, M_f]$ .

In Section 7, for every face of each chamber, we compare the boundary traces of  $f$  taken from the adjacent regions: the *interior trace* is computed by approaching the face from within the given chamber, while the *exterior trace* is obtained by approaching the same face from the neighboring chamber that shares the same face. Those boundary values are demonstrated to be the same. This allows us to show that the function  $f$  is globally Sobolev and that its Sobolev semi-norm is controlled by  $\|\cdot\|_{\mathcal{L}_p(L_2(\mathbb{R}^N, w(x)dx))}$ -norm of the commutator  $[R_{\text{Dunkl},j}, M_f]$  (i.e., we prove Theorem 1.1 (ii)).

**1.3. Notation.** Conventionally, we let  $\mathbb{N}$  denote the set of positive integers and set  $\mathbb{Z}_+ := \{0\} \cup \mathbb{N}$ . Throughout the paper, the symbols  $C$  and  $c$  denote positive constants (possibly varying from line to line) that may depend only on  $N$ ,  $p$ , and the structural data  $(R, k)$  of the Dunkl setting, but are independent of the main parameters (e.g., the symbol  $b$  and auxiliary decay factors). We write  $A \lesssim B$  to mean that  $A \leq CB$  for some constant  $C > 0$ , and  $A \sim B$  to mean that both  $A \lesssim B$  and  $B \lesssim A$  hold. For  $\lambda > 0$  and a cube  $Q$  with sidelength  $\ell(Q)$ , write  $\lambda Q$  for the cube concentric with  $Q$  and sidelength  $\lambda \ell(Q)$ . Similarly, for every ball  $B = B(x, r)$ , set  $\lambda B := B(x, \lambda r)$ . For any  $1 \leq p \leq \infty$ , we denote by  $p'$  the conjugate of  $p$ , which satisfies  $\frac{1}{p} + \frac{1}{p'} = 1$ . We denote the average of a function  $f$  over a  $\mu$ -measurable set  $E$  by

$$(1.1) \quad (f)_E := \int_E f(x) d\mu(x) := \frac{1}{\mu(E)} \int_E f(x) d\mu(x).$$

For a subset  $E \subset \mathbb{R}^N$ , we denote by  $\chi_E$  its characteristic function.

Denote by  $\{e_j\}_{j=1}^N$  the standard orthonormal basis on  $\mathbb{R}^N$ .

## 2. PRELIMINARIES

**2.1. Harmonic analysis in the Dunkl setting.** Let  $\mathbb{R}^N$  be the Euclidean space equipped with the usual scalar product  $\langle \cdot, \cdot \rangle$  and the induced norm  $\| \cdot \|$ . For a nonzero vector  $v \in \mathbb{R}^N$ , the reflection  $\sigma_v$  with respect to the hyperplane orthogonal to a normalised vector  $v$  is defined by

$$\sigma_v x := x - 2\langle x, v \rangle v.$$

A finite set  $R \subset \mathbb{R}^N \setminus \{0\}$  is called a root system if  $\sigma_v(R) = R$  for every  $v \in R$ . A root system  $R$  is called a normalized reduced root system if  $|v| = 1$  for every  $v \in R$ . The finite group  $G$  generated by the reflections  $\{\sigma_v\}_{v \in R}$  is called the Weyl group (reflection group) of the root system  $R$ . Let  $k : R \rightarrow [0, \infty)$  be a multiplicity function, which is a  $G$ -invariant function fixed throughout the paper.

Define the Dunkl measure  $w(x)dx$  on  $\mathbb{R}^N$  by

$$w(x) = \prod_{v \in R} |\langle x, v \rangle|^{k(v)}.$$

Let  $B(x, r) := \{y \in \mathbb{R}^N : |x - y| < r\}$  be the Euclidean ball centred at  $x$  with radius  $r > 0$ . Furthermore,

$$(2.1) \quad w(B(x, r)) \sim r^N \prod_{v \in R} (|\langle x, v \rangle| + r)^{k(v)},$$

where the implicit constant only depends on  $N, R$  and  $k$ . It follows from (2.1) that there exists a constant  $C \geq 1$  such that for every  $x \in \mathbb{R}^N$ ,

$$(2.2) \quad C^{-1} \left( \frac{r_2}{r_1} \right)^N \leq \frac{w(B(x, r_2))}{w(B(x, r_1))} \leq C \left( \frac{r_2}{r_1} \right)^{N'}, \text{ for } 0 < r_1 < r_2,$$

so the numbers  $N'$  and  $N$  are called the upper and lower homogeneous dimensions, respectively.

The Dunkl operators  $T_\xi$  are defined by

$$T_\xi f(x) := \partial_\xi f(x) + \sum_{v \in R} \frac{k(v)}{2} \langle v, \xi \rangle \frac{f(x) - f(\sigma_v(x))}{\langle v, x \rangle},$$

where  $\partial_\xi f$  denotes the directional derivative of  $f$  in the direction  $\xi$ . Furthermore, the Dunkl Laplacian is defined by

$$\Delta_{\text{Dunkl}} f(x) := \sum_{j=1}^N T_{e_j}^2 f(x) = \Delta_{\mathbb{R}^N} f(x) + \sum_{v \in R} k(v) \delta_v f(x),$$

where  $\Delta_{\mathbb{R}^N}$  is the Laplacian on  $\mathbb{R}^N$  and  $\delta_v$  is defined by

$$\delta_v f(x) := \frac{\partial_v f(x)}{\langle v, x \rangle} - \frac{f(x) - f(\sigma_v(x))}{\langle v, x \rangle^2}.$$

In this article, we consider the Dunkl-Riesz transforms defined by  $R_{\text{Dunkl}, j} := T_{e_j} (-\Delta_{\text{Dunkl}})^{-1/2}$ ,  $j = 1, 2, \dots, N$ .

**Definition 2.1.** The Weyl chambers are defined as the closures of connected components of the set

$$(2.3) \quad \{x \in \mathbb{R}^N : \langle x, v \rangle \neq 0 \text{ for all } v \in R\}.$$

Define

$$d(x, y) = \min_{\sigma \in G} |x - \sigma(y)|, \quad x, y \in \mathbb{R}^N.$$

Recall from [5, Chapter VII, proof of Theorem 2.12] that

$$(2.4) \quad d(x, y) = |x - y| \text{ if and only if } x, y \in \Gamma \text{ for some chamber } \Gamma.$$

Denote by  $K_{\text{Dunkl},j}(x, y)$  the integral kernel of  $R_{\text{Dunkl},j}$ . We recall the kernel estimates for the Riesz transforms in the Dunkl setting proved in [4]. Although these estimates may not be sharp (see the comment following Theorem 2.1 in [4, p. 10]), they suffice for our purposes.

**Lemma 2.2.** *There exists a constant  $C$  such that for  $j \in \{1, 2, \dots, N\}$  and every  $x, y$  with  $d(x, y) \neq 0$ ,*

$$|K_{\text{Dunkl},j}(x, y)| \leq \frac{d(x, y)}{|x - y|} \cdot \frac{C}{w(B(x, d(x, y)))}.$$

*Proof.* The proof is given in [4, Theorem 1.1].  $\square$

**Lemma 2.3.** *We have*

$$d(x, y)^2 \cdot w(x)w(y) \leq Cw^2(B(x, d(x, y))), \quad x, y \in \mathbb{R}^N.$$

*Proof.* Define equivalence relation  $\sim$  on  $R$  by saying  $v_1 \sim v_2$  if there exists  $g \in G$  with  $v_1 = gv_2$ . Let us split  $R$  into equivalence classes with respect to  $\sim$ . Since  $k$  is  $G$ -invariant, it follows that  $k$  is constant on each equivalence class. Note that  $G$  leaves each equivalence class invariant.

We, therefore, write

$$w(x) = \prod_{A \in R/\sim} \left( \prod_{v \in A} |\langle x, v \rangle| \right)^{k|A|}.$$

On the other hand,

$$w(B(x, r)) \geq c_{R,k}^{(1)} r^N \prod_{A \in R/\sim} \left( \prod_{v \in A} (r + |\langle x, v \rangle|) \right)^{k|A|}.$$

Let  $x \in \Gamma$  and  $y \in g\Gamma$ . Set  $z = g^{-1}y \in \Gamma$ . Note that  $d(x, y) = |x - z|$  and, therefore,

$$|\langle x, v \rangle| + d(x, y) = |\langle x, v \rangle| + |x - z| \geq 2^{-\frac{1}{2}} |\langle z, v \rangle|.$$

Thus,

$$\begin{aligned} \prod_{v \in A} (|\langle x, v \rangle| + d(x, y)) &\geq \prod_{v \in A} 2^{-\frac{1}{2}} |\langle z, v \rangle| = 2^{-\frac{1}{2}|A|} \prod_{v \in A} |\langle g^{-1}y, v \rangle| = \\ &= 2^{-\frac{1}{2}|A|} \prod_{v \in A} |\langle y, gv \rangle| = 2^{-\frac{1}{2}|A|} \prod_{v \in gA} |\langle y, v \rangle| = 2^{-\frac{1}{2}|A|} \prod_{v \in A} |\langle y, v \rangle|. \end{aligned}$$

Thus,

$$\begin{aligned} w(B(x, d(x, y))) &\geq c_{R,k}^{(1)} d^N(x, y) \prod_{A \in R/\sim} \left( \prod_{v \in A} (d(x, y) + |\langle x, v \rangle|) \right)^{k|A|} \geq \\ &\geq c_{R,k}^{(1)} d^N(x, y) \prod_{A \in R/\sim} \left( 2^{-\frac{1}{2}|A|} \prod_{v \in A} |\langle y, v \rangle| \right)^{k|A|} = c_{R,k}^{(2)} d^N(x, y) w(y). \end{aligned}$$

On the other hand,

$$w(B(x, d(x, y))) \geq c_{R,k}^{(1)} d^N(x, y) w(x).$$

Multiplying the last two equations, we complete the proof.  $\square$

**Lemma 2.4.** *Let  $1 \leq j \leq N$ . If  $x_j \geq y_j$ , then*

$$K_{\text{Dunkl},j}(x, y) \geq c_{R,k} \frac{x_j - y_j}{|x - y|} \cdot \frac{1}{w(B(x, |x - y|))}.$$

*If  $x_j \leq y_j$ , then*

$$K_{\text{Dunkl},j}(x, y) \leq c_{R,k} \frac{x_j - y_j}{|x - y|} \cdot \frac{1}{w(B(x, |x - y|))}.$$

*Proof.* The lower bound is identical to that in [4, Theorem 1.2].  $\square$

## 2.2. Schatten class. [Zhijie will add this later]

**2.3. Nearly weakly orthonormal sequences.** Rochberg and Semmes [11] introduced the notion of *nearly weakly orthonormal (NWO)* sequences of functions. This notion has since proved highly effective in characterizing Schatten class membership of singular integral commutators in a variety of contexts (see [3, 7, 10, 1, 2, 9, 11, 13, 12]). The properties of NWO sequences were originally investigated in the Euclidean setting (equipped with Lebesgue measure), but the extension to spaces of homogeneous type is straightforward once a system of dyadic cubes is available in this general framework (see [7, Section 13] for a complete proof). For our purposes, we adapt the extension to the Dunkl setting  $(\mathbb{R}^N, \|\cdot\|, w)$  rather than recalling the general definition of NWO sequences and the general extension to spaces of homogeneous type.

**Definition 2.5.** *Let  $\mathcal{D}$  be the dyadic lattice in  $\mathbb{R}^N$ . A sequence of functions  $\{e_Q\}_{Q \in \mathcal{D}}$  is called *nearly weakly orthonormal (NWO, in short) sequence* if*

- (i)  $\text{supp}(e_Q) \subset cQ$  for some positive constant  $c$  independent of  $Q$ .
- (ii)  $\|e_Q\|_{L_\infty(\mathbb{R}^N, w(x)dx)} \leq w(Q)^{-\frac{1}{2}}$ .

**Lemma 2.6.** *Let  $1 < p < \infty$  and let  $\mathcal{D}$  be the dyadic lattice in  $\mathbb{R}^N$ . If  $T \in \mathcal{L}_p(L_2(\mathbb{R}^N, w(x)dx))$ , then*

$$\left\| \left\{ \langle T e_Q, f_Q \rangle_{L_2(\mathbb{R}^N, w(x)dx)} \right\}_{Q \in \mathcal{D}} \right\|_{l_p(\mathcal{D})} \leq c_{N,p,w,c_1,c_2} \|T\|_{\mathcal{L}_p(L_2(\mathbb{R}^N, w(x)dx))}$$

*for every NWO systems  $\{e_Q\}_{Q \in \mathcal{D}}$  and  $\{f_Q\}_{Q \in \mathcal{D}}$  (with respective constants  $c_1$  and  $c_2$ ).*

*Proof.* reference to [11] is inappropriate. Rochberg and Semmes do not consider weighted setting. we absolutely must provide correct reference. This property can be found in [11, (1.10)] (see also [7, Section 7]).  $\square$

## 3. COMPARISON OF SOBOLEV NORMS

**Theorem 3.1.** *For every  $f \in \dot{W}^{\frac{N}{p},p}(\mathbb{R}^N)$ ,  $p > N$ , we have*

$$\left( \int_{\mathbb{R}^N \times \mathbb{R}^N} \left( \frac{d(x,y)}{|x-y|} \right)^p |f(x) - f(y)|^p \frac{dxdy}{d(x,y)^{2N}} \right)^{\frac{1}{p}} \leq c_{p,R} \|f\|_{\dot{W}^{\frac{N}{p},p}(\mathbb{R}^N)}.$$

**Lemma 3.2.** *For every convex polyhedral cone  $\Gamma$ , we have*

$$\int_{\Gamma} \frac{dx}{|x-y|^{2N}} \geq c_{\Gamma} \text{dist}(y, \Gamma)^{-N}, \quad y \notin \Gamma.$$

*Proof.* Find a point  $z \in \Gamma$  such that

$$|y - z| = \text{dist}(y, \Gamma).$$

We have  $z + \Gamma \subset \Gamma$ . Thus,

$$\int_{\Gamma} \frac{dx}{|x - y|^{2N}} \geq \int_{z+\Gamma} \frac{dx}{|x - y|^{2N}} = \int_{\Gamma} \frac{dx}{|x - (y - z)|^{2N}}.$$

Clearly,

$$|x - (y - z)| \leq |x| + |y - z|.$$

Thus,

$$\int_{\Gamma} \frac{dx}{|x - y|^{2N}} \geq \int_{\Gamma} \frac{dx}{(|x| + |y - z|)^{2N}} = |y - z|^{-N} \int_{\Gamma} \frac{dx}{(|x| + 1)^{2N}}.$$

□

**Lemma 3.3.** *If  $p \in (N, 2N)$ , then*

$$\sup_{a>0} \int_{\mathbb{R}^N} \frac{|x - ae_N|^{p-2N} dx}{(1 + |x|)^p} < \infty.$$

*Proof. Step 1:* we claim that, for every  $a > 0$ ,

$$\begin{aligned} & \int_{\mathbb{R}^N} \frac{|x - ae_N|^{p-2N} dx}{(1 + |x|)^p} \leq \\ & \leq 4^p \int_{\mathbb{R}^{N-1} \times (-a, a)} \frac{|x|^{p-2N} dx}{(1 + |x'| + a)^p} + 3^{2N-p} \int_{\mathbb{R}^N} \frac{|x|^{p-2N} dx}{(1 + |x|)^p}. \end{aligned}$$

We clearly have

$$\begin{aligned} & \int_{\mathbb{R}^N} \frac{|x - ae_N|^{p-2N} dx}{(1 + |x|)^p} = \\ & = \int_{\mathbb{R}^{N-1} \times (\frac{a}{2}, \frac{3a}{2})} \frac{|x - ae_N|^{p-2N} dx}{(1 + |x|)^p} + \int_{\mathbb{R}^{N-1} \times \mathbb{R} \setminus (\frac{a}{2}, \frac{3a}{2})} \frac{|x - ae_N|^{p-2N} dx}{(1 + |x|)^p}. \end{aligned}$$

If  $x_N \notin (\frac{a}{2}, \frac{3a}{2})$ , then  $|x_N - a| \geq \frac{1}{3}|x_N|$ . Hence,  $|x - ae_N| \geq \frac{1}{3}|x|$ . Thus,

$$\begin{aligned} & \int_{\mathbb{R}^{N-1} \times \mathbb{R} \setminus (\frac{a}{2}, \frac{3a}{2})} \frac{|x - ae_N|^{p-2N} dx}{(1 + |x|)^p} \leq \\ & \leq \int_{\mathbb{R}^{N-1} \times \mathbb{R} \setminus (\frac{a}{2}, \frac{3a}{2})} \frac{3^{2N-p} |x|^{p-2N} dx}{(1 + |x|)^p} \leq \int_{\mathbb{R}^N} \frac{3^{2N-p} |x|^{p-2N} dx}{(1 + |x|)^p}. \end{aligned}$$

If  $x_N \in (\frac{a}{2}, \frac{3a}{2})$ , then (here, we using the notation  $x' = (x_1, \dots, x_{N-1})$ )

$$1 + |x| \geq 1 + \max\{|x'|, \frac{1}{2}a\} \geq \frac{1}{4}(1 + |x'| + a).$$

Thus,

$$\begin{aligned} & \int_{\mathbb{R}^{N-1} \times (\frac{a}{2}, \frac{3a}{2})} \frac{|x - ae_N|^{p-2N} dx}{(1 + |x|)^p} \leq 4^p \int_{\mathbb{R}^{N-1} \times (\frac{a}{2}, \frac{3a}{2})} \frac{|x - ae_N|^{p-2N} dx}{(1 + |x'| + a)^p} = \\ & = 4^p \int_{\mathbb{R}^{N-1} \times (-\frac{a}{2}, \frac{a}{2})} \frac{|x|^{p-2N} dx}{(1 + |x'| + a)^p} \leq 4^p \int_{\mathbb{R}^{N-1} \times (-a, a)} \frac{|x|^{p-2N} dx}{(1 + |x'| + a)^p}. \end{aligned}$$

Combining all preceding estimates, we infer the claim in Step 1.

**Step 2:** we claim that

$$\sup_{a>0} \int_{\mathbb{R}^{N-1} \times (-a,a)} \frac{|x|^{p-2N} dx}{(1+|x'|+a)^p} < \infty.$$

Passing to polar coordinates in  $\mathbb{R}^{N-1}$ , we equivalently rewrite the claim as

$$\sup_{a>0} \int_{\mathbb{R}_+ \times (0,a)} \frac{(r^2 + x_N^2)^{\frac{p}{2}-N} r^{N-2} dr dx_N}{(1+r+a)^p} < \infty.$$

If  $0 < a < 1$ , then

$$\int_{\mathbb{R}_+ \times (0,a)} \frac{(r^2 + x_N^2)^{\frac{p}{2}-N} r^{N-2} dr dx_N}{(1+r+a)^p} \leq \int_{\mathbb{R}_+ \times (0,1)} \frac{(r^2 + x_N^2)^{\frac{p}{2}-N} r^{N-2} dr dx_N}{(1+r)^p}.$$

If  $a \geq 1$ , then

$$\begin{aligned} \int_{\mathbb{R}_+ \times (0,a)} \frac{(r^2 + x_N^2)^{\frac{p}{2}-N} r^{N-2} dr dx_N}{(1+r+a)^p} &\leq \int_{\mathbb{R}_+ \times (0,a)} \frac{(r^2 + x_N^2)^{\frac{p}{2}-N} r^{N-2} dr dx_N}{(r+a)^p} = \\ &= \int_{(a,\infty) \times (0,a)} \frac{(r^2 + x_N^2)^{\frac{p}{2}-N} r^{N-2} dr dx_N}{(r+a)^p} + \\ &+ \int_0^a \int_0^r \frac{(r^2 + x_N^2)^{\frac{p}{2}-N} r^{N-2} dx_N dr}{(r+a)^p} + \int_0^a \int_0^{x_N} \frac{(r^2 + x_N^2)^{\frac{p}{2}-N} r^{N-2} dr dx_N}{(r+a)^p} \leq \\ &\leq \int_{(a,\infty) \times (0,a)} \frac{r^{p-2N} r^{N-2} dr dx_N}{r^p} + \\ &+ \int_0^a \int_0^r \frac{r^{p-2N} r^{N-2} dx_N dr}{a^p} + \int_0^a \int_0^{x_N} \frac{x_N^{p-2N} r^{N-2} dr dx_N}{a^p} = \\ &= \int_a^\infty r^{-N-1} dr + a^{-p} \int_0^a r^{p-N-1} dr + \frac{1}{N-1} a^{-p} \int_0^a x_N^{p-N-1} dx_N = \\ &= \frac{1}{N} a^{-N} + \frac{1}{p-N} a^{-N} + \frac{1}{(N-1)(p-N)} a^{-N} \leq \frac{1}{N} + \frac{1}{p-N} + \frac{1}{(N-1)(p-N)}. \end{aligned}$$

This yields the claim in Step 2.

Combining Steps 1 and 2, we complete the proof of the lemma.  $\square$

**Lemma 3.4.** *If  $p \in (N, 2N)$ , then*

$$\int_{\mathbb{R}^{N-1} \times \mathbb{R}_+} \frac{|x-y|^{p-2N} dx}{|x+z|^p} \leq c_{p,N} z_N^{-N}, \quad y, z \in \mathbb{R}^{N-1} \times \mathbb{R}_+.$$

*Proof.* By dilation invariance, we may assume without loss of generality that  $z_N = 1$ . By translation invariance in  $\mathbb{R}^{N-1}$ , we may assume without loss of generality that  $z_1 = \dots = z_{N-1} = 0$ . We, therefore, have

$$|x+z| = |x+e_N| \geq \max\{1, |x|\} \geq \frac{1}{2}(1+|x|), \quad x \in \mathbb{R}^{N-1} \times \mathbb{R}_+.$$

Thus,

$$\begin{aligned} \int_{\mathbb{R}^{N-1} \times \mathbb{R}_+} \frac{|x-y|^{p-2N} dx}{|x+z|^p} &\leq 2^p \int_{\mathbb{R}^{N-1} \times \mathbb{R}_+} \frac{|x-y|^{p-2N} dx}{(1+|x|)^p} \leq \\ &= 2^p \int_{\mathbb{R}^N} \frac{|x-y|^{p-2N} dx}{(1+|x|)^p} = 2^p \int_{\mathbb{R}^N} \frac{|x-y|^{p-2N} dx}{(1+|x|)^p}. \end{aligned}$$

---

<sup>1</sup>It is exactly at this point we are using the assumption  $p < 2N$ .

The assertion follows from Lemma 3.3.  $\square$

We need the following elementary version of the Hahn-Banach theorem.

**Lemma 3.5.** *Let  $K \subset \mathbb{R}^N$  be a convex set and let  $y \notin K$ . Let  $z \in K$  be the point closest to  $y$ . Affine hyperplane  $P$  passing through  $z$  and orthogonal to  $y - z$  splits the space into two half-spaces, one of them containing  $y$ , the other one containing  $K$ .*

*Proof.* By translation invariance, we may assume without loss of generality that  $z = 0$ . By rotation and dilation invariance, we may assume without loss of generality that  $y = e_N$ . By assumption,  $\text{dist}(e_N, K) = 1$ . We need to show that  $K$  is contained in the (closed) lower half-plane.

Assume the contrary: let  $x \in K$  be such that  $x_N > 0$ . Since  $0 \in K$  and since  $K$  is convex, it follows that  $\epsilon x \in K$  for every  $\epsilon \in (0, 1)$ . Thus,

$$\text{dist}(e_N, K) \leq |e_N - \epsilon x|, \quad \epsilon \in (0, 1).$$

Since  $\epsilon \in (0, 1)$  is arbitrary, it follows that

$$\text{dist}(e_N, K) \leq \inf_{\epsilon \in (0, 1)} |e_N - \epsilon x| = \inf_{\epsilon \in (0, 1)} \left( \epsilon^2 \sum_{k=1}^{N-1} x_k^2 + (1 - \epsilon x_N)^2 \right)^{\frac{1}{2}}.$$

Since  $x_N > 0$ , it follows that infimum is strictly less than 1. Thus,  $\text{dist}(e_N, K) < 1$ . However, it is granted in the lemma that  $\text{dist}(e_N, K) = 1$ . This contradiction shows that such an  $x$  cannot exist. This completes the proof.  $\square$

**Lemma 3.6.** *If  $p \in (N, 2N)$ , then*

$$\int_{\Gamma} \frac{|x - y|^{p-2N} dx}{|x - gy|^p} \leq c_p \text{dist}(gy, \Gamma)^{-N}, \quad y \in \Gamma.$$

*Proof.* Find a point  $z \in \Gamma$  such that

$$|gy - z| = \text{dist}(gy, \Gamma).$$

By Lemma 3.5, affine hyperplane passing through  $z$  and orthogonal to  $gy - z$  splits the space into two half-spaces, one of them containing  $gy$ , the other one containing  $\Gamma$  (denote this one by  $H$ ).

We have  $y \in H$ ,  $\Gamma \subset H$  and  $gy \notin H$ . So,

$$\int_{\Gamma} \frac{|x - y|^{p-2N} dx}{|x - gy|^p} \leq \int_H \frac{|x - y|^{p-2N} dx}{|x - gy|^p}.$$

Using Lemma 3.4, we write

$$\int_H \frac{|x - y|^{p-2N} dx}{|x - gy|^p} \leq c_{p,N} \text{dist}(gy, H)^{-N}.$$

Since  $z \in \partial H$  and since  $gy - z$  is orthogonal to  $\partial H$ , it follows that  $\text{dist}(gy, H) = |gy - z| = \text{dist}(gy, \Gamma)$ .  $\square$

*Proof of Theorem 3.1. Case 1:* Let  $p \in (N, 2N)$ .

We have

$$\begin{aligned} & \left( \int_{\Gamma \times g\Gamma} \left( \frac{d(x, y)}{|x - y|} \right)^p |f(x) - f(y)|^p \frac{dxdy}{d(x, y)^{2N}} \right)^{\frac{1}{p}} = \\ & = \left( \int_{\Gamma \times \Gamma} \left( \frac{|x - y|}{|x - gy|} \right)^p |f(x) - f(gy)|^p \frac{dxdy}{|x - y|^{2N}} \right)^{\frac{1}{p}} \leq \end{aligned}$$

$$\begin{aligned}
&\leq \left( \int_{\Gamma \times \Gamma} \left( \frac{|x-y|}{|x-gy|} \right)^p |f(x) - f(y)|^p \frac{dxdy}{|x-y|^{2N}} \right)^{\frac{1}{p}} + \\
&+ \left( \int_{\Gamma \times \Gamma} \left( \frac{|x-y|}{|x-gy|} \right)^p |f(y) - f(gy)|^p \frac{dxdy}{|x-y|^{2N}} \right)^{\frac{1}{p}} \leq \\
&\leq \left( \int_{\Gamma \times \Gamma} |f(x) - f(y)|^p \frac{dxdy}{|x-y|^{2N}} \right)^{\frac{1}{p}} + \\
&+ \left( \int_{\Gamma} |f(y) - f(gy)|^p \left( \int_{\Gamma} \frac{|x-y|^{p-2N}}{|x-gy|^p} dx \right) dy \right)^{\frac{1}{p}}.
\end{aligned}$$

Obviously, the first summand on the right hand side is less than  $\|f\|_{\dot{W}^{\frac{N}{p},p}(\mathbb{R}^N)}$ . By Lemmas 3.6 and 3.2,

$$\begin{aligned}
&\left( \int_{\Gamma \times g\Gamma} \left( \frac{d(x,y)}{|x-y|} \right)^p |f(x) - f(y)|^p \frac{dxdy}{d(x,y)^{2N}} \right)^{\frac{1}{p}} \leq \|f\|_{\dot{W}^{\frac{N}{p},p}(\mathbb{R}^N)} + \\
&+ c_{p,R}^{(1)} \left( \int_{\Gamma} |f(y) - f(gy)|^p \left( \int_{\Gamma} \frac{dx}{|x-gy|^{2N}} \right) dy \right)^{\frac{1}{p}} = \\
&= \|f\|_{\dot{W}^{\frac{N}{p},p}(\mathbb{R}^N)} + c_{p,R}^{\frac{1}{p}} \left( \int_{\Gamma \times \Gamma} \frac{|f(y) - f(gy)|^p dxdy}{|x-gy|^{2N}} \right)^{\frac{1}{p}} \leq \\
&\leq \|f\|_{\dot{W}^{\frac{N}{p},p}(\mathbb{R}^N)} + c_{p,R}^{(1)} \left( \int_{\Gamma \times \Gamma} \frac{|f(x) - f(gy)|^p dxdy}{|x-gy|^{2N}} \right)^{\frac{1}{p}} + \\
&+ c_{p,R}^{(1)} \left( \int_{\Gamma \times \Gamma} \frac{|f(x) - f(y)|^p dxdy}{|x-gy|^{2N}} \right)^{\frac{1}{p}}.
\end{aligned}$$

Now,

$$\left( \int_{\Gamma \times \Gamma} \frac{|f(x) - f(gy)|^p dxdy}{|x-gy|^{2N}} \right)^{\frac{1}{p}} = \left( \int_{\Gamma \times g\Gamma} \frac{|f(x) - f(y)|^p dxdy}{|x-y|^{2N}} \right)^{\frac{1}{p}} \leq \|f\|_{\dot{W}^{\frac{N}{p},p}(\mathbb{R}^N)}$$

and

$$\left( \int_{\Gamma \times \Gamma} \frac{|f(x) - f(y)|^p dxdy}{|x-gy|^{2N}} \right)^{\frac{1}{p}} \leq \left( \int_{\Gamma \times \Gamma} \frac{|f(x) - f(y)|^p dxdy}{|x-y|^{2N}} \right)^{\frac{1}{p}} \leq \|f\|_{\dot{W}^{\frac{N}{p},p}(\mathbb{R}^N)}.$$

Thus,

$$\left( \int_{\Gamma \times g\Gamma} \left( \frac{d(x,y)}{|x-y|} \right)^p |f(x) - f(y)|^p \frac{dxdy}{d(x,y)^{2N}} \right)^{\frac{1}{p}} \leq (1 + 2c_{p,R}^{(1)}) \|f\|_{\dot{W}^{\frac{N}{p},p}(\mathbb{R}^N)}.$$

Summing over  $\Gamma$  and  $1 \neq g \in G$ , we obtain

$$\left( \int_{\mathbb{R}^N \times \mathbb{R}^N} \left( \frac{d(x,y)}{|x-y|} \right)^p |f(x) - f(y)|^p \frac{dxdy}{d(x,y)^{2N}} \right)^{\frac{1}{p}} \leq |G|^2 (1 + 2c_{p,R}^{(1)}) \|f\|_{\dot{W}^{\frac{N}{p},p}(\mathbb{R}^N)}.$$

**Case 2:** Let  $p \geq 2N$ . We have

$$\begin{aligned}
&\left( \int_{\Gamma \times g\Gamma} \left( \frac{d(x,y)}{|x-y|} \right)^p |f(x) - f(y)|^p \frac{dxdy}{d(x,y)^{2N}} \right)^{\frac{1}{p}} \leq \\
&\leq \left( \int_{\Gamma \times g\Gamma} \left( \frac{d(x,y)}{|x-y|} \right)^{2N} |f(x) - f(y)|^p \frac{dxdy}{d(x,y)^{2N}} \right)^{\frac{1}{p}} = \|f\|_{\dot{W}^{\frac{N}{p},p}(\mathbb{R}^N)}.
\end{aligned}$$

□

## 4. PROOF OF THEOREM 1.1 (i)

Let  $(X, \mu)$  be a measure space and let  $L_p(L_{q,\infty})^{\text{symm}}(X \times X, \mu \times \mu)$  denote the mixed-norm space consisting of all  $\mu \times \mu$ -measurable functions  $f : X \times X \rightarrow \mathbb{C}$  such that

$$\int_X \|f(x, \cdot)\|_{L_{q,\infty}(X,\mu)}^p d\mu(x), \int_X \|f(\cdot, y)\|_{L_{q,\infty}(X,\mu)}^p d\mu(y) < \infty.$$

We equip  $L_p(L_{q,\infty})^{\text{symm}}(X \times X, \mu \times \mu)$  with its natural norm.

**Lemma 4.1.** *Let  $(X, \mu)$  be a measure space and let  $2 < p < \infty$ . Suppose that  $K \in L_p(L_{p',\infty})^{\text{symm}}(X \times X, \mu \times \mu)$ . The integral operator*

$$T_K f(x) := \int_X K(x, y) f(y) d\mu(y)$$

*belongs to  $S_{p,\infty}(L_2(X, d\mu))$ . Moreover, we have*

$$\|T_K\|_{S_{p,\infty}(L_2(X, d\mu))} \leq c_p \|K\|_{L_p(L_{p',\infty})^{\text{symm}}(X \times X, \mu \times \mu)}.$$

*Proof.* The proof is given in [8, Lemma 2] (see also [7, Corollary 11.2]).  $\square$

**Lemma 4.2.** *Let  $p > 2$  and let  $\frac{1}{q} = 1 - \frac{2}{p}$ . We have*

$$\|F_1 F_2\|_{L_p(L_{p',\infty})(X \times X, \mu \times \mu)} \leq 2^{\frac{1}{pp'}} \|F_1\|_{L_p(X \times X, \mu \times \mu)} \|F_2\|_{L_\infty(L_{q,\infty})(X \times X, \mu \times \mu)}.$$

*Proof.* We clearly have

$$LHS = \left( \int_X \|F_1(x, \cdot) F_2(x, \cdot)\|_{L_{p',\infty}(X,\mu)}^p d\mu(x) \right)^{\frac{1}{p}}.$$

Clearly,  $\frac{1}{p'} = \frac{1}{p} + \frac{1}{q}$ . Thus,

$$\begin{aligned} \|F_1(x, \cdot) F_2(x, \cdot)\|_{L_{p',\infty}(X,\mu)} &\leq 2^{\frac{1}{p'}} \|F_1(x, \cdot)\|_{L_{p,\infty}(X,\mu)} \|F_2(x, \cdot)\|_{L_{q,\infty}(X,\mu)} \\ &\leq 2^{\frac{1}{p'}} \|F_1(x, \cdot)\|_{L_p(X,\mu)} \|F_2(x, \cdot)\|_{L_{q,\infty}(X,\mu)}. \end{aligned}$$

Hence,

$$\begin{aligned} LHS &\leq 2^{\frac{1}{pp'}} \left( \int_X \|F_1(x, \cdot)\|_{L_p(X,\mu)}^p \|F_2(x, \cdot)\|_{L_{q,\infty}(X,\mu)}^p d\mu(x) \right)^{\frac{1}{p}} \leq \\ &\leq 2^{\frac{1}{pp'}} \left( \int_X \|F_1(x, \cdot)\|_{L_p(X,\mu)}^p d\mu(x) \right)^{\frac{1}{p}} \times \sup_{x \in X} \|F_2(x, \cdot)\|_{L_{q,\infty}(X,\mu)} = \\ &= 2^{\frac{1}{pp'}} \|F_1\|_{L_p(X \times X, \mu \times \mu)} \|F_2\|_{L_\infty(L_{q,\infty})(X \times X, \mu \times \mu)}. \end{aligned}$$

$\square$

**Lemma 4.3.** *Let  $(X, \mu)$  be a measure space and let  $2 < p < \infty$ . Suppose that  $F \in L_p(X \times X, H^2(\mu \times \mu))$  and  $H \in L_\infty(L_{1,\infty})^{\text{symm}}(X \times X, \mu \times \mu)$ . We have*

$$\|T_{FH}\|_{S_{p,\infty}(L_2(X, d\mu))} \leq c_p \|F\|_{L_p(X \times X, H^2(\mu \times \mu))} \|H\|_{L_\infty(L_{1,\infty})^{\text{symm}}(X \times X, \mu \times \mu)}^{1-\frac{2}{p}}.$$

*Proof.* By Lemma 4.1,

$$\|T_{FH}\|_{S_{p,\infty}(L_2(X, d\mu))} \leq c_p \|FH\|_{L_p(L_{p',\infty})^{\text{symm}}(X \times X, \mu \times \mu)}.$$

Let  $\frac{1}{q} = 1 - \frac{2}{p}$ . It follows from Lemma 4.2 that

$$\|F_1 F_2\|_{L_p(L_{p',\infty})^{\text{symm}}(X \times X, \mu \times \mu)} \leq$$

$$\leq 2^{\frac{1}{pp'}} \|F_1\|_{L_p(X \times X, \mu \times \mu)} \|F_2\|_{L_\infty(L_{q,\infty})^{\text{symm}}(X \times X, \mu \times \mu)}.$$

Setting  $F_1 = FH^{\frac{2}{p}}$  and  $F_2 = H^{\frac{1}{q}}$ , we obtain

$$\begin{aligned} & \|FH\|_{L_p(L_{p',\infty})^{\text{symm}}(X \times X, \mu \times \mu)} \leq \\ & \leq 2^{\frac{1}{pp'}} \|FH^{\frac{2}{p}}\|_{L_p(X \times X, \mu \times \mu)} \|H^{\frac{1}{q}}\|_{L_\infty(L_{q,\infty})^{\text{symm}}(X \times X, \mu \times \mu)} = \\ & = 2^{\frac{1}{pp'}} \|F\|_{L_p(X \times X, H^2(\mu \times \mu))} \|H\|_{L_\infty(L_{1,\infty})^{\text{symm}}(X \times X, \mu \times \mu)} = \\ & = 2^{\frac{1}{pp'}} \|F\|_{L_p(X \times X, H^2(\mu \times \mu))} \|H\|_{L_\infty(L_{1,\infty})^{\text{symm}}(X \times X, \mu \times \mu)}^{1-\frac{2}{p}}. \end{aligned}$$

□

**Lemma 4.4.** *Let  $(X, \mu)$  be a measure space and let  $2 < p < \infty$ . Suppose that  $F \in L_p(X \times X, GH^2(\mu \times \mu))$  and  $H \in L_\infty(L_{1,\infty})^{\text{symm}}(X \times X, \mu \times \mu)$ . We have*

$$\|T_{FH}\|_{S_p(L_2(X, d\mu))} \leq c_{p,H} \|F\|_{L_p(X \times X, H^2(\mu \times \mu))}.$$

*Proof.* The assertion follows from Lemma 4.3 by real interpolation. □

The following proposition, strictly speaking, is not part of the proof of Theorem 1.1 (i). Instead, it features in the proof of Theorem 1.1 (ii) in the next section. However, in terms of the employed technique, it certainly belongs to this section.

**Proposition 4.5.** *Let  $p > 2$ , let  $\Gamma$  be a chamber and let  $f \in \dot{W}^{\frac{N}{p},p}(\Gamma)$ . Let  $V$  be an integral operator whose integral kernel  $K$  satisfies the condition*

$$|K(x, y)| \leq \frac{1}{w(B(x, |x - y|))}.$$

*For every  $1 \leq j \leq N$ , we have*

$$\|[V, M_f]\|_{\mathcal{L}_p(L_2(\Gamma, w(x)dx))} \leq c_{R,k} \|f\|_{\dot{W}^{\frac{N}{p},p}(\Gamma)}.$$

*Proof.* In what follows, let  $X = \Gamma$  and  $d\mu(x) = w(x)dx$ . Set

$$F(x, y) = w(B(x, |x - y|))K(x, y)(f(x) - f(y)), \quad x \neq y,$$

$$H(x, y) = \frac{1}{w(B(x, |x - y|))}, \quad x \neq y.$$

By [7, Lemma 5.7],  $H \in L_\infty(L_{1,\infty})^{\text{symm}}(\Gamma \times \Gamma, \mu \times \mu)$ . By assumption on the kernel  $K$ ,

$$\|F\|_{L_p(\Gamma \times \Gamma, G^2(\mu \times \mu))} \leq \left( \int_{\Gamma \times \Gamma} |f(x) - f(y)|^p \frac{w(x)w(y)dxdy}{w^2(B(x, |x - y|))} \right)^{\frac{1}{p}}.$$

By (2.1),

$$w(B(x, |x - y|)) \geq c'_{R,k} |x - y|^N w^{\frac{1}{2}}(x) w^{\frac{1}{2}}(y), \quad x, y \in \mathbb{R}^N.$$

Thus,

$$\begin{aligned} & \|F\|_{L_p(\Gamma \times \Gamma, H^2(\mu \times \mu))} \leq \\ & \leq (c'_{R,k})^{-\frac{2}{p}} \left( \int_{\Gamma \times \Gamma} \frac{|f(x) - f(y)|^p}{|x - y|^{2N}} dxdy \right)^{\frac{1}{p}} = (c'_{R,k})^{-\frac{2}{p}} \|f\|_{\dot{W}^{\frac{N}{p},p}(\Gamma)}. \end{aligned}$$

Obviously, the integral kernel of the operator  $[V, M_f]$  is  $FH$ . The assertion follows now from Lemma 4.4. □

*Proof of Theorem 1.1 (i).* In what follows, let  $X = \mathbb{R}^N$  and  $d\mu(x) = w(x)dx$ . Set

$$F(x, y) = w(B(x, d(x, y)))K_{\text{Dunkl},j}(x, y)(f(x) - f(y)), \quad x \neq y,$$

$$H(x, y) = \frac{1}{w(B(x, d(x, y)))}, \quad x \neq y.$$

By [7, Lemma 11.7],  $H \in L_\infty(L_{1,\infty})^{\text{symm}}(X \times X, \mu \times \mu)$ . By Lemma 2.2,

$$\begin{aligned} & \|F\|_{L_p(X \times X, H^2(\mu \times \mu))} \leq \\ & \leq c_{R,k}^{(1)} \left( \int_{\mathbb{R}^N \times \mathbb{R}^N} \left( \frac{d(x, y)}{|x - y|} \right)^p |f(x) - f(y)|^p \frac{w(x)w(y)dx dy}{w(B(x, d(x, y)))^2} \right)^{\frac{1}{p}}. \end{aligned}$$

Using Lemma 2.3, we conclude that

$$\|F\|_{L_p(X \times X, H^2(\mu \times \mu))} \leq c_{R,k}^{(2)} \left( \int_{\mathbb{R}^N \times \mathbb{R}^N} \left( \frac{d(x, y)}{|x - y|} \right)^p |f(x) - f(y)|^p \frac{dx dy}{d(x, y)^{2N}} \right)^{\frac{1}{p}}.$$

Using Theorem 3.1, we conclude that

$$\|F\|_{L_p(X \times X, H^2(\mu \times \mu))} \leq c_{R,k} \|f\|_{\dot{W}^{\frac{N}{p}, p}(\mathbb{R}^N)}.$$

Obviously, the integral kernel of the operator  $[R_{\text{Dunkl},j}, M_f]$  is  $FH$ . The assertion follows now from Lemma 4.4.  $\square$

## 5. PROOF OF RESTRICTED VERSION OF THEOREM 1.1 (ii)

**Theorem 5.1.** *If  $[R_{\text{Dunkl},j}, M_f] \in \mathcal{L}_p(L_2(\mathbb{R}^N, w(x)dx))$ , then  $f \in \dot{W}^{\frac{N}{p}, p}(\Gamma)$  for every chamber  $\Gamma$ . Furthermore,*

$$\|f\|_{\dot{W}^{\frac{N}{p}, p}(\Gamma)} \leq c_{R,k} \|[R_{\text{Dunkl},j}, M_f]\|_{\mathcal{L}_p(L_2(\mathbb{R}^N, w(x)dx))}.$$

**Lemma 5.2.** *Let  $\Gamma$  be a chamber and let  $x_0 \in \Gamma$ . We have*

$$\sup_{x \in B(x_0, r) \cap \Gamma} w(x) \leq \frac{c_{k,\Gamma}}{m(B(x_0, r) \cap \Gamma)} \int_{B(x_0, r) \cap \Gamma} w(x) dx.$$

*Proof.* If  $x = x_0 + y$ ,  $y \in B(0, r)$ , then

$$|\langle x, v \rangle| \leq |\langle x_0, v \rangle| + r|\langle y, v \rangle| \leq |\langle x_0, v \rangle| + r\sqrt{2}.$$

Thus,

$$w(x) = \prod_{v \in R} (|\langle x_0, v \rangle| + |\langle y, v \rangle|)^{\epsilon \cdot k(v)} \leq \prod_{v \in R} (|\langle x_0, v \rangle| + r\sqrt{2})^{k(v)}.$$

In particular,

$$LHS \leq \prod_{v \in R} (|\langle x_0, v \rangle| + r\sqrt{2})^{k(v)}.$$

By convexity,  $x_0 + \Gamma \subset \Gamma$ . Hence,

$$x_0 + (B(0, r) \cap \Gamma) \subset B(x_0, r) \cap \Gamma.$$

So,

$$\int_{B(x_0, r) \cap \Gamma} w(x) dx \geq \int_{x_0 + (B(0, r) \cap \Gamma)} w(x) dx = \int_{B(0, r) \cap \Gamma} w(x_0 + y) dy.$$

If  $y \in \Gamma$ , then

$$|\langle x_0 + y, v \rangle| = |\langle x_0, v \rangle| + |\langle y, v \rangle|, \quad v \in R.$$

Thus,

$$\begin{aligned} w(x_0 + y) &= \prod_{v \in R} (|\langle x_0, v \rangle| + |\langle y, v \rangle|)^{k(v)} \geq \\ &\geq \max\left\{ \prod_{v \in R} |\langle x_0, v \rangle|^{k(v)}, \prod_{v \in R} |\langle y, v \rangle|^{k(v)} \right\}. \end{aligned}$$

Hence,

$$\begin{aligned} \int_{B(x_0, r) \cap \Gamma} w(x) dx &\geq \int_{B(0, r) \cap \Gamma} w(x_0 + y) dy \geq \\ &\geq \max\left\{ \int_{B(0, r) \cap \Gamma} \prod_{v \in R} |\langle x_0, v \rangle|^{k(v)} dy, \int_{B(0, r) \cap \Gamma} \prod_{v \in R} |\langle y, v \rangle|^{k(v)} dy \right\}. \end{aligned}$$

Obviously,

$$\begin{aligned} \int_{B(0, r) \cap \Gamma} \prod_{v \in R} |\langle x_0, v \rangle|^{k(v)} dy &= r^N \prod_{v \in R} |\langle x_0, v \rangle|^{k(v)} \int_{B(0, 1) \cap \Gamma} dy, \\ \int_{B(0, r) \cap \Gamma} \prod_{v \in R} |\langle y, v \rangle|^{k(v)} dy &= r^{N + \sum_{v \in R} k(v)} \int_{B(0, 1) \cap \Gamma} \prod_{v \in R} |\langle y, v \rangle|^{k(v)} dy. \end{aligned}$$

Thus,

$$\int_{B(x_0, r) \cap \Gamma} w(x) dx \geq c_{k, \Gamma}^{(1)} \max\left\{ r^N \prod_{v \in R} |\langle x_0, v \rangle|^{k(v)}, r^{N + \sum_{v \in R} k(v)} \right\}.$$

Combining this with the preceding paragraph, we complete the proof.  $\square$

**Lemma 5.3.** *Let  $\Gamma$  be a chamber and let  $x_0 \in \Gamma$ . Suppose  $1 < t < 1 + \frac{1}{|R| \cdot \max_{v \in R} k(v)}$ . For every  $r > 0$ , we have*

$$\left( \frac{\int_{B(x_0, r) \cap \Gamma} w^{1-t}(x) dx}{\int_{B(x_0, r) \cap \Gamma} w(x) dx} \right)^{\frac{1}{t}} \leq c_{t, k, \Gamma} \frac{m(B(x_0, r) \cap \Gamma)}{\int_{B(x_0, r) \cap \Gamma} w(x) dx}.$$

*Proof.* If  $v \in R$  is such that

$$2\sqrt{2}r \leq |\langle x_0, v \rangle|,$$

then

$$|\langle x, v \rangle| \geq |\langle x_0, v \rangle| - |\langle x - x_0, v \rangle| \geq |\langle x_0, v \rangle| - r\sqrt{2} \geq \frac{1}{2} |\langle x_0, v \rangle|, \quad x \in B(x_0, r).$$

In this case,

$$\begin{aligned} \int_{B(x_0, r)} |\langle x, v \rangle|^{(1-t)|R|k(v)} dx &\leq 2^{(t-1)|R|k(v)} |\langle x_0, v \rangle|^{(1-t)|R|k(v)} \cdot \int_{B(x_0, r)} dx = \\ &= 2^{(t-1)|R|k(v)} |\langle x_0, v \rangle|^{(1-t)|R|k(v)} \cdot \text{Vol}(\mathbb{B}^N) r^N. \end{aligned}$$

If  $v \in R$  is such that

$$|\langle x_0, v \rangle| \leq 2\sqrt{2}r,$$

then let  $y_0 = \langle x_0, v \rangle$  and  $z_0 = x_0 - \frac{1}{2}y_0v_0$ . Note that  $|y_0| \leq 2\sqrt{2}r$ . Using new variables  $y = \langle x, v \rangle$  and  $z = x - \frac{1}{2}yv$ , we write

$$\begin{aligned} \int_{B(x_0, r)} |\langle x, v \rangle|^{(1-t)|R|k(v)} dx &= \int_{|y-y_0|^2 + |z-z_0|^2 \leq r^2} |y|^{(1-t)|R|k(v)} dy dz \leq \\ &\leq \int_{|y-y_0|^2, |z-z_0|^2 \leq r^2} |y|^{(1-t)|R|k(v)} dy dz = \end{aligned}$$

$$\begin{aligned}
&= r^{N-1} \text{Vol}(\mathbb{B}^{N-1}) \cdot \int_{|y-y_0| \leq r} |y|^{(1-t)|R|k(v)} dy \leq \\
&= r^{N-1} \text{Vol}(\mathbb{B}^{N-1}) \cdot \int_{|y| \leq (1+2\sqrt{2})r} |y|^{(1-t)|R|k(v)} dy = \\
&= r^{N+(1-t)|R|k(v)} \text{Vol}(\mathbb{B}^{N-1}) \cdot \int_{|y| \leq 1+2\sqrt{2}} |y|^{(1-t)|R|k(v)} dy.
\end{aligned}$$

Combining the estimates in the preceding paragraphs, we write

$$\int_{B(x_0, r)} |\langle x, v \rangle|^{(1-t)|R|k(v)} dx \leq c_1 R^n (\max\{|\langle x_0, v \rangle|, r\})^{(1-t)|R|k(v)}.$$

By Hölder inequality,

$$\begin{aligned}
\int_{B(x_0, r)} w^{1-t}(x) dx &\leq \prod_{v \in R} \left( \int_{B(x_0, r)} |\langle x, v \rangle|^{(1-t)|R|k(v)} dx \right)^{\frac{1}{|R|}} \leq \\
&\leq c_2 R^N \prod_{v \in R} (\max\{|\langle x_0, v \rangle|, r\})^{(1-t)k(v)}.
\end{aligned}$$

The assertion follows easily.  $\square$

**Lemma 5.4.** *Let  $\Gamma$  be a chamber equipped with Euclidean distance. Measures  $dx$  and  $w(x)dx$  and operator  $T_{\Gamma, j} = M_{\chi_\Gamma} R_{\text{Dunkl}, j} M_{\chi_\Gamma}$  satisfy **on  $\Gamma$**  the conditions in Theorem 1.4 in [6].*

*Proof. Step 1:* We claim that measures  $dx$  and  $w(x)dx$  are  $A_\infty$ -equivalent (in the sense of Proposition 1.2 in [6]). This verifies Assumption (i) in Theorem 1.4 in [6].

Let  $x_0 \in \Gamma$ . By Lemmas 5.2 and 5.3,

$$\begin{aligned}
\left( \frac{1}{m(B \cap \Gamma)} \int_{B \cap \Gamma} w^t(x) dx \right)^{\frac{1}{t}} &\leq \frac{c_{k, \Gamma}}{m(B \cap \Gamma)} \int_{B \cap \Gamma} w(x) dx, \quad t > 0. \\
\left( \frac{\int_{B \cap \Gamma} w^{1-t}(x) dx}{\int_{B \cap \Gamma} w(x) dx} \right)^{\frac{1}{t}} &\leq c_G \frac{m(B \cap \Gamma)}{\int_{B \cap \Gamma} w(x) dx}, \quad 1 < t < 1 + \frac{1}{|R| \cdot \max_{v \in R} k(v)}.
\end{aligned}$$

The assertion follows now from Proposition 3.6 in [6].

**Step 2:** Assumption (ii) in Theorem 1.4 in [6] is obvious.

**Step 3:** Integral kernel  $K_j$  of the operator  $T_{\Gamma, j}$  satisfies the estimates

$$\begin{aligned}
|K_j(x, y)| &\leq \frac{c_G}{w(B(x, |x-y|) \cap \Gamma)}, \quad x, y \in \Gamma, \\
|K_j(x, y) - K_j(x', y)| &\leq \frac{|x-x'|}{|x-y|} \cdot \frac{c_G}{w(B(x, |x-y|))}, \quad |x-x'| \leq \frac{1}{2}|x-y|, \quad x, x', y \in \Gamma, \\
|K_j(x, y) - K_j(x, y')| &\leq \frac{|y-y'|}{|x-y|} \cdot \frac{c_G}{w(B(x, |x-y|))}, \quad |y-y'| \leq \frac{1}{2}|x-y|, \quad x, y, y' \in \Gamma.
\end{aligned}$$

In other words,  $K_j$  is a standard kernel for the measure  $w(x)dx$ .

This follows from Theorem 1.1 in [4].

**Step 4:** Kernel  $K_j$  is non-degenerate in the following sense: for every  $x \in \Gamma$  and  $r > 0$ , there exists  $y \in \Gamma$  with  $|x-y| = r$  and

$$|K_j(x, y)| \geq \frac{c_G}{w(B(x, |x-y|) \cap \Gamma)}.$$

Let

$$c_\Gamma = \left( \max_{z \in \Gamma, |z|=1} |z_j| \right)^{-1}.$$

Choose  $z \in \Gamma$  such that  $|z_j| = c_\Gamma^{-1}$  and  $|z| = 1$ . Set  $y = x + rz$ . Since  $x, z \in \Gamma$ , it follows that  $y \in \Gamma$ . Clearly,  $|(x - y)_j| \geq rc_\Gamma^{-1}$  and  $|x - y| = r$ . The claim follows from **the proof of the** Theorem 1.2 in [4].

**Step 4:** Lebesgue measure on  $\Gamma$  is, obviously,  $N$ -regular and possesses the  $(1, 1)$  Poincare inequality.  $\square$

*Proof of Theorem 5.1.* This follows immediately from Lemma 5.4 and Theorem 1.4 in [6].  $\square$

## 6. USEFUL PART OF ZHIJIE ARGUMENT

**Theorem 6.1.** *If  $[R_{\text{Dunkl},j}, M_f] \in \mathcal{L}_p(L_2(\mathbb{R}^N, w(x)dx))$ , then  $f \in B^p(\mathbb{R}^N, w(x)dx)$ .*

**Lemma 6.2.** *Let  $Q \in \mathcal{D}$ . We have*

$$K_j(x, y) \geq \frac{c_k}{w(Q)}, \quad x \in 3Q, \quad y \in Q + 3s(Q)e_j.$$

*Proof.* **TO BE DONE.** Follows from Lemma 2.4.  $\square$

**Lemma 6.3.** *Let real-valued  $f \in L_1^{\text{loc}}(\mathbb{R}^N)$  be such that the commutator  $[R_{\text{Dunkl},j}, M_f]$  is bounded on  $L_2(\mathbb{R}^N, w(x)dx)$ . For every  $Q \in \mathcal{D}$ , there exist  $\xi_Q$  and  $\eta_Q$  supported in  $7Q$  such that  $\|\xi_Q\|_\infty, \|\eta_Q\|_\infty \leq m(Q)^{-\frac{1}{2}}$  and such that*

$$\frac{1}{w(3Q)} \int_{3Q} |f(y) - f_{3Q}| w(y) dy \leq c_k |\langle [R_{\text{Dunkl},j}, M_f] \xi_Q, \eta_Q \rangle_{L_2(\mathbb{R}^N, w(x)dx)}|.$$

*Proof.* Let  $\alpha$  be the median of  $f$  on  $3Q$  and  $\beta$  be the median of  $f$  on  $Q + 3s(Q)e_j$ . Assume for definiteness  $\alpha \geq \beta$ . Set

$$\begin{aligned} A_1 &= \{x \in 3Q : f(x) \geq \alpha\}, & B_1 &= \{x \in Q + 3s(Q)e_j : f(x) \leq \beta\}, \\ A_2 &= \{x \in 3Q : f(x) \leq \beta\}, & B_2 &= \{x \in Q + 3s(Q)e_j : f(x) \geq \alpha\}, \\ \xi_{Q,s} &= w(Q)^{-\frac{1}{2}} \chi_{A_s}, & \eta_{Q,s} &= w(Q)^{-\frac{1}{2}} \chi_{B_s}, \quad s = 1, 2. \end{aligned}$$

We have

$$\begin{aligned} &\langle [R_{\text{Dunkl},j}, M_f] \xi_{Q,s}, \eta_{Q,s} \rangle_{L_2(\mathbb{R}^N, w(x)dx)} = \\ &= \frac{1}{w(Q)} \int_{B_s \times A_s} K_j(x, y) (f(y) - f(x)) w(x) w(y) dx dy. \end{aligned}$$

If  $x \in B_1$  and  $y \in A_1$ , then

$$\begin{aligned} f(y) - f(x) &= (f(y) - \alpha) + (\alpha - \beta) + (\beta - f(x)) = \\ &= |f(y) - \alpha| + |\alpha - \beta| + |f(x) - \beta| \geq |f(y) - \alpha| + |\alpha - \beta| \geq 0. \end{aligned}$$

If  $x \in B_2$  and  $y \in A_2$ , then

$$\begin{aligned} f(y) - f(x) &= (f(y) - \beta) + (\beta - f(x)) = \\ &= -|f(y) - \beta| - |f(x) - \beta| \leq -|f(y) - \beta| \leq 0. \end{aligned}$$

Thus,

$$\begin{aligned} &\langle [R_{\text{Dunkl},j}, M_f] \xi_{Q,1}, \eta_{Q,1} \rangle_{L_2(\mathbb{R}^N, w(x)dx)} \geq \\ &\geq \frac{1}{w(Q)} \int_{B_1 \times A_1} K_j(x, y) (|f(y) - \alpha| + |\alpha - \beta|) w(x) w(y) dx dy \geq \\ &\geq \frac{c_k}{w^2(Q)} \int_{B_1 \times A_1} (|f(y) - \alpha| + |\alpha - \beta|) w(x) w(y) dx dy = \end{aligned}$$

$$\begin{aligned}
&= \frac{c_k}{w^2(Q)} \cdot \left( w(B_1) \cdot \int_{A_1} |f(y) - \alpha| w(y) dy + |\alpha - \beta| w(B_1) w(A_1) \right) \geq \\
&= \frac{c_k}{w^2(Q)} \cdot \left( \frac{1}{2} w(Q) \cdot \int_{A_1} |f(y) - \alpha| w(y) dy + |\alpha - \beta| \cdot \frac{1}{4} w^2(Q) \right) \geq \\
&\geq \frac{c_k}{4w(Q)} \cdot \left( \int_{A_1} |f(y) - \alpha| w(y) dy + |\alpha - \beta| w(Q) \right)
\end{aligned}$$

and

$$\begin{aligned}
& -\langle [R_{\text{Dunkl},j}, M_f] \xi_{Q,2}, \eta_{Q,2} \rangle_{L_2(\mathbb{R}^N, w(x) dx)} \geq \\
& \geq \frac{1}{w(Q)} \int_{B_2 \times A_2} K_j(x, y) |f(y) - \beta| w(x) w(y) dx dy \geq \\
& \geq \frac{c_k}{w^2(Q)} \int_{B_2 \times A_2} |f(y) - \beta| w(x) w(y) dx dy = \\
& = \frac{c_k}{w^2(Q)} \cdot w(B_2) \cdot \int_{A_2} |f(y) - \beta| w(y) dy \geq \\
& = \frac{c_k}{2w(Q)} \int_{A_2} |f(y) - \beta| w(y) dy.
\end{aligned}$$

It remains to note that

$$\begin{aligned}
& \int_{3Q} |f(y) - \alpha| w(y) dy = \int_{A_1} |f(y) - \alpha| w(y) dy + \\
& + \int_{A_2} |f(y) - \alpha| w(y) dy + \int_{3Q \setminus (A_1 \cup A_2)} |f(y) - \alpha| w(y) dy \leq \\
& \leq \int_{A_1} |f(y) - \alpha| w(y) dy + \int_{A_2} |f(y) - \beta| w(y) dy \\
& + \int_{A_2} |\alpha - \beta| w(y) dy + \int_{3Q \setminus (A_1 \cup A_2)} |\alpha - \beta| w(y) dy \leq \\
& \leq \int_{A_1} |f(y) - \alpha| w(y) dy + \int_{A_2} |f(y) - \beta| w(y) dy + 2|\alpha - \beta| w(3Q).
\end{aligned}$$

Combining the last 3 equations, we conclude

$$\frac{c'_k}{w(Q)} \int_{3Q} |f(y) - \alpha| w(y) dy \leq \max_{s=1,2} |\langle [R_{\text{Dunkl},j}, M_f] \xi_{Q,s}, \eta_{Q,s} \rangle_{L_2(\mathbb{R}^N, w(x) dx)}|.$$

If

$|\langle [R_{\text{Dunkl},j}, M_f] \xi_{Q,1}, \eta_{Q,1} \rangle_{L_2(\mathbb{R}^N, w(x) dx)}| \geq |\langle [R_{\text{Dunkl},j}, M_f] \xi_{Q,2}, \eta_{Q,2} \rangle_{L_2(\mathbb{R}^N, w(x) dx)}|$ ,  
then we set  $\xi_Q = \xi_{Q,1}$  and  $\eta_Q = \eta_{Q,1}$ . Otherwise, set  $\xi_Q = \xi_{Q,2}$  and  $\eta_Q = \eta_{Q,2}$ . Thus,

$$\frac{c'_k}{w(Q)} \int_{3Q} |f(y) - \alpha| w(y) dy \leq |\langle [R_{\text{Dunkl},j}, M_f] \xi_Q, \eta_Q \rangle_{L_2(\mathbb{R}^N, w(x) dx)}|.$$

Hence,

$$c'_k |f_{3Q} - \alpha| \leq |\langle [R_{\text{Dunkl},j}, M_f] \xi_Q, \eta_Q \rangle_{L_2(\mathbb{R}^N, w(x) dx)}|$$

and, finally,

$$\frac{c'_k}{w(Q)} \int_{3Q} |f(y) - f_{3Q}| w(y) dy \leq 2 |\langle [R_{\text{Dunkl},j}, M_f] \xi_Q, \eta_Q \rangle_{L_2(\mathbb{R}^N, w(x) dx)}|.$$

□

**Lemma 6.4.** *Suppose that  $1 < p < \infty$  and  $f \in L_1^{\text{loc}}(\mathbb{R}^N)$ . If  $[R_{\text{Dunkl},j}, M_f] \in \mathcal{L}_p(L_2(\mathbb{R}^N, w(x)dx))$  for some  $j \in \{1, 2, \dots, N\}$ , then*

$$\left\{ \frac{1}{w(3Q)} \int_{3Q} |f(x) - f_{3Q}| w(x) dx \right\}_{Q \in \mathcal{D}} \in l_p.$$

*Proof.* Let  $\{\xi_Q\}_{Q \in \mathcal{D}}$  and  $\{\eta_Q\}_{Q \in \mathcal{D}}$  be functions constructed in Lemma 6.3. By Lemma 2.6,

$$\left\{ \langle [R_{\text{Dunkl},j}, M_f] \xi_Q, \eta_Q \rangle_{L_2(\mathbb{R}^N, w(x)dx)} \right\}_{Q \in \mathcal{D}} \in l_p.$$

The assertion follows now from Lemma 6.3.  $\square$

**Lemma 6.5.** *Suppose that  $1 \leq p < \infty$  and  $f \in L_1^{\text{loc}}(\mathbb{R}^N, w(x)dx)$ . We have*

$$\begin{aligned} \|f\|_{B^p(\mathbb{R}^N, w(x)dx)}^p &\leq c_{p,k} \sum_{Q \in \mathcal{D}} \frac{1}{w(Q)} \int_Q |f(x) - f_Q|^p w(x) dx + \\ &+ c_{p,k} \sum_{Q \in \mathcal{D}} \left( \frac{1}{w(3Q)} \int_{3Q} |f(x) - f_{3Q}| w(x) dx \right)^p. \end{aligned}$$

*Proof.* We first decompose  $\mathbb{R}^N \times \mathbb{R}^N$  into dyadic shells as follows.

$$\begin{aligned} &\int_{\mathbb{R}^N \times \mathbb{R}^N} \frac{|f(x) - f(y)|^p}{w(B(x, |x-y|))^2} w(x) w(y) dx dy \leq \\ &\leq \sum_{m \in \mathbb{Z}} \int_{2^{-m-1} < |x-y| < 2^{-m}} \frac{|f(x) - f(y)|^p}{w(B(x, |x-y|))^2} w(x) w(y) dx dy \leq \\ &\leq \sum_{m \in \mathbb{Z}} \frac{1}{w(B(x, 2^{-m-1}))^2} \int_{\substack{x, y \in \mathbb{R}^N \\ |x-y| < 2^{-m}}} |f(x) - f(y)|^p w(x) w(y) dx dy. \end{aligned}$$

For  $x, y \in \mathbb{R}^N$  with  $2^{-m-1} < |x-y| < 2^{-m}$ , fix  $Q \in \mathcal{D}$  with  $x \in Q$  and  $s(Q) = 2^{-m}$ . We have  $|x-y|_\infty < 2^{-m}$  and  $|x - c_Q|_\infty < 2^{-m-1}$ . Thus,  $|y - c_Q|_\infty < \frac{3}{2} \cdot 2^{-m}$ . In other words,  $y \in 3Q$ . Consequently, there exists  $Q \in \mathcal{D}$  with  $s(Q) = 2^{-m}$ ,  $x, y \in 3Q$ . Thus,

$$\begin{aligned} &\sum_{m \in \mathbb{Z}} \frac{1}{w(B(x, 2^{-m-1}))^2} \int_{\substack{x, y \in \mathbb{R}^N \\ |x-y| < 2^{-m}}} |f(x) - f(y)|^p w(x) w(y) dx dy \leq \\ &\leq \sum_{m \in \mathbb{Z}} \frac{1}{w(B(x, 2^{-m-1}))^2} \sum_{\substack{Q \in \mathcal{D} \\ s(Q) = 2^{-m}}} \int_{3Q \times 3Q} |f(x) - f(y)|^p w(x) w(y) dx dy \leq \\ &\leq 2^p \sum_{m \in \mathbb{Z}} \frac{1}{w(B(x, 2^{-m-1}))^2} \sum_{\substack{Q \in \mathcal{D} \\ s(Q) = 2^{-m}}} \int_{3Q \times 3Q} |f(x) - f_{3Q}|^p w(x) w(y) dx dy + \\ &+ 2^p \sum_{m \in \mathbb{Z}} \frac{1}{w(B(x, 2^{-m-1}))^2} \sum_{\substack{Q \in \mathcal{D} \\ s(Q) = 2^{-m}}} \int_{3Q \times 3Q} |f(y) - f_{3Q}|^p w(x) w(y) dx dy \leq \\ &\leq c_{p,k} \sum_{Q \in \mathcal{D}} \frac{1}{w^2(3Q)} \int_{3Q \times 3Q} |f(x) - f_{3Q}|^p w(x) w(y) dx dy = \\ &= c_{p,k} \sum_{Q \in \mathcal{D}} \frac{1}{w(3Q)} \int_{3Q} |f(x) - f_{3Q}|^p w(x) dx. \end{aligned}$$

Next,

$$\begin{aligned} \int_{3Q} |f(x) - f_{3Q}|^p w(x) dx &= \sum_{\epsilon \in \{-1, 0, 1\}^d} \int_{Q+\epsilon s(Q)} |f(x) - f_{3Q}|^p w(x) dx \leq \\ &\leq 2^{p-1} \sum_{\epsilon \in \{-1, 0, 1\}^d} \int_{Q+\epsilon s(Q)} |f(x) - f_{Q+\epsilon s(Q)}|^p w(x) dx + \\ &+ 2^{p-1} \sum_{\epsilon \in \{-1, 0, 1\}^d} \int_{Q+\epsilon s(Q)} |f_{Q+\epsilon s(Q)} - f_{3Q}|^p w(x) dx. \end{aligned}$$

Obviously,

$$\begin{aligned} \int_{Q+\epsilon s(Q)} |f_{Q+\epsilon s(Q)} - f_{3Q}|^p w(x) dx &= |f_{Q+\epsilon s(Q)} - f_{3Q}|^p \cdot w(Q + \epsilon s(Q)) \leq \\ &\leq \left( \int_{Q+\epsilon s(Q)} |f(x) - f_{3Q}| w(x) dx \right)^p \cdot w^{1-p}(Q + \epsilon s(Q)) \leq \\ &\leq \left( \int_{3Q} |f(x) - f_{3Q}| w(x) dx \right)^p \cdot w^{1-p}(Q + \epsilon s(Q)) \leq \\ &\leq c_{p,k} \left( \int_{3Q} |f(x) - f_{3Q}| w(x) dx \right)^p \cdot w^{1-p}(3Q). \end{aligned}$$

Thus,

$$\begin{aligned} &\int_{\mathbb{R}^N \times \mathbb{R}^N} \frac{|f(x) - f(y)|^p}{w(B(x, |x - y|))^2} w(x) w(y) dx dy \leq \\ &\leq c_{p,k} \sum_{Q \in \mathcal{D}} \sum_{\epsilon \in \{-1, 0, 1\}^d} \frac{1}{w(Q + \epsilon s(Q))} \int_{Q+\epsilon s(Q)} |f(x) - f_{Q+\epsilon s(Q)}|^p w(x) dx + \\ &+ c_{p,k} \sum_{Q \in \mathcal{D}} \left( \frac{1}{w(3Q)} \int_{3Q} |f(x) - f_{3Q}| w(x) dx \right)^p. \end{aligned}$$

This completes the proof.  $\square$

**Lemma 6.6.** *Suppose that  $1 \leq p < \infty$  and  $f \in L_1^{\text{loc}}(\mathbb{R}^N, w(x) dx)$ . We have*

$$\sum_{Q \in \mathcal{D}} \frac{1}{w(Q)} \int_Q |f(x) - f_Q|^p w(x) dx \leq c_{p,k} \sum_{Q \in \mathcal{D}} |f_Q - f_{\text{parent}(Q)}|^p.$$

*Proof.* Obviously,

$$\begin{aligned} \sum_{Q \in \mathcal{D}} \frac{1}{w(Q)} \int_Q |f(x) - f_Q|^p w(x) dx &= \sum_{m \in \mathbb{Z}} \sum_{\substack{Q \in \mathcal{D} \\ s(Q) = 2^{-m}}} \frac{1}{w(Q)} \int_Q |f(x) - f_Q|^p w(x) dx \leq \\ &\leq c_{p,k} \sum_{m \in \mathbb{Z}} \sum_{\substack{Q \in \mathcal{D} \\ s(Q) = 2^{-m}}} \int_Q \frac{|f(x) - f_Q|^p w(x) dx}{w(B(x, 2^{-m}))} = \\ &= c_{p,k} \sum_{m \in \mathbb{Z}} \int_{\mathbb{R}^N} \frac{|f(x) - (E_m f)(x)|^p w(x) dx}{w(B(x, 2^{-m}))}. \end{aligned}$$

Note that

$$f - E_m f = \sum_{l \geq m} (E_{l+1} f - E_l f).$$

By triangle inequality,

$$\begin{aligned} & \left( \int_{\mathbb{R}^N} \frac{|f(x) - (E_m f)(x)|^p w(x) dx}{w(B(x, 2^{-m}))} \right)^{\frac{1}{p}} \leq \\ & \leq \sum_{l \geq m} \left( \int_{\mathbb{R}^N} \frac{|(E_{l+1} f)(x) - (E_l f)(x)|^p w(x) dx}{w(B(x, 2^{-m}))} \right)^{\frac{1}{p}}. \end{aligned}$$

Note that

$$w(B(x, 2^{-m})) \geq c_k 2^{N(l-m)} w(B(x, 2^{-l})).$$

Thus,

$$\begin{aligned} & \left( \int_{\mathbb{R}^N} \frac{|f(x) - (E_m f)(x)|^p w(x) dx}{w(B(x, 2^{-m}))} \right)^{\frac{1}{p}} \leq \\ & \leq c_{p,k} \sum_{l \geq m} 2^{\frac{N(m-l)}{p}} \left( \int_{\mathbb{R}^N} \frac{|(E_{l+1} f)(x) - (E_l f)(x)|^p w(x) dx}{w(B(x, 2^{-l}))} \right)^{\frac{1}{p}} \end{aligned}$$

and, therefore,

$$\begin{aligned} & \sum_{m \in \mathbb{Z}} \int_{\mathbb{R}^N} \frac{|f(x) - (E_m f)(x)|^p w(x) dx}{w(B(x, 2^{-m}))} \leq \\ & \leq c_{p,k} \sum_{m \in \mathbb{Z}} \left( \sum_{l \geq m} 2^{\frac{N(m-l)}{p}} \left( \int_{\mathbb{R}^N} \frac{|(E_{l+1} f)(x) - (E_l f)(x)|^p w(x) dx}{w(B(x, 2^{-l}))} \right)^{\frac{1}{p}} \right)^p. \end{aligned}$$

Using the discrete Young's inequality

$$\|a * b\|_p \leq \|a\|_1 \|b\|_p$$

with

$$a(l) = 2^{\frac{Nl}{p}} \chi_{\mathbb{Z}_-}(l), \quad b(l) = \left( \int_{\mathbb{R}^N} \frac{|(E_{l+1} f)(x) - (E_l f)(x)|^p w(x) dx}{w(B(x, 2^{-l}))} \right)^{\frac{1}{p}},$$

we obtain

$$\begin{aligned} & \sum_{m \in \mathbb{Z}} \left( \sum_{l \geq m} 2^{\frac{N(m-l)}{p}} \left( \int_{\mathbb{R}^N} \frac{|(E_{l+1} f)(x) - (E_l f)(x)|^p w(x) dx}{w(B(x, 2^{-l}))} \right)^{\frac{1}{p}} \right)^p \leq \\ & \leq c_{N,p} \sum_{l \in \mathbb{Z}} \int_{\mathbb{R}^N} \frac{|(E_{l+1} f)(x) - (E_l f)(x)|^p w(x) dx}{w(B(x, 2^{-l}))}. \end{aligned}$$

Thus,

$$\begin{aligned} & \sum_{m \in \mathbb{Z}} \int_{\mathbb{R}^N} \frac{|f(x) - (E_m f)(x)|^p w(x) dx}{w(B(x, 2^{-m}))} \leq \\ & \leq c_{p,k} \sum_{l \in \mathbb{Z}} \int_{\mathbb{R}^N} \frac{|(E_{l+1} f)(x) - (E_l f)(x)|^p w(x) dx}{w(B(x, 2^{-l}))} \leq \\ & \leq c_{p,k} \sum_{l \in \mathbb{Z}} \sum_{\substack{Q \in \mathcal{D} \\ s(Q)=2^{-l-1}}} \frac{1}{w(Q)} \int_Q |(E_{l+1} f)(x) - (E_l f)(x)|^p w(x) dx = \\ & = c_{p,k} \sum_{l \in \mathbb{Z}} \sum_{\substack{Q \in \mathcal{D} \\ s(Q)=2^{-l-1}}} |f_Q - f_{\text{parent}(Q)}|^p. \end{aligned}$$

This completes the proof.  $\square$

In what follows, conditional expectations are with respect to the measure  $w(x)dx$ .

**Lemma 6.7.** *Suppose that  $1 \leq p < \infty$  and  $f \in L_1^{\text{loc}}(\mathbb{R}^N, w(x)dx)$ . We have*

$$\|f\|_{B^p(\mathbb{R}^N, w(x)dx)}^p \leq c_{p,k} \sum_{Q \in \mathcal{D}} \left( \frac{1}{w(3Q)} \int_{3Q} |f(x) - f_{3Q}| w(x) dx \right)^p.$$

*Proof.* Combining Lemmas 6.5 and 6.6, we arrive at

$$\begin{aligned} \|f\|_{B^p(\mathbb{R}^N, w(x)dx)}^p &\leq c_{p,k} \sum_{Q \in \mathcal{D}} \frac{1}{w(Q)} \int_Q |f(x) - f_Q|^p w(x) dx + \\ &\quad + c_{p,k} \sum_{Q \in \mathcal{D}} |f_Q - f_{\text{parent}(Q)}|^p. \end{aligned}$$

Obviously,

$$\begin{aligned} |f_Q - f_{\text{parent}(Q)}| &\leq \frac{1}{w(Q)} \int_{\text{parent}(Q)} |f(x) - f_Q| w(x) dx \leq \\ &\leq \frac{1}{w(Q)} \int_{3Q} |f(x) - f_Q| w(x) dx \leq \frac{c_k}{w(3Q)} \int_{3Q} |f(x) - f_Q| w(x) dx \leq \\ &\leq \frac{2c_k}{w(3Q)} \int_{3Q} |f(x) - f_Q| w(x) dx. \end{aligned}$$

This completes the proof.  $\square$

*Proof of Theorem 6.1.* The assertion follows by combining Lemma 6.4 and Lemma 6.7.  $\square$

## 7. PROOF OF THEOREM 1.1 (ii)

**Theorem 7.1.** *Let  $f \in L_1^{\text{loc}}(\mathbb{R}^N)$  be such that*

- (i) *for every chamber  $\Gamma$ , we have  $f \in \dot{W}^{\frac{N}{p}, p}(\Gamma)$ ;*
- (ii)  *$f$  belongs to the space  $B^p(\mathbb{R}^N, w(x)dx)$ ;*

*It follows that  $f \in \dot{W}^{\frac{N}{p}, p}(\mathbb{R}^N)$ .*

**Lemma 7.2.** *Denote  $H_+ = \mathbb{R}^{N-1} \times \mathbb{R}_+$  and  $H_- = \mathbb{R}^{N-1} \times \mathbb{R}_-$ . Let  $z \in \mathbb{R}^N$  be such that  $z_N = 0$ . Suppose  $f \in \dot{W}^{\frac{N}{p}, p}(H_+ \cap B(z, 2\epsilon))$  and  $f \in \dot{W}^{\frac{N}{p}, p}(H_- \cap B(z, 2\epsilon))$ . If*

$$\int_{\substack{x \in H_+ \cap B(z, 2\epsilon) \\ y \in H_- \cap B(z, 2\epsilon)}} \frac{|f(x) - f(y)|^p}{|x - y|^{2N+2K}} |x_N|^K |y_N|^K dx dy < \infty,$$

*then*

$$\int_{B(z, \epsilon)} \frac{|f(x_1, \dots, x_{N-1}, x_N) - f(x_1, \dots, x_{N-1}, -x_N)|^p dx}{|x_N|^N} < \infty.$$

*Proof.* Let  $f_1(x) = f(x_1, \dots, x_{N-1}, |x_N|)$ ,  $x \in B(z, 2\epsilon)$ . It is immediate **we, probably, need reference** that  $f_1 \in \dot{W}^{\frac{N}{p}, p}(B(z, 2\epsilon))$ . In other words,

$$\int_{B(z, 2\epsilon) \times B(z, 2\epsilon)} \frac{|f_1(x) - f_1(y)|^p}{|x - y|^{2N}} dx dy < \infty.$$

In particular,

$$\int_{\substack{x \in H_+ \cap B(z, 2\epsilon) \\ y \in H_- \cap B(z, 2\epsilon)}} \frac{|f_1(x) - f_1(y)|^p}{|x - y|^{2N}} dx dy < \infty.$$

Since  $|x_N|, |y_N| \leq |x - y|$  for  $x \in H_+$ ,  $y \in H_-$ , it follows that

$$\int_{\substack{x \in H_+ \cap B(z, 2\epsilon) \\ y \in H_- \cap B(z, 2\epsilon)}} \frac{|f_1(x) - f_1(y)|^p}{|x - y|^{2N+2K}} |x_N|^K |y_N|^K dx dy < \infty.$$

Set  $f_2 = f - f_1$ . By triangle inequality,

$$\int_{\substack{x \in H_+ \cap B(z, 2\epsilon) \\ y \in H_- \cap B(z, 2\epsilon)}} \frac{|f_2(x) - f_2(y)|^p}{|x - y|^{2N+2K}} |x_N|^K |y_N|^K dx dy < \infty.$$

Note that  $f_1 = f$  on  $H_+ \cap B(z, 2\epsilon)$  and, therefore,  $f_2 = 0$  on  $H_+ \cap B(z, 2\epsilon)$ . We, therefore, have

$$\int_{\substack{x \in H_+ \cap B(z, 2\epsilon) \\ y \in H_- \cap B(z, 2\epsilon)}} \frac{|f_2(y)|^p}{|x - y|^{2N+2K}} |x_N|^K |y_N|^K dx dy < \infty.$$

In other words,

$$\int_{H_- \cap B(z, 2\epsilon)} |f_2|^p(y) \left( \int_{H_+ \cap B(z, 2\epsilon)} \frac{|x_N|^K |y_N|^K dx}{|x - y|^{2N+2K}} \right) dy < \infty.$$

If  $y \in B(z, \epsilon)$ , then  $B(y', \epsilon) \subset B(z, 2\epsilon)$ . Thus,

$$\int_{H_- \cap B(z, \epsilon)} |f_2|^p(y) \left( \int_{H_+ \cap B(y', \epsilon)} \frac{|x_N|^K |y_N|^K dx}{|x - y|^{2N+2K}} \right) dy < \infty.$$

Clearly,

$$\int_{H_+ \cap B(y', \epsilon)} \frac{|x_N|^K |y_N|^K dx}{|x - y|^{2N+2K}} = \int_{H_+ \cap B(0, \epsilon)} \frac{|x_N|^K |y_N|^K dx}{(|x'|^2 + (x_N + |y_N|)^2)^{N+K}}.$$

Thus,

$$\int_{H_- \cap B(z, \epsilon)} |f_2|^p(y) \left( \int_{H_+ \cap B(0, \epsilon)} \frac{|x_N|^K |y_N|^K dx}{\max\{|x'|, x_N, |y_N|\}^{2N+2K}} \right) dy < \infty.$$

Clearly,

$$\begin{aligned} \int_{H_+ \cap B(0, \epsilon)} \frac{|x_N|^K |y_N|^K dx}{\max\{|x'|, x_N, |y_N|\}^{2N+2K}} &\geq \int_{\substack{|x'| < \frac{1}{2}|y_N| \\ x_N \in (0, \frac{1}{2}|y_N|)}} \frac{|x_N|^K |y_N|^K dx}{|y_N|^{2N+2K}} = \\ &= |y_N|^{-2N-K} \cdot \int_{|x'| < \frac{1}{2}|y_N|} dx' \cdot \int_0^{\frac{1}{2}|y_N|} dx_N = |y_N|^{-N} \cdot 2^{-N} \text{Vol}(\mathbb{B}^{N-1}). \end{aligned}$$

Hence,

$$\int_{H_- \cap B(z, \epsilon)} \frac{|f_2|^p(y) dy}{|y_N|^N} < \infty.$$

This completes the proof.  $\square$

**Lemma 7.3.** Denote  $H_+ = \mathbb{R}^{N-1} \times \mathbb{R}_+$  and  $H_- = \mathbb{R}^{N-1} \times \mathbb{R}_-$ . Let  $z \in \mathbb{R}^N$  be such that  $z_N = 0$ . Suppose  $f \in \dot{W}^{\frac{N}{p}, p}(H_+ \cap B(z, \epsilon))$  and  $f \in \dot{W}^{\frac{N}{p}, p}(H_- \cap B(z, \epsilon))$ . If

$$\int_{B(z, \epsilon)} \frac{|f(x_1, \dots, x_{N-1}, x_N) - f(x_1, \dots, x_{N-1}, -x_N)|^p dx}{|x_N|^N} < \infty,$$

then  $f \in \dot{W}^{\frac{N}{p}, p}(B(z, \epsilon))$ .

*Proof.* Set  $r(x) = (x_1, \dots, x_{N-1}, -x_N)$ ,  $x \in B(z, \epsilon)$  and let  $f_1 = f + f \circ r$ ,  $f_2 = f - f \circ r$ . It is known that [provide reference](#) that  $f_1 \in \dot{W}^{\frac{N}{p}, p}(B(z, \epsilon))$ . Hence, we may assume without loss of generality that  $f = -f \circ r$  and

$$\int_{B(z, \epsilon)} \frac{|f|^p(x) dx}{|x_N|^N} < \infty.$$

We have

$$\begin{aligned} & \int_{B(z, \epsilon) \times B(z, \epsilon)} \frac{|f(x) - f(y)|^p dx dy}{|x - y|^{2N}} \leq \\ & \leq \int_{\substack{x \in H_+ \cap B(z, \epsilon) \\ y \in H_+ \cap B(z, \epsilon)}} \frac{|f(x) - f(y)|^p dx dy}{|x - y|^{2N}} + \int_{\substack{x \in H_- \cap B(z, \epsilon) \\ y \in H_- \cap B(z, \epsilon)}} \frac{|f(x) - f(y)|^p dx dy}{|x - y|^{2N}} + \\ & + 2^{p-1} \int_{\substack{x \in H_+ \cap B(z, \epsilon) \\ y \in H_- \cap B(z, \epsilon)}} \frac{|f|^p(x) dx dy}{|x - y|^{2N}} + 2^{p-1} \int_{\substack{x \in H_+ \cap B(z, \epsilon) \\ y \in H_- \cap B(z, \epsilon)}} \frac{|f|^p(y) dx dy}{|x - y|^{2N}} + \\ & + 2^{p-1} \int_{\substack{x \in H_- \cap B(z, \epsilon) \\ y \in H_+ \cap B(z, \epsilon)}} \frac{|f|^p(x) dx dy}{|x - y|^{2N}} + 2^{p-1} \int_{\substack{x \in H_- \cap B(z, \epsilon) \\ y \in H_+ \cap B(z, \epsilon)}} \frac{|f|^p(y) dx dy}{|x - y|^{2N}}. \end{aligned}$$

It remains to note that, for every  $x \in H_+$ ,

$$\int_{y \in H_- \cap B(z, \epsilon)} \frac{dy}{|x - y|^{2N}} \leq \int_{y \in \mathbb{R}^N} \frac{dy}{(|y|^2 + (|y_N| + |x_N|)^2)^N} = c_N |x_N|^{-N}.$$

Thus,

$$\begin{aligned} & \int_{B(z, \epsilon) \times B(z, \epsilon)} \frac{|f(x) - f(y)|^p dx dy}{|x - y|^{2N}} \leq \\ & \leq \int_{\substack{x \in H_+ \cap B(z, \epsilon) \\ y \in H_+ \cap B(z, \epsilon)}} \frac{|f(x) - f(y)|^p dx dy}{|x - y|^{2N}} + \int_{\substack{x \in H_- \cap B(z, \epsilon) \\ y \in H_- \cap B(z, \epsilon)}} \frac{|f(x) - f(y)|^p dx dy}{|x - y|^{2N}} + \\ & + 2^{p-1} c_N \int_{x \in H_+ \cap B(z, \epsilon)} \frac{|f|^p(x) dx}{|x_N|^N} + 2^{p-1} c_N \int_{x \in H_- \cap B(z, \epsilon)} \frac{|f|^p(x) dx}{|x_N|^N} + \\ & + 2^{p-1} c_N \int_{y \in H_+ \cap B(z, \epsilon)} \frac{|f|^p(y) dy}{|y_N|^N} + 2^{p-1} c_N \int_{y \in H_- \cap B(z, \epsilon)} \frac{|f|^p(y) dy}{|y_N|^N}. \end{aligned}$$

Since the right hand side is finite, the assertion follows.  $\square$

**Lemma 7.4.** *Let  $\Gamma_1$  and  $\Gamma_2$  be distinct chambers sharing the same face  $P \subset \{\langle x, v \rangle = 0\}$ . Let  $z$  be in the relative interior of  $P$ . If  $f$  is as in Theorem [7.1](#), then  $f \in \dot{W}^{\frac{N}{p}, p}(B(z, \epsilon))$  for every sufficiently small  $\epsilon > 0$ .*

*Proof.* To simplify the notations, let  $v = e_N$ . Denote for brevity,  $k_{e_N} = K$ . Let  $z$

For every  $\epsilon > 0$ , we have

$$\int_{B(z, 2\epsilon) \times B(z, 2\epsilon)} \frac{|f(x) - f(y)|^p}{w(B(x, |x - y|))^2} w(x) w(y) dx dy < \infty.$$

Now, let  $\epsilon > 0$  be so small that  $|\langle x, \tilde{v} \rangle| \geq c_{z, \epsilon} > 0$  for every  $v \neq \tilde{v} \in R$  and for every  $x \in B(z, 2\epsilon)$ . It follows that

$$w(x) \geq c_{z, \epsilon, k} |x_N|^K, \quad w(y) \geq c_{z, \epsilon, k} |y_N|^K, \quad x, y \in B(z, 2\epsilon).$$

Also,

$$w(B(x, |x - y|)) \leq c_{z, \epsilon, k} |x - y|^N \cdot (|x - y| + |x_N|)^K, \quad x, y \in B(z, 2\epsilon).$$

Consequently,

$$w(B(x, |x - y|)) \leq c_{z, \epsilon, k} |x - y|^{N+K}, \quad x \in H_+ \cap B(z, 2\epsilon), \quad y \in H_- \cap B(z, 2\epsilon).$$

We, therefore, have

$$\int_{\substack{x \in H_+ \cap B(z, 2\epsilon) \\ y \in H_- \cap B(z, 2\epsilon)}} \frac{|f(x) - f(y)|^p}{|x - y|^{2N+2K}} |x_N|^K |y_N|^K dx dy < \infty.$$

On the other hand,  $H_+ \cap B(z, 2\epsilon) \subset \Gamma_1$  and  $H_- \cap B(z, 2\epsilon) \subset \Gamma_2$ . Thus,  $f \in \dot{W}^{\frac{N}{p}, p}(H_+ \cap B(z, 2\epsilon))$  and  $f \in \dot{W}^{\frac{N}{p}, p}(H_- \cap B(z, 2\epsilon))$ . By Lemma 7.2 and Lemma 7.3,  $f \in \dot{W}^{\frac{N}{p}, p}(B(z, \epsilon))$ .  $\square$

**Lemma 7.5.** *Suppose  $f \in \dot{W}^{\frac{N}{p}, p}(\Gamma)$  for every chamber  $\Gamma$ . Suppose that, for every face of every chamber and for every  $z$  in the relative interior of that face, we have  $f \in \dot{W}^{\frac{N}{p}, p}(B(z, \epsilon))$  for every sufficiently small  $\epsilon > 0$ . It follows that  $f \in \dot{W}^{\frac{N}{p}, p}(\mathbb{R}^N)$ . Furthermore, we have*

$$\|f\|_{\dot{W}^{\frac{N}{p}, p}(\mathbb{R}^N)} \leq c_R \sum_{\Gamma} \|f\|_{\dot{W}^{\frac{N}{p}, p}(\Gamma)}.$$

*Proof.* [find reference](#).  $\square$

*Proof of Theorem 7.1.* The assertion follows by combining Lemmas 7.4 and 7.5.  $\square$

*Proof of Theorem 1.1 (ii).* The assertion follows by combining Theorem 5.1, Theorem 6.1 and Theorem 7.1.  $\square$

## 8. CONSTANT CHARACTERIZATION

**Theorem 8.1.** *If  $f \in L_1^{\text{loc}}(\mathbb{R}^N, w(x)dx)$  is such that*

$$[R_{\text{Dunkl}, j}, M_f] \in \mathcal{L}_p(L_2(\mathbb{R}^N, w(x)dx))$$

*for some  $p \leq d$ , then  $f$  is constant.*

*Proof.* By Lemma 5.4, measures  $dx$  and  $w(x)dx$  and operator  $T_{\Gamma, j} = M_{\chi_{\Gamma}} R_{\text{Dunkl}, j} M_{\chi_{\Gamma}}$  satisfy [on  \$\Gamma\$](#)  the assumptions in Theorem 1.4 in [6]. By assumption,  $[T_{\Gamma, j}, M_f] \in \mathcal{L}_p(L_2(\Gamma, w(x)dx))$ . By Theorem 1.4 in [6],  $f$  is constant on  $\Gamma$ .

Hence,  $f$  is constant on every chamber. Let  $f = c_1$  on  $\Gamma_1$  and  $f = c_2$  on  $\Gamma_2$ . On the other hand,  $[R_{\text{Dunkl}, j}, M_f] \in \mathcal{L}_{2N}(L_2(\mathbb{R}^N, w(x)dx))$ . By Theorem 1.1 (ii),  $f \in \dot{W}^{\frac{1}{2}, 2N}(\mathbb{R}^N)$ . We have

$$\begin{aligned} \int_{\Gamma_1 \times \Gamma_2} \frac{|c_1 - c_2|^{2N} dx dy}{|x - y|^{2N}} &= \int_{\Gamma_1 \times \Gamma_2} \frac{|f(x) - f(y)|^{2N} dx dy}{|x - y|^{2N}} \leq \\ &\leq \int_{\mathbb{R}^N \times \mathbb{R}^N} \frac{|f(x) - f(y)|^{2N} dx dy}{|x - y|^{2N}} = \|f\|_{\dot{W}^{\frac{1}{2}, 2N}(\mathbb{R}^N)}^{2N} < \infty. \end{aligned}$$

It follows that  $c_1 = c_2$ . Since chambers  $\Gamma_1$  and  $\Gamma_2$  are arbitrary, the assertion follows.  $\square$

## 9. GLOBAL RESULT FROM HYTONEN

**Lemma 9.1.** *Let  $1 < p < \infty$ . Then, for every measurable function  $b$  on  $\mathbb{R}^N$ ,*

$$\|b\|_{B^p(\mathbb{R}^N, dw)} \sim \|b\|_{B^p(\mathbb{R}^N, dx)},$$

where

$$\|b\|_{B^p(\mathbb{R}^N, d\mu)} := \left( \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|b(x) - b(y)|^p}{\mu(B(x, \|x - y\|))^2} d\mu(x) d\mu(y) \right)^{1/p}.$$

*Proof.* Write

$$h_k(x) := \prod_{v \in R} |\langle x, v \rangle|^{k(v)}, \quad dw(x) = h_k(x) dx.$$

$dx$  is obvious doubling, and  $dw$  is doubling by (2.2).

We first verify that  $dx$  and  $dw$  satisfy the  $A_\infty$ -relation used in [6, Proposition 1.2]. Fix  $s > 1$ , and set

$$dw_s(x) := h_k(x)^s dx = \prod_{v \in R} |\langle x, v \rangle|^{sk(v)} dx.$$

We have

$$w_s(B(x, r)) \sim r^N \prod_{v \in R} (|\langle x, v \rangle| + r)^{sk(v)}.$$

On the other hand, (2.1) gives

$$w(B(x, r)) \sim r^N \prod_{v \in R} (|\langle x, v \rangle| + r)^{k(v)}.$$

Since  $|B(x, r)| \sim r^N$ , we obtain

$$\begin{aligned} \left( \int_{B(x, r)} h_k(z)^s dz \right)^{1/s} &= \left( \frac{w_s(B(x, r))}{|B(x, r)|} \right)^{1/s} \\ &\sim \prod_{v \in R} (|\langle x, v \rangle| + r)^{k(v)} \\ &\sim \frac{w(B(x, r))}{|B(x, r)|} = \int_{B(x, r)} h_k(z) dz. \end{aligned}$$

Thus  $h_k \in RH_s(dx)$ . Hence, by [6, Proposition 3.6], the two doubling measures  $dx$  and  $dw = h_k dx$  satisfy the  $A_\infty$ -relation

$$|E| \leq \varepsilon |B| \implies w(E) \leq (1 - \delta) w(B)$$

for some  $\varepsilon, \delta \in (0, 1)$ , every Euclidean ball  $B$ , and every measurable set  $E \subset B$ .

Applying [6, Proposition 1.2] with  $\mu = dx$ ,  $\nu = dw$ , and  $q = p$ , we get

$$\|b\|_{\text{Osc}^{p,p}(\mathbb{R}^N, dx)} \sim \|b\|_{\text{Osc}^{p,p}(\mathbb{R}^N, dw)}.$$

Finally, since both

$$(\mathbb{R}^N, \|\cdot\|, dx) \quad \text{and} \quad (\mathbb{R}^N, \|\cdot\|, dw)$$

are doubling metric measure spaces, [7, Proposition 11.5] yields

$$\|b\|_{B^p(\mathbb{R}^N, dx)} \sim \|b\|_{\text{Osc}^{p,p}(\mathbb{R}^N, dx)}$$

and

$$\|b\|_{B^p(\mathbb{R}^N, dw)} \sim \|b\|_{\text{Osc}^{p,p}(\mathbb{R}^N, dw)}.$$

Combining the last three equivalences gives

$$\|b\|_{B^p(\mathbb{R}^N, dw)} \sim \|b\|_{B^p(\mathbb{R}^N, dx)}.$$

This completes the proof.  $\square$

### Acknowledgements:

Z. Fan is supported by the Guangdong Basic and Applied Basic Research Foundation (No. 2023A1515110879). J. Li is supported by the Australian Research Council (ARC) through the research grant DP220100285.

### REFERENCES

- [1] Z. Fan, M. Lacey, and J. Li. Schatten classes and commutators of Riesz transform on Heisenberg group and applications. *J. Fourier Anal. Appl.*, 29(2):Paper No. 17, 28, 2023. [5](#)
- [2] Z. Fan, M. Lacey, J. Li, M.N. Vempati, and B.D. Wick. Besov space, Schatten classes, and commutators of Riesz transforms associated with the Neumann Laplacian. *Michigan Math. J.*, 75(1):199–224, 2025. [5](#)
- [3] Z. Fan, M.T. Lacey, J. Li, and X. Xiong. Schatten–Lorentz characterization of Riesz transform commutator associated with Bessel operators. Preprint available at <https://arxiv.org/abs/2403.08249>, 2024. [5](#)
- [4] Y. Han, M. Lee, J. Li, and B.D. Wick. Riesz transforms and commutators in the Dunkl setting. *Anal. Math. Phys.*, 14(3):Paper No. 46, 32, 2024. [4](#), [5](#), [14](#), [15](#)
- [5] S. Helgason. *Differential geometry, Lie groups, and symmetric spaces*, volume 80 of *Pure and Applied Mathematics*. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York-London, 1978. [4](#)
- [6] T. Hytonen.  $A_\infty$ -invariance of oscillatory norms and Schatten characterisations of commutators. [2](#), [14](#), [15](#), [23](#), [24](#)
- [7] T. Hytonen. Schatten properties of commutators on metric spaces. Preprint available at <https://arxiv.org/abs/2411.02613>, 2024. [5](#), [10](#), [11](#), [12](#), [24](#)
- [8] S. Janson and T.H. Wolff. Schatten classes and commutators of singular integral operators. *Ark. Mat.*, 20(2):301–310, 1982. [10](#)
- [9] M.T. Lacey, J. Li, and B.D. Wick. Schatten classes and commutator in the two weight setting, I. Hilbert transform. *Potential Anal.*, 60(2):875–894, 2024. [5](#)
- [10] J. Li, X. Xiong, and F. Yang. Schatten properties of Calderón–Zygmund singular integral commutator on stratified Lie groups. *J. Math. Pures Appl. (9)*, 188:73–113, 2024. [5](#)
- [11] R. Rochberg and S. Semmes. Nearly weakly orthonormal sequences, singular value estimates, and Calderon–Zygmund operators. *J. Funct. Anal.*, 86(2):237–306, 1989. [5](#)
- [12] Z. Wei and H. Zhang. Complex median method and Schatten class membership of commutators. Preprint available at <https://arxiv.org/abs/2411.05810>, 2024. [5](#)
- [13] Z. Wei and H. Zhang. On the boundedness and Schatten class property of noncommutative martingale paraproducts and operator-valued commutators. Preprint available at <https://arxiv.org/abs/2407.14988>, 2024. [5](#)

ZHIJIE FAN, SCHOOL OF MATHEMATICAL SCIENCES, SOUTH CHINA NORMAL UNIVERSITY, GUANGZHOU 510631, CHINA

*E-mail address:* [fanzhj3@mail2.sysu.edu.cn](mailto:fanzhj3@mail2.sysu.edu.cn)

Ji LI, DEPARTMENT OF MATHEMATICS, MACQUARIE UNIVERSITY, NSW 2109, AUSTRALIA

*E-mail address:* [ji.li@mq.edu.au](mailto:ji.li@mq.edu.au)

SCHOOL OF MATHEMATICS AND STATISTICS, CENTRAL SOUTH UNIVERSITY, CHANGSHA 410075, PEOPLE’S REPUBLIC OF CHINA

*E-mail address:* [d.zanin@csu.edu.cn](mailto:d.zanin@csu.edu.cn)