WEYL–CHAMBER GEOMETRY, POINCARÉ INEQUALITY MEETS SCHATTEN CLASS PROPERTIES OF DUNKL–RIESZ TRANSFORM COMMUTATORS

ZHIJIE FAN, JI LI, AND DMITRIY ZANIN

Abstract.

Contents

1. Introduction 1
1.1. Background and motivation 1
1.2. Notation 2
2. Preliminaries 2
2.1. Harmonic analysis in the Dunkl setting 2
2.2. Schatten class 4
2.3. Nearly weakly orthonormal sequences 5
3. Comparison of Sobolev norms 5
4. Proof of Theorem 1.1 (i) 10
5. Expected dyadic result 12
6. Simplified version of Zhijie's argument 13
7. Proof of Theorem 1.1 (ii) 17
8. Useful part of Zhijie argument 20
9. Constant characterization 27
References 27

1. Introduction

1.1. Background and motivation.

[Zhijie will write the background in the Dunkl setting.]

Theorem 1.1. *Let R be a normalised root system in $\mathbb{R}^N$ and let G be the corresponding finite reflection group. Let $k : R \to \mathbb{R}_+$ be G-invariant multiplicity function. Let $R_{\mathrm{Dunkl},j}$ be the j-th Dunkl-Riesz transform corresponding to R and k. Let w be the Dunkl weight corresponding to R and k. Let $f \in L_1^{\mathrm{loc}}(\mathbb{R}^N)$.*

(i) *If $f \in \dot{W}^{\frac{N}{p},p}(\mathbb{R}^N)$, $p > N$, then $[R_{\mathrm{Dunkl},j}, M_f] \in \mathcal{L}_p(L_2(\mathbb{R}^N, w(x)dx))$. Furthermore,*

$$\|[R_{\mathrm{Dunkl},j}, M_f]\|_{\mathcal{L}_p(L_2(\mathbb{R}^N,w(x)dx))} \le c_{p,R,k}\|f\|_{\dot{W}^{\frac{N}{p},p}(\mathbb{R}^N)};$$

Date: May 6, 2026.
2010 *Mathematics Subject Classification.* 47B10, 42B20, 43A85.
Key words and phrases. Schatten class, Riesz transform commutators, Dunkl operator, Besov space, Nearly weakly orthogonal sequence.

(ii) *If* $[R_{\mathrm{Dunkl},j}, M_f] \in \mathcal{L}_p(L_2(\mathbb{R}^N, w(x)dx))$, $p > N$, *then* $f \in \dot{W}^{\frac{N}{p},p}(\mathbb{R}^N)$. *Furthermore,*

$$\|[R_{\mathrm{Dunkl},j}, M_f]\|_{\mathcal{L}_p(L_2(\mathbb{R}^N,w(x)dx))} \le c_{p,R,k}^{-1}\|f\|_{\dot{W}^{\frac{N}{p},p}(\mathbb{R}^N)};$$

(iii) *If* $f \in \dot{W}^{1,N}(\mathbb{R}^N)$, *then* $[R_{\mathrm{Dunkl},j}, M_f] \in \mathcal{L}_{N,\infty}(L_2(\mathbb{R}^N, w(x)dx))$. *Furthermore,*

$$\|[R_{\mathrm{Dunkl},j}, M_f]\|_{\mathcal{L}_{N,\infty}(L_2(\mathbb{R}^N,w(x)dx))} \le c_{R,k}\|f\|_{\dot{W}^{1,N}(\mathbb{R}^N)};$$

(iv) *If* $[R_{\mathrm{Dunkl},j}, M_f] \in \mathcal{L}_{N,\infty}(L_2(\mathbb{R}^N, w(x)dx))$, *then* $f \in \dot{W}^{1,N}(\mathbb{R}^N)$. *Furthermore,*

$$\|[R_{\mathrm{Dunkl},j}, M_f]\|_{\mathcal{L}_{N,\infty}(L_2(\mathbb{R}^N,w(x)dx))} \ge c_{R,k}^{-1}\|f\|_{\dot{W}^{1,N}(\mathbb{R}^N)};$$

1.2. **Notation.** Conventionally, we let $\mathbb{N}$ denote the set of positive integers and set $\mathbb{Z}_+ := \{0\} \cup \mathbb{N}$. Throughout the paper, the symbols C and c denote positive constants (possibly varying from line to line) that may depend only on N, p, and the structural data (R, k) of the Dunkl setting, but are independent of the main parameters (e.g., the symbol b and auxiliary decay factors). We write $A \lesssim B$ to mean that $A \le CB$ for some constant $C > 0$, and $A \sim B$ to mean that both $A \lesssim B$ and $B \lesssim A$ hold. For $\lambda > 0$ and a cube Q with sidelength $\ell(Q)$, write λQ for the cube concentric with Q and sidelength $\lambda\,\ell(Q)$. Similarly, for every ball $B = B(x, r)$, set $\lambda B := B(x, \lambda r)$. For any $1 \le p \le \infty$, we denote by p' the conjugate of p, which satisfies $\frac{1}{p} + \frac{1}{p'} = 1$. We denote the average of a function f over a μ-measurable set E by

$$(f)_E := \fint_E f(x)d\mu(x) := \frac{1}{\mu(E)}\int_E f(x)d\mu(x). \tag{1.1}$$

For a subset $E \subset \mathbb{R}^N$, we denote by χ_E its characteristic function.

Denote by $\{e_j\}_{j=1}^N$ the standard orthonormal basis on $\mathbb{R}^N$.

2. Preliminaries

2.1. **Harmonic analysis in the Dunkl setting.** Let $\mathbb{R}^N$ be the Euclidean space equipped with the usual scalar product $\langle\cdot,\cdot\rangle$ and the induced norm $\|\cdot\|$. For a nonzero vector $v \in \mathbb{R}^N$, the reflection σ_v with respect to the hyperplane orthogonal to a normalised vector v is defined by

$$\sigma_v x := x - 2\langle x, v\rangle v.$$

A finite set $R \subset \mathbb{R}^N\backslash\{0\}$ is called a root system if $\sigma_v(R) = R$ for every $v \in R$. A root system R is called a normalized reduced root system if $|v| = 1$ for every $v \in R$. The finite group G generated by the reflections $\{\sigma_v\}_{v\in R}$ is called the Weyl group (reflection group) of the root system R. Let $k : R \to [0, \infty)$ be a multiplicity function, which is a G-invariant function fixed throughout the paper.

Define the Dunkl measure $w(x)dx$ on $\mathbb{R}^N$ by

$$w(x) = \prod_{v\in R} |\langle x, v\rangle|^{k(v)}.$$

Let $B(x, r) := \{y \in \mathbb{R}^N : \|x - y\| < r\}$ be the Euclidean ball centred at x with radius $r > 0$. Let $\mathbf{N} := N + \sum_{v\in R} k(v)$. Note that

$$w(B(tx, tr)) = t^{\mathbf{N}} w(B(x, r)), \text{ for } x \in \mathbb{R}^N,\ t, r > 0.$$

Furthermore,

$$w(B(x,r)) \sim r^N \prod_{v\in R} (|\langle x,v\rangle| + r)^{k(v)}, \tag{2.1}$$

where the implicit constant only depends on N, R and k. It follows from (2.1) that there exists a constant $C \geq 1$ such that for every $x \in \mathbb{R}^N$,

$$C^{-1}\left(\frac{r_2}{r_1}\right)^N \leq \frac{w(B(x,r_2))}{w(B(x,r_1))} \leq C\left(\frac{r_2}{r_1}\right)^{\mathbf{N}}, \text{ for } 0 < r_1 < r_2, \tag{2.2}$$

so the numbers $\mathbf{N}$ and N are called the upper and lower homogeneous dimensions, respectively.

The Dunkl operators T_ξ are defined by

$$T_\xi f(x) := \partial_\xi f(x) + \sum_{v\in R} \frac{k(v)}{2}\langle v,\xi\rangle \frac{f(x) - f(\sigma_v(x))}{\langle v,x\rangle},$$

where $\partial_\xi f$ denotes the directional derivative of f in the direction ξ. Furthermore, the Dunkl Laplacian is defined by

$$\Delta_{\mathrm{Dunkl}} f(x) := \sum_{j=1}^N T_{e_j}^2 f(x) = \Delta_{\mathbb{R}^N} f(x) + \sum_{v\in R} k(v)\delta_v f(x),$$

where $\Delta_{\mathbb{R}^N}$ is the Laplacian on $\mathbb{R}^N$ and δ_v is defined by

$$\delta_v f(x) := \frac{\partial_v f(x)}{\langle v,x\rangle} - \frac{f(x) - f(\sigma_v(x))}{\langle v,x\rangle^2}.$$

In this article, we consider the Dunkl-Riesz transforms defined by $R_{\mathrm{Dunkl},j} := T_{e_j}(-\Delta_{\mathrm{Dunkl}})^{-1/2}$, $j = 1, 2, \ldots, N$.

Definition 2.1. The Weyl chambers are defined as the closures of connected components of the set

$$\left\{x \in \mathbb{R}^N : \langle x,v\rangle \neq 0 \ \ \textit{for all } v \in R\right\}. \tag{2.3}$$

Define

$$d(x,y) = \min_{\sigma\in G} |x - \sigma(y)|, \quad x,y \in \mathbb{R}^N.$$

Recall from [5, Chapter VII, proof of Theorem 2.12] that

$$d(x,y) = \|x-y\| \text{ if and only if } x,y \in \Gamma \tag{2.4}$$

for some closed Weyl chamber Γ.

Denote by $K_{\mathrm{Dunkl},j}(x,y)$ the integral kernel of $R_{\mathrm{Dunkl},j}$. We recall the kernel estimates for the Riesz transforms in the Dunkl setting proved in [4]. Although these estimates may not be sharp (see the comment following Theorem 2.1 in [4, p. 10]), they suffice for our purposes.

Lemma 2.2. *There exists a constant C such that for $j \in \{1, 2, \cdots, N\}$ and every x, y with $d(x,y) \neq 0$,*

$$|K_{\mathrm{Dunkl},j}(x,y)| \leq \frac{d(x,y)}{|x-y|} \cdot \frac{C}{w(B(x,d(x,y)))}.$$

Proof. The proof is given in [4, Theorem 1.1]. □

Lemma 2.3. *We have*

$$d(x,y)^2 \cdot w(x)w(y) \le Cw^2(B(x,d(x,y))), \quad x,y \in \mathbb{R}^N.$$

Proof. Define equivalence relation $\sim$ on R by saying $v_1 \sim v_2$ if there exists $g \in G$ with $v_1 = gv_2$. Let us split R into equivalence classes with respect to $\sim$. Since k is G-invariant, it follows that k is constant on each equivalence class. Note that G leaves each equivalence class invariant.

We, therefore, write

$$w(x) = \prod_{A \in R/\sim} \big(\prod_{v \in A} |\langle x, v\rangle| \big)^{k|_A}.$$

On the other hand,

$$w(B(x,r)) \ge c^{(1)}_{R,k} r^N \prod_{A \in R/\sim} \big(\prod_{v \in A} (r + |\langle x, v\rangle|) \big)^{k|_A}.$$

Let $x \in \Gamma$ and $y \in g\Gamma$. Set $z = g^{-1}y \in \Gamma$. Note that $d(x,y) = |x-z|$ and, therefore,

$$|\langle x, v\rangle| + d(x,y) = |\langle x, v\rangle| + |x-z| \ge 2^{-\frac{1}{2}} |\langle z, v\rangle|.$$

Thus,

$$\prod_{v \in A} (|\langle x, v\rangle| + d(x,y)) \ge \prod_{v \in A} 2^{-\frac{1}{2}} |\langle z, v\rangle| = 2^{-\frac{1}{2}|A|} \prod_{v \in A} |\langle g^{-1}y, v\rangle| =$$

$$= 2^{-\frac{1}{2}|A|} \prod_{v \in A} |\langle y, gv\rangle| = 2^{-\frac{1}{2}|A|} \prod_{v \in gA} |\langle y, v\rangle| = 2^{-\frac{1}{2}|A|} \prod_{v \in A} |\langle y, v\rangle|.$$

Thus,

$$w(B(x,d(x,y))) \ge c^{(1)}_{R,k} d^N(x,y) \prod_{A \in R/\sim} \big(\prod_{v \in A} (d(x,y) + |\langle x, v\rangle|) \big)^{k|_A} \ge$$

$$\ge c^{(1)}_{R,k} d^N(x,y) \prod_{A \in R/\sim} \big(2^{-\frac{1}{2}|A|} \prod_{v \in A} |\langle y, v\rangle| \big)^{k|_A} = c^{(2)}_{R,k} d^N(x,y) w(y).$$

On the other hand,

$$w(B(x,d(x,y))) \ge c^{(1)}_{R,k} d^N(x,y) w(x).$$

Multiplying the last two equations, we complete the proof. □

Lemma 2.4. *Let $1 \le j \le N$. If $x_j \ge y_j$, then*

$$K_{\mathrm{Dunkl},j}(x,y) \ge c_{R,k} \frac{x_j - y_j}{|x-y|} \cdot \frac{1}{w(B(x,|x-y|))}.$$

If $x_j \le y_j$, then

$$K_{\mathrm{Dunkl},j}(x,y) \le c_{R,k} \frac{x_j - y_j}{|x-y|} \cdot \frac{1}{w(B(x,|x-y|))}.$$

Proof. The lower bound is identical to that in [4, Theorem 1.2]. □

2.2. **Schatten class.** [Zhijie will add this later]

2.3. **Nearly weakly orthonormal sequences.** Rochberg and Semmes [11] introduced the notion of *nearly weakly orthonormal (NWO)* sequences of functions. This notion has since proved highly effective in characterizing Schatten class membership of singular integral commutators in a variety of contexts (see [3, 7, 10, 1, 2, 9, 11, 13, 12]). The properties of NWO sequences were originally investigated in the Euclidean setting (equipped with Lebesgue measure), but the extension to spaces of homogeneous type is straightforward once a system of dyadic cubes is available in this general framework (see [7, Section 13] for a complete proof). For our purposes, we adapt the extension to the Dunkl setting $(\mathbb{R}^N, \|\cdot\|, w)$ rather than recalling the general definition of NWO sequences and the general extension to spaces of homogeneous type.

Let $\mathscr{D}$ be a system of dyadic cubes. A sequence of functions $\{e_Q\}_{Q\in\mathscr{D}}$ is an NWO sequence if

(1) $\operatorname{supp} e_Q \subseteq cQ$ *for some positive constant c independent of Q*
(2) $\|e_Q\|_{L^s(\mathbb{R}^N,dw)} \leq w(Q)^{1/s-1/2}$ *for some $s > 2$.*

Lemma 2.5. *Let $1 < p < \infty$, and let $\mathscr{D}$ be a system of dyadic cubes over $\mathbb{R}^N$. For any compact operator T on $L^2(\mathbb{R}^N, dw)$, we have*

$$\Big(\sum_{Q\in\mathscr{D}} |\langle Te_Q, f_Q\rangle_{L^2(\mathbb{R}^N,dw)}|^p\Big)^{1/p} \leq C_{N,p,w}\|T\|_{S^p},$$

where $\{e_Q\}_{Q\in\mathscr{D}}$ and $\{f_Q\}_{Q\in\mathscr{D}}$ are NWO sequences.

Proof. This property can be found in [11, (1.10)] (see also [7, Section 7]). □

Lemma 2.6. *Let $0 < p < \infty$, and let $\mathscr{D}$ be a system of dyadic cubes over $\mathbb{R}^N$. Assume that*

$$A = \sum_{Q\in\mathscr{D}} \lambda_Q \langle\cdot,\, e_Q\rangle_{L^2(\mathbb{R}^N,dw)} f_Q,$$

where $\{e_Q\}_{Q\in\mathscr{D}}$ and $\{f_Q\}_{Q\in\mathscr{D}}$ are NWO sequences and $\{\lambda_Q\}_{Q\in\mathscr{D}}$ is a sequence of scalars. We have

$$\|A\|_{S^p} \leq C_{N,p,w}\|\{\lambda_Q\}_Q\|_{\ell^p}. \tag{2.5}$$

Proof. This property can be found in [11, (1.9)] (see also [7, Section 7]). □

3. Comparison of Sobolev norms

Theorem 3.1. *For every $f \in \dot{W}^{\frac{N}{p},p}(\mathbb{R}^N)$, $p > N$, we have*

$$\Big(\int_{\mathbb{R}^N\times\mathbb{R}^N} \Big(\frac{d(x,y)}{|x-y|}\Big)^p |f(x)-f(y)|^p \frac{dxdy}{d(x,y)^{2N}}\Big)^{\frac{1}{p}} \leq c_{p,R}\|f\|_{\dot{W}^{\frac{N}{p},p}(\mathbb{R}^N)}.$$

Lemma 3.2. *For every convex polyhedral cone Γ, we have*

$$\int_\Gamma \frac{dx}{|x-y|^{2N}} \geq c_\Gamma \operatorname{dist}(y,\Gamma)^{-N}, \quad y \notin \Gamma.$$

Proof. Find a point $z \in \Gamma$ such that

$$|y-z| = \operatorname{dist}(y,\Gamma).$$

We have $z + \Gamma \subset \Gamma$. Thus,

$$\int_\Gamma \frac{dx}{|x-y|^{2N}} \geq \int_{z+\Gamma} \frac{dx}{|x-y|^{2N}} = \int_\Gamma \frac{dx}{|x-(y-z)|^{2N}}.$$

Clearly,

$$|x-(y-z)| \le |x|+|y-z|.$$

Thus,

$$\int_\Gamma \frac{dx}{|x-y|^{2N}} \ge \int_\Gamma \frac{dx}{(|x|+|y-z|)^{2N}} = |y-z|^{-N}\int_\Gamma \frac{dx}{(|x|+1)^{2N}}.$$

□

Lemma 3.3. *If $p\in(N,2N)$, then*

$$\sup_{a>0}\int_{\mathbb{R}^N}\frac{|x-ae_N|^{p-2N}dx}{(1+|x|)^p}<\infty.$$

Proof. **Step 1:** we claim that, for every $a>0$,

$$\int_{\mathbb{R}^N}\frac{|x-ae_N|^{p-2N}dx}{(1+|x|)^p} \le$$

$$\le 4^p\int_{\mathbb{R}^{N-1}\times(-a,a)}\frac{|x|^{p-2N}dx}{(1+|x'|+a)^p}+3^{2N-p}\int_{\mathbb{R}^N}\frac{|x|^{p-2N}dx}{(1+|x|)^p}.$$

We clearly have

$$\int_{\mathbb{R}^N}\frac{|x-ae_N|^{p-2N}dx}{(1+|x|)^p} =$$

$$=\int_{\mathbb{R}^{N-1}\times(\frac{a}{2},\frac{3a}{2})}\frac{|x-ae_N|^{p-2N}dx}{(1+|x|)^p}+\int_{\mathbb{R}^{N-1}\times\mathbb{R}\setminus(\frac{a}{2},\frac{3a}{2})}\frac{|x-ae_N|^{p-2N}dx}{(1+|x|)^p}.$$

If $x_N\notin(\frac{a}{2},\frac{3a}{2})$, then $|x_N-a|\ge\frac{1}{3}|x_N|$. Hence, $|x-ae_N|\ge\frac{1}{3}|x|$. Thus,

$$\int_{\mathbb{R}^{N-1}\times\mathbb{R}\setminus(\frac{a}{2},\frac{3a}{2})}\frac{|x-ae_N|^{p-2N}dx}{(1+|x|)^p} \le$$

$$\le\int_{\mathbb{R}^{N-1}\times\mathbb{R}\setminus(\frac{a}{2},\frac{3a}{2})}\frac{3^{2N-p}|x|^{p-2N}dx}{(1+|x|)^p}\le\int_{\mathbb{R}^N}\frac{3^{2N-p}|x|^{p-2N}dx}{(1+|x|)^p}.$$

If $x_N\in(\frac{a}{2},\frac{3a}{2})$, then (here, we using the notation $x'=(x_1,\cdots,x_{N-1})$)

$$1+|x|\ge 1+\max\{|x'|,\frac{1}{2}a\}\ge\frac{1}{4}(1+|x'|+a).$$

Thus,

$$\int_{\mathbb{R}^{N-1}\times(\frac{a}{2},\frac{3a}{2})}\frac{|x-ae_N|^{p-2N}dx}{(1+|x|)^p}\le 4^p\int_{\mathbb{R}^{N-1}\times(\frac{a}{2},\frac{3a}{2})}\frac{|x-ae_N|^{p-2N}dx}{(1+|x'|+a)^p}=$$

$$=4^p\int_{\mathbb{R}^{N-1}\times(-\frac{a}{2},\frac{a}{2})}\frac{|x|^{p-2N}dx}{(1+|x'|+a)^p}\le 4^p\int_{\mathbb{R}^{N-1}\times(-a,a)}\frac{|x|^{p-2N}dx}{(1+|x'|+a)^p}.$$

Combining all preceding estimates, we infer the claim in Step 1.

Step 2: we claim that

$$\sup_{a>0}\int_{\mathbb{R}^{N-1}\times(-a,a)}\frac{|x|^{p-2N}dx}{(1+|x'|+a)^p}<\infty.$$

Passing to polar coordinates in $\mathbb{R}^{N-1}$, we equivalently rewrite the claim as

$$\sup_{a>0}\int_{\mathbb{R}_+\times(0,a)}\frac{(r^2+x_N^2)^{\frac{p}{2}-N}r^{N-2}drdx_N}{(1+r+a)^p}<\infty.$$

If $0 < a < 1$, then

$$\int_{\mathbb{R}_+\times(0,a)} \frac{(r^2+x_N^2)^{\frac{p}{2}-N} r^{N-2} dr dx_N}{(1+r+a)^p} \le \int_{\mathbb{R}_+\times(0,1)} \frac{(r^2+x_N^2)^{\frac{p}{2}-N} r^{N-2} dr dx_N}{(1+r)^p}.$$

If $a \ge 1$, then

$$\int_{\mathbb{R}_+\times(0,a)} \frac{(r^2+x_N^2)^{\frac{p}{2}-N} r^{N-2} dr dx_N}{(1+r+a)^p} \le \int_{\mathbb{R}_+\times(0,a)} \frac{(r^2+x_N^2)^{\frac{p}{2}-N} r^{N-2} dr dx_N}{(r+a)^p} =$$

$$= \int_{(a,\infty)\times(0,a)} \frac{(r^2+x_N^2)^{\frac{p}{2}-N} r^{N-2} dr dx_N}{(r+a)^p} +$$

$$+ \int_0^a \int_0^r \frac{(r^2+x_N^2)^{\frac{p}{2}-N} r^{N-2} dx_N dr}{(r+a)^p} + \int_0^a \int_0^{x_N} \frac{(r^2+x_N^2)^{\frac{p}{2}-N} r^{N-2} dr dx_N}{(r+a)^p} \le$$

$$\le {}^{1} \int_{(a,\infty)\times(0,a)} \frac{r^{p-2N} r^{N-2} dr dx_N}{r^p} +$$

$$+ \int_0^a \int_0^r \frac{r^{p-2N} r^{N-2} dx_N dr}{a^p} + \int_0^a \int_0^{x_N} \frac{x_N^{p-2N} r^{N-2} dr dx_N}{a^p} =$$

$$= \int_a^\infty r^{-N-1} dr + a^{-p} \int_0^a r^{p-N-1} dr + \frac{1}{N-1} a^{-p} \int_0^a x_N^{p-N-1} dx_N =$$

$$= \frac{1}{N} a^{-N} + \frac{1}{p-N} a^{-N} + \frac{1}{(N-1)(p-N)} a^{-N} \le \frac{1}{N} + \frac{1}{p-N} + \frac{1}{(N-1)(p-N)}.$$

This yields the claim in Step 2.

Combining Steps 1 and 2, we complete the proof of the lemma. □

Lemma 3.4. *If $p \in (N, 2N)$, then*

$$\int_{\mathbb{R}^{N-1}\times\mathbb{R}_+} \frac{|x-y|^{p-2N} dx}{|x+z|^p} \le c_{p,N} z_N^{-N}, \quad y, z \in \mathbb{R}^{N-1}\times\mathbb{R}_+.$$

Proof. By dilation invariance, we may assume without loss of generality that $z_N = 1$. By translation invariance in $\mathbb{R}^{N-1}$, we may assume without loss of generality that $z_1 = \cdots = z_{N-1} = 0$. We, therefore, have

$$|x+z| = |x+e_N| \ge \max\{1, |x|\} \ge \frac{1}{2}(1+|x|), \quad x \in \mathbb{R}^{N-1}\times\mathbb{R}_+.$$

Thus,

$$\int_{\mathbb{R}^{N-1}\times\mathbb{R}_+} \frac{|x-y|^{p-2N} dx}{|x+z|^p} \le 2^p \int_{\mathbb{R}^{N-1}\times\mathbb{R}_+} \frac{|x-y|^{p-2N} dx}{(1+|x|)^p} \le$$

$$= 2^p \int_{\mathbb{R}^N} \frac{|x-y|^{p-2N} dx}{(1+|x|)^p} = 2^p \int_{\mathbb{R}^N} \frac{\big|x-|y|e_N\big|^{p-2N} dx}{(1+|x|)^p}.$$

The assertion follows from Lemma 3.3. □

We need the following elementary version of the Hahn-Banach theorem.

Lemma 3.5. *Let $K \subset \mathbb{R}^N$ be a convex set and let $y \notin K$. Let $z \in K$ be the point closest to y. Affine hyperplane P passing through z and orthogonal to $y - z$ splits the space into two half-spaces, one of them containing y, the other one containing K.*

[1] It is exactly at this point we are using the assumption $p < 2N$.

Proof. By translation invariance, we may assume without loss of generality that $z = 0$. By rotation and dilation invariance, we may assume without loss of generality that $y = e_N$. By assumption, $\mathrm{dist}(e_N, K) = 1$. We need to show that K is contained in the (closed) lower half-plane.

Assume the contrary: let $x \in K$ be such that $x_N > 0$. Since $0 \in K$ and since K is convex, it follows that $\epsilon x \in K$ for every $\epsilon \in (0,1)$. Thus,

$$\mathrm{dist}(e_N, K) \leq |e_N - \epsilon x|, \quad \epsilon \in (0,1).$$

Since $\epsilon \in (0,1)$ is arbitrary, it follows that

$$\mathrm{dist}(e_N, K) \leq \inf_{\epsilon \in (0,1)} |e_N - \epsilon x| = \inf \big(\epsilon^2 \sum_{k=1}^{N-1} x_k^2 + (1 - \epsilon x_N)^2\big)^{\frac{1}{2}}.$$

Since $x_N > 0$, it follows that infimum is strictly less than 1. Thus, $\mathrm{dist}(e_N, K) < 1$. However, it is granted in the lemma that $\mathrm{dist}(e_N, K) = 1$. This contradiction shows that such an x cannot exist. This completes the proof. □

Lemma 3.6. *If $p \in (N, 2N)$, then*

$$\int_\Gamma \frac{|x-y|^{p-2N} dx}{|x - gy|^p} \leq c_p \mathrm{dist}(gy, \Gamma)^{-N}, \quad y \in \Gamma.$$

Proof. Find a point $z \in \Gamma$ such that

$$|gy - z| = \mathrm{dist}(gy, \Gamma).$$

By Lemma 3.5, affine hyperplane passing through z and orthogonal to $gy - z$ splits the space into two half-spaces, one of them containing gy, the other one containing Γ (denote this one by H).

We have $y \in H$, $\Gamma \subset H$ and $gy \notin H$. So,

$$\int_\Gamma \frac{|x-y|^{p-2N} dx}{|x - gy|^p} \leq \int_H \frac{|x-y|^{p-2N} dx}{|x - gy|^p}.$$

Using Lemma 3.4, we write

$$\int_H \frac{|x-y|^{p-2N} dx}{|x - gy|^p} \leq c_{p,N} \mathrm{dist}(gy, H)^{-N}.$$

Since $z \in \partial H$ and since $gy - z$ is orthogonal to ∂H, it follows that $\mathrm{dist}(gy, H) = |gy - z| = \mathrm{dist}(gy, \Gamma)$. □

Proof of Theorem 3.1. **Case 1:** Let $p \in (N, 2N)$.

We have

$$\Big(\int_{\Gamma \times g\Gamma} \Big(\frac{d(x,y)}{|x-y|}\Big)^p |f(x) - f(y)|^p \frac{dxdy}{d(x,y)^{2N}}\Big)^{\frac{1}{p}} =$$

$$= \Big(\int_{\Gamma \times \Gamma} \Big(\frac{|x-y|}{|x-gy|}\Big)^p |f(x) - f(gy)|^p \frac{dxdy}{|x-y|^{2N}}\Big)^{\frac{1}{p}} \leq$$

$$\leq \Big(\int_{\Gamma \times \Gamma} \Big(\frac{|x-y|}{|x-gy|}\Big)^p |f(x) - f(y)|^p \frac{dxdy}{|x-y|^{2N}}\Big)^{\frac{1}{p}} +$$

$$+ \Big(\int_{\Gamma \times \Gamma} \Big(\frac{|x-y|}{|x-gy|}\Big)^p |f(y) - f(gy)|^p \frac{dxdy}{|x-y|^{2N}}\Big)^{\frac{1}{p}} \leq$$

$$\leq \Big(\int_{\Gamma \times \Gamma} |f(x) - f(y)|^p \frac{dxdy}{|x-y|^{2N}}\Big)^{\frac{1}{p}} +$$

$$+\Big(\int_\Gamma |f(y)-f(gy)|^p\Big(\int_\Gamma \frac{|x-y|^{p-2N}}{|x-gy|^p}dx\Big)dy\Big)^{\frac{1}{p}}.$$

Obviously, the first summand on the right hand side is less than $\|f\|_{\dot{W}^{\frac{N}{p},p}(\mathbb{R}^N)}$. By Lemmas 3.6 and 3.2,

$$\Big(\int_{\Gamma\times g\Gamma}\Big(\frac{d(x,y)}{|x-y|}\Big)^p|f(x)-f(y)|^p\frac{dxdy}{d(x,y)^{2N}}\Big)^{\frac{1}{p}}\leq \|f\|_{\dot{W}^{\frac{N}{p},p}(\mathbb{R}^N)}+$$

$$+c^{(1)}_{p,R}\Big(\int_\Gamma |f(y)-f(gy)|^p\Big(\int_\Gamma \frac{dx}{|x-gy|^{2N}}\Big)dy\Big)^{\frac{1}{p}}=$$

$$=\|f\|_{\dot{W}^{\frac{N}{p},p}(\mathbb{R}^N)}+c^{\frac{1}{p}}_{p,R}\Big(\int_{\Gamma\times\Gamma}\frac{|f(y)-f(gy)|^p dxdy}{|x-gy|^{2N}}\Big)^{\frac{1}{p}}\leq$$

$$\leq \|f\|_{\dot{W}^{\frac{N}{p},p}(\mathbb{R}^N)}+c^{(1)}_{p,R}\Big(\int_{\Gamma\times\Gamma}\frac{|f(x)-f(gy)|^p dxdy}{|x-gy|^{2N}}\Big)^{\frac{1}{p}}+$$

$$+c^{(1)}_{p,R}\Big(\int_{\Gamma\times\Gamma}\frac{|f(x)-f(y)|^p dxdy}{|x-gy|^{2N}}\Big)^{\frac{1}{p}}.$$

Now,

$$\Big(\int_{\Gamma\times\Gamma}\frac{|f(x)-f(gy)|^p dxdy}{|x-gy|^{2N}}\Big)^{\frac{1}{p}}=\Big(\int_{\Gamma\times g\Gamma}\frac{|f(x)-f(y)|^p dxdy}{|x-y|^{2N}}\Big)^{\frac{1}{p}}\leq \|f\|_{\dot{W}^{\frac{N}{p},p}(\mathbb{R}^N)}$$

and

$$\Big(\int_{\Gamma\times\Gamma}\frac{|f(x)-f(y)|^p dxdy}{|x-gy|^{2N}}\Big)^{\frac{1}{p}}\leq\Big(\int_{\Gamma\times\Gamma}\frac{|f(x)-f(y)|^p dxdy}{|x-y|^{2N}}\Big)^{\frac{1}{p}}\leq \|f\|_{\dot{W}^{\frac{N}{p},p}(\mathbb{R}^N)}.$$

Thus,

$$\Big(\int_{\Gamma\times g\Gamma}\Big(\frac{d(x,y)}{|x-y|}\Big)^p|f(x)-f(y)|^p\frac{dxdy}{d(x,y)^{2N}}\Big)^{\frac{1}{p}}\leq (1+2c^{(1)}_{p,R})\|f\|_{\dot{W}^{\frac{N}{p},p}(\mathbb{R}^N)}.$$

Summing over Γ and $1\neq g\in G$, we obtain

$$\Big(\int_{\mathbb{R}^N\times\mathbb{R}^N}\Big(\frac{d(x,y)}{|x-y|}\Big)^p|f(x)-f(y)|^p\frac{dxdy}{d(x,y)^{2N}}\Big)^{\frac{1}{p}}\leq |G|^2(1+2c^{(1)}_{p,R})\|f\|_{\dot{W}^{\frac{N}{p},p}(\mathbb{R}^N)}.$$

Case 2: Let $p\geq 2N$. We have

$$\Big(\int_{\Gamma\times g\Gamma}\Big(\frac{d(x,y)}{|x-y|}\Big)^p|f(x)-f(y)|^p\frac{dxdy}{d(x,y)^{2N}}\Big)^{\frac{1}{p}}\leq$$

$$\leq\Big(\int_{\Gamma\times g\Gamma}\Big(\frac{d(x,y)}{|x-y|}\Big)^{2N}|f(x)-f(y)|^p\frac{dxdy}{d(x,y)^{2N}}\Big)^{\frac{1}{p}}=\|f\|_{\dot{W}^{\frac{N}{p},p}(\mathbb{R}^N)}.$$

□

4. Proof of Theorem 1.1 (i)

Let (X,μ) be a measure space and let $L_p(L_{q,\infty})^{\rm symm}(X\times X,\mu\times\mu)$ denote the mixed-norm space consisting of all $\mu\times\mu$-measurable functions $f:X\times X\to\mathbb{C}$ such that

$$\int_X \|f(x,\cdot)\|^p_{L_{q,\infty}(X,\mu)}d\mu(x),\ \int_X \|f(\cdot,y)\|^p_{L_{q,\infty}(X,\mu)}d\mu(y)<\infty.$$

We equip $L_p(L_{q,\infty})^{\rm symm}(X\times X,\mu\times\mu)$ with its natural norm.

Lemma 4.1. *Let (X,μ) be a measure space and let $2<p<\infty$. Suppose that $K\in L_p(L_{p',\infty})^{\rm symm}(X\times X,\mu\times\mu)$. The integral operator*

$$T_Kf(x):=\int_X K(x,y)f(y)d\mu(y)$$

belongs to $S_{p,\infty}(L_2(X,d\mu))$. Moreover, we have

$$\|T_K\|_{S_{p,\infty}(L_2(X,d\mu))}\le c_p\|K\|_{L_p(L_{p',\infty})^{\rm symm}(X\times X,\mu\times\mu)}.$$

Proof. The proof is given in [8, Lemma 2] (see also [7, Corollary 11.2]). □

Lemma 4.2. *Let $p>2$ and let $\frac{1}{q}=1-\frac{2}{p}$. We have*

$$\|F_1F_2\|_{L_p(L_{p',\infty})(X\times X,\mu\times\mu)}\le 2^{\frac{1}{pp'}}\|F_1\|_{L_p(X\times X,\mu\times\mu)}\|F_2\|_{L_\infty(L_{q,\infty})(X\times X,\mu\times\mu)}.$$

Proof. We clearly have

$$LHS=\Big(\int_X \|F_1(x,\cdot)F_2(x,\cdot)\|^p_{L_{p',\infty}(X,\mu)}d\mu(x)\Big)^{\frac{1}{p}}.$$

Clearly, $\frac{1}{p'}=\frac{1}{p}+\frac{1}{q}$. Thus,

$$\|F_1(x,\cdot)F_2(x,\cdot)\|_{L_{p',\infty}(X,\mu)}\le 2^{\frac{1}{p'}}\|F_1(x,\cdot)\|_{L_{p,\infty}(X,\mu)}\|F_2(x,\cdot)\|_{L_{q,\infty}(X,\mu)}\le$$
$$\le 2^{\frac{1}{p'}}\|F_1(x,\cdot)\|_{L_p(X,\mu)}\|F_2(x,\cdot)\|_{L_{q,\infty}(X,\mu)}.$$

Hence,

$$LHS\le 2^{\frac{1}{pp'}}\Big(\int_X \|F_1(x,\cdot)\|^p_{L_p(X,\mu)}\|F_2(x,\cdot)\|^p_{L_{q,\infty}(X,\mu)}d\mu(x)\Big)^{\frac{1}{p}}\le$$
$$\le 2^{\frac{1}{pp'}}\Big(\int_X \|F_1(x,\cdot)\|^p_{L_p(X,\mu)}d\mu(x)\Big)^{\frac{1}{p}}\times\sup_{x\in X}\|F_2(x,\cdot)\|_{L_{q,\infty}(X,\mu)}=$$
$$=2^{\frac{1}{pp'}}\|F_1\|_{L_p(X\times X,\mu\times\mu)}\|F_2\|_{L_\infty(L_{q,\infty})(X\times X,\mu\times\mu)}.$$

□

Lemma 4.3. *Let (X,μ) be a measure space and let $2<p<\infty$. Suppose that $F\in L_p(X\times X,H^2(\mu\times\mu))$ and $H\in L_\infty(L_{1,\infty})^{\rm symm}(X\times X,\mu\times\mu)$. We have*

$$\|T_{FH}\|_{S_{p,\infty}(L_2(X,d\mu))}\le c_p\|F\|_{L_p(X\times X,H^2(\mu\times\mu))}\|H\|^{1-\frac{2}{p}}_{L_\infty(L_{1,\infty})^{\rm symm}(X\times X,\mu\times\mu)}.$$

Proof. By Lemma 4.1,

$$\|T_{FH}\|_{S_{p,\infty}(L_2(X,d\mu))}\le c_p\|FH\|_{L_p(L_{p',\infty})^{\rm symm}(X\times X,\mu\times\mu)}.$$

Let $\frac{1}{q}=1-\frac{2}{p}$. It follows from Lemma 4.2 that

$$\|F_1F_2\|_{L_p(L_{p',\infty})^{\rm symm}(X\times X,\mu\times\mu)}\le$$

$$\leq 2^{\frac{1}{pp'}}\|F_1\|_{L_p(X\times X,\mu\times\mu)}\|F_2\|_{L_\infty(L_{q,\infty})^{\rm symm}(X\times X,\mu\times\mu)}.$$

Setting $F_1 = FH^{\frac{2}{p}}$ and $F_2 = H^{\frac{1}{q}}$, we obtain

$$\|FH\|_{L_p(L_{p',\infty})^{\rm symm}(X\times X,\mu\times\mu)} \leq$$

$$\leq 2^{\frac{1}{pp'}}\|FH^{\frac{2}{p}}\|_{L_p(X\times X,\mu\times\mu)}\|H^{\frac{1}{q}}\|_{L_\infty(L_{q,\infty})^{\rm symm}(X\times X,\mu\times\mu)} =$$

$$= 2^{\frac{1}{pp'}}\|F\|_{L_p(X\times X,H^2(\mu\times\mu))}\|H\|^{\frac{1}{q}}_{L_\infty(L_{1,\infty})^{\rm symm}(X\times X,\mu\times\mu)} =$$

$$= 2^{\frac{1}{pp'}}\|F\|_{L_p(X\times X,H^2(\mu\times\mu))}\|H\|^{1-\frac{2}{p}}_{L_\infty(L_{1,\infty})^{\rm symm}(X\times X,\mu\times\mu)}.$$

□

Lemma 4.4. *Let (X,μ) be a measure space and let $2 < p < \infty$. Suppose that $F \in L_p(X\times X, GH^2(\mu\times\mu))$ and $H \in L_\infty(L_{1,\infty})^{\rm symm}(X\times X,\mu\times\mu)$. We have*

$$\|T_{FH}\|_{S_p(L_2(X,d\mu))} \leq c_{p,H}\|F\|_{L_p(X\times X,H^2(\mu\times\mu))}.$$

Proof. The assertion follows from Lemma 4.3 by real interpolation. □

The following proposition, strictly speaking, is not part of the proof of Theorem 1.1 (i). Instead, it features in the proof of Theorem 1.1 (ii) in the next section. However, in terms of the employed technique, it certainly belongs to this section.

Proposition 4.5. *Let $p > 2$, let Γ be a chamber and let $f \in \dot{W}^{\frac{N}{p},p}(\Gamma)$. Let V be an integral operator whose integral kernel K satisfies the condition*

$$|K(x,y)| \leq \frac{1}{w(B(x,|x-y|))}.$$

For every $1 \leq j \leq N$, we have

$$\|[V,M_f]\|_{\mathcal{L}_p(L_2(\Gamma,w(x)dx))} \leq c_{R,k}\|f\|_{\dot{W}^{\frac{N}{p},p}(\Gamma)}.$$

Proof. In what follows, let $X = \Gamma$ and $d\mu(x) = w(x)dx$. Set

$$F(x,y) = w(B(x,|x-y|))K(x,y)(f(x)-f(y)), \quad x \neq y,$$

$$H(x,y) = \frac{1}{w(B(x,|x-y|))}, \quad x \neq y.$$

By [7, Lemma 5.7], $H \in L_\infty(L_{1,\infty})^{\rm symm}(\Gamma\times\Gamma,\mu\times\mu)$. By assumption on the kernel K,

$$\|F\|_{L_p(\Gamma\times\Gamma,G^2(\mu\times\mu))} \leq \Big(\int_{\Gamma\times\Gamma}|f(x)-f(y)|^p\frac{w(x)w(y)dxdy}{w^2(B(x,|x-y|))}\Big)^{\frac{1}{p}}.$$

By (2.1),

$$w(B(x,|x-y|)) \geq c'_{R,k}|x-y|^N w^{\frac{1}{2}}(x)w^{\frac{1}{2}}(y), \quad x,y \in \mathbb{R}^N.$$

Thus,

$$\|F\|_{L_p(\Gamma\times\Gamma,H^2(\mu\times\mu))} \leq$$

$$\leq (c'_{R,k})^{-\frac{2}{p}}\Big(\int_{\Gamma\times\Gamma}\frac{|f(x)-f(y)|^p}{|x-y|^{2N}}dxdy\Big)^{\frac{1}{p}} = (c'_{R,k})^{-\frac{2}{p}}\|f\|_{\dot{W}^{\frac{N}{p},p}(\Gamma)}.$$

Obviously, the integral kernel of the operator $[V,M_f]$ is FH. The assertion follows now from Lemma 4.4. □

Proof of Theorem 1.1 (i). In what follows, let $X = \mathbb{R}^N$ and $d\mu(x) = w(x)dx$. Set

$$F(x,y) = w(B(x,d(x,y)))K_{\mathrm{Dunkl},j}(x,y)(f(x)-f(y)), \quad x \neq y,$$

$$H(x,y) = \frac{1}{w(B(x,d(x,y)))}, \quad x \neq y.$$

By [7, Lemma 11.7], $H \in L_\infty(L_{1,\infty})^{\mathrm{symm}}(X \times X, \mu \times \mu)$. By Lemma 2.2,

$$\|F\|_{L_p(X\times X, H^2(\mu\times\mu))} \leq$$

$$\leq c^{(1)}_{R,k}\left(\int_{\mathbb{R}^N\times\mathbb{R}^N}\left(\frac{d(x,y)}{|x-y|}\right)^p |f(x)-f(y)|^p \frac{w(x)w(y)dxdy}{w(B(x,d(x,y)))^2}\right)^{\frac{1}{p}}.$$

Using Lemma 2.3, we conclude that

$$\|F\|_{L_p(X\times X, H^2(\mu\times\mu))} \leq c^{(2)}_{R,k}\left(\int_{\mathbb{R}^N\times\mathbb{R}^N}\left(\frac{d(x,y)}{|x-y|}\right)^p |f(x)-f(y)|^p \frac{dxdy}{d(x,y)^{2N}}\right)^{\frac{1}{p}}.$$

Using Theorem 3.1, we conclude that

$$\|F\|_{L_p(X\times X, H^2(\mu\times\mu))} \leq c_{R,k}\|f\|_{\dot{W}^{\frac{N}{p},p}(\mathbb{R}^N)}.$$

Obviously, the integral kernel of the operator $[R_{\mathrm{Dunkl},j}, M_f]$ is FH. The assertion follows now from Lemma 4.4. □

5. Expected dyadic result

I expect the following result to be true.

Theorem 5.1. *If*

$$\left\{\int_Q \frac{|f|(y)dy}{|y_N|^N}\right\}_{\substack{Q\in\mathcal{D}\\ |(c_Q)_N|\geq c\cdot s(Q)}} \in l_p$$

for some $c > 0$, then

$$\int_{\mathbb{R}^N} \frac{|f|^p(y)dy}{|y_N|^N}dx < \infty.$$

It is immediate that

$$\left\{\int_{3Q} \frac{|f|(y)dy}{|y_N|^N}\right\}_{\substack{Q\in\mathcal{D}\\ |(c_Q)_N|\geq (c+1)\cdot s(Q)}} \in l_p.$$

Let $x \in Q$ and let $t \in (\frac{s(Q)}{2}, s(Q))$. We have

$$B(x,t) = x + B(0,t) \subset x + B(0,s(Q)) \subset x + [-s(Q), s(Q)]^d \subset$$

$$\subset Q + [-s(Q), s(Q)]^d = 3Q.$$

Thus,

$$\int_{B(x,t)} \frac{|f|(y)dy}{|y_N|^N} \leq \int_{3Q} \frac{|f|(y)dy}{|y_N|^N}.$$

Set

$$h(x,t) = \int_{B(x,t)} \frac{|f|(y)dy}{|y_N|^N}, \quad x \in [0,1]^N, \quad t \in (0,1).$$

So, we have

$$\sup_{\substack{x\in Q\\ t\in(\frac{s(Q)}{2}, s(Q))}} h(x,t) \leq \int_{3Q} \frac{|f|(y)dy}{|y_N|^N}.$$

In particular,

$$\int_{Q\times(\frac{s(Q)}{2},s(Q))} h^p(x,t)\frac{dxdt}{t^{N+1}} \leq 2^N\Big(\int_{3Q}|f|\Big)^p.$$

Consequently,

$$\sum_{\substack{Q\in\mathcal{D}\\ |(c_Q)_N|\geq(c+1)\cdot s(Q)}} \int_{Q\times(\frac{s(Q)}{2},s(Q))} h^p(x,t)\frac{dxdt}{t^{N+1}} \leq$$

$$\leq 2^N \sum_{\substack{Q\in\mathcal{D}\\ |(c_Q)_N|\geq(c+1)\cdot s(Q)}} \Big(\int_{3Q}\frac{|f|(y)dy}{|y_N|^N}\Big)^p.$$

Suppose $(x,t)\in\mathbb{R}^N\times\mathbb{R}_+$ is such that $|x_N|\geq(2c+4)t$. Find $m\in\mathbb{Z}$ such that $t\in(2^{-m-1},2^{-m}]$. Find $Q\in\mathcal{D}$ such that $s(Q)=2^{-m}$ and such that $x\in Q$. We have

$$|(c_Q)_N|\geq|x_N|-|(x-c_Q)_N|\geq|x_N|-s(Q)\geq|x_N|-2t\geq$$
$$\geq(2c+4)t-2t=(2c+2)t\geq(c+1)s(Q).$$

Hence,

$$\int_{\substack{(x,t)\in\mathbb{R}^N\times\mathbb{R}_+\\ |x_N|\geq(2c+4)t}} h^p(x,t)\frac{dxdt}{t^{N+1}} \leq 2^N \sum_{\substack{Q\in\mathcal{D}\\ Q\subset Q_0\\ |(c_Q)_N|\geq(c+1)\cdot s(Q)}} \Big(\int_{3Q}\frac{|f|(y)dy}{|y_N|^N}\Big)^p.$$

So,

$$\int_{\substack{(x,t)\in\mathbb{R}^N\times\mathbb{R}_+\\ |x_N|\geq(2c+4)t}} h^p(x,t)\frac{dxdt}{t^{N+1}} < \infty.$$

Furthermore, it is easy to see that do we need details here?

$$\int_{\substack{x\in\mathbb{R}^N\\ t\in\mathbb{R}_+\\ |x_N|\geq t}} h^p(x,t)\frac{dxdt}{t^{N+1}} < \infty.$$

from this, we should be able to derive that our weighted L_p-norm is finite.

6. Simplified version of Zhijie's argument

Lemma 6.1. *Let $\mathcal{D}$ be the standard system of dyadic cubes. Let $O_v\in O(N,\mathbb{R})$ be such $O_ve_N=v$ and let $g\in G$. We have*

$$\Big\|\Big\{w(O_vQ)^{-1}\langle T\chi_{gO_vQ},\chi_{O_vQ}\rangle_{L_2(\mathbb{R}^N,w(x)dx)}\Big\}_{Q\in\mathcal{D}}\Big\|_{l_p(\mathcal{D})} \leq c_{g,v}\|T\|_{\mathcal{L}_p(L_2(\mathbb{R}^N,w(x)dx))}.$$

Proof. to be done later. □

It is decidedly inconvenient that integral kernel K_j of the operator $R_{\mathrm{Dunkl},j}$ is not positive.

Lemma 6.2. *Integral kernel of $R_{\mathrm{Dunkl},j}$ has the same sign as $(x-y)_j$.*

Proof. This should follow from Lemma 3.3 in refer to Dziubanski-Hejna paper Ji refers to. □

Consider the operator B_j I do not know whether this operator is bounded whose integral kernel is given by the formula

$$(x,y)\to|K_j|(x,y),\quad x,y\in\mathbb{R}^N.$$

Lemma 6.3. *Suppose* $M_{f\chi_\Gamma} R_{\mathrm{Dunkl},j} M_{\chi_{g\Gamma}} \in \mathcal{L}_p(L_2(\mathbb{R}^N, w(x)dx))$. *It follows that*

$$M_{|f|\chi_\Gamma} B_j M_{\chi_{g\Gamma}} \in \mathcal{L}_p(L_2(\Gamma, w(x)dx))$$

Proof. Multiplying by $\mathrm{sgn}(f)$ on the left, we may assume without loss of generality that $f \geq 0$. By Lemma 6.2, $K_j(x,y)$ has the same sign as $(x-y)_j$. Thus,

$$M_{f\chi_\Gamma} B_j M_{\chi_{g\Gamma}} = \mathfrak{S}_{\mathrm{sgn}((x-y)_j)}(M_{f\chi_\Gamma} R_{\mathrm{Dunkl},j} M_{\chi_{g\Gamma}}).$$

Schur multiplier $\mathfrak{S}_{\mathrm{sgn}((x-y)_j)}$ is easily seen to be bounded on $\mathcal{L}_p(L_2(\Gamma, w))$. This completes the proof. □

Let $g \in \Gamma$ be a reflection with respect to vector $v_0 \in R$.

Lemma 6.4. *Let* $Q_0 \in \mathcal{D}$ *be such that*

(i) $Q_0 \subset O_{v_0}^{-1}\Gamma$ *touches the plane* $\{x_N = 0\}$.
(ii) $O_{v_0} Q_0$ *is separated from any plane* $\{\langle x, v\rangle = 0\}$, $v_0 \neq v \in R$.

If $Q \in \mathcal{D}$ *is such that* $Q \subset Q_0$ *and*[2]

$$|(c_Q)_N| \geq 2N^{\frac{1}{2}} s(Q), \quad s(Q) \leq \mathrm{const}_{Q_0}.$$

(i) *We have*

$$|\langle O_{v_0} c_Q, v\rangle| \geq 2N^{\frac{1}{2}} s(Q), \quad |\langle g O_{v_0} c_Q, v\rangle| \geq 2N^{\frac{1}{2}} s(Q), \quad v \in \mathbb{R}.$$

(ii) *If* $x \in O_{v_0} Q$ *and* $y \in g O_{v_0} Q$, *then*

$$\frac{1}{2}|\langle O_{v_0} c_Q, v\rangle| \leq |\langle x, v\rangle| \leq \frac{3}{2}|\langle O_{v_0} c_Q, v\rangle|,$$
$$\frac{1}{2}|\langle g O_{v_0} c_Q, v\rangle| \leq |\langle y, v\rangle| \leq \frac{3}{2}|\langle g O_{v_0} c_Q, v\rangle|.$$

(iii) *If* $x \in O_{v_0} Q$ *and* $y \in g O_{v_0} Q$, *then*

$$c_k^{-1} w(O_{v_0} c_Q) \leq w(x), w(y) \leq c_k w(O_{v_0} c_Q).$$

(iv) *If* $x \in O_{v_0} Q$ *and* $y \in g O_{v_0} Q$, *then*

$$|(c_Q)_N| \leq |x - y| \leq 3|(c_Q)_N|.$$

(v) *If* $x \in O_{v_0} Q$ *and* $y \in g O_{v_0} Q$, *then*

$$c_k^{-1} |(c_Q)_N|^N \cdot w(O_{v_0} c_Q) \leq w(B(x, |x-y|)) \leq c_k |(c_Q)_N|^N \cdot w(O_{v_0} c_Q).$$

(vi) *We have*

$$c_k^{-1} m(Q) w(O_{v_0} Q) \leq w(O_{v_0} Q) \leq c_k m(Q) w(O_{v_0} Q).$$

Proof. Since $c_Q \in Q_0$, it follows that

$$|\langle O_{v_0} c_Q, v\rangle| \geq \inf_{z \in Q_0} |\langle O_{v_0} z, v| \geq 2N^{\frac{1}{2}} \mathrm{const}_{Q_0} \geq 2N^{\frac{1}{2}} s(Q), \quad v_0 \neq v \in R,$$

$$|\langle g O_{v_0} c_Q, v\rangle| \geq \inf_{z \in Q_0} |\langle g O_{v_0} z, v| \geq 2N^{\frac{1}{2}} \mathrm{const}_{Q_0} \geq 2N^{\frac{1}{2}} s(Q), \quad v_0 \neq v \in R.$$

On the other hand, we have

$$|\langle g O_{v_0} c_Q, v_0\rangle| = |\langle O_{v_0} c_Q, v_0\rangle| = |\langle O_{v_0} c_Q, O_{v_0} e_N\rangle| = |\langle c_Q, e_N\rangle| = |(c_Q)_N|.$$

[2]Here, we use the shorthand

$$\mathrm{const}_{Q_0} = \frac{1}{2N^{\frac{1}{2}}} \min\{ \inf_{\substack{x \in Q_0 \\ v_0 \neq v \in R}} |\langle x, v\rangle|, \inf_{\substack{x \in Q_0 \\ v_0 \neq v \in R}} |\langle gx, v\rangle| \}.$$

This yields (i).

If $x \in O_{v_0}Q$ and $y \in gO_{v_0}Q$, then

$$|\langle x, v\rangle| \geq |\langle O_{v_0}c_Q, v\rangle| - |\langle (x - O_{v_0}c_Q), v\rangle| \geq |\langle O_{v_0}c_Q, v\rangle| - |x - O_{v_0}c_Q| \geq$$

$$\geq |\langle O_{v_0}c_Q, v\rangle| - N^{\frac{1}{2}}s(Q) \geq \frac{1}{2}|\langle O_{v_0}c_Q, v\rangle|, \quad x \in O_vQ.$$

The other inequalities in (ii) are proved similarly.

The assertion (iii) follows immediately from (ii).

If $x \in O_{v_0}Q$ and $y \in gO_{v_0}Q$, then

$$|x - y| \leq |x - O_{v_0}c_Q| + |y - gO_{v_0}c_Q| + |O_{v_0}c_Q - gO_{v_0}c_Q| \leq$$

$$\leq 2N^{\frac{1}{2}}s(Q) + |O_{v_0}c_Q - (O_{v_0}c_Q - 2\langle O_{v_0}c_Q, v_0\rangle v_0)| =$$

$$= 2N^{\frac{1}{2}}s(Q) + 2|\langle O_{v_0}c_Q, v_0\rangle| \leq 3|\langle O_{v_0}c_Q, v_0\rangle| = 3|(c_Q)_N|.$$

The other inequality in (iv) is proved similarly.

Note that

$$w(B(x, |x - y|)) \geq c_k^{-1}|x - y|^N \cdot w(x).$$

If $x \in O_{v_0}Q$ and $y \in gO_{v_0}Q$, then the it follows from (iii) and (iv) that

$$w(B(x, |x - y|)) \geq c_k^{-1}|\langle O_{v_0}c_Q, v_0\rangle|^N \cdot w(O_{v_0}c_Q).$$

Thus, we obtained the left hand side estimate in (v).

Note that

$$w(B(x, |x - y|)) \leq c_k|x - y|^N \cdot \prod_{v \in R}(|x - y| + |\langle x, v\rangle|)^{k_v}.$$

If $x \in O_{v_0}Q$ and $y \in gO_{v_0}Q$, then it follows from (ii) and (iv) that

$$w(B(x, |x - y|)) \leq c_k|\langle O_{v_0}c_Q, v_0\rangle|^N \cdot \prod_{v \in R}(|\langle O_{v_0}c_Q, v_0\rangle| + |\langle O_{v_0}c_Q, v\rangle|)^{k_v}.$$

By assumption on Q_0,

$$|\langle z, v_0\rangle| \leq \mathrm{const}'_{Q_0}|\langle z, v\rangle|, \quad z \in Q_0, \quad v \in R.$$

Since $O_{v_0}c_Q \in Q_0$, it follows that

$$|\langle O_{v_0}c_Q, v_0\rangle| \leq \mathrm{const}'_{Q_0}|\langle O_{v_0}c_Q, v\rangle|, \quad v \in R.$$

Hence,

$$w(B(x, |x - y|)) \leq c_k|\langle O_{v_0}c_Q, v_0\rangle|^N \cdot \prod_{v \in R}((1 + \mathrm{const}'_{Q_0})|\langle O_{v_0}c_Q, v\rangle|)^{k_v}.$$

Thus, we obtained the right hand side estimate in (v).

Finally, the assertion (vi) follows immediately from (iii). □

Lemma 6.5. *Suppose that v_0 is not orthogonal to e_j. Suppose $Q \in \mathcal{D}$ is such that*

$$|(c_Q)_N| \cdot |(v_0)_j| \geq 2N^{\frac{1}{2}}s(Q).$$

We have

$$\int_{gO_{v_0}Q} |(x - y)_j|dy \geq m(Q) \cdot |(c_Q)_N| \cdot |(v_0)_j|.$$

Proof. Clearly,

$$\int_{gO_{v_0}Q} |(x-y)_j|dy \geq \max\Big\{\int_{gO_{v_0}Q}(x-y)_j dy, \int_{gO_{v_0}Q}(y-x)_j dy\Big\}.$$

Next,

$$\int_{gO_{v_0}Q}(x-y)_j dy = (x-gO_{v_0}c_Q)_j \cdot m(Q)$$

and

$$|(x-gO_{v_0}c_Q)_j| \geq |(O_{v_0}c_Q)_j - (gO_{v_0}c_Q)_j| - |(x-O_{v_0c_Q})_j| \geq$$
$$\geq |(O_{v_0}c_Q - gO_{v_0}c_Q)_j| - N^{\frac{1}{2}}s(Q) =$$
$$= 2|\langle O_{v_0}c_Q, v_0\rangle| \cdot |(v_0)_j| - N^{\frac{1}{2}}s(Q) \geq |\langle O_{v_0}c_Q, v_0\rangle| \cdot |(v_0)_j|.$$

Combining these inequalities, we complete the proof. □

Lemma 6.6. *Suppose $M_{f\chi_\Gamma}R_{\mathrm{Dunkl},j}M_{\chi_{g\Gamma}} \in \mathcal{L}_p(L_2(\mathbb{R}^N, w(x)dx))$. Let Q_0 be as in Lemma 6.4. If v_0 is not orthogonal to e_j, then*

$$\Big\{\int_Q \frac{|f|(O_{v_0}^{-1}x)dx}{|x_N|^N}\Big\}_{\substack{Q\in\mathcal{D}\\ Q\subset Q_0\\ |(c_Q)_N|\geq c\cdot s(Q)}} \in l_p, \quad c = \frac{2N^{\frac{1}{2}}}{|(v_0)_j|}.$$

Proof. Combining Lemma 6.1 and Lemma 6.3, we obtain

$$\Big\{w(O_{v_0}Q)^{-1}\langle |f|\cdot B_j(\chi_{gO_{v_0}Q}), \chi_{O_{v_0}Q}\rangle_{L_2(\mathbb{R}^N,w(x)dx)}\Big\}_{\substack{Q\in\mathcal{D}\\ O_{v_0}Q\subset\Gamma}} \in l_p.$$

In particular,

$$\Big\{w(O_{v_0}Q)^{-1}\langle |f|\cdot B_j(\chi_{gO_{v_0}Q}), \chi_{O_{v_0}Q}\rangle_{L_2(\mathbb{R}^N,w(x)dx)}\Big\}_{\substack{Q\in\mathcal{D}\\ Q\subset Q_0\\ s(Q)\leq \mathrm{const}_{Q_0}\\ |(c_Q)_N|\geq c\cdot s(Q)}} \in l_p.$$

Here, we use the shorthands

$$\mathrm{const}_{Q_0} = \frac{1}{2N^{\frac{1}{2}}}\min\{\inf_{\substack{x\in Q_0\\ v_0\neq v\in R}}|\langle x,v\rangle|, \inf_{\substack{x\in Q_0\\ v_0\neq v\in R}}|\langle gx,v\rangle|\}, \quad c = \frac{2N^{\frac{1}{2}}}{|(v_0)_j|}.$$

Using Lemmas 2.4, 6.4 and 6.5, we write

$$w(O_{v_0}Q)^{-1}\langle |f|\cdot B_j(\chi_{gO_{v_0}Q}), \chi_{O_{v_0}Q}\rangle_{L_2(\mathbb{R}^N,w(x)dx)} =$$
$$= w(O_{v_0}Q)^{-1}\int_{O_{v_0}Q\times gO_{v_0}Q}|f|(x)|K_j|(x,y)w(x)w(y)dxdy \geq$$
$$\overset{L.2.4}{\geq} c_k^{(1)}w(O_{v_0}Q)^{-1}\int_{O_{v_0}Q\times gO_{v_0}Q}|f|(x)\frac{|(x-y)_j|}{|x-y|}\cdot\frac{w(x)w(y)}{w(B(x,|x-y|))}dxdy \geq$$
$$\overset{L.6.4(\mathrm{iii})}{\geq} c_k^{(2)}w(O_{v_0}Q)^{-1}\int_{O_{v_0}Q\times gO_{v_0}Q}|f|(x)\frac{|(x-y)_j|}{|x-y|}\cdot\frac{w^2(O_{v_0}c_Q)}{w(B(x,|x-y|))}dxdy \geq$$
$$\overset{L.6.4(\mathrm{iv})}{\geq} c_k^{(3)}w(O_{v_0}Q)^{-1}\int_{O_{v_0}Q\times gO_{v_0}Q}|f|(x)\frac{|(x-y)_j|}{|(c_Q)_N|}\cdot\frac{w^2(O_{v_0}c_Q)}{w(B(x,|x-y|))}dxdy \geq$$
$$\overset{L.6.4(\mathrm{vi})}{\geq} c_k^{(4)}m(Q)^{-1}\int_{O_{v_0}Q\times gO_{v_0}Q}|f|(x)\frac{|(x-y)_j|}{|(c_Q)_N|}\cdot\frac{w(O_{v_0}c_Q)}{w(B(x,|x-y|))}dxdy \geq$$

$$\overset{L.6.4(v)}{\geq} c^{(5)}_{k,Q_0} m(Q)^{-1} \int_{O_{v_0}Q\times gO_{v_0}Q} |f|(x)|(x-y)_j| \cdot \frac{1}{|(c_Q)_N|^{N+1}} dxdy =$$

$$= c^{(5)}_{k,Q_0} \frac{1}{m(Q)\cdot |(c_Q)_N|^{N+1}} \int_{O_{v_0}Q} |f|(x)\Big(\int_{gO_{v_0}Q} |(x-y)_j| dy\Big) dx \geq$$

$$\overset{L.6.5}{\geq} c^{(6)}_{k,Q_0} \frac{1}{|(c_Q)_N|^N} \int_{O_{v_0}Q} |f|(x)dx \geq c^{(7)}_{k,Q_0} \int_Q \frac{|f|(O^{-1}_{v_0}x)dx}{|x_N|^N}.$$

As the condition $s(Q) \leq \mathrm{const}_{Q_0}$ excludes only finitely many terms, we can drop this condition. □

In case v_0 is orthogonal to e_j, we similarly obtain

$$\Big\{\frac{1}{|(c_Q)_N|^{N+1}} \int_{O_{v_0}Q} |f|(x)|(x-O_{v_0}c_Q)_j| dx\Big\}_{Q\in\mathcal{D}_{v_0,Q_0}} \in l_p.$$

Of course, in this case, we should omit $|(v_0)_j|$ from the definition of $\mathcal{D}_{v_0,Q_0}$.

7. Proof of Theorem 1.1 (ii)

Lemma 7.1. *Let Γ be a chamber and let $x_0 \in \Gamma$. We have*

$$\sup_{x\in B(x_0,r)\cap\Gamma} w(x) \leq \frac{c_{k,\Gamma}}{m(B(x_0,r)\cap\Gamma)} \int_{B(x_0,r)\cap\Gamma} w(x)dx.$$

Proof. If $x = x_0 + y$, $y \in B(0,r)$, then

$$|\langle x, v\rangle| \leq |\langle x_0, v\rangle| + r|\langle y, v\rangle| \leq |\langle x_0, v\rangle| + r\sqrt{2}.$$

Thus,

$$w(x) = \prod_{v\in R} (|\langle x_0, v\rangle| + |\langle y, v\rangle|)^{\epsilon\cdot k(v)} \leq \prod_{v\in R} (|\langle x_0, v\rangle| + r\sqrt{2})^{k(v)}.$$

In particular,

$$LHS \leq \prod_{v\in R} (|\langle x_0, v\rangle| + r\sqrt{2})^{k(v)}.$$

By convexity, $x_0 + \Gamma \subset \Gamma$. Hence,

$$x_0 + (B(0,r)\cap\Gamma) \subset B(x_0,r)\cap\Gamma.$$

So,

$$\int_{B(x_0,r)\cap\Gamma} w(x)dx \geq \int_{x_0+(B(0,r)\cap\Gamma)} w(x)dx = \int_{B(0,r)\cap\Gamma} w(x_0+y)dy.$$

If $y \in \Gamma$, then

$$|\langle x_0 + y, v\rangle| = |\langle x_0, v\rangle| + |\langle y, v\rangle|, \quad v \in R.$$

Thus,

$$w(x_0+y) = \prod_{v\in R} (|\langle x_0, v\rangle| + |\langle y, v\rangle|)^{k(v)} \geq$$

$$\geq \max\{\prod_{v\in R} |\langle x_0, v\rangle|^{k(v)}, \prod_{v\in R} |\langle y, v\rangle|^{k(v)}\}.$$

Hence,

$$\int_{B(x_0,r)\cap\Gamma} w(x)dx \geq \int_{B(0,r)\cap\Gamma} w(x_0+y)dy \geq$$

$$\geq \max\Big\{\int_{B(0,r)\cap\Gamma}\prod_{v\in R}|\langle x_0,v\rangle|^{k(v)}dy, \int_{B(0,r)\cap\Gamma}\prod_{v\in R}|\langle y,v\rangle|^{k(v)}dy\Big\}.$$

Obviously,

$$\int_{B(0,r)\cap\Gamma}\prod_{v\in R}|\langle x_0,v\rangle|^{k(v)}dy = r^N\prod_{v\in R}|\langle x_0,v\rangle|^{k(v)}\int_{B(0,1)\cap\Gamma}dy,$$

$$\int_{B(0,r)\cap\Gamma}\prod_{v\in R}|\langle y,v\rangle|^{k(v)}dy = r^{N+\sum_{v\in R}k(v)}\int_{B(0,1)\cap\Gamma}\prod_{v\in R}|\langle y,v\rangle|^{k(v)}dy.$$

Thus,

$$\int_{B(x_0,r)\cap\Gamma}w(x)dx \geq c^{(1)}_{k,\Gamma}\max\{r^N\prod_{v\in R}|\langle x_0,v\rangle|^{k(v)}, r^{N+\sum_{v\in R}k(v)}\}.$$

Combining this with the preceding paragraph, we complete the proof. □

Lemma 7.2. *Let Γ be a chamber and let $x_0\in\Gamma$. Suppose $1<t<1+\frac{1}{|R|\cdot\max_{v\in R}k(v)}$. For every $r>0$, we have*

$$\Big(\frac{\int_{B(x_0,r)\cap\Gamma}w^{1-t}(x)dx}{\int_{B(x_0,r)\cap\Gamma}w(x)dx}\Big)^{\frac{1}{t}} \leq c_{t,k,\Gamma}\frac{m(B(x_0,r)\cap\Gamma)}{\int_{B(x_0,r)\cap\Gamma}w(x)dx}.$$

Proof. If $v\in R$ is such that

$$2\sqrt{2}r\leq|\langle x_0,v\rangle|,$$

then

$$|\langle x,v\rangle|\geq|\langle x_0,v\rangle|-|\langle x-x_0,v\rangle|\geq|\langle x_0,v\rangle|-r\sqrt{2}\geq\frac{1}{2}|\langle x_0,v\rangle|,\quad x\in B(x_0,r).$$

In this case,

$$\int_{B(x_0,r)}|\langle x,v\rangle|^{(1-t)|R|k(v)}dx\leq 2^{(t-1)|R|k(v)}|\langle x_0,v\rangle|^{(1-t)|R|k(v)}\cdot\int_{B(x_0,r)}dx=$$

$$=2^{(t-1)|R|k(v)}|\langle x_0,v\rangle|^{(1-t)|R|k(v)}\cdot\mathrm{Vol}(\mathbb{B}^N)r^N.$$

If $v\in R$ is such that

$$|\langle x_0,v\rangle|\leq 2\sqrt{2}r,$$

then let $y_0=\langle x_0,v\rangle$ and $z_0=x_0-\frac{1}{2}y_0v_0$. Note that $|y_0|\leq 2\sqrt{2}r$. Using new variables $y=\langle x,v\rangle$ and $z=x-\frac{1}{2}yv$, we write

$$\int_{B(x_0,r)}|\langle x,v\rangle|^{(1-t)|R|k(v)}dx=\int_{|y-y_0|^2+|z-z_0|^2\leq r^2}|y|^{(1-t)|R|k(v)}dydz\leq$$

$$\leq\int_{|y-y_0|^2,|z-z_0|^2\leq r^2}|y|^{(1-t)|R|k(v)}dydz=$$

$$=r^{N-1}\mathrm{Vol}(\mathbb{B}^{N-1})\cdot\int_{|y-y_0|\leq r}|y|^{(1-t)|R|k(v)}dy\leq$$

$$=r^{N-1}\mathrm{Vol}(\mathbb{B}^{N-1})\cdot\int_{|y|\leq(1+2\sqrt{2})r}|y|^{(1-t)|R|k(v)}dy=$$

$$=r^{N+(1-t)|R|k(v)}\mathrm{Vol}(\mathbb{B}^{N-1})\cdot\int_{|y|\leq 1+2\sqrt{2}}|y|^{(1-t)|R|k(v)}dy.$$

Combining the estimates in the preceding paragraphs, we write

$$\int_{B(x_0,r)} |\langle x,v\rangle|^{(1-t)|R|k(v)}dx \le c_1 R^n \big(\max\{|\langle x_0,v\rangle|,r\}\big)^{(1-t)|R|k(v)}.$$

By Hölder inequality,

$$\int_{B(x_0,r)} w^{1-t}(x)dx \le \prod_{v\in R}\Big(\int_{B(x_0,r)} |\langle x,v\rangle|^{(1-t)|R|k(v)}dx\Big)^{\frac{1}{|R|}} \le$$

$$\le c_2 R^N \prod_{v\in R}\big(\max\{|\langle x_0,v\rangle|,r\}\big)^{(1-t)k(v)}.$$

The assertion follows easily. □

Lemma 7.3. *Let Γ be a chamber equipped with Euclidean distance. Measures dx and $w(x)dx$ and operator $T_{\Gamma,j} = M_{\chi_\Gamma} R_{\mathrm{Dunkl},j} M_{\chi_\Gamma}$ satisfy on Γ the conditions in Theorem 1.4 in [6].*

Proof. **Step 1:** We claim that measures dx and $w(x)dx$ are A_∞-equivalent (in the sense of Proposition 1.2 in [6]). This verifies Assumption (i) in Theorem 1.4 in [6].

Let $x_0 \in \Gamma$. By Lemmas 7.1 and 7.2,

$$\Big(\frac{1}{m(B\cap\Gamma)}\int_{B\cap\Gamma} w^t(x)dx\Big)^{\frac{1}{t}} \le \frac{c_{k,\Gamma}}{m(B\cap\Gamma)}\int_{B\cap\Gamma} w(x)dx, \quad t>0.$$

$$\Big(\frac{\int_{B\cap\Gamma} w^{1-t}(x)dx}{\int_{B\cap\Gamma} w(x)dx}\Big)^{\frac{1}{t}} \le c_G \frac{m(B\cap\Gamma)}{\int_{B\cap\Gamma} w(x)dx}, \quad 1<t<1+\frac{1}{|R|\cdot\max_{v\in R}k(v)}.$$

The assertion follows now from Proposition 3.6 in [6].

Step 2: Assumption (ii) in Theorem 1.4 in [6] is obvious.

Step 3: Integral kernel K_j of the operator $T_{\Gamma,j}$ satisfies the estimates

$$|K_j(x,y)| \le \frac{c_G}{w(B(x,|x-y|)\cap\Gamma)}, \quad x,y\in\Gamma,$$

$$|K_j(x,y)-K_j(x',y)| \le \frac{|x-x'|}{|x-y|}\cdot\frac{c_G}{w(B(x,|x-y|))}, \quad |x-x'|\le\frac{1}{2}|x-y|, \quad x,x',y\in\Gamma,$$

$$|K_j(x,y)-K_j(x,y')| \le \frac{|y-y'|}{|x-y|}\cdot\frac{c_G}{w(B(x,|x-y|))}, \quad |y-y'|\le\frac{1}{2}|x-y|, \quad x,y,y'\in\Gamma.$$

In other words, K_j is a standard kernel for the measure $w(x)dx$.

This follows from Theorem 1.1 in [4].

Step 4: Kernel K_j is non-degenerate in the following sense: for every $x\in\Gamma$ and $r>0$, there exists $y\in\Gamma$ with $|x-y|=r$ and

$$|K_j(x,y)| \ge \frac{c_G}{w(B(x,|x-y|)\cap\Gamma)}.$$

Let

$$c_\Gamma = (\max_{z\in\Gamma,\ |z|=1}|z_j|)^{-1}.$$

Choose $z\in\Gamma$ such that $|z_j| = c_\Gamma^{-1}$ and $|z|=1$. Set $y = x+rz$. Since $x,z\in\Gamma$, it follows that $y\in\Gamma$. Clearly, $|(x-y)_j|\ge rc_\Gamma^{-1}$ and $|x-y|=r$. The claim follows from the proof of the Theorem 1.2 in [4].

Step 4: Lebesgue measure on Γ is, obviously, N-regular and possesses the $(1,1)$ Poincare inequality. □

Lemma 7.4. *If* $[R_{\mathrm{Dunkl},j}, M_f] \in \mathcal{L}_p(L_2(\mathbb{R}^N, w(x)dx))$, *then* $f \in \dot{W}^{\frac{N}{p},p}(\Gamma)$ *for every chamber* Γ. *Furthermore,*

$$\|f\|_{\dot{W}^{\frac{N}{p},p}(\Gamma)} \leq c_{R,k} \|[R_{\mathrm{Dunkl},j}, M_f]\|_{\mathcal{L}_p(L_2(\mathbb{R}^N, w(x)dx))}.$$

Proof. This follows immediately from Lemma 7.3 and Theorem 1.4 in [6]. □

Let $U_g : L_2(\mathbb{R}^N, w(x)dx) \to L_2(\mathbb{R}^N, w(x)dx)$ be given by the formula $U_g\xi = \xi \circ g^{-1}$. Denote for brevity

$$A_{g,\Gamma,j} = M_{\chi_\Gamma} R_{\mathrm{Dunkl},j} M_{\chi_{g\Gamma}} U_g.$$

Lemma 7.5. *If* $[R_{\mathrm{Dunkl},j}, M_f] \in \mathcal{L}_p(L_2(\mathbb{R}^N, w(x)dx))$, *then*[3]

$$M_{(f-f\circ g)\chi_\Gamma} R_{\mathrm{Dunkl},j} M_{\chi_{g\Gamma}} \in \mathcal{L}_p(L_2(\Gamma, w(x)dx))$$

for every $g \in G$ *and for every chamber* Γ. *Furthermore,*

$$\|M_{(f-f\circ g)\chi_\Gamma} R_{\mathrm{Dunkl},j} M_{\chi_{g\Gamma}}\|_{\mathcal{L}_p(L_2(\Gamma, w(x)dx))} \leq c_{R,k} \|[R_{\mathrm{Dunkl},j}, M_f]\|_{\mathcal{L}_p(L_2(\mathbb{R}^N, w(x)dx))}.$$

Proof. By assumption,

$$M_{\chi_\Gamma}[R_{\mathrm{Dunkl},j}, M_f] M_{\chi_{g\Gamma}} U_g \in \mathcal{L}_p(L_2(\mathbb{R}^N, w(x)dx)).$$

We, therefore, have

$$A_{g,\Gamma,j} M_{f\circ g} - M_f A_{g,\Gamma,j} \in \mathcal{L}_p(L_2(\Gamma, w(x)dx)). \tag{7.1}$$

Let us denote by $K_{g,\Gamma,j}$ the integral kernel of the operator $A_{g,\Gamma,j}$. By Lemma 2.2,

$$|K_{g,\Gamma,j}|(x,y) \leq \frac{c_G}{w(B(x,|x-y|))}, \quad x, y \in \Gamma.$$

We know from Lemma 7.4 that $f \in \dot{W}^{\frac{N}{p},p}(\Gamma)$. We now infer from the Proposition 4.5 that

$$A_{g,\Gamma,j} M_{f\circ g} - M_{f\circ g} A_{g,\Gamma,j} \in \mathcal{L}_p(L_2(\Gamma, w(x)dx)). \tag{7.2}$$

Combining (7.1) and (7.2), we write

$$M_{f-f\circ g} A_{g,\Gamma,j} \in \mathcal{L}_p(L_2(\Gamma, w(x)dx)).$$

Multiplying by U_g^{-1} on the right, we complete the proof. □

8. Useful part of Zhijie argument

Lemma 8.1. *Let* $P, Q \subset \mathbb{R}^N$ *be measurable and such that* $P \subset Q$. *There is mean zero*[4] *function* $H_{P,Q}$ *on* Q *with*

$$\|H_{P,Q}\|_\infty = \max\left\{\frac{w(Q\backslash P)^{\frac{1}{2}}}{w(P)^{\frac{1}{2}} w(Q)^{\frac{1}{2}}}, \frac{w(P)^{\frac{1}{2}}}{w(Q)^{\frac{1}{2}} w(Q\backslash P)^{\frac{1}{2}}}\right\}$$

and such that

$$\langle b, H_{P,Q}\rangle_{L^2(\mathbb{R}^N, dw)} = \frac{w(P)^{\frac{1}{2}} w(Q)^{\frac{1}{2}}}{w(Q\backslash P)^{\frac{1}{2}}}(f_P - f_Q)$$

for every $f \in L_1^{\mathrm{loc}}(\mathbb{R}^N)$.

[3] this footnote is temporary, for the authors' use only. the combination of assertions of Lemmas 7.4 and 7.5 yield $[R_{\mathrm{Dunkl},j}, M_f] \in \mathcal{L}_p(L_2(\mathbb{R}^N, w(x)dx))$.

[4] With respect to measure $w(x)dx$.

Proof. Set

$$H_{P,Q} = \frac{w(Q\backslash P)^{\frac{1}{2}}}{w(P)^{\frac{1}{2}}w(Q)^{\frac{1}{2}}}\chi_P - \frac{w(P)^{\frac{1}{2}}}{w(Q)^{\frac{1}{2}}w(Q\backslash P)^{\frac{1}{2}}}\chi_{Q\backslash P}.$$

Clearly, $H_{P,Q}$ satisfies the expected estimate in the uniform norm and is mean zero on Q.

We have

$$\langle f, H_{P,Q}\rangle_{L_2(\mathbb{R}^N,w(x)dx)} =$$

$$= \frac{w(Q\backslash P)^{\frac{1}{2}}}{w(P)^{\frac{1}{2}}w(Q)^{\frac{1}{2}}}\int_P f(x)w(x)dx - \frac{w(P)^{\frac{1}{2}}}{w(Q)^{\frac{1}{2}}w(Q\backslash P)^{\frac{1}{2}}}\int_{Q\backslash P} f(x)w(x)dx =$$

$$= \frac{w(Q\backslash P)^{\frac{1}{2}}}{w(P)^{\frac{1}{2}}w(Q)^{\frac{1}{2}}}\int_P f(x)w(x)dx - \frac{w(P)^{\frac{1}{2}}}{w(Q)^{\frac{1}{2}}w(Q\backslash P)^{\frac{1}{2}}}\int_Q f(x)w(x)dx+$$

$$+\frac{w(P)^{\frac{1}{2}}}{w(Q)^{\frac{1}{2}}w(Q\backslash P)^{\frac{1}{2}}}\int_P f(x)w(x)dx =$$

$$= \frac{w(Q\backslash P)^{\frac{1}{2}}}{w(P)^{\frac{1}{2}}w(Q)^{\frac{1}{2}}}\cdot w(P)f_P - \frac{w(P)^{\frac{1}{2}}}{w(Q)^{\frac{1}{2}}w(Q\backslash P)^{\frac{1}{2}}}\cdot w(Q)f_Q+$$

$$+\frac{w(P)^{\frac{1}{2}}}{w(Q)^{\frac{1}{2}}w(Q\backslash P)^{\frac{1}{2}}}\cdot w(P)f_P =$$

$$= \frac{w(P)^{\frac{1}{2}}w(Q\backslash P)^{\frac{1}{2}}}{w(Q)^{\frac{1}{2}}}f_P - \frac{w(P)^{\frac{1}{2}}w(Q)^{\frac{1}{2}}}{w(Q\backslash P)^{\frac{1}{2}}}f_Q + \frac{w(P)^{\frac{3}{2}}}{w(Q)^{\frac{1}{2}}w(Q\backslash P)^{\frac{1}{2}}}f_P =$$

$$= \frac{w(P)^{\frac{1}{2}}\cdot w(Q\backslash P)}{w(Q)^{\frac{1}{2}}w(Q\backslash P)^{\frac{1}{2}}}f_P - \frac{w(P)^{\frac{1}{2}}w(Q)^{\frac{1}{2}}}{w(Q\backslash P)^{\frac{1}{2}}}f_Q + \frac{w(P)^{\frac{1}{2}}\cdot w(P)}{w(Q)^{\frac{1}{2}}w(Q\backslash P)^{\frac{1}{2}}}f_P =$$

$$= \frac{w(P)^{\frac{1}{2}}\cdot w(Q)}{w(Q)^{\frac{1}{2}}w(Q\backslash P)^{\frac{1}{2}}}f_P - \frac{w(P)^{\frac{1}{2}}w(Q)^{\frac{1}{2}}}{w(Q\backslash P)^{\frac{1}{2}}}f_Q =$$

$$= \frac{w(P)^{\frac{1}{2}}w(Q)^{\frac{1}{2}}}{w(Q\backslash P)^{\frac{1}{2}}}f_P - \frac{w(P)^{\frac{1}{2}}w(Q)^{\frac{1}{2}}}{w(Q\backslash P)^{\frac{1}{2}}}f_Q.$$

This completes the proof. □

Lemma 8.2. *Let $Q \in \mathcal{D}$. We have*

$$K_j(x,y) \geq \frac{c_k}{w(Q)}, \quad x \in 3Q, \quad y \in Q + 3s(Q)e_j.$$

Proof. TO BE DONE. Follows from Lemma 2.4.

□

Lemma 8.3. *Let real-valued $f \in L_1^{\mathrm{loc}}(\mathbb{R}^N)$ be such that the commutator $[R_{\mathrm{Dunkl},j}, M_f]$ is bounded on $L_2(\mathbb{R}^N, w(x)dx)$. For every $Q \in \mathcal{D}$, there exist e_Q and f_Q supported in $7Q$ such that $\|e_Q\|_\infty, \|f_Q\|_\infty \leq m(Q)^{-\frac{1}{2}}$ and such that*

$$\frac{1}{w(Q)}\int_{3Q}|f(y) - f_{3Q}|w(y)dy \leq c_k|\langle [R_{\mathrm{Dunkl},j}, M_f]e_Q, f_Q\rangle_{L_2(\mathbb{R}^N,w(x)dx)}|.$$

Proof. Let α be the median of f on $3Q$ and β be the median of f on $Q+3s(Q)e_j$. Assume for definiteness $\alpha \geq \beta$. Set

$$A_1 = \{x \in 3Q : \ f(x) \geq \alpha\}, \quad B_1 = \{x \in Q+3s(Q)e_j : \ f(x) \leq \beta\},$$
$$A_2 = \{x \in 3Q : \ f(x) \leq \beta\}, \quad B_2 = \{x \in Q+3s(Q)e_j : \ f(x) \geq \beta\},$$
$$e_{Q,s} = w(Q)^{-\frac{1}{2}}\chi_{A_s}, \quad f_{Q,s} = w(Q)^{-\frac{1}{2}}\chi_{B_s}, \quad s=1,2.$$

We have

$$\langle [R_{\mathrm{Dunkl},j}, M_f]e_{Q,s}, f_{Q,s}\rangle_{L_2(\mathbb{R}^N,w(x)dx)} =$$
$$= \frac{1}{w(Q)}\int_{B_s\times A_s} K_j(x,y)(f(y)-f(x))w(x)w(y)dxdy.$$

If $x \in B_1$ and $y \in A_1$, then

$$f(y)-f(x) = (f(y)-\alpha)+(\alpha-\beta)+(\beta-f(x)) =$$
$$= |f(y)-\alpha|+|\alpha-\beta|+|f(x)-\beta| \geq |f(y)-\alpha|+|\alpha-\beta| \geq 0.$$

If $x \in B_2$ and $y \in A_2$, then

$$f(y)-f(x) = (f(y)-\beta)+(\beta-f(x)) =$$
$$= -|f(y)-\beta|-|f(x)-\beta| \leq -|f(y)-\beta| \leq 0.$$

Thus,

$$\langle [R_{\mathrm{Dunkl},j}, M_f]e_{Q,1}, f_{Q,1}\rangle_{L_2(\mathbb{R}^N,w(x)dx)} \geq$$
$$\geq \frac{1}{w(Q)}\int_{B_1\times A_1} K_j(x,y)\big(|f(y)-\alpha|+|\alpha-\beta|\big)w(x)w(y)dxdy \geq$$
$$\geq \frac{c_k}{w^2(Q)}\int_{B_1\times A_1} \big(|f(y)-\alpha|+|\alpha-\beta|\big)w(x)w(y)dxdy =$$
$$= \frac{c_k}{w^2(Q)}\cdot\Big(w(B_1)\cdot\int_{A_1}|f(y)-\alpha|w(y)dy+|\alpha-\beta|w(B_1)w(A_1)\Big) \geq$$
$$= \frac{c_k}{w^2(Q)}\cdot\Big(\frac{1}{2}w(Q)\cdot\int_{A_1}|f(y)-\alpha|w(y)dy+|\alpha-\beta|\cdot\frac{1}{4}w^2(Q)\Big) \geq$$
$$\geq \frac{c_k}{4w(Q)}\cdot\Big(\int_{A_1}|f(y)-\alpha|w(y)dy+|\alpha-\beta|w(Q)\Big)$$

and

$$-\langle [R_{\mathrm{Dunkl},j}, M_f]e_{Q,2}, f_{Q,2}\rangle_{L_2(\mathbb{R}^N,w(x)dx)} \geq$$
$$\geq \frac{1}{w(Q)}\int_{B_2\times A_2} K_j(x,y)|f(y)-\beta|w(x)w(y)dxdy \geq$$
$$\geq \frac{c_k}{w^2(Q)}\int_{B_2\times A_2}|f(y)-\beta|w(x)w(y)dxdy =$$
$$= \frac{c_k}{w^2(Q)}\cdot w(B_2)\cdot\int_{A_2}|f(y)-\beta|w(y)dy \geq$$
$$= \frac{c_k}{2w(Q)}\int_{A_2}|f(y)-\beta|w(y)dy.$$

It remains to note that

$$\int_{3Q}|f(y)-\alpha|w(y)dy = \int_{A_1}|f(y)-\alpha|w(y)dy+$$
$$+\int_{A_2}|f(y)-\alpha|w(y)dy+\int_{3Q\setminus(A_1\cup A_2)}|f(y)-\alpha|w(y)dy \leq$$

$$\leq \int_{A_1} |f(y)-\alpha| w(y)dy + \int_{A_2} |f(y)-\beta| w(y)dy$$
$$+ \int_{A_2} |\alpha-\beta| w(y)dy + \int_{3Q\setminus(A_1\cup A_2)} |\alpha-\beta| w(y)dy \leq$$
$$\leq \int_{A_1} |f(y)-\alpha| w(y)dy + \int_{A_2} |f(y)-\beta| w(y)dy + 2|\alpha-\beta| w(3Q).$$

Combining the last 3 equations, we conclude

$$\frac{c_k'}{w(Q)} \int_{3Q} |f(y)-\alpha| w(y)dy \leq \max_{s=1,2} |\langle [R_{\mathrm{Dunkl},j}, M_f] e_{Q,s}, f_{Q,s}\rangle_{L_2(\mathbb{R}^N, w(x)dx)}|.$$

If

$$|\langle [R_{\mathrm{Dunkl},j}, M_f] e_{Q,1}, f_{Q,1}\rangle_{L_2(\mathbb{R}^N, w(x)dx)}| \geq |\langle [R_{\mathrm{Dunkl},j}, M_f] e_{Q,2}, f_{Q,2}\rangle_{L_2(\mathbb{R}^N, w(x)dx)}|,$$

then we set $e_Q = e_{Q,1}$ and $f = f_{Q,1}$. Otherwise, set $e_Q = e_{Q,2}$ and $f_Q = f_{Q,2}$. Thus,

$$\frac{c_k'}{w(Q)} \int_{3Q} |f(y)-\alpha| w(y)dy \leq |\langle [R_{\mathrm{Dunkl},j}, M_f] e_Q, f_Q\rangle_{L_2(\mathbb{R}^N, w(x)dx)}|.$$

Hence,

$$c_k' |f_{3Q}-\alpha| \leq |\langle [R_{\mathrm{Dunkl},j}, M_f] e_Q, f_Q\rangle_{L_2(\mathbb{R}^N, w(x)dx)}|$$

and, finally,

$$\frac{c_k'}{w(Q)} \int_{3Q} |f(y)-f_{3Q}| w(y)dy \leq 2|\langle [R_{\mathrm{Dunkl},j}, M_f] e_Q, f_Q\rangle_{L_2(\mathbb{R}^N, w(x)dx)}|.$$

□

Lemma 8.4. *Suppose that $1 < p < \infty$ and $f \in L_1^{\mathrm{loc}}(\mathbb{R}^N)$. If $[R_{\mathrm{Dunkl},j}, M_f] \in \mathcal{L}_p(L_2(\mathbb{R}^N, w(x)dx))$ for some $j \in \{1, 2, \cdots, N\}$, then*

$$\Big\{ \frac{1}{w(Q)} \int_{3Q} |f(x)-f_{3Q}| w(x)dx \Big\}_{Q\in\mathcal{D}} \in l_p.$$

Proof. Let $\{e_Q\}_{Q\in\mathcal{D}}$ and $\{f_Q\}_{Q\in\mathcal{D}}$ be functions constructed in Lemma 8.3. By some version of Rochberg-Semmes theorem PROOF OR REFERENCE IS NEEDED,

$$\Big\{ \langle [R_{\mathrm{Dunkl},j}, M_f] e_Q, f_Q\rangle_{L_2(\mathbb{R}^N, w(x)dx)} \Big\}_{Q\in\mathcal{D}} \in l_p.$$

The assertion follows now from Lemma 8.3. □

In what follows, conditional expectations are with respect to the measure $w(x)dx$.

Lemma 8.5. *Suppose that $1 < p < \infty$ and $f \in L_1^{\mathrm{loc}}(\mathbb{R}^N, w(x)dx)$. We have*

$$\|f\|^p_{B^p(\mathbb{R}^N, w(x)dx)} \leq c_{p,k} \sum_{Q\in\mathcal{D}} \Big(\frac{1}{w(3Q)} \int_{3Q} |f(x)-f_{3Q}| w(x)dx \Big)^p.$$

Proof. We first decompose $\mathbb{R}^N \times \mathbb{R}^N$ into dyadic shells as follows.

$$\int_{\mathbb{R}^N\times\mathbb{R}^N} \frac{|f(x)-f(y)|^p}{w(B(x,|x-y|))^2} w(x)w(y)dxdy \leq$$
$$\leq \sum_{m\in\mathbb{Z}} \int_{\substack{x,y\in\mathbb{R}^N \\ 2^{-m-1}<|x-y|<2^{-m}}} \frac{|f(x)-f(y)|^p}{w(B(x,|x-y|))^2} w(x)w(y)dxdy \leq$$

$$\leq \sum_{m\in\mathbb{Z}} \frac{1}{w(B(x,2^{-m-1}))^2} \int_{\substack{x,y\in\mathbb{R}^N\\ |x-y|<2^{-m}}} |f(x)-f(y)|^p w(x)w(y)dxdy.$$

For $x, y \in \mathbb{R}^N$ with $2^{-m-1} < |x-y| < 2^{-m}$, fix $Q \in \mathcal{D}$ with $x \in Q$ and $s(Q) = 2^{-m}$. We have $|x-y|_\infty < 2^{-m}$ and $|x - c_Q|_\infty < 2^{-m-1}$. Thus, $|y - c_Q|_\infty < \frac{3}{2} \cdot 2^{-m}$. In other words, $y \in 3Q$. Consequently, there exists $Q \in \mathcal{D}$ with $s(Q) = 2^{-m}$, $x, y \in 3Q$. Thus,

$$\sum_{m\in\mathbb{Z}} \frac{1}{w(B(x,2^{-m-1}))^2} \int_{\substack{x,y\in\mathbb{R}^N\\ |x-y|<2^{-m}}} |f(x)-f(y)|^p w(x)w(y)dxdy \leq$$

$$\leq \sum_{m\in\mathbb{Z}} \frac{1}{w(B(x,2^{-m-1}))^2} \sum_{\substack{Q\in\mathcal{D}\\ s(Q)=2^{-m}}} \int_{3Q\times 3Q} |f(x)-f(y)|^p w(x)w(y)dxdy \leq$$

$$\leq 2^p \sum_{m\in\mathbb{Z}} \frac{1}{w(B(x,2^{-m-1}))^2} \sum_{\substack{Q\in\mathcal{D}\\ s(Q)=2^{-m}}} \int_{3Q\times 3Q} |f(x)-f_{3Q}|^p w(x)w(y)dxdy +$$

$$+2^p \sum_{m\in\mathbb{Z}} \frac{1}{w(B(x,2^{-m-1}))^2} \sum_{\substack{Q\in\mathcal{D}\\ s(Q)=2^{-m}}} \int_{3Q\times 3Q} |f(y)-f_{3Q}|^p w(x)w(y)dxdy \leq$$

$$\leq c_{p,k} \sum_{Q\in\mathcal{D}} \frac{1}{w^2(3Q)} \int_{3Q\times 3Q} |f(x)-f_{3Q}|^p w(x)w(y)dxdy =$$

$$= c_{p,k} \sum_{Q\in\mathcal{D}} \frac{1}{w(3Q)} \int_{3Q} |f(x)-f_{3Q}|^p w(x)dx.$$

Next,

$$\int_{3Q} |f(x)-f_{3Q}|^p w(x)dx = \sum_{\epsilon\in\{-1,0,1\}^d} \int_{Q+\epsilon s(Q)} |f(x)-f_{3Q}|^p w(x)dx \leq$$

$$\leq 2^{p-1} \sum_{\epsilon\in\{-1,0,1\}^d} \int_{Q+\epsilon s(Q)} |f(x)-f_{Q+\epsilon s(Q)}|^p w(x)dx +$$

$$+2^{p-1} \sum_{\epsilon\in\{-1,0,1\}^d} \int_{Q+\epsilon s(Q)} |f_{Q+\epsilon s(Q)}-f_{3Q}|^p w(x)dx.$$

Obviously,

$$\int_{Q+\epsilon s(Q)} |f_{Q+\epsilon s(Q)}-f_{3Q}|^p w(x)dx = |f_{Q+\epsilon s(Q)}-f_{3Q}|^p \cdot w(Q+\epsilon s(Q)) \leq$$

$$\leq \Big(\int_{Q+\epsilon s(Q)} |f(x)-f_{3Q}|w(x)dx\Big)^p \cdot w^{1-p}(Q+\epsilon s(Q)) \leq$$

$$\leq \Big(\int_{3Q} |f(x)-f_{3Q}|w(x)dx\Big)^p \cdot w^{1-p}(Q+\epsilon s(Q)) \leq$$

$$\leq c_{p,k}\Big(\int_{3Q} |f(x)-f_{3Q}|w(x)dx\Big)^p \cdot w^{1-p}(3Q).$$

Thus,

$$\int_{\mathbb{R}^N\times\mathbb{R}^N} \frac{|f(x)-f(y)|^p}{w(B(x,|x-y|))^2} w(x)w(y)dxdy \leq$$

$$\leq c_{p,k}\sum_{Q\in\mathcal{D}}\sum_{\epsilon\in\{-1,0,1\}^d}\frac{1}{w(Q+\epsilon s(Q))}\int_{Q+\epsilon s(Q)}|f(x)-f_{Q+\epsilon s(Q)}|^p w(x)dx+$$
$$+c_{p,k}\sum_{Q\in\mathcal{D}}\Big(\frac{1}{w(3Q)}\int_{3Q}|f(x)-f_{3Q}|w(x)dx\Big)^p.$$

In other words,

$$\|f\|^p_{B^p(\mathbb{R}^N,w(x)dx)}\leq c_{p,k}\sum_{Q\in\mathcal{D}}\frac{1}{w(Q)}\int_Q|f(x)-f_Q|^p w(x)dx+$$
$$+c_{p,k}\sum_{Q\in\mathcal{D}}\Big(\frac{1}{w(3Q)}\int_{3Q}|f(x)-f_{3Q}|w(x)dx\Big)^p.$$

Obviously,

$$\sum_{Q\in\mathcal{D}}\frac{1}{w(Q)}\int_Q|f(x)-f_Q|^p w(x)dx=\sum_{m\in\mathbb{Z}}\sum_{\substack{Q\in\mathcal{D}\\ s(Q)=2^{-m}}}\frac{1}{w(Q)}\int_Q|f(x)-f_Q|^p w(x)dx\leq$$
$$\leq c_{p,k}\sum_{m\in\mathbb{Z}}\sum_{\substack{Q\in\mathcal{D}\\ s(Q)=2^{-m}}}\int_Q\frac{|f(x)-f_Q|^p w(x)dx}{w(B(x,2^{-m}))}=$$
$$=c_{p,k}\sum_{m\in\mathbb{Z}}\int_{\mathbb{R}^N}\frac{|f(x)-(E_mf)(x)|^p w(x)dx}{w(B(x,2^{-m}))}.$$

Note that

$$f-E_mf=\sum_{l\geq m}(E_{l+1}f-E_lf).$$

By triangle inequality,

$$\Big(\int_{\mathbb{R}^N}\frac{|f(x)-(E_mf)(x)|^p w(x)dx}{w(B(x,2^{-m}))}\Big)^{\frac{1}{p}}\leq$$
$$\leq\sum_{l\geq m}\Big(\int_{\mathbb{R}^N}\frac{|(E_{l+1}f)(x)-(E_lf)(x)|^p w(x)dx}{w(B(x,2^{-m}))}\Big)^{\frac{1}{p}}.$$

Note that

$$w(B(x,2^{-m}))\geq c_k2^{N(l-m)}w(B(x,2^{-l})).$$

Thus,

$$\Big(\int_{\mathbb{R}^N}\frac{|f(x)-(E_mf)(x)|^p w(x)dx}{w(B(x,2^{-m}))}\Big)^{\frac{1}{p}}\leq$$
$$\leq c_{p,k}\sum_{l\geq m}2^{\frac{N(m-l)}{p}}\Big(\int_{\mathbb{R}^N}\frac{|(E_{l+1}f)(x)-(E_lf)(x)|^p w(x)dx}{w(B(x,2^{-l}))}\Big)^{\frac{1}{p}}$$

and, therefore,

$$\sum_{m\in\mathbb{Z}}\int_{\mathbb{R}^N}\frac{|f(x)-(E_mf)(x)|^p w(x)dx}{w(B(x,2^{-m}))}\leq$$
$$\leq c_{p,k}\sum_{m\in\mathbb{Z}}\Big(\sum_{l\geq m}2^{\frac{N(m-l)}{p}}\Big(\int_{\mathbb{R}^N}\frac{|(E_{l+1}f)(x)-(E_lf)(x)|^p w(x)dx}{w(B(x,2^{-l}))}\Big)^{\frac{1}{p}}\Big)^p.$$

Using the discreet Young's inequality

$$\|a*b\|_p\leq\|a\|_1\|b\|_p$$

with

$$a(l) = 2^{-\frac{Nl}{p}}\chi_{\mathbb{Z}_+}(l), \quad b(l) = \Big(\int_{\mathbb{R}^N} \frac{|(E_{l+1}f)(x) - (E_l f)(x)|^p w(x)dx}{w(B(x,2^{-l}))}\Big)^{\frac{1}{p}},$$

we obtain

$$\sum_{m\in\mathbb{Z}} \Big(\sum_{l\geq m} 2^{\frac{N(m-l)}{p}} \Big(\int_{\mathbb{R}^N} \frac{|(E_{l+1}f)(x) - (E_l f)(x)|^p w(x)dx}{w(B(x,2^{-l}))}\Big)^{\frac{1}{p}}\Big)^p \leq$$

$$\leq c_{N,p} \sum_{l\in\mathbb{Z}} \int_{\mathbb{R}^N} \frac{|(E_{l+1}f)(x) - (E_l f)(x)|^p w(x)dx}{w(B(x,2^{-l}))}.$$

Thus,

$$\sum_{m\in\mathbb{Z}} \int_{\mathbb{R}^N} \frac{|f(x) - (E_m f)(x)|^p w(x)dx}{w(B(x,2^{-m}))} \leq$$

$$\leq c_{p,k} \sum_{l\in\mathbb{Z}} \int_{\mathbb{R}^N} \frac{|(E_{l+1}f)(x) - (E_l f)(x)|^p w(x)dx}{w(B(x,2^{-l}))} \leq$$

$$\leq c_{p,k} \sum_{l\in\mathbb{Z}} \sum_{\substack{Q\in\mathcal{D}\\ s(Q)=2^{-l-1}}} \frac{1}{w(Q)} \int_Q |(E_{l+1}f)(x) - (E_l f)(x)|^p w(x)dx =$$

$$= c_{p,k} \sum_{l\in\mathbb{Z}} \sum_{\substack{Q\in\mathcal{D}\\ s(Q)=2^{-l-1}}} |f_Q - f_{\mathrm{parent}(Q)}|^p.$$

Obviously,

$$|f_Q - f_{\mathrm{parent}(Q)}| \leq \frac{1}{w(Q)} \int_{\mathrm{parent}(Q)} |f(x) - f_Q| w(x)dx \leq$$

$$\leq \frac{1}{w(Q)} \int_{3Q} |f(x) - f_Q| w(x)dx \leq \frac{c_k}{w(3Q)} \int_{3Q} |f(x) - f_Q| w(x)dx \leq$$

$$\leq \frac{3c_k}{w(3Q)} \int_{3Q} |f(x) - f_Q| w(x)dx.$$

Hence,

$$\sum_{m\in\mathbb{Z}} \int_{\mathbb{R}^N} \frac{|f(x) - (E_m f)(x)|^p w(x)dx}{w(B(x,2^{-m}))} \leq$$

$$\leq c_{p,k} \sum_{l\in\mathbb{Z}} \sum_{\substack{Q\in\mathcal{D}\\ s(Q)=2^{-l-1}}} \Big(\frac{1}{w(3Q)} \int_{3Q} |f(x) - f_Q| w(x)dx\Big)^p =$$

$$= c_{p,k} \sum_{Q\in\mathcal{D}} \Big(\frac{1}{w(3Q)} \int_{3Q} |f(x) - f_Q| w(x)dx\Big)^p.$$

This completes the proof. □

9. Constant characterization

Theorem 9.1. *If $f \in L_1^{\mathrm{loc}}(\mathbb{R}^N, w(x)dx)$ is such that*

$$[R_{\mathrm{Dunkl},j}, M_f] \in \mathcal{L}_p(L_2(\mathbb{R}^N, w(x)dx))$$

for some $p \le d$, then f is constant.

Proof. By Lemma 7.3, measures dx and $w(x)dx$ and operator $T_{\Gamma,j} = M_{\chi_\Gamma} R_{\mathrm{Dunkl},j} M_{\chi_\Gamma}$ satisfy on Γ the assumptions in Theorem 1.4 in [6]. By assumption, $[T_{\Gamma,j}, M_f] \in \mathcal{L}_p(L_2(\Gamma, w(x)dx))$. By Theorem 1.4 in [6], f is constant on Γ.

Hence, f is constant on every chamber. Let $f = c_1$ on Γ_1 and $f = c_2$ on Γ_2. On the other hand, $[R_{\mathrm{Dunkl},j}, M_f] \in \mathcal{L}_{2N}(L_2(\mathbb{R}^N, w(x)dx))$. By Theorem 1.1 (ii), $f \in \dot{W}^{\frac{1}{2},2N}(\mathbb{R}^N)$. We have

$$\int_{\Gamma_1\times\Gamma_2} \frac{|c_1 - c_2|^{2N} dxdy}{|x-y|^{2N}} = \int_{\Gamma_1\times\Gamma_2} \frac{|f(x) - f(y)|^{2N} dxdy}{|x-y|^{2N}} \le$$

$$\le \int_{\mathbb{R}^N\times\mathbb{R}^N} \frac{|f(x) - f(y)|^{2N} dxdy}{|x-y|^{2N}} = \|f\|_{\dot{W}^{\frac{1}{2},2N}(\mathbb{R}^N)} < \infty.$$

It follows that $c_1 = c_2$. Since chambers Γ_1 and Γ_2 are arbitrary, the assertion follows. □

Acknowledgements:

Z. Fan is supported by the Guangdong Basic and Applied Basic Research Foundation (No. 2023A1515110879). J. Li is supported by the Australian Research Council (ARC) through the research grant DP220100285.

References

[1] Z. Fan, M. Lacey, and J. Li. Schatten classes and commutators of Riesz transform on Heisenberg group and applications. *J. Fourier Anal. Appl.*, 29(2):Paper No. 17, 28, 2023. 5

[2] Z. Fan, M. Lacey, J. Li, M.N. Vempati, and B.D. Wick. Besov space, Schatten classes, and commutators of Riesz transforms associated with the Neumann Laplacian. *Michigan Math. J.*, 75(1):199–224, 2025. 5

[3] Z. Fan, M.T. Lacey, J. Li, and X. Xiong. Schatten–Lorentz characterization of Riesz transform commutator associated with Bessel operators. Preprint available at https://arxiv.org/abs/2403.08249, 2024. 5

[4] Y. Han, M. Lee, J. Li, and B.D. Wick. Riesz transforms and commutators in the Dunkl setting. *Anal. Math. Phys.*, 14(3):Paper No. 46, 32, 2024. 3, 4, 19

[5] S. Helgason. *Differential geometry, Lie groups, and symmetric spaces*, volume 80 of *Pure and Applied Mathematics*. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York-London, 1978. 3

[6] T. Hytonen. A_∞-invariance of oscillatory norms and schatten characterisations of commutators. 19, 20, 27

[7] T. Hytonen. Schatten properties of commutators on metric spaces. Preprint available at https://arxiv.org/abs/2411.02613, 2024. 5, 10, 11, 12

[8] S. Janson and T.H. Wolff. Schatten classes and commutators of singular integral operators. *Ark. Mat.*, 20(2):301–310, 1982. 10

[9] M.T. Lacey, J. Li, and B.D. Wick. Schatten classes and commutator in the two weight setting, I. Hilbert transform. *Potential Anal.*, 60(2):875–894, 2024. 5

[10] J. Li, X. Xiong, and F. Yang. Schatten properties of Calderón–Zygmund singular integral commutator on stratified Lie groups. *J. Math. Pures Appl. (9)*, 188:73–113, 2024. 5

[11] R. Rochberg and S. Semmes. Nearly weakly orthonormal sequences, singular value estimates, and Calderon-Zygmund operators. *J. Funct. Anal.*, 86(2):237–306, 1989. 5

[12] Z. Wei and H. Zhang. Complex median method and Schatten class membership of commutators. Preprint available at https://arxiv.org/abs/2411.05810, 2024. 5

[13] Z. Wei and H. Zhang. On the boundedness and Schatten class property of noncommutative martingale paraproducts and operator-valued commutators. Preprint available at https://arxiv.org/abs/2407.14988, 2024. 5

Zhijie Fan, School of Mathematical Sciences, South China Normal University, Guangzhou 510631, China
Email address: fanzhj3@mail2.sysu.edu.cn

Ji Li, Department of Mathematics, Macquarie University,NSW2109, Australia
Email address: ji.li@mq.edu.au

Dmitriy Zanin, University of New South Wales, Kensington, NSW 2052, Australia
Email address: d.zanin@unsw.edu.au