Com S 574 Spring Semester 2021

Machine Learning

Homework 5

Instructor: Forrest Bao Submitted by: Ibne Farabi Shihab

1 no question

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{bmatrix}$$

$$B = \begin{bmatrix} 0.5 & 0.1 & 0.3 \\ -1 & -20 & 1.5 \end{bmatrix}$$

$$A \circ B = \begin{bmatrix} 1 \times 0.5 & 2 \times 0.1 & 3 \times 0.3 \\ 3 \times -1 & 2 \times -20 & 1 \times 1.5 \end{bmatrix}$$

$$A \circ B = \begin{bmatrix} 0.5 & 0.2 & 0.9 \\ -3 & -40 & 1.5 \end{bmatrix}$$

2 no Question

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{bmatrix}$$

$$B^T = \begin{bmatrix} 0.5 & -1 \\ 0.1 & -20 \\ 0.3 & 1.5 \end{bmatrix}$$

$$AB^T = \begin{bmatrix} 1.6 & -36.5 \\ 2 & -41.5 \end{bmatrix}$$

1

$$B = \begin{bmatrix} 0.5 & 0.1 & 0.3 \\ -1 & -20 & 1.5 \end{bmatrix}$$

$$A^T = \begin{bmatrix} 1 & 3 \\ 2 & 2 \\ 3 & 1 \end{bmatrix}$$

$$BA^T = \begin{bmatrix} 1.6 & 2\\ -36.5 & -41.5 \end{bmatrix}$$

3 no Question

Product of A and B is not possible as in matrix multiplication number of rows of 1st matrix must be equal to the number of columns of the second one.

4 no Question

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{bmatrix}$$

$$B^T = \begin{bmatrix} 0.5 & -1\\ 0.1 & -20\\ 0.3 & 1.5 \end{bmatrix}$$

$$AB^T = \begin{bmatrix} 1.6 & -36.5 \\ 2 & -41.5 \end{bmatrix}$$

$$f(x) = x + 1$$

$$f(AB^T) = AB^T + 1$$

$$= \begin{bmatrix} 2.6 & -35.5 \\ 3 & -40.5 \end{bmatrix}$$

no Question

We can get it in the following way:

$$\begin{split} \frac{\partial E}{\partial \tilde{y}} &= 2(\tilde{y} - y) \\ \frac{\partial W^T X}{\partial \tilde{w}_i} &= x \text{ (one sample)} \\ \frac{\partial \tilde{y}}{\partial W^T X} &= 1 \\ \frac{\partial E}{\partial w_i} &= 2(\tilde{y} - y) * x \end{split}$$

no Question 6

Given,
$$x = \begin{bmatrix} x_0, x_1, x_2, x_3 \end{bmatrix}$$

$$= \begin{bmatrix} 1, 0, 1, 0 \end{bmatrix}$$

$$w = \begin{bmatrix} w_0 \\ w_1 \\ w_2 \\ w_3 \end{bmatrix} = \begin{bmatrix} 5 \\ 4 \\ 6 \\ 1 \end{bmatrix}$$

Prediction -

$$\begin{split} \hat{y} &= \phi(W^T X) \\ &= \phi(W_0 X_0 + W_1 X_1 + W_2 X_2 + W_3 X_3)) \\ &= \phi(11) \\ &= 11^2 = 121 \end{split}$$

7 no Question

= 0

$$\begin{split} &\frac{\partial E}{\partial X_1} = \frac{\partial}{\partial X_1} (\hat{y} - y) \\ &= \frac{\partial}{\partial X_1} (W_0 X_0 + W_1 X_1 + W_2 X_2 + W_3 X_3)^2 - \frac{\partial}{\partial X_1} (y) \\ &= 2 \times (W_0 X_0 + W_1 X_1 + W_2 X_2 + W_3 X_3) \times W_1 \\ &= 2 \times (11) \times 4 - 0 \\ &= 88 \end{split}$$

$$&\frac{\partial E}{\partial W_1} = \frac{\partial}{\partial W_1} (\hat{y} - y) \\ &= 2 \times (W_0 X_0 + W_1 X_1 + W_2 X_2 + W_3 X_3) \times X_1 \end{split}$$

8 no Question

$$\frac{\partial E}{\partial X} = \begin{bmatrix} \frac{\partial E}{\partial X_0} \\ \frac{\partial E}{\partial X_1} \\ \frac{\partial E}{\partial X_2} \\ \frac{\partial E}{\partial X_3} \end{bmatrix} = \begin{bmatrix} 22 \times 5 \\ 22 \times 4 \\ 22 \times 6 \\ 22 \times 1 \end{bmatrix} = \begin{bmatrix} 110 \\ 88 \\ 132 \\ 22 \end{bmatrix}$$

$$\frac{\partial E}{\partial W} = \begin{bmatrix} \frac{\partial E}{\partial W_0} \\ \frac{\partial E}{\partial W_1} \\ \frac{\partial E}{\partial W_2} \\ \frac{\partial E}{\partial W_3} \end{bmatrix} = \begin{bmatrix} 22 \times 1 \\ 22 \times 0 \\ 22 \times 1 \\ 22 \times 0 \end{bmatrix} = \begin{bmatrix} 22 \\ 0 \\ 22 \\ 0 \end{bmatrix}$$