


28 sept 202	1	·Hon	nework #	-3			
			& linear.				
1			Arth A		Tro. F.		
	Problem	1			4,7	les .	
				ş.,			
a)	Lim	2- cos Q		Sin20+C	10(20 = 1		
)	Lim Ø→0	02					
			and the same		Salver !	Fair Him	
	lim	sin <sup>2</sup> O		10.2			
-	0->0	8 <sup>2</sup>	Set 31.2	-4 5 2		Table 1	
		And the second			The street		
<b>-</b> 2	Usin a	L'Hopital 1	Rule.				
	J			A Section of Section	- 5 <u>5</u>	WIJ 2	
	lim	2 sin 0 cos 8	ψ <sup>come tie</sup> ,	at hy	The second secon		
	0-0	20		- 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1			
					~		
		7 Sin 20	apply	L'Hopital	Bule a	egain.	
	0-0	20	manuscript of a second electronic of the	A Section 1	377 1. 2.1. X = 26	0.3036-1.5	
				- May	2		
	lim	2(05 20			<i>3</i> %		
<b>2</b> )	0->0	2	5	The second	7 - V 3		
<u></u>	*_14-747	C		-	- 1	The same to the	
	$-\frac{2\cos(2.0)}{2\cos(2.0)} = \frac{2\times 1}{2} = \boxed{1}$						
	-	2	2	2 - 1 ( - 1	1	~	
,D.			1	-		and - w	
3				- 7 No. %			
			040				
		William Francisco					
		d the little like			, 3	-	



Vertical Asyptotes: 322 +-12x +9 denominator =0. so n² -5 = 0  $n^2 = 5$   $n = \pm \sqrt{5}$ Horizontal Asymptotes y=3 Vertical Asymptotes n=+J5, n=J5





Problem 3

a)  $(os(n) = e^{x} + x + z$ 

applying intermediate Value Theorem

 $y = e^{n} + n - (os(n) + 2)$ 

f(n) = ex +2 +05x +2.

f(1) = e + 1 + 3 - cos(1)= [5.18]

 $f(-1) = \frac{1}{2} + (-1) + 2 - (03(2))$ 

= 0.83

Value of f(n) acrosses at n = 1 and n = 1So it is ploven with IVT mat that  $\cos(n) = 2e^{n} + n + 2 \quad \text{is having of least}$ one solution.

$$f(n) = e^{x} + x + 2$$
  
 $f'(n) = xe^{x} + 1 + 0$ 

applying Intermediate Value Incorum

$$f'(-1) = 1/e + 1 = 1.37$$

$$f'(0) = e' + 1 = 2$$

hence me roof lies between n=-1, n=-3