

BÉZOUT INEQUALITY AND APPLICATIONS

FARHAD BABAEE AND TIEN-CUONG DINH

ABSTRACT.

1. INTRODUCTION

Let X be a complex manifold of dimension k . Let Z_1, \dots, Z_n be analytic subsets of X of dimension q_1, \dots, q_n respectively. When $q_1 + \dots + q_n < k$, the intersection $Z_1 \cap \dots \cap Z_n$ is expected to be empty. Otherwise, it is expected to have dimension $q_1 + \dots + q_n - k$ when it is non-empty. More precisely, $Z_1 \cap \dots \cap Z_n$ is an analytic subset of X whose irreducible components has a dimension at least $q_1 + \dots + q_n - k$. When this dimension is strictly larger than $q_1 + \dots + q_n - k$ we refer to the "dimension excess" phenomenon. We will define to each component Y of $Z_1 \cap \dots \cap Z_n$ a multiplicity m_Y which is a positive integer.

Here is our main result where the degree $\|Y\|$ of an analytic subset of pure dimension of X is defined by

$$\|Y\| := \int_Y \omega^{\dim Y}.$$

Theorem 1.1 (Bézout inequality). *Let X be a compact Kähler manifold of dimension k and ω be a Kähler form on X . Let Z_1, \dots, Z_n be analytic subsets of X of dimension q_1, \dots, q_n respectively. Let q be an integer. Denote by Y_1, \dots, Y_l the irreducible components of dimension q of $Z_1 \cap \dots \cap Z_n$ and m_1, \dots, m_l their multiplicities. Then there is a constant $A > 0$ depending only on (X, ω) such that*

$$\sum_{i=1}^l m_i \|Y_i\| \leq A \|Z_1\| \dots \|Z_n\|.$$

This is an application of the density theory for currents.

Applying this result to the intersection of $\Gamma_n \cap \pi_1^{-1}(Z)$ and Δ where Γ_n is the graph of f^n and Δ is the diagonal of $X \times X$.

Corollary 1.2. *Let X be a compact Kähler manifold of dimension k and ω be a Kähler form on X . Let f be a dominant meromorphic correspondence from X to X . Let P_n be the set of isolated periodic points of period n of f where we count points with multiplicities. Then there is a constant A depending only on (X, ω) such that if Z is any analytic subset of pure dimension q of X we have*

$$\#(P_n \cap Z) \leq A \|Z\| \max_{0 \leq p \leq q} d_{n,p}, \quad \text{where} \quad d_{n,p} := \int_X (f^n)^*(\omega^p) \wedge \omega^{k-p}.$$

Recall that the limit

$$d_p := \lim_{n \rightarrow \infty} (d_{n,p})^{1/n}$$

exists and is called the dynamical degree of order p of f . It follows that the RHS of the inequality in the corollary can be bounded by a constant times $(\max d_p + \epsilon)^n$ for each $\epsilon > 0$.

The important fact in the corollary is that the estimate is uniform on Z . The case of endomorphism of \mathbb{P}^k without multiplicities was obtained in [DY25] and has been used to get the exponential rate of equidistribution for periodic points. In this case, we have $d_{n,p} = d^{pn}$, where d is the algebraic degree of the map.

2010 *Mathematics Subject Classification.* XXXXX (primary), XXXXX (secondary).

Corollary 1.3. *Growth of the indeterminacy sets of f^n ? Other applications ? How to get a good estimate for periodic points of a Henon map on a curve ?*

REFERENCES

[DY25] Tien-Cuong Dinh and Jit Wu Yap. *Unpublished note*, 2025.

SCHOOL OF MATHEMATICS, UNIVERSITY OF BRISTOL

E-mail address: `farhad.babae@bristol.ac.uk`

NATIONAL UNIVERSITY OF SINGAPORE, LOWER KENT RIDGE ROAD 10, SINGAPORE 119076, SINGAPORE

E-mail address: `matdtc@nus.edu.sg`