

AVERAGING ANALYTIC VARITIES

FARHAD BABAEE AND TIEN CUONG DINH

ABSTRACT.

CONTENTS

1. Introduction	1
2. Questions	1
References	1

1. INTRODUCTION

2. QUESTIONS

Theorem 2.1. Let $V \subseteq T_N \subseteq X_\Sigma$, be an algebraic subvariety compatible with the toric variety X_Σ . Define

$$\overline{V}_\mu := \int_{x \in (S^1)_N} [\overline{V}] d\mu(x),$$

where μ is a Haar measure on S^1_N . The current \overline{V}_μ has a continuous superpotential.

Question 2.2.

- (a) I am not sure if you wrote down the proof for codimension 1 case at some point.
- (b) Do we have any equality of type below, maybe under some homothety? It's

$$\overline{V}_\mu \wedge \overline{V}'_\mu \geq \overline{(V \wedge V')}_\mu$$

- (c) A rather unrelated question: What is the speed of convergence for dynamical tropicalisation? How about the slices?

REFERENCES

SCHOOL OF MATHEMATICS, UNIVERSITY OF BRISTOL
Email address: farhad.babae@bristol.ac.uk

DEPARTMENT OF MATHEMATICS, NATIONAL UNIVERSITY OF SINGAPORE
Email address: matdct@nus.edu.sg