

ON THE CONTINUITY OF THE SUPERPOTENTIALS AND SLICES OF TROPICAL CURRENTS

FARHAD BABAEE AND TIEN CUONG DINH

ABSTRACT. We study the question of the continuity of slices of currents and explain how it relates to several seemingly unrelated problems in tropical geometry. On the one hand, through this lens, we show that the continuity of superpotentials constitutes a very general instance of stable intersection theory in tropical geometry. On the other hand, questions concerning tropicalisation with respect to a non-trivial valuation and the commutativity of tropicalisation with intersections offer insights to the problem of the continuity of slices of tropical currents with integration currents. Within this framework, we reformulate and reprove known theorems in tropical geometry and derive new results and techniques in both tropical geometry and the theory of currents.

CONTENTS

1. Introduction	1
2. Tools from Superpotential and Slicing Theory of Currents	4
3. Tropical Varieties, Tori, Tropical Currents	11
4. Tropical Currents	13
5. Continuity of superpotentials	15
6. Slicing tropical currents	23
7. Dynamical tropicalisation and Intersections	28
References	34

1. INTRODUCTION

Let X be a complex manifold of dimension d , and p, q non-negative integers with $d = p + q$. We denote by $\mathcal{C}^q(X) = \mathcal{C}_p(X)$ the cone of positive closed bidegree (q, q) , or bidimension (p, p) -currents on X . We also consider $\mathcal{D}^q(X) = \mathcal{D}_p(X)$, the \mathbb{R} -vector space spanned by $\mathcal{C}^q(X)$. It is well-known that the intersection of two positive closed currents is not always defined. The main initial progress was due to the works of Federer [Fed69] and Bedford and Taylor in [BT82]. Federer defined the generic slicing theory of currents: for a dominant holomorphic map $f : X \rightarrow Y$, and a positive closed current $\mathcal{T} \in \mathcal{C}_p(X)$, or more generally, a *flat current*, a *slice*

$$\mathcal{T} \wedge [f^{-1}(y)]$$

is well-defined for a generic $y \in Y$. A different but approach was introduced by Bedford and Taylor, and extended by Demailly [Dem12], and Fornaess–Sibony [FS95] which

2020 *Mathematics Subject Classification.* 14T10, 37F80, 32U40.

Key words and phrases. Tropical Geometry - Superpotential Theory - Slicing Theory.

assert the following: if $\mathcal{S} = dd^c u$ is a bidegree $(1, 1)$ -current, then

$$\mathcal{S} \wedge \mathcal{T} := dd^c(u\mathcal{T}),$$

can be defined or it is *admissible*, when

- The potential, u , is bounded
- u is unbounded, but its unbounded locus has a small intersection with $\text{supp}(\mathcal{T})$.

For instance, when $\mathcal{S} = dd^c \log |f|$ and \mathcal{T} are two integration currents, such that their supports intersect in the expected dimension, then

$$\mathcal{S} \wedge \mathcal{T} = \sum c_i [C_i],$$

where each C_i is an irreducible component of the intersection, and c_i is the corresponding vanishing number. This intersection coincides with the slicing of integration currents.

Demaily in [Dem92] asked the question of generalising the intersection theory to the case where \mathcal{T} is of a higher bidegree. In several works, the second-named author and Sibony introduced *superpotential theory* and *density* of currents to answer this question. In [DS09], the situation where X is a homogeneous space was completely resolved, and in [DS10] the authors investigated the intersection theory for currents with continuous superpotentials on Kähler manifolds, which generalises the case of bounded potentials in bidegree $(1, 1)$. In this article, we adopt the superpotential theory and the slicing theory. Let us also note that the works of Anderson, Eriksson, Kalm, Wulcan and Yger [AESK⁺21], and [ASKW22] also consider a non-proper intersection theory.

Once the intersections are defined, one can ask the following *continuity problem*:

Problem A. Let \mathcal{T}_k be a sequence of positive closed currents on X converging to \mathcal{T} . Let \mathcal{S} be also a positive closed current on X . Find sufficient conditions such that

$$\lim_{k \rightarrow \infty} (\mathcal{S} \wedge \mathcal{T}_k) = \mathcal{S} \wedge (\lim_{k \rightarrow \infty} \mathcal{T}_k).$$

Roughly speaking, if \mathcal{S} is a current on a compact Kähler manifold with a *continuous superpotential*, then for a current \mathcal{T} , the wedge product

$$\mathcal{S} \wedge \mathcal{T} := \lim_{n \rightarrow \infty} (\mathcal{S} \wedge \mathcal{T}_n),$$

is independent of the choice of smooth approximation $\mathcal{T}_n \rightarrow \mathcal{T}$. Employing results from [DS10], we can partially answer Problem A:

Proposition B. Let X be a compact Kähler manifold, $\mathcal{T}_k \rightarrow \mathcal{T}$ be a convergent sequence in $\mathcal{D}^p(X)$. If a current \mathcal{S} has a continuous superpotential, then

$$\mathcal{S} \wedge \mathcal{T}_n \rightarrow \mathcal{S} \wedge \mathcal{T}.$$

See Proposition 2.6 for a proof. Problem A becomes more difficult when one considers continuity for slices, particularly when the current \mathcal{S} is an integration current. In this article, we examine this problem in the specific case where

$$\lim_{k \rightarrow \infty} \mathcal{T}_k$$

is a *complex tropical current* [Bab14, BH17]. The complex tropical currents are closed currents on complex tori $(\mathbb{C}^*)^d$ or on a toric variety associated to a *tropical cycle*. Recall that a tropical cycle is a weighted polyhedral complex satisfying the *balancing condition*

(see Definition 3.1). For a tropical cycle $\mathcal{C} \subseteq \mathbb{R}^d$, of dimension p , the associated tropical current $\mathcal{T}_{\mathcal{C}} \in \mathcal{D}_p((\mathbb{C}^*)^d)$, is a closed current with support $\text{Log}^{-1}(\mathcal{C})$, where

$$\text{Log} : (\mathbb{C}^*)^d \longrightarrow \mathbb{R}^d, \quad (z_1, \dots, z_d) \longmapsto (-\log |z_1|, \dots, -\log |z_d|).$$

The tropical current $\mathcal{T}_{\mathcal{C}}$ can be naturally presented as a locally fibration of $\text{Log}^{-1}(\mathcal{C})$, and we say \mathcal{C} is compatible with the fan Σ , if the fibres of $\overline{\mathcal{T}}_{\mathcal{C}}$ intersect the toric invariant divisors of the toric variety X_{Σ} transversely. Here $\overline{\mathcal{T}}_{\mathcal{C}}$ denotes the extension by zero of $\mathcal{T}_{\mathcal{C}}$ to the toric variety X_{Σ} .

Theorem C (See Theorem 5.7). Let X_{Σ} be a smooth projective toric variety, and let \mathcal{C} be a tropical cycle compatible with Σ . Then $\overline{\mathcal{T}}_{\mathcal{C}}$ has a continuous super-potential.

The preceding theorem allows us to define the intersection product of a tropical current with any current on a compatible toric variety. We can then restrict this intersection product to the complex torus $T_N \subseteq X_{\Sigma}$ and, using the isomorphism $T_N \simeq (\mathbb{C}^*)^d$, define the intersection product of two tropical currents in $(\mathbb{C}^*)^d$.

On the tropical geometry side, there exists a *stable intersection theory* for tropical cycles. The term “stable” refers precisely to the continuity of the intersection under generic translations of tropical cycles. This property is reminiscent of the *fan displacement rule* in toric geometry (see [FS97]), and more generally, of the *moving lemma* in algebraic geometry. Following [FS97], the concept of stable intersection theory for fans was studied independently in [Kaz03], [RGST05] and [Mik06], and extended in [JY16]. A new but equivalent approach was developed by [AR10]. In the context of supercurrents, Lagerberg [Lag12] derived an intersection theory for hypersurfaces. We also refer the reader to [Sha13, Gub16, Mih21] for related developments. In our setting, given Theorem C, the stable intersection theory is a very special case of Proposition B.

With the stable intersection and natural addition of tropical cycles, we have the \mathbb{Q} -algebra of the tropical cycles.

Theorem D (See Theorem 5.15). The assignment $\mathcal{C} \longmapsto \mathcal{T}_{\mathcal{C}}$ induces a \mathbb{Q} -algebra isomorphism between

- (a) The \mathbb{Q} -algebra of tropical currents on $(\mathbb{C}^*)^d$ with the usual addition of currents and the wedge product.
- (b) The \mathbb{Q} -algebra generated by the tropical cycles in \mathbb{R}^d with the natural addition (Definition 3.5) and stable intersection (Definition 3.3).

Further, considering the map $\Phi_m : (\mathbb{C}^*)^d \longrightarrow (\mathbb{C}^*)^d, z \longmapsto z^m$. We have the isomorphism of the following:

- (a') The \mathbb{Q} -algebra of Φ_m -invariant tropical currents on $(\mathbb{C}^*)^d$ with the usual addition of currents and the wedge product.
- (b') The \mathbb{Q} -algebra generated by the tropical fans in \mathbb{R}^d with the natural addition and stable intersection.

Works of [FS97], [Kaz03] and [JY16] also imply the isomorphisms of \mathbb{Q} -algebra in (a) with the McMullen algebra of dimensional polytopes in \mathbb{Q}^d as well as the Chow group of the compact space obtained as a direct limit of all complete smooth toric varieties; see Theorem 5.15 for further details.

We also address Problem A in a specific case involving the slicing of currents converging to a tropical current:

Theorem E (See Theorem 6.6). Let $\mathcal{C} \subseteq \mathbb{R}^d$ be a tropical variety, and let $B = A_1 \cap \cdots \cap A_k \subseteq \mathbb{R}^d$ be a complete intersection of rational hyperplanes A_1, \dots, A_k . Assume that \mathcal{C} and B intersect transversely. Let Σ be a smooth, projective fan compatible with $\mathcal{C} + B$, and let $(\bar{\mathcal{S}}_n)$ be a sequence of positive closed currents on X_Σ . Suppose that:

- (a) $\bar{\mathcal{S}}_n \longrightarrow \bar{\mathcal{T}}_{\mathcal{C}}$;
- (b) $\text{supp}(\bar{\mathcal{S}}_n) \longrightarrow \text{supp}(\bar{\mathcal{T}}_{\mathcal{C}})$.

Then, the following limit holds in the smooth projective toric variety X_Σ :

$$\lim_{n \rightarrow \infty} \left(\bar{\mathcal{S}}_n \wedge [\bar{T}^B] \right) = \bar{\mathcal{T}}_{\mathcal{C}} \wedge [\bar{T}^B].$$

The proof relies on a theorem of Berteloot and the second-named author [BD20], which states that the limit of slices satisfies a certain continuity property for harmonic functions. This allows us to use Fourier analysis to prove theorems about tropical currents. We will see that this result, combined with *dynamical tropicalisation* [Bab23], implies analogues of the results in:

- [Jon16] and [MS15], concerning the fundamental theorem of tropical geometry and tropicalisation with respect to a non-trivial valuation (see Theorem 7.8);
- [BJS⁺07] and [OP13], concerning the commutativity of intersection and tropicalisation (see Theorem 7.11).

It is notable that the setting of dynamical tropicalisation allows for the tropicalisation of differential forms and currents, enabling the formulation of theorems that would otherwise be impossible in the classical setting of tropical geometry. For instance:

Theorem F (See Theorem 7.13). Let W and V be two smooth algebraic subvarieties of $(\mathbb{C}^*)^d$, and for $0 < \epsilon < 1$, let $U_\epsilon((S^1)^d)$ be an ϵ -neighbourhood of $(S^1)^d$. Then,

$$\frac{1}{m^{2n-(p+q)}} \int_{(t_1, t_2) \in U_\epsilon((S^1)^{2n})} \Phi_m^*([t_1 V] \wedge [t_2 W]) d\nu(t_1) \otimes d\nu(t_2) \longrightarrow \mathcal{T}_{\text{Trop}_0(V)} \wedge \mathcal{T}_{\text{Trop}_0(W)},$$

as $m \rightarrow \infty$. Here, the $d\nu$ is the normalised Lebesgue measures on $U_\epsilon((S^1)^d)$.

Our results highlight the need for a generalisation of our main theorems in the broader context of dynamical systems, and future works.

Acknowledgements. This project has received funding from the National University of Singapore and MOE of Singapore through the grants A-8002488-00-00, as well as the London Mathematical Society's Research in Pairs ref. 42239.

2. TOOLS FROM SUPERPOTENTIAL AND SLICING THEORY OF CURRENTS

Let (X, ω) be a compact Kähler manifold of dimension d . Assume that \mathcal{S} is either a positive or a negative current of bidegree (q, q) on X . The quantity

$$|\langle \mathcal{S}, \omega^{d-q} \rangle|$$

is referred to as the *total mass* of \mathcal{S} . For $0 \leq r \leq d$, we consider the de Rham cohomology groups $H^r(X, \mathbb{C}) = H^r(X, \mathbb{R}) \otimes_{\mathbb{R}} \mathbb{C}$ with coefficients in \mathbb{C} . Recall that Hodge theory

provides the following decomposition of the de Rham cohomology group into Dolbeault cohomology groups:

$$H^r(X, \mathbb{C}) \simeq \bigoplus_{p+q=r} H^{p,q}(X, \mathbb{C}).$$

We denote by $\mathcal{C}^q(X)$ the cone of positive closed bidegree (q, q) -currents or bidimension $(d - q, d - q)$ in X . We denote by $\mathcal{D}^q(X) = \mathcal{D}_{d-q}(X)$ the \mathbb{R} -vector space spanned by $\mathcal{C}^q(X)$, which is the space of closed real currents of bidegree (q, q) . Every current $\mathcal{T} \in \mathcal{D}^q(X)$ has a cohomology class:

$$[\mathcal{T}] \in H^{q,q}(X, \mathbb{R}) = H^{q,q}(X, \mathbb{C}) \cap H^{2q}(X, \mathbb{R}).$$

We define $\mathcal{D}^{q,0}(X) = \mathcal{D}_{d-q}^0(X)$ to be the subspace of $\mathcal{D}^q(X)$, consisting of currents with vanishing cohomology. The $*$ -topology on $\mathcal{D}^q(X)$ is defined by the norm:

$$\|\mathcal{S}\|_* := \min(\|\mathcal{S}^+\| + \|\mathcal{S}^-\|),$$

where the minimum is taken over positive currents \mathcal{S}^+ and \mathcal{S}^- in $\mathcal{C}^q(X)$ that satisfy $\mathcal{S} = \mathcal{S}^+ - \mathcal{S}^-$. We say that \mathcal{S}_n converges to \mathcal{S} in $\mathcal{D}^q(X)$ if \mathcal{S}_n converges weakly to \mathcal{S} and moreover, $\|\mathcal{S}_n\|_*$ is bounded by a constant independent of n .

Let $h := \dim H^{q,q}(X, \mathbb{R})$, and fix a set of smooth forms $\alpha = (\alpha_1, \dots, \alpha_h)$ such that their cohomology classes $\{\alpha\} = (\{\alpha_1\}, \dots, \{\alpha_h\})$ form a basis for $H^{q,q}(X, \mathbb{R})$. By Poincaré duality, there exists a set of smooth forms $\alpha^\vee = (\alpha_1^\vee, \dots, \alpha_h^\vee)$ such that their cohomology classes $\{\alpha^\vee\}$ form the dual basis of $\{\alpha\}$, with respect to the cup-product. By adding $U_{\mathcal{S}}$ to a suitable combination of α_i^\vee , we can assume that $\langle U_{\mathcal{S}}, \alpha_i \rangle = 0$, for all $i = 1, \dots, h$. In this case, we say that $U_{\mathcal{S}}$ is α -normalised.

Definition 2.1. Let $\mathcal{T} \in \mathcal{D}^q(X)$ and \mathcal{S} be a smooth form in $\mathcal{D}^{d-q+1,0}(X)$.

- (i) The α -normalised super-potential $\mathcal{U}_{\mathcal{T}}$ of \mathcal{T} is given by the function

$$\begin{aligned} \mathcal{U}_{\mathcal{T}} : \{\mathcal{S} \in \mathcal{D}^{d-q+1,0}(X) : \text{smooth}\} &\longrightarrow \mathbb{R} \\ \mathcal{S} &\longmapsto \langle \mathcal{T}, U_{\mathcal{S}} \rangle, \end{aligned}$$

where $U_{\mathcal{S}}$ is the α -normalised potential of \mathcal{S} .

- (ii) We say \mathcal{T} has a *continuous super-potential*, if $\mathcal{U}_{\mathcal{T}}$ can be extended to a function on $\mathcal{D}^{d-q+1,0}(X)$ which is continuous with respect to the $*$ -topology.

In general, consider $\mathcal{T} \in \mathcal{D}^q(X)$ and $\mathcal{S} \in \mathcal{D}^r(X)$. Assume that $q + r \leq d$ and \mathcal{T} has a continuous super-potential. Let $\mathcal{U}_{\mathcal{T}}$ be the α -normalised super-potential of \mathcal{T} . Let $\beta \in \text{Span}_{\mathbb{R}}\{\alpha\}$ such that $\{\beta\} = \{\mathcal{T}\}$. For any compactly supported smooth form φ of bidegree $(d - q - r, d - q - r)$, we define

$$(1) \quad \langle \mathcal{T} \wedge \mathcal{S}, \varphi \rangle := \mathcal{U}_{\mathcal{T}}(\mathcal{S} \wedge dd^c \varphi) + \langle \beta \wedge \mathcal{S}, \varphi \rangle.$$

Now assume that if $f : X \longrightarrow Y$, is a biholomorphism between smooth compact Kähler manifolds, then we have

$$f_* \mathcal{U}_{\mathcal{R}_1} = \mathcal{U}_{f_* \mathcal{R}_1}, \quad f^* \mathcal{U}_{\mathcal{R}_2} = \mathcal{U}_{f^* \mathcal{R}_2},$$

for $\mathcal{R}_1 \in \mathcal{D}^q(X)$ and $\mathcal{R}_2 \in \mathcal{D}^q(Y)$.

Definition 2.2. Let (\mathcal{T}_n) be a sequence of currents in $\mathcal{D}^q(X)$ weakly converging to \mathcal{T} . Let $\mathcal{U}_{\mathcal{T}}$ and $\mathcal{U}_{\mathcal{T}_n}$ be their α -normalised super-potentials. If $\mathcal{U}_{\mathcal{T}_n}$ converges to $\mathcal{U}_{\mathcal{T}}$ uniformly on any $*$ -bounded sets of smooth forms in $\mathcal{D}^{d-q+1,0}(X)$, then the convergence is called *SP-uniform*.

It is shown in [DS10, Proposition 3.2.8] that any current with continuous super-potentials can be SP-uniformly approximated by smooth forms. Moreover, currents with continuous super-potentials have other nice properties:

Theorem 2.3 ([DNV18, Theorem 1.1]). Suppose that \mathcal{T} and \mathcal{T}' are two positive currents in $\mathcal{D}_q(X)$, such that $\mathcal{T} \leq \mathcal{T}'$, i.e., $\mathcal{T}' - \mathcal{T}$ is a positive current. Then, if \mathcal{T}' has a continuous super-potential, then so does \mathcal{T} .

Theorem 2.4. If \mathcal{T}_1 and \mathcal{T}_2 are two positive closed currents, and \mathcal{T}_1 has a continuous superpotentials, then $\mathcal{T}_1 \wedge \mathcal{T}_2$ is well-defined. Moreover, if \mathcal{T}_2 has also a continuous superpotential, then

- (a) [DS10, Proposition 3.3.3] $\mathcal{T}_1 \wedge \mathcal{T}_2$ has a continuous superpotential;
- (b) [DS10, Proposition 3.3.3] This wedge product is continuous with respect to the SP-uniform convergence.
- (c) [DS09, Theorem 4.2.4] $\text{supp}(\mathcal{T}_1 \wedge \mathcal{T}_2) \subseteq \text{supp}(\mathcal{T}_1) \cap \text{supp}(\mathcal{T}_2)$.

Theorem 2.5 ([DS10, Proposition 3.3.4]). Assume that $\mathcal{T}_1, \mathcal{T}_2$ and \mathcal{T}_3 are closed positive currents, and \mathcal{T}_1 and \mathcal{T}_2 have continuous superpotentials. Then,

$$\mathcal{T}_1 \wedge \mathcal{T}_2 = \mathcal{T}_2 \wedge \mathcal{T}_1 \quad \text{and} \quad (\mathcal{T}_1 \wedge \mathcal{T}_2) \wedge \mathcal{T}_3 = \mathcal{T}_1 \wedge (\mathcal{T}_2 \wedge \mathcal{T}_3).$$

Proposition 2.6. Let X be a compact Kähler manifold, $S_n \rightarrow S$ be a convergent sequence in $\mathcal{D}^q(X)$. If a current \mathcal{T} has a continuous superpotential, then

$$\mathcal{T} \wedge S_n \rightarrow \mathcal{T} \wedge S.$$

Proof. The main result of [DS04] implies any current $\mathcal{T} \in \mathcal{D}^p(X)$ can be weakly approximated by a difference of smooth closed positive of bidegree (p, p) -forms. The result then follows from the definition of continuity of super-potentials and Equation (1). \square

Lemma 2.7. Let $\mathcal{T}, \mathcal{T}'$ be positive closed currents such that $\mathcal{T}|_{\Omega} = \mathcal{T}'|_{\Omega}$ in an open subset $\Omega \subseteq X$, and both \mathcal{T} and \mathcal{T}' have continuous super-potentials. Then, for any $\mathcal{S} \in \mathcal{D}^r(X)$,

$$(\mathcal{T} \wedge \mathcal{S})|_{\Omega} = (\mathcal{T}' \wedge \mathcal{S})|_{\Omega}.$$

Proof. In [DS10], for any current \mathcal{S} with continuous super-potential, a family $\{\mathcal{T}_{\theta}\}_{\theta \in \mathbb{C}^*}$ is constructed that \mathcal{T}_{θ} converges SP-uniformly to \mathcal{S} as $|\theta| \rightarrow 0$. Let $\epsilon := |\theta| > 0$, be a small positive number, and $V \subseteq U$ be any open set such that V_{ϵ} , the ϵ -neighbourhood of V , is contained entirely in Ω . Therefore, by the hypothesis of the lemma, we can construct families of smooth forms \mathcal{T}_n and \mathcal{T}'_n converging SP-uniformly to \mathcal{T} and \mathcal{T}' respectively. Moreover,

$$\mathcal{T}_n|_{V_{\epsilon}} = \mathcal{T}'_n|_{V_{\epsilon}}.$$

Now, for any $(d - q - r, d - q - r)$ smooth form φ with compact support on Ω , we can cover the support of φ with an open set of the form V_{ϵ} and deduce

$$(\mathcal{T}_n \wedge \mathcal{S}) \wedge \varphi = (\mathcal{T}'_n \wedge \mathcal{S}) \wedge \varphi.$$

This, together with Theorem 2.4(b) implies the assertion. \square

We also have a very useful local version of Theorem 2.3.

Lemma 2.8. If \mathcal{T} is a positive closed current on a compact Kähler manifold X , which is locally bounded by a product of positive closed bidegree $(1, 1)$ -currents of continuous potentials, resp. Hölder continuous potentials, then \mathcal{T} has continuous superpotentials, respectively Hölder continuous potentials in X .

Proof. Fix a point a in X . In an open neighbourhood of a that we identify with the ball $B(0, 2)$ in \mathbb{C}^d , we have

$$\mathcal{T} \leq dd^c u_1 \wedge \cdots \wedge dd^c u_p$$

with u_i continuous or Hölder continuous. Without loss of generality, we can assume that these functions are strictly negative. On $B(0, 1)$, define

$$u'_i := \max(u_i, A \log \|z\|)$$

with A sufficiently large so that $u'_i = u_i$ on $B(0, 1/2)$. Observe that $u'_i = A \log \|z\|$ near $\partial B(0, 1)$. Hence we can extend it to a function which is smooth in a neighbourhood of $X \setminus B(0, 1)$. Thus, this function is quasi-plurisubharmonic. We have

$$\mathcal{T} \leq (B\omega + dd^c u'_1) \wedge \cdots \wedge (B\omega + dd^c u'_p)$$

in a neighbourhood W_a of a if B is large enough.

Since we can cover X using a finite number of open sets W_{a_k} , we can add up all obtained quasi-plurisubharmonic functions together and obtain a quasi-plurisubharmonic function u . It is clear that

$$\mathcal{T} \leq (C\omega + u)^p$$

if C is large enough. The function u is continuous or Hölder continuous, and we deduce by Theorem 2.3. \square

Theorem 2.9. Let $q : \widehat{X} \rightarrow X$, be the blowing up of the compact Kähler manifold X along a submanifold. Assume that $\mathcal{T} \in \mathcal{D}_p(\widehat{X})$ is such that the support of \mathcal{T} does not intersect the exceptional divisors of \widehat{X} . If the current \mathcal{T} has a continuous superpotential then $q_*\mathcal{T}$ has the same property.

Proof. Assume that $\mathcal{S} \in \mathcal{D}^{d-p+1}(X)$. We need to show that the super-potential of $q_*\mathcal{T}$, which is a function defined on smooth forms in $\mathcal{D}^{d-q+1,0}(X)$, can be extended to a continuous function on $\mathcal{D}^{d-q+1,0}(X)$. Let α be a smooth closed (p, p) -form cohomologous to $q_*\mathcal{T}$ and define $\beta := q^*\alpha$. By hypothesis, we have $q^*q_*\mathcal{T} = \mathcal{T}$. It follows that \mathcal{T} is cohomologous to β . Fix a potential U of $q_*\mathcal{T} - \alpha$ which is smooth near the blowup locus. We have $dd^c U = q_*\mathcal{T} - \alpha$ and therefore $dd^c q^*U = \mathcal{T} - \beta$. The smoothness of U near the blowup locus implies that $q_*(q^*U) = U$.

Since α is smooth, it is enough to show that the superpotential \mathcal{U} of $q_*\mathcal{T} - \alpha$ can be extended to a continuous function on $\mathcal{D}^{d-q+1,0}(X)$. Since the last current has a vanishing cohomology class, its superpotential doesn't depend on the normalization.

Claim 1. Let (\mathcal{S}_n) be a bounded sequence of smooth forms in $\mathcal{D}^{d-p+1}(X)$. Then the sequence $\mathcal{U}(\mathcal{S}_n)$ is bounded.

Proof of Claim 1. Since \mathcal{S}_n is smooth, we have

$$\mathcal{U}(\mathcal{S}_n) = \langle U, \mathcal{S}_n \rangle = \langle q^*(U), q^*\mathcal{S}_n \rangle = \widehat{\mathcal{U}}(q^*\mathcal{S}_n),$$

where $\widehat{\mathcal{U}}$ denotes the superpotential of $\mathcal{T} - \beta$. Since the action of q^* on cohomology is bounded and the mass of a positive closed current only depends on its cohomology class, we see that $q^*\mathcal{S}_n$ is bounded in $\mathcal{D}^{d-p+1,0}(\widehat{X})$. Since \mathcal{T} has a continuous superpotential, we deduce that $\widehat{\mathcal{U}}(q^*\mathcal{S}_n)$ is bounded and hence $\mathcal{U}(\mathcal{S}_n)$ is bounded as claimed. \square

Claim 2. Let (\mathcal{S}_n) be a bounded sequence of smooth forms in $\mathcal{D}^{d-p+1}(X)$ converging to 0. Then $\mathcal{U}(\mathcal{S}_n)$ tends to 0.

Proof of Claim 2. By extracting a subsequence, we can assume that $q^*\mathcal{S}_n$ converges to some current \mathcal{R} supported by the exceptional divisor. By hypothesis, we have $q_*(\mathcal{R}) = 0$. Using the computation in Claim 1, and the fact that $\widehat{\mathcal{U}}$ is continuous, we get

$$\lim \mathcal{U}(\mathcal{S}_n) = \widehat{\mathcal{U}}(\mathcal{R}).$$

Now, since q^*U is smooth near the exceptional divisor which supports \mathcal{R} , we deduce that

$$\widehat{\mathcal{U}}(\mathcal{R}) = \langle q^*U, \mathcal{R} \rangle = \langle U, q_*\mathcal{R} \rangle = 0.$$

This proves the claim. \square

To finish the proof, let \mathcal{S} be any current in $\mathcal{D}^{d-p+1,0}(X)$. Choose a bounded sequence \mathcal{R}_n of smooth forms in $\mathcal{D}^{d-p+1,0}(X)$ converging to \mathcal{S} . By Claim 1, extracting a subsequence allows to assume that $\mathcal{U}(\mathcal{R}_n)$ converges to some real number l . Consider now an arbitrary bounded sequence of (\mathcal{S}_n) of smooth forms in $\mathcal{D}^{d-p+1,0}(X)$ converging to \mathcal{S} . To show that \mathcal{U} extends to a continuous function at \mathcal{S} , it is enough to check that $\mathcal{U}(\mathcal{S}_n)$ converges to l . This is a consequence of Claim 2 applied to the sequence $\mathcal{S}_n - \mathcal{R}_n$. \square

Theorem 2.10. For two complex manifolds X and Y , consider two convergent sequences of currents $\mathcal{T}_n \rightarrow \mathcal{T}$ in $\mathcal{D}^q(X)$ and $\mathcal{S}_n \rightarrow \mathcal{S}$ in $\mathcal{D}^r(Y)$. We have that

$$\mathcal{T}_n \otimes \mathcal{S}_n \rightarrow \mathcal{T} \otimes \mathcal{S},$$

weakly in $\mathcal{D}^{q+r}(X \times Y)$.

Sketch of the proof. Let us denote by (x, y) the coordinates on $X \times Y$. Using local coordinates and a partition of unity and Weierstrass theorem we can approximate any smooth forms on $X \times Y$ with forms with polynomial coefficients in (x, y) . The approximation is in C^∞ . As a result, the convergence, we only need test forms with monomial coefficients. Thus, the variables x, y are separated, and the convergence of the tensor products becomes the convergence of each factor. \square

2.1. Semi-continuity of slices. Let $f : X \rightarrow Y$ be a dominant holomorphic map between complex manifolds, not necessarily compact, of dimension d and m respectively. Let \mathcal{T} be a positive closed current on X of bi-dimension (p, p) with $p \geq m$. Then a slice

$$\mathcal{T}_y^f = \langle \mathcal{T} | f | y \rangle$$

obtained by restricting \mathcal{T} to $f^{-1}(y)$ exists for almost every $y \in Y$; see [Dem, Page 171]. This is a positive closed current of bi-dimension $(p - m, p - m)$ on X with the support

contained $f^{-1}(y)$. If Ω is a smooth form of maximal bi-degree on Y and α a smooth $(p-m, p-m)$ -form with compact support in X , then we have

$$\langle \mathcal{T}, \alpha \wedge f^*(\Omega) \rangle = \int_{y \in Y} \langle \mathcal{T}_y^f, \alpha \rangle \Omega(y).$$

In general, for a fixed f , the equality $\langle \mathcal{T}|f|y \rangle = \langle \mathcal{T}'|f|y \rangle$ for almost every y , does not imply that $\mathcal{T} = \mathcal{T}'$. However, the following is true: let f_1, \dots, f_k be dominant holomorphic maps from X to Y_1, \dots, Y_k . Consider the vector space spanned by all the differential forms of type $\alpha \wedge f_i^*(\Omega_i)$ for some α as above and some smooth form Ω_i on Y_i of maximal degree. Assume this space is equal to the space of all (p, p) -forms of compact support in X . Then if $\langle \mathcal{T}|f_i|y_i \rangle = \langle \mathcal{T}'|f_i|y_i \rangle$ for every i and almost every $y \in Y_i$, we have $\mathcal{T} = \mathcal{T}'$. The proof is a consequence of the above discussion. An important special case is the following:

Lemma 2.11. Assume that \mathcal{T} and \mathcal{T}' are two positive closed currents of bidimension (p, p) on \mathbb{C}^d and

$$\langle \mathcal{T}|f|y \rangle = \langle \mathcal{T}'|f|y \rangle,$$

for all linear maps $f : \mathbb{C}^d \longrightarrow \mathbb{C}^p$ and generic $y \in \mathbb{C}^p$. Then $\mathcal{T} = \mathcal{T}'$.

Proof. Let $z \in \text{supp}(\mathcal{T})$, and consider the local coordinates centered at z . By [Dem, Lemma 1.4], the set of strongly positive forms

$$\beta_s = i\beta_{s,1} \wedge \bar{\beta}_{s,1} \wedge \dots \wedge i\beta_{s,p} \wedge \bar{\beta}_{s,p}, \quad 1 \leq s \leq \binom{d}{p}$$

constitute a basis for $\Lambda^{p,p}(\mathbb{C}^d)$, where $\beta_{s,i}$ are of the type $dz_j \pm dz_k$ or $dz_j \pm idz_k$. These forms can be obtained by the pullback of the standard volume forms by the projections $f : \mathbb{C}^d \longrightarrow \mathbb{C}^p$, where $(z_1, \dots, z_d) \longmapsto (\beta_{s,1}, \dots, \beta_{s,p})$, for all different $\beta_{s,i}$. As a result, for considering all such projections f , the set of the form $\alpha f^*(\Omega)$, where α is a smooth test function, f projections, Ω is the volume form on \mathbb{C}^p span all the test forms of bidegree (p, p) in \mathbb{C}^d and we can conclude by the above discussion. \square

Now let $U \subseteq \mathbb{C}^m$ and $V \subseteq \mathbb{C}^d$ be two bounded open sets. Assume that $\pi_1 : U \times V \longrightarrow U$ and $\pi_2 : U \times V \longrightarrow V$ are the canonical projections. Consider two closed positive currents \mathcal{T} and \mathcal{S} on $U \times V$ of bi-dimension (m, m) and (d, d) respectively. We say that \mathcal{T} horizontal-like if $\pi_2(\text{supp}(\mathcal{T}))$ is relatively compact in V . Similarly, if $\pi_1(\text{supp}(\mathcal{S}))$ is relatively compact in U , \mathcal{S} is called vertical-like.

Theorem 2.12. Let X be a smooth projective variety. Assume that V and W are two analytic subsets of X of dimensions p and q , respectively, with $p+q \geq d$, and that they intersect properly. Then:

- (a) The current $[V] \wedge [W]$ is well-defined and

$$[V] \wedge [W] = \sum c_i [C_i],$$

where the C_i are the irreducible components of $V \cap W$, and the c_i are the corresponding multiplicities.

- (b) If, in addition, V and W are two algebraic varieties, such that for generic $t \in T_N \subseteq X$ any component of $tV \cap W$ contains a point $z_t \in (tV_{\text{reg}}) \cap W_{\text{reg}}$, and the

intersection at this point is transversal. Then the intersection multiplicities c_i coincide with the classical multiplicities from algebraic geometry.

Proof. By the partition of unity, we reduce the problem to the case of the subset $N \subset \mathbb{C}^d$. Using Lemma 2.11, we define $[V] \wedge [W]$ as the unique current satisfying

$$([V] \wedge [W]) \wedge [L] := [V] \wedge ([W] \wedge [L]),$$

for generic linear spaces L of dimension $p+q$ containing $\varphi(z)$. Here, the current $[W] \wedge [L]$ can equivalently be defined by the slicing theory or by noting that it is a complete intersection of hypersurfaces.

Now, identifying

$$[V] \wedge ([W] \wedge [L]) = ([V] \otimes ([W] \wedge [L])) \wedge [\Delta],$$

and noting that the diagonal Δ is smooth, we can construct a submersion

$$\xi : N \times N \rightarrow N,$$

whose fibres are translations of Δ , and which is horizontal-like. Now, Part (a) is deduced from the results [BDR24]. To see part (b), suppose V and W are smooth. On one hand, translating V by an element $t \in T_N \subseteq X$, we obtain the subvariety tV , which is linearly equivalent to V . On the other hand, by Kleiman's theorem in T_N , for generic $t \in T_N$, the intersection $(tV) \cap W \cap L$ is a finite set of points contained in T_N .

In the case of transverse intersection, both algebraic geometry and slicing theory assign multiplicity one to each point of intersection. Therefore, the multiplicities from both theories coincide for $V \cap W$, since slicing theory is invariant under generic translations, and algebraic intersection theory is invariant under linear equivalence. \square

Remark 2.13. We note that the intersection theory of currents even though very general, as it currently stands, cannot coincide with the intersection theory in algebraic geometry, since:

- The self-intersection of integration currents is not defined;
- The intersection of two positive currents is always positive.

In particular, we cannot yet define negative self-intersections of divisors using currents.

Theorem 2.14 ([BD20, Lemma 3.7]). Let $(\mathcal{T}_n) \rightarrow \mathcal{T}$ be a convergent sequence of horizontal-like positive closed currents to a horizontal-like current \mathcal{T} in $U \times V$. Let $a \in U$ and assume that the sequence of measures $(\langle \mathcal{T}_n, \pi_1, a \rangle)_n$ is also convergent. Then,

$$\lim_{n \rightarrow \infty} \langle \mathcal{T}_n | \pi_1 | a \rangle(\phi) \leq \langle \mathcal{T} | \pi_1 | a \rangle(\phi)$$

for every plurisubharmonic function ϕ on \mathbb{C}^d .

There is an simple version of the above theorem for supports which will be useful later.

Lemma 2.15. Assume that \mathcal{T}_n 's, \mathcal{S} and \mathcal{T} are all closed positive currents, and $\mathcal{T}_n \wedge \mathcal{S}$ and $\mathcal{T} \wedge \mathcal{S}$ are well-defined. If we have the following weak convergence, together with the convergence of supports in the Hausdorff metric, that is,

$$\mathcal{T}_n \rightarrow \mathcal{T}, \quad \text{supp}(\mathcal{T}_n) \rightarrow \text{supp}(\mathcal{T}).$$

Then, $\text{supp}(\lim(\mathcal{T}_n \wedge \mathcal{S})) \subseteq \text{supp}(\mathcal{T}) \cap \text{supp}(\mathcal{S})$.

Proof. For a point x outside the support of \mathcal{T} , there exists a sufficiently small radius ϵ , such that for a sufficiently large n , \mathcal{T}_n vanishes on the ball $B_\epsilon(x)$ centred at x . It follows that any limit of $\mathcal{T}_n \wedge \mathcal{S}$ vanishes on $B_\epsilon(x)$. So its support does not contain a . Moreover, its support does not contain any point outside $\text{supp}(\mathcal{S})$. \square

3. TROPICAL VARIETIES, TORI, TROPICAL CURRENTS

In this section, we recall the definition of tropical cycles and note that with the natural addition of tropical cycles and their *stable intersection*, the tropical cycles form a ring.

3.1. Tropical varieties. A linear subspace $H \subseteq \mathbb{R}^d$ is said to be *rational* if there exists a subset of \mathbb{Z}^d that spans H . A *rational polyhedron* is the intersection of finitely many rational half-spaces defined by

$$\{x \in \mathbb{R}^d : \langle m, x \rangle \geq c, \text{ for some } m \in \mathbb{Z}^d, c \in \mathbb{R}\}.$$

A *rational polyhedral complex* is a polyhedral complex consisting solely of rational polyhedra. The polyhedra in a polyhedral complex are also referred to as *cells*. A *fan* is a polyhedral complex whose cells are all cones. If every cone in a fan Σ is contained in another fan Σ' , then Σ is called a *subfan* of Σ . The one-dimensional cones of a fan are often called *rays*. Throughout this article, all fans and polyhedral complexes are assumed to be *rational*.

For a given polyhedron σ , and a finitely generated abelian group N , we denote by

$$\begin{aligned} \text{aff}(\sigma) &:= \text{affine span of } \sigma, \\ H_\sigma &:= \text{translation of } \text{aff}(\sigma) \text{ to the origin,} \\ N_\sigma &:= N \cap H_\sigma, \\ N(\sigma) &:= N/N_\sigma. \end{aligned}$$

Consider τ , a codimension one face of a p -dimensional polyhedron σ , and let $u_{\sigma/\tau}$ be the unique outward generator of the one-dimensional lattice $(\mathbb{Z}^d \cap H_\sigma)/(\mathbb{Z}^d \cap H_\tau)$.

Definition 3.1 (Balancing Condition and Tropical Cycles). Let \mathcal{C} be a p -dimensional polyhedral complex whose p -dimensional cones are equipped with integer weights w_σ . We say that \mathcal{C} satisfies the *balancing condition* at τ if

$$\sum_{\sigma \supset \tau} w(\sigma) u_{\sigma/\tau} = 0, \quad \text{in } \mathbb{Z}^d/(\mathbb{Z}^d \cap H_\tau),$$

where the sum is over all p -dimensional cells σ in \mathcal{C} containing τ as a face. A *tropical cycle* or a *tropical variety* in \mathbb{R}^d is a weighted complex with finitely many cells that satisfies the balancing condition at every cone of dimension $p - 1$. We say a tropical cycle is a *tropical fan* if its support is a fan.

3.2. Stable intersection and addition of tropical cycles. Recall that, generally speaking, the star of a cone in a complex is the extension of the local p -dimensional fan surrounding it. More precisely:

Definition 3.2. Given a polyhedral complex $\Sigma \subseteq \mathbb{R}^d$ and a cell $\tau \in \Sigma$, define the star of τ in Σ , denoted by $\text{star}_\Sigma(\tau)$, is a fan in \mathbb{R}^d . The cones of $\text{star}_\Sigma(\tau)$ are the *extensions* of cells σ that include τ as a face. Here, by extension, we mean

$$\bar{\sigma} = \{\lambda(x - y) : \lambda \geq 0, x \in \sigma, y \in \tau\}.$$

Definition 3.3 (Stable Intersection). (a) Let $\mathcal{C}_1, \mathcal{C}_2 \subseteq \mathbb{R}^d$ be two tropical cycles of dimension p and q , intersecting transversely. That is, the top dimensional cells $\sigma_1 \in \mathcal{C}_1$ and $\sigma_2 \in \mathcal{C}_2$ intersect in dimension $p+q-d$ and in the relative interior of these cells. Then the stable intersection of $\mathcal{C}_1 \cdot \mathcal{C}_2$ is the tropical cycles supported on finitely many cells $\mathcal{C}_1 \cap \mathcal{C}_2$. In this case, the weight of a cell $\sigma_1 \cap \sigma_2$ is defined by

$$w_{\mathcal{C}_1 \cdot \mathcal{C}_2}(\sigma_1 \cap \sigma_2) = w_{\sigma_1} w_{\sigma_2} [N : N_{\sigma_1} + N_{\sigma_2}],$$

where $N = \mathbb{Z}^d$ here.

(b) When \mathcal{C}_1 and \mathcal{C}_2 do not intersect transversely, then $\mathcal{C}_1 \cdot \mathcal{C}_2$ as a set is the Hausdorff limit of

$$\mathcal{C}_1 \cap (\epsilon b + \mathcal{C}_2), \quad \text{as } \epsilon \rightarrow 0,$$

for a fixed generic $b \in \mathbb{R}^d$, and the weights are the sum of all the tropical multiplicities of the cells in the transversal intersection $\mathcal{C}_1 \cap (\epsilon b + \mathcal{C}_2)$ which converge to the same $(p+q-d)$ -dimensional cell in the Hausdorff metric. Equivalently, for top dimensional cones $\sigma_1 \in \mathcal{C}_1$ and $\sigma_2 \in \mathcal{C}_2$

$$w_{\mathcal{C}_1 \cdot \mathcal{C}_2}(\sigma_1 \cap \sigma_2) = \sum_{\tau_1, \tau_2} w_{\tau_1} w_{\tau_2} [N : N_{\tau_1} + N_{\tau_2}],$$

where the sum is taken over all $\tau_1 \in \text{star}_{\mathcal{C}_1}(\sigma_1 \cap \sigma_2), \tau_2 \in \text{star}_{\mathcal{C}_2}(\sigma_1 \cap \sigma_2)$ with $\tau_1 \cap (\epsilon b + \tau_2) \neq \emptyset$, for some fixed generic vector $b \in \mathbb{R}^d$.

(c) When $p+q < d$, then the stable intersection of \mathcal{C}_1 and \mathcal{C}_2 is the empty set.

The following result is proved in tropical geometry; see [MS15, Lemmas 3.6.4 and 3.6.9]. We will revisit its proof later through the lens of superpotential theory.


Theorem 3.4. When $p+q \geq d$, the stable intersection, defined above, yields a balanced polyhedral complex of dimension $p+q-d$.

We also need the following for turning the set of tropical cycles into a \mathbb{Z} -algebra.

Definition 3.5 (Addition of Tropical Cycles). For two p -dimensional tropical cycles $\mathcal{C}_1, \mathcal{C}_2$ in \mathbb{R}^d , the addition $\mathcal{C}_1 + \mathcal{C}_2$ is the tropical cycle obtained by the common refinement of the support $|\mathcal{C}_1| \cup |\mathcal{C}_2|$ where the weights of a cone σ in the refinement are determined by $w_{\mathcal{C}_1 + \mathcal{C}_2}(\sigma) = w_{\mathcal{C}_1}(\sigma) + w_{\mathcal{C}_2}(\sigma)$.

Let us end this section with an example of the stable intersection.

Example 3.6. Consider the tropical cycles below. We can easily check that both cycles with the given weights satisfy the balancing condition. The line $\{x_2 = 0\}$ with weight 1 can be considered a tropical cycle that does not properly intersect the blue tropical variety. We can therefore translate by the vector $(0, -\epsilon)$ it to obtain $x_2 = -\epsilon$, and compute the stable intersection by taking the limit as $\epsilon \rightarrow 0$. Calculating the multiplicities and Hausdorff limit gives the stable intersection equal to the origin $(0, 0)$ with multiplicity 2. Note that we can choose any $\epsilon b \in \mathbb{R}^2$ generic for this translation, and thanks to Theorem 3.4 we obtain the same result.



4. TROPICAL CURRENTS

Let us briefly recall the definition of tropical currents from [Bab14, BH17]. To fix the notation,

$$\begin{aligned} T_N &:= \text{the complex algebraic torus } \mathbb{C}^* \otimes_{\mathbb{Z}} N, \\ S_N &:= \text{the compact real torus } S^1 \otimes_{\mathbb{Z}} N, \\ N_{\mathbb{R}} &:= \text{the real vector space } \mathbb{R} \otimes_{\mathbb{Z}} N. \end{aligned}$$

Let \mathbb{C}^* be the group of nonzero complex numbers. As before, the logarithm map is the homomorphism

$$\text{Log} : (\mathbb{C}^*)^d \longrightarrow \mathbb{R}^d, \quad (z_1, \dots, z_d) \longmapsto (-\log |z_1|, \dots, -\log |z_d|),$$

and the *argument map* is

$$\text{Arg} : (\mathbb{C}^*)^d \longrightarrow (S^1)^d, \quad (z_1, \dots, z_d) \longmapsto (z_1/|z_1|, \dots, z_d/|z_d|).$$

For a rational linear subspace $H \subseteq \mathbb{R}^d$ we have the following exact sequences:

$$0 \longrightarrow H \cap \mathbb{Z}^d \longrightarrow \mathbb{Z}^d \longrightarrow \mathbb{Z}^d / (H \cap \mathbb{Z}^d) \longrightarrow 0.$$

Moreover,

$$0 \longrightarrow S_{H \cap \mathbb{Z}^d} \longrightarrow (S^1)^d = S^1 \otimes_{\mathbb{Z}} \mathbb{Z}^d \longrightarrow S_{\mathbb{Z}^d / (H \cap \mathbb{Z}^d)} \longrightarrow 0.$$

Define

$$\pi_H : \text{Log}^{-1}(H) \xrightarrow{\text{Arg}} (S^1)^d \longrightarrow S_{\mathbb{Z}^d / (H \cap \mathbb{Z}^d)}.$$

Similarly,

$$0 \longrightarrow T_{H \cap \mathbb{Z}^d} \longrightarrow (\mathbb{C}^*)^d = \mathbb{C}^* \otimes_{\mathbb{Z}} \mathbb{Z}^d \longrightarrow T_{\mathbb{Z}^d / (H \cap \mathbb{Z}^d)} \longrightarrow 0.$$

We define

$$\Pi_H : (\mathbb{C}^*)^d \simeq \mathbb{C}^* \otimes ((H \cap \mathbb{Z}^d) \oplus \mathbb{Z}^d / (H \cap \mathbb{Z}^d)) \longrightarrow T_{\mathbb{Z}^d / (H \cap \mathbb{Z}^d)}.$$

One has

$$\ker(\Pi_H) = \ker(\pi_H) = T_{H \cap \mathbb{Z}^d} \subseteq (\mathbb{C}^*)^d.$$

As a result, when H is of dimension p , the set $\text{Log}^{-1}(H)$ is naturally foliated by the $\pi_H^{-1}(x) = T_{H \cap \mathbb{Z}^d} \cdot x \simeq (\mathbb{C}^*)^p$ for $x \in S_{\mathbb{Z}^d/(H \cap \mathbb{Z}^d)}$. For a lattice basis u_1, \dots, u_p , of $H \cap \mathbb{Z}^d$, the tori $T_{H \cap \mathbb{Z}^d} \cdot x$ can be parametrised by the monomial map

$$(\mathbb{C}^*)^p \longrightarrow (\mathbb{C}^*)^d, \quad z \longmapsto x \cdot z^{[u_1, \dots, u_p]^t}$$

where $U = [u_1, \dots, u_p]$ is the matrix with column vectors u_1, \dots, u_p , and z^{U^t} denotes that $z \in (\mathbb{C}^*)^p$ is taken to have the exponents with rows of the matrix U . Accordingly, one can easily check that

$$T_{H \cap \mathbb{Z}^d} \cdot x = \{z \in (\mathbb{C}^*)^d : z^{m_i} = x^{m_i}, i = 1, \dots, d-p\}.$$

for any choice of a \mathbb{Z} -basis $\{m_1, \dots, m_{d-p}\}$ of $\mathbb{Z}^d/(H \cap \mathbb{Z}^d)$.

Definition 4.1. Let H be a rational subspace of dimension p , and μ be the Haar measure of mass 1 on $S_{\mathbb{Z}^d/(H \cap \mathbb{Z}^d)}$. We define a (p, p) -dimensional closed current \mathcal{T}_H on $(\mathbb{C}^*)^d$ by

$$\mathcal{T}_H := \int_{x \in S_{\mathbb{Z}^d/(H \cap \mathbb{Z}^d)}} [\pi_H^{-1}(x)] d\mu(x).$$

When A is a rational affine subspace of \mathbb{R}^d parallel to the linear subspace $H = A - a$ for $a \in A$, we define \mathcal{T}_A by translation of \mathcal{T}_H . Namely, we define the submersion π_A as the composition

$$\pi_A : \text{Log}^{-1}(A) \xrightarrow{e^a} \text{Log}^{-1}(H) \xrightarrow{\pi_H} S_{\mathbb{Z}^d/(H \cap \mathbb{Z}^d)}.$$

We will call $T^A := \pi_A^{-1}(1) = \ker \pi_A = e^{-a} T_{H \cap \mathbb{Z}^d}$, the *distinguished fibre* of \mathcal{T}_A .

Definition 4.2. Let \mathcal{C} be a weighted polyhedral complex of dimension p . The tropical current $\mathcal{T}_{\mathcal{C}}$ associated to \mathcal{C} is given by

$$\mathcal{T}_{\mathcal{C}} = \sum_{\sigma} w_{\sigma} \mathbb{1}_{\text{Log}^{-1}(\sigma)} \mathcal{T}_{\text{aff}(\sigma)},$$

where the sum runs over all p -dimensional cells σ of \mathcal{C} .

Theorem 4.3 ([Bab14]). A weighted complex \mathcal{C} is balanced, if and only if, $\mathcal{T}_{\mathcal{C}}$ is closed.

Theorem 4.4 ([Bab14]). Any tropical current $\mathcal{T}_{\mathcal{C}} \in \mathcal{D}_{d-1}((\mathbb{C}^*)^d)$ is of the form $dd^c[\mathfrak{q} \circ \text{Log}]$, where $\mathfrak{q} : \mathbb{R}^d \rightarrow \mathbb{R}$, is a tropical Laurent polynomial, that is $\mathfrak{q}(x) = \max_{\alpha \in A} \{c_{\alpha} + \langle \alpha, x \rangle\}$, for $A \subseteq \mathbb{Z}^d$ a finite subset and $c_{\alpha} \in \mathbb{R}$.

Remark 4.5. Note that the support of $dd^c[\mathfrak{q} \circ \text{Log}]$, is given by $\text{Log}^{-1}(\text{Trop}(\mathfrak{q}))$, where $\text{Trop}(\mathfrak{q})$ is the set of points $x \in \mathbb{R}^d$ where \mathfrak{q} is not smooth at x . This set can be balanced with natural weights which coincides with the weights of the closed current $dd^c[\mathfrak{q} \circ \text{Log}]$ and it is called the tropical variety associated to \mathfrak{q} , and we can write $\mathcal{T}_{\text{Trop}(\mathfrak{q})} = dd^c[\mathfrak{q} \circ \text{Log}]$.

Proposition 4.6 ([Bab23, Proposition 4.6]). Assume that $\mathcal{T} \in \mathcal{D}_p((\mathbb{C}^*)^d)$ is a closed positive $(S^1)^d$ -invariant current whose support is given by $\text{Log}^{-1}(|\mathcal{C}|)$, for a polyhedral complex $\mathcal{C} \subseteq \mathbb{R}^d$ of pure dimension p . Then \mathcal{T} is a tropical current.

5. CONTINUITY OF SUPERPOTENTIALS

Let $q : \mathbb{R}^d \rightarrow \mathbb{R}$, be a tropical polynomial function, and $\text{Log} : (\mathbb{C}^*)^d \rightarrow \mathbb{R}^d$, as before. The current $dd^c[q \circ \text{Log}] \in \mathcal{D}_{d-1}((\mathbb{C}^*)^d)$ has a bounded potential on any relatively compact open set, and by Bedford–Taylor theory, for any positive closed current $\mathcal{T} \in \mathcal{D}_p(\mathbb{C}^*)^d$, the product

$$dd^c[q \circ \text{Log}] \wedge \mathcal{T} = dd^c([q \circ \text{Log}] \mathcal{T}),$$

is well-defined. See [Dem, Section III.3]. In higher codimensions though, to prove that any two tropical currents have a well-defined wedge product, we utilise the superpotential theory [DS09] on a compact Kähler manifold, and as a result, we extend the tropical currents to smooth compact toric varieties.

5.1. Tropical Currents on Toric Varieties. In a toric variety X_Σ , for a cone $\sigma \in \Sigma$, we denote by \mathcal{O}_σ , the toric orbit associated with σ . We have

$$X_\Sigma = \bigcup_{\sigma \in \Sigma} \mathcal{O}_\sigma.$$

We also set D_σ to be the closure of \mathcal{O}_σ in the X_Σ , and $\Sigma(p)$ the p -dimensional skeleton of Σ , that is, the union of p -dimensional cells of Σ . Fibres of tropical currents are algebraic varieties with finite degrees and can be extended by zero to any toric variety, in consequence, any tropical current can be extended by zero to toric varieties. Moreover, with the following compatibility condition, we can ask for the extension of the fibres to intersect the toric invariant divisors transversally.

Definition 5.1. (i) For a polyhedron σ , its *recession cone* is the convex polyhedral cone

$$\text{rec}(\sigma) = \{b \in \mathbb{R}^d : \sigma + b \subseteq \sigma\} \subseteq H_\sigma.$$

- (ii) Let \mathcal{C} be a p -dimensional balanced weighted complex in \mathbb{R}^d , and Σ a p -dimensional fan. We say that \mathcal{C} is *compatible* with Σ , if $\text{rec}(\sigma) \in \Sigma$ for all $\sigma \in \mathcal{C}$.
- (iii) We say the tropical current $\mathcal{T}_\mathcal{C}$ is *compatible* with X_Σ , if all the closures of the fibers $\pi_{\text{aff}(\sigma)}^{-1}(x)$ in X_Σ of $\mathcal{T}_\mathcal{C}$ intersect the torus invariant divisors of X_Σ transversely.

Theorem 5.2 ([BH17, Lemma 4.10]). Let \mathcal{C} be a p -dimensional tropical cycle and Σ be a fan. Assume that $\sigma \in \mathcal{C}$ is a p -dimensional polyhedron and $\rho \in \Sigma$ is a one-dimensional cone. Then

- (a) The intersection $D_\rho \cap \overline{\pi_{\text{aff}(\sigma)}^{-1}(x)}$ is non-empty and transverse, if and only if, $\rho \in \text{rec}(\sigma)$. Here $\overline{\pi_{\text{aff}(\sigma)}^{-1}(x)}$ corresponds the closure of a fiber of $\mathcal{T}_{\text{aff}(\sigma)}$ in the toric variety X_Σ .
- (b) In particular, \mathcal{C} is compatible with Σ , if and only if, $\mathcal{T}_\mathcal{C}$ is compatible with X_Σ .

For a tropical current $\mathcal{T}_\mathcal{C} \in \mathcal{D}_p((\mathbb{C}^*)^d)$, and given a toric variety X_Σ we denote its extension by zero $\overline{\mathcal{T}}_\mathcal{C} \in \mathcal{D}_p(X_\Sigma)$.

Proposition 5.3. For every tropical variety \mathcal{C} , there exists a smooth projective toric fan Σ compatible with a subdivision of \mathcal{C} .

Proof. By [BS11], for \mathcal{C} there is a refinement \mathcal{C}' , and a complete fan $\Sigma_1 \subseteq \mathbb{R}^d$ such that \mathcal{C}' is compatible with Σ_1 . Applying the toric Chow lemma [CLS11, Theorem 6.1.18] and the toric resolution of singularities [CLS11, Theorem 11.1.9] we can find a fan Σ which is a refinement of Σ_1 that defines a smooth projective variety X_Σ . The tropical variety \mathcal{C}'' which is the refinement of \mathcal{C}' induced by Σ , satisfies the statement. \square

Remark 5.4. When \mathcal{C}' is a refinement of a tropical variety \mathcal{C} , then \mathcal{C}' is a tropical variety with natural induced weights. It is also easy to check that we have the equality of currents $\mathcal{T}_{\mathcal{C}} = \mathcal{T}_{\mathcal{C}'}$ in $(\mathbb{C}^*)^d$; see [BH17, Section 2.6].

Lemma 5.5. Let $\mathbf{q} : \mathbb{R}^d \rightarrow \mathbb{R}$ be a tropical Laurent polynomial and X_Σ be a smooth projective toric variety compatible with a subdivision of $\text{Trop}(\mathbf{q})$. Let $\rho \in \Sigma(1)$. Assume that $\zeta_0 \in D_\rho \cap \text{supp}(\overline{dd^c[\mathbf{q} \circ \text{Log}]})$, and Ω is a sufficiently small neighbourhood of ζ_0 . Then, $\mathbf{q} \circ \text{Log} \in \text{PSH}(\Omega \setminus D_\rho) \cap \mathcal{C}^0(\Omega \setminus D_\rho)$ can be extended to a function $u : \Omega \rightarrow \mathbb{R} \cup \{+\infty\}$, such that

- (a) In Ω , $u = g + \kappa \log |f|$, where g is a continuous function, f is the local equation for D_ρ , and κ is a negative integer.
- (b) Restricted to Ω , we have $dd^c u = \overline{\mathcal{T}}_{\text{Trop}(\mathbf{q})} + \kappa [D_\rho]$.
- (c) In Ω , we have $\overline{\mathcal{T}}_{\mathcal{C}} = dd^c g$. In particular, $\overline{\mathcal{T}}_{\text{Trop}(\mathbf{q})}$ has a continuous superpotential.

Proof. Assume that $\mathbf{q} = \max_{\alpha \in A} \{c_\alpha + \langle \alpha, x \rangle\}$. Recall that

$$\text{Log} = (-\log |\cdot|, \dots, -\log |\cdot|).$$

We write

$$\mathbf{q} \circ \text{Log} = \log \max_{\alpha} \{e^{c_\alpha} z^{-\alpha}\}.$$

Assume that near ζ_0 , $\mathbf{q} \circ \text{Log}$ is given by $\max\{|e^{c_\beta} z^{-\beta}|, |e^{c_\gamma} z^{-\gamma}|\}$. This implies that in $\text{Log}(\Omega \setminus D_\rho)$, \mathbf{q} is given by $\max\{c_\beta + \langle \beta, x \rangle, c_\gamma + \langle \gamma, x \rangle\}$. For $\mathbf{q} = \max_{\alpha \in A} \{c_\alpha + \langle \alpha, x \rangle\}$ we set $\text{rec}(\mathbf{q}) = \max_{\alpha \in A} \{\langle \alpha, x \rangle\}$. It is not hard to check that

$$\text{rec}(\text{Trop}(\mathbf{q})) = \text{Trop}(\text{rec}(\mathbf{q}));$$

see [MS15, Page 132].

We now show that by extending each $z^{-\alpha}$ as a rational function to X_Σ , the compatibility condition implies that $\mathbf{q} \circ \text{Log}$ extends to X_Σ . By [CLS11, Proposition 4.1.2] the divisor of the extension of a character z^α in X_Σ is given by

$$(2) \quad \text{Div}(z^\alpha) = \sum_{\rho \in \Sigma(1)} \langle \alpha, n_\rho \rangle D_\rho,$$

where n_ρ is the minimal generator of ρ . By assumption,

$$D_\rho \cap \text{supp}(\overline{dd^c[\mathbf{q} \circ \text{Log}]}) \neq \emptyset.$$

Theorem 5.2 implies that

$$n_\rho \in \text{rec}(V_{\text{Trop}(\mathbf{q})}).$$

Moreover, if $\zeta_1 \in D_\rho \cap \text{supp}(\overline{dd^c[\text{rec}(\mathbf{q}) \circ \text{Log}]})$, then in a small neighbourhood of $\text{Log}(\zeta_1)$, $\text{rec}(\mathbf{q})(x) = \max\{\langle \beta, x \rangle, \langle \gamma, x \rangle\}$. By definition

$$n_\rho \in \text{rec}(\text{Trop}(\mathbf{q})) \quad \text{if and only if} \quad \kappa := \langle \beta, n_\rho \rangle = \langle \gamma, n_\rho \rangle.$$

This, together with Equation (2) implies that the extension of $z^{-\beta}$ and $z^{-\gamma}$ as rational functions to X_Σ have the same vanishing order along D_ρ , and we write $z^{-\beta} = f^\kappa \frac{g_1}{h_1}$ and $z^{-\gamma} = f^\kappa \frac{g_2}{h_2}$. Now note that in $\Omega \setminus D_\rho$,

$$\mathfrak{q} \circ \text{Log} = \max \log \{|e^{c_\beta} z^{-\beta}|, |e^{c_\gamma} z^{-\gamma}|\} = \kappa \log |f| + \max \{|e^{c_\beta} \frac{g_1}{h_1}|, |e^{c_\gamma} \frac{g_2}{h_2}|\},$$

we must have $\kappa < 0$, otherwise $q \circ \text{Log} = -\infty$ in $\Omega \setminus D_\rho$. Consequently, $\mathfrak{q} \circ \text{Log} : \Omega \setminus D_\rho \rightarrow \mathbb{R}$, can be extended to

$$u := \kappa \log |f| + \max \{|e^{c_\beta} \frac{g_1}{h_1}|, |e^{c_\gamma} \frac{g_2}{h_2}|\}$$

on Ω . Setting

$$g = \max \{|e^{-c_\beta} \frac{g_1}{h_1}|, |e^{-c_\gamma} \frac{g_2}{h_2}|\},$$

implies (a).

We have

$$dd^c[\mathfrak{q} \circ \text{Log}]|_{\Omega \setminus D_\rho} = (dd^c \log |f|^\kappa dd^c \log |g|)|_{\Omega \setminus D_\rho} = dd^c \log |g|_{|\Omega \setminus D_\rho},$$

since $dd^c \log |f|^\kappa$ is holomorphic in $\Omega \setminus D_\rho$. As a result of compatibility with X_Σ , $\overline{dd^c[\mathfrak{q} \circ \text{Log}]}$ does not charge any mass in D_ρ , and we obtain

$$\overline{dd^c[\mathfrak{q} \circ \text{Log}]} = dd^c \log |g|.$$

This together with Theorem 4.4 implies (c) and (b). \square

Lemma 5.6. Assume that σ is p -dimensional and $\text{aff}(\sigma) = H_1 \cap \cdots \cap H_{d-p}$, is given as the transversal intersection hyperplanes $H_i \subseteq \mathbb{R}^d$. If Σ is a smooth projective fan compatible with $\bigcup_i H_i$, then

$$\overline{\mathcal{T}_{\text{aff}(\sigma)}} \leq \overline{\mathcal{T}_{H_1} \wedge \cdots \wedge \mathcal{T}_{H_{d-p}}} = \overline{\mathcal{T}_{H_1}} \wedge \cdots \wedge \overline{\mathcal{T}_{H_{d-p}}}.$$

Proof. By the definition of tropical currents, we have the inequality

$$\mathcal{T}_{\text{aff}(\sigma)} \leq \mathcal{T}_{H_1} \wedge \cdots \wedge \mathcal{T}_{H_{d-p}},$$

as currents in $(\mathbb{C}^*)^d$, since the right-hand side might have multiplicities but the currents have the same support. Now, the wedge products in X_Σ are well-defined by Lemma 5.5 and Theorem 2.4. As both currents on both sides of the equation coincide on $(\mathbb{C}^*)^d$, the support of the current on the right-hand side contains the closure of the support of $\mathcal{T}_{\text{aff}(\sigma)}$ in X_Σ . For the equality, note that compatibility with Σ , implies that $\overline{\mathcal{T}_{H_i}}$ has a zero mass in $X_\Sigma \setminus T_N$. \square

Theorem 5.7. Let \mathcal{C} be a positively weighted tropical cycle of dimension p compatible with a smooth, projective fan Σ , then $\overline{\mathcal{T}_{\mathcal{C}}}$ has a continuous superpotential in X_Σ .

We need the following definition.

Definition 5.8. We define the affine extension p -dimensional a tropical cycle \mathcal{C} , by as the addition of tropical cycles

$$\widehat{\mathcal{C}} := \sum_{\sigma \in \mathcal{C}} w_\sigma \text{aff}(\sigma).$$

It is clear that if \mathcal{C} is a positively weighted tropical cycle, then $\mathcal{T}_{\widehat{\mathcal{C}}} - \mathcal{T}_{\mathcal{C}} \geq 0$.

Proof of 5.7. Let $\widehat{\mathcal{C}}$ be the affine extension of \mathcal{C} , and $\widehat{\Sigma}$ be a smooth projective fan which is a refinement of Σ and compatible with $\widehat{\mathcal{C}}$. By the preceding lemma and repeated application of Theorem 2.4 for any $\sigma \in \mathcal{C}$, $\overline{\mathcal{T}}_{\text{aff}(\sigma)}$ has a bounded superpotential, which implies this property for $\overline{\mathcal{T}}_{\widehat{\mathcal{C}}}$. Now, since $\mathcal{T}_{\widehat{\mathcal{C}}} - \mathcal{T}_{\mathcal{C}}$ is a positive closed tropical current in $(\mathbb{C}^*)^d$,

$$\overline{\mathcal{T}_{\widehat{\mathcal{C}}} - \mathcal{T}_{\mathcal{C}}} = \overline{\mathcal{T}}_{\widehat{\mathcal{C}}} - \overline{\mathcal{T}}_{\mathcal{C}} \geq 0$$

in $X_{\widehat{\Sigma}}$. Continuity of the superpotential of $\overline{\mathcal{T}}_{\mathcal{C}}$ in $X_{\widehat{\Sigma}}$ follows from Theorem 2.3.

We now show that $\overline{\mathcal{T}}_{\mathcal{C}}$ has also a continuous super-potential on X_{Σ} as well. We consider the proper map $f : X_{\widehat{\Sigma}} \rightarrow X_{\Sigma}$, which can be understood as a composition of multiple blow-ups along toric points with exceptional divisors D_{ρ} for any ray $\rho \in \widehat{\Sigma} \setminus \Sigma$. These divisors satisfy $D_{\rho} \cap \text{supp}(\overline{\mathcal{T}}_{\mathcal{C}}) = \emptyset$. We conclude the proof by Theorem 2.9. \square

Remark 5.9. When \mathcal{C} is a tropical cycle which is not positively weighted, then there exist positively weighted tropical cycles \mathcal{C}_1 and \mathcal{C}_2 such that

$$\mathcal{C} = \mathcal{C}_1 - \mathcal{C}_2.$$

Therefore, if Σ is compatible with both \mathcal{C}_1 and \mathcal{C}_2 , then by preceding lemma, $\overline{\mathcal{T}}_{\mathcal{C}}$ has a continuous superpotential in X_{Σ} .

Proposition 5.10. Let \mathcal{C}_1 and \mathcal{C}_2 be two positively weighted tropical cycles. If X_{Σ} is compatible $\mathcal{C}_1 + \mathcal{C}_2$, then

$$\overline{\mathcal{T}_{\mathcal{C}_1} \wedge \mathcal{T}_{\mathcal{C}_2}} = \overline{\mathcal{T}}_{\mathcal{C}_1} \wedge \overline{\mathcal{T}}_{\mathcal{C}_2}.$$

Proof. The proof is clear since both $\overline{\mathcal{T}}_{\mathcal{C}_1}$ and $\overline{\mathcal{T}}_{\mathcal{C}_2}$ have continuous superpotentials with no mass on the boundary divisors $X_{\Sigma} \setminus T_N$. \square

Proposition 5.11. For any two tropical currents \mathcal{C}_1 and \mathcal{C}_2 , the intersection product

$$\mathcal{T}_{\mathcal{C}_1} \wedge \mathcal{T}_{\mathcal{C}_2} := \overline{\mathcal{T}}_{\mathcal{C}_1} \wedge \overline{\mathcal{T}}_{\mathcal{C}_2}|_{(\mathbb{C}^*)^d},$$

does not depend on the choice of a smooth projective toric variety of the fan Σ compatible with $\mathcal{C}_1 + \mathcal{C}_2$, where $(\mathbb{C}^*)^d$ is identified with $T_N \subseteq X_{\Sigma}$. Moreover, this product coincides with the definition of wedge products with bi-degree $(1, 1)$ tropical currents in Bedford–Taylor Theory in $(\mathbb{C}^*)^d$.

Proof. This is a consequence of Lemma 2.7, and the fact that intersection product with a bidegree $(1, 1)$ current in super-potential theory, in an open set of compact Kähler manifold, coincides with the Bedford–Taylor theory. \square

Proposition 5.12. Stable intersection of tropical cycles is associative and commutative

Proof. This is the application of Theorem 5.7 and Theorem 2.5. \square

5.2. Proof of $\mathcal{T}_{\mathcal{C}_1} \wedge \mathcal{T}_{\mathcal{C}_2} = \mathcal{T}_{\mathcal{C}_1 \cdot \mathcal{C}_2}$. We prove our main intersection theorems here, and it is not hard to visit the proof of Theorem 3.4 using tools from superpotential theory.

Theorem 5.13. For two tropical varieties \mathcal{C} and \mathcal{C}' of dimension p and q , respectively, we have

$$\mathcal{T}_{\mathcal{C}_1} \wedge \mathcal{T}_{\mathcal{C}_2} = \mathcal{T}_{\mathcal{C}_1 \cdot \mathcal{C}_2}$$

where the $\mathcal{C} \cdot \mathcal{C}'$ is the stable intersection \mathcal{C}_1 and \mathcal{C}_2 , defined in Definition 3.3. Moreover, the $\mathcal{C}_1 \cdot \mathcal{C}_2$ is a balanced polyhedral complex of dimension $p + q - d$.

Proposition 5.14 ([Kat09, Propositions 6.1]). Let $H_1, H_2 \subseteq \mathbb{R}^d$ be two rational planes of dimension p and q with $p + q = d$ that intersect transversely. Then, the complex tori $T_{H_1 \cap \mathbb{Z}^d}$ and $T_{H_2 \cap \mathbb{Z}^d}$ intersect at $[N : N_{H_1} + N_{H_2}]$ distinct points.

Proof of Theorem 5.13. Note that $\mathcal{T}_{\mathcal{C}_1} \wedge \mathcal{T}_{\mathcal{C}_2}$ is well-defined by Proposition 5.11. Assume that \mathcal{C}_1 and \mathcal{C}_2 are two tropical cycles of dimension p and q , respectively. Note that when $p + q < d$, both sides of the equality are zero. Therefore, we assume that $p + q \geq d$. We proceed with the following steps:

- (a) In the transversal case $\text{supp}(\mathcal{T}_{\mathcal{C}_1} \wedge \mathcal{T}_{\mathcal{C}_2}) = \text{Log}^{-1}(\mathcal{C}_1 \cdot \mathcal{C}_2) = \text{Log}^{-1}(\mathcal{C}_1 \cap \mathcal{C}_2)$.
- (b) When $p + q = d$, $\mathcal{T}_{\mathcal{C}_1} \wedge \mathcal{T}_{\mathcal{C}_2} = \mathcal{T}_{\mathcal{C}_1 \cdot \mathcal{C}_2}$, in the transversal case.
- (c) When $p + q > d$, $\mathcal{T}_{\mathcal{C}_1} \wedge \mathcal{T}_{\mathcal{C}_2} = \mathcal{T}_{\mathcal{C}_1 \cdot \mathcal{C}_2}$, in the transversal case.
- (d) $\text{Log}(\text{supp}(\mathcal{T}_{\mathcal{C}_1} \wedge \mathcal{T}_{\mathcal{C}_2})) = \mathcal{C}_1 \cdot \mathcal{C}_2$ when $p + q = d$, also in the non-transversal case.
- (e) $\text{Log}(\text{supp}(\mathcal{T}_{\mathcal{C}_1} \wedge \mathcal{T}_{\mathcal{C}_2})) = \mathcal{C}_1 \cdot \mathcal{C}_2$ when $p + q > d$, also in the non-transversal case.

To see (a), note the by Theorem 2.4(c), $\text{supp}(\mathcal{T}_{\mathcal{C}_1} \wedge \mathcal{T}_{\mathcal{C}_2}) \subseteq \text{Log}^{-1}(\mathcal{C}_1 \cdot \mathcal{C}_2) = \text{Log}^{-1}(\mathcal{C}_1 \cap \mathcal{C}_2)$. Moreover, on $\mathcal{C}_1 \cap \mathcal{C}_2$, the fibres of $\mathcal{T}_{\mathcal{C}_1}$ and $\mathcal{T}_{\mathcal{C}_2}$ have a non-zero intersection.

To prove (b), let $a \in \mathcal{C}_1 \cap \mathcal{C}_2$ be an isolated point of intersection. We can choose a small ball $B_\epsilon(a) \subseteq \mathbb{R}^d$ such that a is an isolated point of intersection $\sigma_1 \cap \sigma_2 \cap B$, where σ_1 and σ_2 are cells of dimension p and q in \mathcal{C}_1 and \mathcal{C}_2 respectively. For any rational polyhedron σ , let

$$\begin{aligned} N_\sigma &:= N \cap \text{aff}(\sigma), \\ S^1(\sigma) &:= S^1 \otimes_{\mathbb{Z}} (\mathbb{Z}^d / (\mathbb{Z}^d \cap \text{aff}(\sigma))), \\ \pi_\sigma &:= \pi_{\text{aff}(\sigma)}, \end{aligned}$$

where $\pi_{\text{aff}(\sigma)}$ was defined after Definition 4.1. By Lemma 2.7

$$\begin{aligned} \mathcal{T}_{\mathcal{C}_1} \wedge \mathcal{T}_{\mathcal{C}_2}|_{\text{Log}^{-1}(B)} &= w_{\sigma_1} w_{\sigma_2} \mathbb{I}_{\text{Log}^{-1}(B)} \\ &\quad \int_{(x_1, x_2) \in S^1(\sigma_1) \times S^1(\sigma_2)} [\pi_{\sigma_1}^{-1}(x_1)] \wedge [\pi_{\sigma_2}^{-1}(x_2)] d\mu_{\sigma_1}(x_1) \otimes d\mu_{\sigma_2}(x_2). \end{aligned}$$

Transversality of the fibres implies

$$[\pi_{\sigma_1}^{-1}(x_1)] \wedge [\pi_{\sigma_2}^{-1}(x_2)] = [\pi_{\sigma_1}^{-1}(x_1) \cap \pi_{\sigma_2}^{-1}(x_2)].$$

By Proposition 5.14, we have $\kappa = [\mathbb{Z}^d : \mathbb{Z}_{\text{aff}(\sigma)}^d + \mathbb{Z}_{\text{aff}(\sigma')}^d]$ distinct intersection points covering $\text{Log}^{-1}(a) \simeq S^1(\sigma_1) \times S^1(\sigma_2)$. When $(x_1, x_2) \in \text{Log}^{-1}(a) \simeq$ vary with respect to the normalised Haar measure, these κ points cover $S^1(\sigma_1) \times S^1(\sigma_2) \simeq (S^1)^d$ with speed

κ . As a result,

$$\begin{aligned} \int_{(x_1, x_2) \in S^1(\sigma_1) \times S^1(\sigma_2)} [\pi_{\sigma_1}^{-1}(x)] \wedge [\pi_{\sigma_2}^{-1}(x')] d\mu_{\sigma_1}(x) \otimes d\mu_{\sigma_2}(x') \\ = \int_{y \in (S^1)^d} \kappa[\pi_{\sigma_1 \cap \sigma_2}^{-1}(y)] d\mu_{\sigma_1 \cap \sigma_2}(y). \end{aligned}$$

This proves (b).

To deduce (c), let $\sigma = \sigma_1 \cap \sigma_2$ be a $(p + q - d)$ -dimensional cell in the intersection. Assume that $0 \in \sigma$, by a translation, and $L := \text{aff}(\sigma)^\perp$. By Proposition 5.12,

$$(\mathcal{T}_{\mathcal{C}_1} \wedge \mathcal{T}_{\mathcal{C}_2}) \wedge \mathcal{T}_L = \mathcal{T}_{\mathcal{C}_1} \wedge (\mathcal{T}_{\mathcal{C}_2} \wedge \mathcal{T}_L).$$

Note that $[N : N_\sigma + N_L] = 1$. Assume that $w_{\sigma_1} = w_{\sigma_2} = 1$. As a result, if the multiplicity of $\mathcal{T}_{\mathcal{C}_1} \wedge \mathcal{T}_{\mathcal{C}_2}$ at σ equals κ , we have that the multiplicity of $(\mathcal{T}_{\mathcal{C}_1} \wedge \mathcal{T}_{\mathcal{C}_2}) \wedge \mathcal{T}_L$ at the origin is also κ . We have that $[N : N_{\sigma_1} + N_{\sigma_2 \cap L}] = [N/N_\sigma : (N_{\sigma_1} + N_{\sigma_2 \cap L})/N_\sigma]$, which equals $[N/N_\sigma : N_{\sigma_1 \cap L} + N_{\sigma_2 \cap L}] = [N : N_{\sigma_1} + N_{\sigma_2}]$. As a consequence, the intersection multiplicity induced on σ by $\mathcal{T}_{\mathcal{C}_1} \wedge \mathcal{T}_{\mathcal{C}_2}$ equals the intersection multiplicity in Definition 3.3.

To prove (d) note that if $\mathcal{C}_1 + \epsilon b$ is the translation of the the tropical variety, where $b \in \mathbb{R}^d$ and $\epsilon \in \mathbb{R}_{\geq 0}$, then $(e^{\epsilon b})^* \mathcal{T}_{\mathcal{C}_1} = \mathcal{T}_{\mathcal{C}_1 + \epsilon b}$. Moreover, we have the SP-convergence of currents with continuous superpotentials.

$$(e^{\epsilon b})^* \mathcal{T}_{\mathcal{C}_1} \longrightarrow \mathcal{T}_{\mathcal{C}_1}, \quad \text{as } \epsilon \rightarrow 0.$$

Therefore, by Theorem 2.4,

$$(3) \quad (e^{\epsilon b})^* \mathcal{T}_{\mathcal{C}_1} \wedge \mathcal{T}_{\mathcal{C}_2} = \mathcal{T}_{\mathcal{C}_1 + \epsilon b} \wedge \mathcal{T}_{\mathcal{C}_2} \longrightarrow \mathcal{T}_{\mathcal{C}_1} \wedge \mathcal{T}_{\mathcal{C}_2}, \quad \text{as } \epsilon \rightarrow 0.$$

Considering the support, we obtain the Hausdorff limit

$$\limsup((e^{\epsilon b})^* \mathcal{T}_{\mathcal{C}_1} \wedge \mathcal{T}_{\mathcal{C}_2}) \supseteq \text{supp}(\mathcal{T}_{\mathcal{C}_1} \wedge \mathcal{T}_{\mathcal{C}_2}).$$

We now note that for all ϵ , the number of intersection points in $(\mathcal{C}_1 + \epsilon b) \cap \mathcal{C}_2$ is uniformly bounded by the number of p -dimensional cells in \mathcal{C}_1 and q -dimensional cells in \mathcal{C}_2 , the Hausdorff limit of $(\mathcal{C}_1 + \epsilon b) \cap \mathcal{C}_2$ is also zero dimensional. Now, by definition of $\mathcal{C}_1 \cdot \mathcal{C}_2$ it suffices to show that

$$\limsup((e^{\epsilon b})^* \mathcal{T}_{\mathcal{C}_1} \wedge \mathcal{T}_{\mathcal{C}_2}) = \text{supp}(\mathcal{T}_{\mathcal{C}_1} \wedge \mathcal{T}_{\mathcal{C}_2}),$$

for any fixed generic b . This is also easy. Let $a_\epsilon \in (\mathcal{C}_1 + \epsilon b) \cap \mathcal{C}_2$. Since the translation by ϵb does not change the slopes of the cells, as $a_\epsilon \rightarrow a$, the multiplicity for all a_ϵ remains constant for $\epsilon > 0$, therefore $\lim(e^{\epsilon b})^* \mathcal{T}_{\mathcal{C}_1} \wedge \mathcal{T}_{\mathcal{C}_2}$ has a non-zero mass at $\text{Log}^{-1}(a)$.

For Part (e), first observe that any $\lim(\mathcal{C}_1 + \epsilon b) \cap \mathcal{C}_2$ is obtained by a translation ϵb , as $\epsilon \rightarrow 0$, of finitely many $(p + q - d)$ -dimensional cells. Therefore, $\mathcal{C}_1 \cdot \mathcal{C}_2$ is also of dimension $p + q - d$, and the SP-convergence readily implies that the limit is independent of generic b . Now, it only remains to show that

$$\limsup((e^{\epsilon b})^* \mathcal{T}_{\mathcal{C}_1} \wedge \mathcal{T}_{\mathcal{C}_2}) = \text{supp}(\mathcal{T}_{\mathcal{C}_1} \wedge \mathcal{T}_{\mathcal{C}_2}).$$

Let σ be a $p + q - d$ dimensional cell in $\mathcal{C}_1 \cdot \mathcal{C}_2$, and by translation, assume that $0 \in \sigma$. Let $L = \text{aff}(\sigma)^\perp$. By Proposition 2.6,

$$((e^{cb})^* \mathcal{T}_{\mathcal{C}_1} \wedge \mathcal{T}_{\mathcal{C}_2}) \wedge \mathcal{T}_L \longrightarrow (\mathcal{T}_{\mathcal{C}_1} \wedge \mathcal{T}_{\mathcal{C}_2}) \wedge \mathcal{T}_L, \quad \text{as } \epsilon \rightarrow 0.$$

By Part (d), the left-hand-side has mass at the origin. This show (e).

To deduce Part (f), recall that in Equation (3) since we can choose b generically and the previous discussion. The balancing condition is also deduced by the fact that $\mathcal{T}_{\mathcal{C}_1} \wedge \mathcal{T}_{\mathcal{C}_2}$ is closed and Theorem 4.3. \square

Let us recall a few notable theorems from [McM89], [FS97], and [JY16], and add the tropical currents into the picture.

Theorem 5.15. The assignment $\mathcal{C} \mapsto \mathcal{T}_{\mathcal{C}}$ induces a \mathbb{Q} -algebra isomorphism between

- (a) The \mathbb{Q} -algebra of tropical currents on $(\mathbb{C}^*)^d$ with the usual addition of currents and the wedge product.
- (b) The \mathbb{Q} -algebra generated by the tropical cycles in \mathbb{R}^d with the natural addition (Definition 3.5) and stable intersection (Definition 3.3).

Further, considering the map $\Phi_m : (\mathbb{C}^*)^d \longrightarrow (\mathbb{C}^*)^d$, $z \mapsto z^m$. We have the isomorphism of the following:

- (a') The \mathbb{Q} -algebra of Φ_m -invariant tropical currents on $(\mathbb{C}^*)^d$ with the usual addition of currents and the wedge product.
- (b') The \mathbb{Q} -algebra generated by the tropical fans in \mathbb{R}^d with the natural addition and stable intersection.
- (c') The \mathbb{Q} -algebra generated by tropical hypersurfaces.
- (d') The McMullen algebra of dimensional polytopes in \mathbb{Q}^d .
- (e') Chow group of the compact space obtained as a direct limit of all complete toric varieties.

Proof. By Proposition 5.12, Items (a) and (a') form a \mathbb{Q} -algebra. In Item (b), the fact that the tropical cycles in \mathbb{R}^d generate an algebra is the content of [JY16, Theorem 2.6], which can also be deduced from Theorem 5.13 and Proposition 5.12, then the isomorphism of (a) and (b) is also clear.

For the isomorphism between (a') and (b'), we observe that a tropical current of bidimension (p, p) , denoted $\mathcal{T}_{\mathcal{C}}$, is invariant under the map Φ , that is,

$$m^{p-d} \Phi_m^* \mathcal{T}_{\mathcal{C}} = \mathcal{T}_{\mathcal{C}}$$

if and only if \mathcal{C} is a tropical fan.

The isomorphism between (b') and (e') is discussed in [FS97] and can also be deduced directly from [BH17, Theorem 4.7]. The isomorphism between (d') and (e') is proved in [FS97, Theorem 4.2], while that between (b') and (d') appears in [JY16, Theorem 5.1]. The equality of (b') and (c') is treated independently in [Kaz03, Proposition 9] and [JY16, Corollary 5.2]. \square

A generalisation of the isomorphism between items (a') and (d') to a homeomorphism in dimension 2, along with its applications to dynamical systems, is presented in [DR25].

5.2.1. *Calculating Intersection Multiplicities Using Monge-Ampère Measures.* Using the equality of the supports in the previous section, we only need to prove the intersection multiplicities in the transversal case locally.

5.2.2. *The Real Monge-Ampère Measures.* Let $\Omega \subseteq \mathbb{R}^d$ be an open subset and $u : \Omega \rightarrow \mathbb{R}$ be a convex (hence continuous) function. The *generalised gradient* of u at $x_0 \in \Omega$ is defined by

$$\nabla u(x_0) = \{\xi \in (\mathbb{R}^d)^* : u(x) - u(x_0) \geq \langle \xi, x - x_0 \rangle, \text{ for all } x \in \Omega\}.$$

In the above, $\langle \cdot, \cdot \rangle$ denotes the standard inner product in \mathbb{R}^d , and $(\mathbb{R}^d)^*$ is the dual. The real Monge-Ampère measure associated to a convex function u on a Borel set $E \subseteq \Omega$, is given by

$$\text{MA}[u](E) = \mu\left(\bigcup_{y \in E} \nabla u(y)\right),$$

where μ is the Lebesgue measure on $(\mathbb{R}^d)^*$.

It is interesting that for the tropical polynomials, one can compute the associate real Monge-Ampère measures explicitly. Recall that, for any tropical polynomial, there is a natural subdivision of its Newton polytope which is dual to the tropical variety of it. See Figure for an example and [BS14, MS15] for details.

Lemma 5.16 ([Yge13, Page 59], [BGPS14, Proposition 2.7.4]). Let $\mathbf{q} : \mathbb{R}^d \rightarrow \mathbb{R}$ be a tropical polynomial associated tropical variety $\mathcal{C} = V_{\text{trop}}(\mathbf{q})$, one has

$$\text{MA}[\mathbf{q}] = \sum_{a \in \mathcal{C}(0)} \text{Vol}(\{a\}^*) \delta_a,$$

where $\mathcal{C}(0)$ is the 0-dimensional skeleton of \mathcal{C} , and $\{a\}^*$ is the dual of the vertex $a \in \mathcal{C}(0)$.

A detailed discussion of the preceding theorem can be also found in [Bab14].

5.3. **Polarisation.** For n convex functions $u_1, \dots, u_d : \mathbb{R}^d \rightarrow \mathbb{R}$, their *mixed Monge-Ampère measure* is defined by

$$\widetilde{\text{MA}}[u_1, \dots, u_d] = \frac{1}{d!} \sum_{k=1}^d \sum_{1 \leq j_1 < \dots < j_k \leq d} (-1)^{d-k} \text{MA}[u_{j_1} + \dots + u_{j_k}].$$

Recall that this is how the *mixed volume* of n convex bodies can be defined from the n -dimensional volume. Moreover, it is easy to check that for a convex function $u : \mathbb{R}^d \rightarrow \mathbb{R}$, $\text{MA}[u] = \widetilde{\text{MA}}[u, \dots, u]$.

It is customary at this point to prove the Bernstein-Khovanskii-Kushnirenko Theorem, which is simply obtained by taking the total mass from Lemma 5.16.

Proposition 5.17. Let $\mathbf{q}, \mathbf{q}_1, \dots, \mathbf{q}_d : \mathbb{R}^d \rightarrow \mathbb{R}$ be tropical polynomials. We have the following facts:

- (a) $\text{MA}[\mathbf{q}](\mathbb{R}^d) = \text{Vol}_d(\Delta_{\mathbf{q}})$, where $\Delta_{\mathbf{q}}$ is the Newton polytope of \mathbf{q} .
- (b) (Tropical Bernstein-Khovanskii-Kushnirenko Theorem) $\widetilde{\text{MA}}[\mathbf{q}_1, \dots, \mathbf{q}_d](\mathbb{R}^d) = \widetilde{\text{Vol}}(\Delta_{\mathbf{q}_1}, \dots, \Delta_{\mathbf{q}_d})$, where $\widetilde{\text{Vol}}$ is the mixed volume.

Proposition 5.18. Assume that $\alpha_i, \beta_i \in \mathbb{Z}^d$ for $i = 1, \dots, d$. Let $\mathbf{q}_i = \max\{\langle \alpha_i, x \rangle, \langle \beta_i, x \rangle\}$ be n tropical polynomials. Then,

$$d! \widetilde{\text{MA}}[\mathbf{q}_1, \dots, \mathbf{q}_d] = \kappa \delta_0,$$

where κ is given by the volume *zonotope* of the Minkowski sum of the vectors $\sum_{i=1}^d [\alpha_i - \beta_i]$. In addition, these multiplicities coincide with the intersection multiplicities in Definition 3.3 for d tropical hypersurfaces.

Proof. Note that $\Delta_{\mathbf{q}_i}$ is the line segment between α_i and β_i . Moreover, in the definition of $\widetilde{\text{MA}}[\mathbf{q}_1, \dots, \mathbf{q}_d]$ only $\text{Vol}(\sum_{i=1}^d [\alpha_i - \beta_i])$ possibly has a non-zero n -dimensional volume. Finally, the origin is the only 0-dimensional cell of the tropical variety of polynomial $\mathbf{q}_1 + \dots + \mathbf{q}_d$, if and only if, $\{\alpha_1 - \beta_1, \dots, \alpha_d - \beta_d\}$ forms a linearly independent set. Therefore, $d! \text{MA}[\mathbf{q}_1 + \dots + \mathbf{q}_d] = \kappa \delta_0$. To prove the second statement, we first claim that any Borel subset $\Omega \subseteq \mathbb{R}^n$,

$$\int_{\text{Log}^{-1}(U)} \mathcal{T}_{\text{Trop}(\mathbf{q}_1)} \wedge \dots \wedge \mathcal{T}_{\text{Trop}(\mathbf{q}_d)} = d! \widetilde{\text{MA}}[\mathbf{q}_1, \dots, \mathbf{q}_d](\Omega).$$

To see this, recall that $\mathcal{T}_{\text{Trop}(\mathbf{q}_i)} = dd^c[\mathbf{q}_i \circ \text{Log}]$, by Theorem 4.4 and Remark 4.5. Further, by [Ras01]

$$\int_{\text{Log}^{-1}(U)} (dd^c[\mathbf{q} \circ \text{Log}])^d = d! \text{MA}[\mathbf{q}](U).$$

Polarising both sides proves the claim. This implies that the tropical intersection multiplicities coincide with the multiplicities in this proposition for hypersurfaces. \square

6. SLICING TROPICAL CURRENTS

Proposition 6.1. Let \mathcal{C} be a p -dimensional positively weighted tropical cycle in \mathbb{R}^d with $p \geq 1$. Assume that $S \subseteq (\mathbb{C}^*)^d$ is an algebraic hypersurface with transversal intersection with $\mathcal{T}_{\mathcal{C}}$. Then, $[S] \wedge \mathcal{T}_{\mathcal{C}}$ is admissible and it is a closed positive current of bidimension $(p-1, p-1)$ given by

$$[S] \wedge \mathcal{T}_{\mathcal{C}} = \sum_{\sigma \in \mathcal{C}} w_{\sigma} \int_{x \in S_{N(\sigma)}} \mathbb{1}_{\text{Log}^{-1}(\sigma^{\circ})} [S \cap \pi_{\text{aff}(\sigma)}^{-1}(x)] d\mu(x).$$

Proof. The idea of the proof is similar to that of [BH17, Proposition 4.11]. Let f be a polynomial with vanishing S in $(\mathbb{C}^*)^d$. Assume that $\text{Log}^{-1}(\sigma^{\circ}) \cap S \neq \emptyset$, for a p -dimensional cell $\sigma \in \mathcal{C}$, then for each fibre, $\pi_{\sigma}^{-1}(x) := \pi_{\text{aff}(\sigma)}^{-1}(x)$, the transversality assumption allows for the application of the Lelong–Poincaré formula to deduce

$$dd^c(\log |f| \mathbb{1}_{\text{Log}^{-1}(\sigma^{\circ})} [\pi_{\sigma}^{-1}(x)]) = \mathbb{1}_{\text{Log}^{-1}(\sigma^{\circ})} [S \cap \pi_{\sigma}^{-1}(x)] + \mathcal{R}_{\sigma}(x),$$

where $\mathcal{R}_{\sigma}(x)$ is a $(p-1, p-1)$ -bidimensional current. The support of $\mathcal{R}_{\sigma}(x)$ lies in the boundary of $\text{Log}^{-1}(\sigma)$, as $\mathcal{R}_{\sigma}(x)$ is the difference of two currents that coincide in any set of form $\text{Log}^{-1}(B)$, where $B \subseteq \mathbb{R}^d$ is a small ball with

$$B \cap \sigma' = \emptyset, \quad \text{for a } p\text{-dimensional cell } \sigma' \in \mathcal{C}, \sigma' \neq \sigma,$$

and both vanish outside $\text{Log}^{-1}(\sigma)$. Integrating along the fibers, and adding for all p -dimensional cells $\sigma \in \mathcal{C}$, we obtain

$$[S] \wedge \mathcal{T}_{\mathcal{C}} = \sum_{\sigma \in \mathcal{C}} w_{\sigma} \int_{x \in S_{N(\sigma)}} \mathbb{1}_{\text{Log}^{-1}(\sigma^{\circ})} [S \cap \pi_{\text{aff}(\sigma)}^{-1}(x)] d\mu(x) + \mathcal{R}_{\mathcal{C}},$$

where $\mathcal{R}_{\mathcal{C}}$ is $(p-1, p-1)$ -dimensional current. We claim that $\mathcal{R}_{\mathcal{C}}$ is *normal*, i.e. $\mathcal{R}_{\mathcal{C}}$ and $d\mathcal{R}_{\mathcal{C}}$ have measure coefficients; $\mathcal{R}_{\mathcal{C}}$ is a difference of two normal currents, where the first current $[S] \wedge \mathcal{T}_{\mathcal{C}}$ is a positive closed current, and the second current is an addition of normal pieces. Moreover, the support of $\mathcal{R}_{\mathcal{C}}$ is a subset of S as it is a difference of two currents that both vanish outside S . As a result, the current $\mathcal{R}_{\mathcal{C}}$ is supported on $S \cap \bigcup_{\sigma} \partial \text{Log}(\sigma)$. This set is a real manifold of Cauchy–Riemann dimension less than $p-1$, therefore by Demailly’s first theorem of support the normal current $\mathcal{R}_{\mathcal{C}}$ vanishes; see also the discussion following [BH17, Proposition 4.11]. \square

Corollary 6.2. Let $H \subseteq \mathbb{R}^d$ be a rational plane of dimension $r \geq d-p$ and $A := a + H$, a translation of H for $a \in \mathbb{R}^d$. Assume also that $\mathcal{C} \subseteq \mathbb{R}^d$ is a tropical variety of dimension p that intersects A transversely. Then

$$[(e^{-a})T_{H \cap \mathbb{Z}^d}] \wedge \mathcal{T}_{\mathcal{C}}$$

can be viewed as a tropical current of dimension $p - (d-r)$ in the complex subtorus $T^A := (e^{-a})T_{H \cap \mathbb{Z}^d} \subseteq (\mathbb{C}^*)^d$.

Proof. Note that the hypothesis implies that the intersection $T^A \cap \pi_{\text{aff}(\sigma)}^{-1}(x)$ is transversal for any $x \in S_{N(\sigma)}$. By translation, it is sufficient to prove the statement for $a = 0$. By the preceding theorem,

$$[T^A] \wedge \mathcal{T}_{\mathcal{C}} = \sum_{\sigma \in \mathcal{C}} w_{\sigma} \int_{x \in S_{N(\sigma)}} \mathbb{1}_{\text{Log}^{-1}(\sigma^{\circ})} [T^A \cap \pi_{\text{aff}(\sigma)}^{-1}(x)] d\mu(x).$$

The sets $T^A \cap \pi_{\text{aff}(\sigma)}^{-1}(x)$ can be understood as the toric sets in T^A . We can conclude since each every $d\mu$ associated to each σ is a Haar measure. \square

Theorem 6.3. Let $M \subseteq (\mathbb{C}^*)^{d-p}$ and $N \subseteq (\mathbb{C}^*)^p$ be two bounded open subsets such that N contains the real torus $(S^1)^p$. Let $\pi : M \times N \rightarrow M$ be the canonical projection. Let \mathcal{T}_n be a sequence of positive closed (p, p) -bidimensional currents on $M \times N$ such that $\overline{\text{supp}(\mathcal{T}_n)} \cap (M \times \partial \bar{N}) = \emptyset$. Assume that $\mathcal{T}_n \rightarrow \mathcal{T}$ and $\text{supp}(\mathcal{T}) \subseteq M \times (S^1)^p$. Then we have the following convergence of slices

$$\langle \mathcal{T}_n | \pi | x \rangle \rightarrow \langle \mathcal{T} | \pi | x \rangle \quad \text{for every } x \in M.$$

Note that all the above slices are well-defined for all $x \in M$.

Proof. Since all the currents \mathcal{T}_n and \mathcal{T} are horizontal-like, the slices are well-defined, and we prove that the slices have the same cluster value. Let \mathcal{S} be any cluster value of $\langle \mathcal{T}_n | \pi | x \rangle$. Note that such \mathcal{S} always exists by Banach–Alaoglu theorem. As both measures \mathcal{S} and $\langle \mathcal{T} | \pi | x \rangle$, are supported $\{x\} \times (S^1)^p$ to prove their equality, it suffices to prove that they have the same Fourier coefficients. By Theorem 2.14, we have

$$\langle \mathcal{S}, \phi \rangle \leq \langle \mathcal{T} | \pi | x \rangle(\phi),$$

for every plurisubharmonic function ϕ on \mathbb{C}^d , and the mass of \mathcal{S} coincides with the mass of $\langle \mathcal{T} | \pi | x \rangle$. Now, note that if ϕ is pluriharmonic, then $-\phi$ and ϕ are plurisubharmonic. As a result,

$$\langle \mathcal{S}, \phi \rangle = \langle \mathcal{T} | \pi | x \rangle(\phi),$$

for every pluriharmonic function. Recall that if f is a holomorphic function, then $\operatorname{Re}(f)$ and $\operatorname{Im}(f)$ are pluriharmonic. We now consider the elements of the Fourier basis $f(\theta) = \exp 2\pi i \langle \nu, \theta \rangle$ for $\nu \in \mathbb{Z}^d$. Then we have the equality

$$\langle \mathcal{S}, f \rangle = \langle \mathcal{T} | \pi | x \rangle(f).$$

This implies that the Fourier measure coefficients of both \mathcal{S} and $\langle \mathcal{T} | \pi | x \rangle$ coincide. \square

Lemma 6.4. Let $\mathcal{C} \subseteq \mathbb{R}^d$ be a tropical variety of dimension p , and L be a rational $(d-p)$ -dimensional plane such that L is transversal to all the affine extensions $\operatorname{aff}(\sigma)$ for $\sigma \in \mathcal{C}$. Assume that \mathcal{T} is a positive closed current of bidimension (p, p) on a smooth projective toric variety X_Σ compatible with $\mathcal{C} + L$ such that $\operatorname{supp}(\mathcal{T}) \subseteq \operatorname{supp}(\mathcal{T}_\mathcal{C})$. Further, for all $a \in \mathbb{R}^d$,

$$\overline{\mathcal{T}}_{L+a} \wedge \mathcal{T} = \overline{\mathcal{T}}_{L+a} \wedge \overline{\mathcal{T}}_\mathcal{C}.$$

Then $\mathcal{T} = \mathcal{T}_\mathcal{C}$ in T_N .

Proof. Let us first remark that $\operatorname{rec}(L + a) = \operatorname{rec}(L)$ for all $a \in \mathbb{R}^d$ and therefore, all \mathcal{T}_{a+L} are compatible with X_Σ and have a continuous super-potential in X_Σ and as a result, all the above wedge products are well-defined. By Demailly's second theorem of support [Dem, III.2.13], there are measures $\mu_\sigma^\mathcal{T}$ such that

$$\mathcal{T} = \sum_\sigma \int_{x \in S(\mathbb{Z}^d \cap H_\sigma)} \mathbb{1}_{\operatorname{Log}^{-1}(\sigma^\circ)}[\pi_\sigma^{-1}(x)] d\mu_\sigma^\mathcal{T}(x).$$

By repeated application of Proposition 6.1,

$$\mathcal{T}_L \wedge \mathcal{T} = \sum_\sigma \int_{(x,y) \in S(\mathbb{Z}^d \cap H_L) \times S(\mathbb{Z}^d \cap H_\sigma)} [\pi_H^{-1}(x) \cap \pi_\sigma^{-1}(y)] d\mu_L(x) \otimes \mu_\sigma^\mathcal{T}(y).$$

Applying both sides of the equality $\mathcal{T}_L \wedge \mathcal{T} = \mathcal{T}_L \wedge \mathcal{T}_\mathcal{C}$ on test-functions of the form

$$\omega_\nu = \exp(-i \langle \nu, \theta \rangle) \rho(r)$$

where $\rho : \mathbb{R}^d \rightarrow \mathbb{R}$ is a smooth function with compact support of $r \in \mathbb{R}^d$ and $\theta \in [0, 2\pi)^d$, and $\nu \in \mathbb{Z}^d$, completely determines the Fourier coefficients of $\mu_\sigma^\mathcal{T}$ which have to coincide with the normalised Haar measures multiplied by the weight of σ , i.e., $\mu_\sigma^\mathcal{T} = w_\sigma \mu_\sigma$. \square

Note that any subtorus of $(\mathbb{C}^*)^d$, can be understood as a fibre of a tropical current. We have the following slicing theorem.

Proposition 6.5. Let $\mathcal{C} \subseteq \mathbb{R}^d$ be a tropical variety and $A \subseteq \mathbb{R}^d$ be a rational hyperplane intersecting \mathcal{C} transversely. Let Σ be a fan compatible with $\mathcal{C} + A$. Assume that $\overline{\mathcal{S}}_n$ is a sequence of positive closed currents on X_Σ , and denote by \mathcal{S}_n the restriction to T_N . Further, assume that

- (a) $\overline{\mathcal{S}}_n \rightarrow \overline{\mathcal{T}}_\mathcal{C}$;
- (b) $\operatorname{supp}(\overline{\mathcal{S}}_n) \rightarrow \operatorname{supp}(\overline{\mathcal{T}}_\mathcal{C})$,

then

$$\lim_{n \rightarrow \infty} (\mathcal{S}_n \wedge [T^A]) = \mathcal{T}_\mathcal{C} \wedge [T^A],$$

as currents on $T_N \subseteq X_\Sigma$.

Proof. Assume that $L \subseteq \mathbb{R}^d$ is an $(d - p - 1)$ -dimensional affine plane intersecting all $\text{aff}(\sigma)$ for all $\sigma \in \mathcal{C} \cap A$ transversely. Then, on a projective smooth toric variety $X_{\Sigma'}$ compatible with $\mathcal{C} + L + A$ the tropical currents $\bar{\mathcal{T}}_{a+L}$, $a \in \mathbb{R}^d$ have continuous super-potentials. Therefore, by Proposition 2.6, we have

$$\lim_{n \rightarrow \infty} (\bar{\mathcal{S}}_n \wedge \bar{\mathcal{T}}_{a+L}) = \bar{\mathcal{T}}_\mathcal{C} \wedge \bar{\mathcal{T}}_{a+L}.$$

Now, for any $x \in \mathcal{C} \cap L \cap A$, let $B \subseteq \mathbb{R}^d$ containing x be a bounded open set containing only x as an isolated point of the intersection. By a translation we can assume that $x = 0$. Consider the isomorphism

$$\xi : (\mathbb{C}^*)^d \xrightarrow{\sim} T_{\mathbb{Z}^d/(\mathbb{Z}^d \cap A)} \times T_{\mathbb{Z}^d \cap A},$$

and let π_1 and π_2 be the respective projections. Note that $\pi_1^{-1}(1) = T^A$. We now set

$$\begin{aligned} U &:= \pi_1 \circ \xi(\text{Log}^{-1}(U) \cap \text{supp}(\mathcal{T}_\mathcal{C} \wedge \mathcal{T}_{a+L})), \\ V &:= \pi_2 \circ \xi(\text{Log}^{-1}(U) \cap T^A), \\ \mathcal{T}_n &:= \xi_*(\mathcal{S}_n \wedge \mathcal{T}_{a+L}), \text{ in } T_N, \\ \mathcal{T} &:= \xi_*(\mathcal{T}_\mathcal{C} \wedge \mathcal{T}_{a+L}). \end{aligned}$$

Note that $\mathcal{T}_\mathcal{C}$ are horizontal-like as in the setting of Theorem 6.3. Assumption (b) now implies that \mathcal{T}_n for a large n , is also horizontal-like. Thus we obtain

$$\lim_{n \rightarrow \infty} (\mathcal{S}_n \wedge [T^A]) \wedge \mathcal{T}_{a+L} = \mathcal{T}_\mathcal{C} \wedge [T^A] \wedge \mathcal{T}_{a+L},$$

for every a . We now deduce the convergence on T_N by Lemma 6.4. \square

Theorem 6.6. Let $\mathcal{C} \subseteq \mathbb{R}^d$ be a tropical variety, and let $B = A_1 \cap \dots \cap A_k \subseteq \mathbb{R}^d$ be a complete intersection of rational hyperplanes A_1, \dots, A_k . Assume that \mathcal{C} and B intersect transversely. Let Σ be a smooth, projective fan compatible with $\mathcal{C} + B$, and let $(\bar{\mathcal{S}}_n)$ be a sequence of positive closed currents on X_Σ . Suppose that:

- (a) $\bar{\mathcal{S}}_n \rightarrow \bar{\mathcal{T}}_\mathcal{C}$;
- (b) $\text{supp}(\bar{\mathcal{S}}_n) \rightarrow \text{supp}(\bar{\mathcal{T}}_\mathcal{C})$.

Then, the following limit holds in the smooth projective toric variety X_Σ :

$$\lim_{n \rightarrow \infty} (\bar{\mathcal{S}}_n \wedge [\bar{T}^B]) = \bar{\mathcal{T}}_\mathcal{C} \wedge [\bar{T}^B].$$

We need a simple observation:

Lemma 6.7. Let $U \subseteq \mathbb{C}^d$ be an open subset and D be an analytic subset of \mathbb{C}^d . Assume that we have the convergence of closed positive currents $\mathcal{V}_n \rightarrow \mathcal{V}$ in $U \setminus D$, and \mathcal{V}_n 's and \mathcal{V} have a uniformly bounded local masses near D . Further, assume that for any cluster value \mathcal{W} of the sequence $\{\bar{\mathcal{V}}_n\}_n$, we have

- (a) $\text{supp}(\mathcal{W}) \subseteq \text{supp}(\bar{\mathcal{V}})$,
- (b) $\text{supp}(\bar{\mathcal{V}}) \cap D$ has the expected Cauchy–Riemann dimension.

Then,

$$\bar{\mathcal{V}}_n \longrightarrow \bar{\mathcal{V}}.$$

Proof. $\mathcal{W} - \bar{\mathcal{V}}$ is a positive closed current with the Cauchy–Riemann dimension less than or equal to p , therefore, it must be zero by Demailly’s first theorem of support [Dem, Theorem III.2.10]. \square

Proof of Theorem 6.6. Applying Theorem 5.2 (or [OP13, Proposition 3.3.2] to each fibre of $\bar{\mathcal{T}}_{\mathcal{C}}$ separately), we obtain $\text{supp}(\bar{\mathcal{T}}_{\mathcal{C}}) \cap \bar{T}^B \cap [D_{\rho}]$ has the expected Cauchy–Riemann dimension $p - k - 1$. By Demailly’s first theorem of support [Dem, Theorem III.2.10],

$$\bar{\mathcal{S}}_{\mathcal{C}} \wedge [\bar{T}^B] = \overline{\mathcal{T}_{\mathcal{C}} \wedge [T^B]}.$$

By assumption $\bar{\mathcal{S}}_n \longrightarrow \bar{\mathcal{T}}_{\mathcal{C}}$ and $\text{supp}(\mathcal{T}_n) \longrightarrow \text{supp}(\bar{\mathcal{T}}_{\mathcal{C}})$. The observation in Lemma 2.15,

$$\lim_{n \rightarrow \infty} \text{supp}(\bar{\mathcal{S}}_n \wedge [\bar{T}^B]) \subseteq \text{supp}(\bar{\mathcal{T}}_{\mathcal{C}} \wedge [\bar{T}^B]).$$

Therefore, any cluster value of $\bar{\mathcal{S}}_n \wedge [\bar{T}^B] \leq \bar{\mathcal{S}}_n \wedge [\bar{T}^B]$ has a support in $\text{supp}(\bar{\mathcal{T}}_{\mathcal{C}} \wedge [\bar{T}^B])$. Now, we set

- (a) $\mathcal{V}_n := \mathcal{S}_n \wedge [T^B]$,
- (b) $\mathcal{V} := \mathcal{T}_{\mathcal{C}} \wedge [T^B]$,
- (c) \mathcal{W} a cluster value of $\overline{\mathcal{T}_n \wedge [T^B]}$.

A repeated application of Proposition 6.5 for $B = A_1 \cap \cdots \cap A_k$, we are in the situation of Lemma 6.7, and conclude. \square

Lemma 6.8. Let X_{Σ} be a smooth projective toric variety, and $\bar{\Delta} \subseteq X_{\Sigma} \times X_{\Sigma}$ be the diagonal. Let \mathcal{S} and \mathcal{T} be two positive currents on X . Then, for any ray $\rho \in \Sigma$,

$$\text{supp}(\mathcal{S}) \cap \text{supp}(\mathcal{T}) \cap D_{\rho} \subseteq X_{\Sigma}$$

has a Cauchy–Riemann dimension ℓ , if and only if,

$$\text{supp}(\mathcal{S} \otimes \mathcal{T}) \cap \bar{\Delta} \cap D_{(0,\rho)} \subseteq X_{\Sigma} \times X_{\Sigma},$$

has a Cauchy–Riemann dimension ℓ , where $D_{(0,\rho)}$ is the toric invariant divisor corresponding to the ray $(0, \rho)$ in $\Sigma \times \Sigma$.

Proof. The fan of $X_{\Sigma} \times X_{\Sigma}$ is $\Sigma \times \Sigma$, we have that $D_{(0,\rho)} \simeq X_{\Sigma} \times D_{\rho}$ and the assertion follows. \square

Theorem 6.9. Let $\mathcal{C}_1, \mathcal{C}_2 \subseteq \mathbb{R}^d$ be two tropical cycles intersecting properly. Assume that X_{Σ} is a smooth toric projective variety compatible with $\mathcal{C}_1 + \mathcal{C}_2$. If moreover, for two sequence of positive closed currents $\bar{\mathcal{V}}_n$ and $\bar{\mathcal{W}}_n$ we have

- (a) $\bar{\mathcal{W}}_n \longrightarrow \bar{\mathcal{T}}_{\mathcal{C}_1}$, and $\bar{\mathcal{V}}_n \longrightarrow \bar{\mathcal{T}}_{\mathcal{C}_2}$;
- (b) $\text{supp}(\bar{\mathcal{W}}_n) \longrightarrow \text{supp}(\bar{\mathcal{T}}_{\mathcal{C}_1})$, and $\text{supp}(\bar{\mathcal{V}}_n) \longrightarrow \text{supp}(\bar{\mathcal{T}}_{\mathcal{C}_2})$;
- (c) For any large n , $\bar{\mathcal{W}}_n \wedge \bar{\mathcal{V}}_n$ is well-defined,

then

$$\bar{\mathcal{W}}_n \wedge \bar{\mathcal{V}}_n \longrightarrow \bar{\mathcal{T}}_{\mathcal{C}_1} \wedge \bar{\mathcal{T}}_{\mathcal{C}_2}, \quad \text{as } m \rightarrow \infty.$$

Proof. For two closed currents \mathcal{S} and \mathcal{T} on X_Σ we naturally identify

$$\mathcal{S} \wedge \mathcal{T} = \pi_*(\mathcal{S} \otimes \mathcal{T} \wedge [\bar{\Delta}]),$$

where $\pi : X_\Sigma \times X_\Sigma \rightarrow X_\Sigma$ is the projection onto the first factor. Let \mathcal{W}_n and \mathcal{V}_n be the restriction of $\bar{\mathcal{W}}_n$ and $\bar{\mathcal{V}}_n$ to $T_N \subseteq X_\Sigma$. We can define the tropical current $\mathcal{T}_\mathcal{C} := \mathcal{T}_{\mathcal{C}_1} \otimes \mathcal{T}_{\mathcal{C}_2}$. By Theorem 4.3, $\mathcal{C}_1 \times \mathcal{C}_2 := \mathcal{C}$ is a balanced tropical variety. Let us denote the diagonal $\Delta_\mathbb{R} \subseteq \mathbb{R}^d \times \mathbb{R}^n$. It is not hard to see that the transversality of \mathcal{C}_1 and \mathcal{C}_2 is equivalent to the transversality of the intersection

$$(\mathcal{C}_1 \times \mathcal{C}_2) \cap \Delta_\mathbb{R} \subseteq \mathbb{R}^d \times \mathbb{R}^d.$$

Now, fixing the coordinates $((z_1, \dots, z_d), (z'_1, \dots, z'_d)) \in (\mathbb{C}^*)^d \times (\mathbb{C}^*)^d$, the diagonal of $(\mathbb{C}^*)^d \times (\mathbb{C}^*)^d$ is given by the complete intersection of the tori $z_i = z'_i$, $i = 1, \dots, n$. Therefore, by refining $\Sigma \times \Sigma$ to a fan compatible also with $\Delta_\mathbb{R}$, and setting $B = \Delta$ in Proposition 6.5, we obtain

$$\mathcal{T}_n \wedge [\Delta] \rightarrow \mathcal{T}_\mathcal{C} \wedge [\Delta], \quad \text{i.e., } \mathcal{W}_n \wedge \mathcal{V}_n \rightarrow \mathcal{T}_{\mathcal{C}_1} \wedge \mathcal{T}_{\mathcal{C}_2},$$

in the open torus $T_N \times T_N$. Now, we can employ the compatibility of $\mathcal{C}_1 + \mathcal{C}_2$ and X_Σ together with Lemma 6.8, imply that for any ray $\rho \in \Sigma$,

$$\text{supp}(\bar{\mathcal{T}}_{\mathcal{C}_1} \otimes \bar{\mathcal{T}}_{\mathcal{C}_2}) \cap [\bar{\Delta}] \cap D_\rho$$

has the expected Cauchy–Riemann dimension, and finally conclude by Lemma 6.7. \square

7. DYNAMICAL TROPICALISATION AND INTERSECTIONS

7.1. Dynamical tropicalisation with a non-trivial valuation. Recall that for a field \mathbb{K} , $\nu : \mathbb{K} \rightarrow \mathbb{R} \cup \{\infty\}$, is called a valuation if it satisfies the following properties for every $a, b \in \mathbb{K}$:

- (a) $\nu(a) = \infty$ if and only if $a = 0$;
- (b) $\nu(ab) = \nu(a) + \nu(b)$;
- (c) $\nu(a + b) \geq \min\{\nu(a), \nu(b)\}$.

A valuation is called *trivial*, if the valuation of any non-zero element is 0. For an element $a \in \mathbb{K}$, we denote by \bar{a} its image in the residue field. We are interested in the case where $\mathbb{K} = \mathbb{C}((t))$, is the field of *formal Laurent series* with the variable t , with the usual valuation. That is, for $g(t) = \sum_{j \geq k} a_j t^j$, with $a_k \neq 0$, the valuation equals the minimal exponent $\nu(g) = k \in \mathbb{Z}$.

Definition 7.1. (a) Let $f = \sum_{\alpha \in A} c_\alpha z^\alpha \in \mathbb{K}[z^{\pm 1}]$, be a Laurent polynomial in d variables, where $A \subseteq \mathbb{Z}^d$ is a finite subset. The tropicalisation of f with respect to ν ,

$$\begin{aligned} \text{trop}_\nu(f) : \mathbb{R}^d &\rightarrow \mathbb{R}, \\ x &\mapsto \max\{-\nu(c_\alpha) + \langle x, \alpha \rangle\}. \end{aligned}$$

- (b) Let $I \subseteq \mathbb{K}[z^{\pm 1}]$ be an ideal. The tropical variety associated to I , as a set, is defined as

$$\text{Trop}_\nu(I) := \bigcap_{f \in I} \text{Trop}(\text{trop}_\nu(f)),$$

where $\text{Trop}(\text{trop}_\nu(f))$ is the set of points where $\text{trop}_\nu(f)$ is not differentiable; see Remark 4.5.

- (c) For an algebraic subvariety of the torus $Z \subseteq (\mathbb{K}^*)^d$, with the associated ideal $\mathbb{I}(Z)$, the tropicalisation of Z , as a set, is $\text{Trop}_\nu(Z) := \text{Trop}_\nu(\mathbb{I}(Z))$.
- (d) In all the situations above, trop_0 denotes the tropicalisation with respect to the trivial valuation.

We need to relate a non-trivial valuation to the trivial valuation. Compare to [BJS⁺07, Lemma 1.1].

Lemma 7.2. Consider the ideal $I \subseteq \mathbb{C}[t^{\pm 1}, z^{\pm 1}] \xhookrightarrow{\iota} \mathbb{C}((t))[z^{\pm 1}]$. Assume that (u, x) are the coordinates in $\mathbb{R} \times \mathbb{R}^d$. Then, we have the following equality of sets

$$\text{Trop}_0(I) \cap \{u = -1\} = \text{Trop}_\nu(\iota(I)),$$

where ν is the usual valuation in $\mathbb{C}((t))$. In other words, the tropicalisation of I as an ideal in $\mathbb{C}[t^{\pm 1}, z^{\pm 1}]$ with respect to the trivial valuation intersected with $\{u = -1\}$ coincides with the tropicalisation of $I = \iota(I)$ with respect to the usual valuation in $\mathbb{C}((t))$.

The proof of the lemma becomes clear with the following example.

Example 7.3. Let

$$f(x, t) = 4(t^3 + t^{-1})z_1z_2 + (1 + t + t^2)z_1.$$

Then, the tropicalisation of $f \in \mathbb{C}[t^{\pm 1}, z^{\pm 1}]$, with respect to the trivial valuation equals:

$$\text{trop}_0(f) = \max \left\{ \max\{3u + x_1 + x_2, -u + x_1 + x_2\}, \max\{x_1, u + x_1, 2u + x_1\} \right\}.$$

Letting $u := -1$, $\text{trop}_0(f)(-1, x) = \max\{1 + x_1 + x_2, x_1\}$. The latter equals $\text{trop}_\nu(f)$ as an element of $\mathbb{C}((t))[z^{\pm 1}]$.

Proof of Lemma 7.2. If f is a monomial in $\mathbb{C}[t][z]$, then it is clear that

$$\text{trop}_0(f)(-1, x) = \text{trop}_\nu(\iota(f)).$$

Therefore, we have the equality for any polynomial in $f \in \mathbb{C}[t, z]$. To prove the main statement, note that

$$\begin{aligned} \text{Trop}_\nu(\iota(I)) &= \bigcap_{f \in \iota(I)} \text{Trop}(\text{trop}_\nu(f)) \\ &= \bigcap_{f \in I} (\text{Trop}(\text{trop}_0(f)) \cap \{u = -1\}) \\ &= \text{Trop}_0(I) \cap \{u = -1\}. \end{aligned}$$

□

Remark 7.4. Bergman in [Ber71], shows that for an algebraic subvariety $Z \subseteq (\mathbb{C}^*)^d$, one has

$$\lim_{t \rightarrow \infty} \text{Log}_t(Z) \subseteq \text{Trop}_0(\mathbb{I}(Z)),$$

and he conjectured the equality. This conjecture was later proved by Bieri and Groves in [BG84]. More precisely, Bieri and Groves prove that $\lim \text{Log}_t(Z) \cap (S^1)^d$ is a polyhedral sphere of real dimension equal to (the complex dimension) $\dim(Z) - 1$. Therefore, the fan $\lim \text{Log}_t(Z)$ is a cone over their spherical complex. See also [MS15, Theorem 1.4.2].

Remark 7.5. The above lemma is related to the results of Markwig and Ren in [MR20]. They considered the tropicalisation of an ideal $J \subseteq R[[t]][z]$, where R is the ring of integers of a discrete valuation ring \mathbb{K} , which is non-trivially valued. To obtain finiteness properties, however, the authors consider the associated tropical variety in the half-space $\mathbb{R}_{\leq 0} \times \mathbb{R}^d$. Note that such a variety is almost never balanced. The authors also prove that for an ideal $I \subseteq \mathbb{K}[z]$, the tropicalisation of the natural inverse image $\pi^{-1}I \subseteq R[[t]][z]$ with respect to trivial valuation, intersected with $\{u = -1\}$ equals $\text{trop}_\nu(I)$; [MR20, Theorem 4].

Let us also recall the main result of [Bab23].

Theorem 7.6. Let $Z \subseteq (\mathbb{C}^*)^d$ be an irreducible subvariety of dimension p , and \overline{Z} be the closure of Z in a compatible smooth projective toric variety X_Σ . Define $\Phi_m : X_\Sigma \rightarrow X_\Sigma$ to be the unique continuous extension of

$$\begin{aligned} (\mathbb{C}^*)^d &\longrightarrow (\mathbb{C}^*)^d, \\ z &\longmapsto z^m. \end{aligned}$$

Then,

$$\frac{1}{m^{d-p}} \Phi_m^*[\overline{Z}] \longrightarrow \overline{\mathcal{T}}_{\text{Trop}_0(Z)}, \quad \text{as } m \rightarrow \infty,$$

where $\overline{\mathcal{T}}_{\text{Trop}_0(Z)}$ is the extension by zero of $\mathcal{T}_{\text{Trop}_0(Z)}$ to X_Σ . Moreover, the supports also converge in Hausdorff metric.

Note that since the limit of a sequence of closed currents is closed, the above theorem implies that $\text{trop}_0(Z)$ can be equipped with weights to become balanced. Note that the compatibility is in the following sense of Tevelev and Sturmfels:

- Theorem 7.7.** (a) The closure \overline{Z} of Z in X_Σ is complete, if and only if, $\text{Trop}_0(Z) \subseteq |\Sigma|$; see [Tev07].
 (b) We have $|\Sigma| = \text{Trop}_0(Z)$, if and only if, for every $\sigma \in \Sigma$ the intersection $\mathcal{O}_\sigma \cap \overline{Z}$ is non-empty and of pure dimension $p - \dim(\sigma)$; see [ST08].

Theorem 7.8. Let $I \subseteq \mathbb{C}[t^{\pm 1}, z^{\pm 1}]$ be an ideal with the associated $(p+1)$ -dimensional algebraic variety $W = \mathbb{V}(I) \subseteq (\mathbb{C}^*)^{d+1}$. Assume that the projection onto the first coordinate $\pi_1 : W \rightarrow \mathbb{C}^*$ is a surjective and Zariski-closed map. Further, $\pi_1(W_{\text{sing}}) \subsetneq \mathbb{C}^*$, the projection of singular points of W is a proper subset of \mathbb{C}^* . If we denote the fibers by $W_t := \pi_1^{-1}(t)$, then

- (a) $\frac{1}{m^{d-p}} \Phi_m^*[W_{e^m}] \rightarrow \mathcal{T}_{\text{Trop}_\nu(I)}$, as $m \rightarrow \infty$, in the sense of currents in $\mathcal{D}_p((\mathbb{C}^*)^d)$, and we have identified $\iota(I) = I$.
- (b) $\text{Trop}_\nu(I)$ can be equipped with weights to become balanced.
- (c) $\limsup(\frac{1}{m^{d-p}} \Phi_m^*[W_{e^m}]) = \text{supp}(\mathcal{T}_{\text{Trop}_\nu(I)})$.
- (d) On a toric variety X_Σ compatible with $\text{trop}_0(W) + \{u = -1\}$,

$$\frac{1}{m^{d-p}} \Phi_m^*[\overline{W}_{e^m}] \longrightarrow \overline{\mathcal{T}}_{\text{Trop}_\nu(I)}, \quad \text{as } m \rightarrow \infty.$$

For the proof, we need the following:

Lemma 7.9. Let $W \subseteq (\mathbb{C}^*)^{d+1}$ be a $(p+1)$ -dimensional subvariety, such that the projection onto the first factor, $\pi_1 : (\mathbb{C}^*)^{d+1} \rightarrow \mathbb{C}^*$ is surjective and a Zariski closed

morphism, and the projection of the singular points are a proper subset: $\pi_1(W_{\text{sing}}) \subsetneq \mathbb{C}^*$. Then, for a sufficiently large $|t_0| \gg 0$

$$[W_{t_0}] = [\pi_1^{-1}(t_0)] = [\{t = t_0\}] \wedge [W].$$

Proof. Let us first fix an ideal associated to $I = \mathbb{I}(W) = \langle f_1, \dots, f_k \rangle \subseteq \mathbb{C}[t^{\pm 1}, z^{\pm 1}]$. At any regular point $w \in W_{\text{reg}}$, $T_w W = \ker J(f)(w) = \left(\frac{\partial f_i}{\partial z_j}(w) \right)_{k \times (d+1)}$ is of dimension $p+1$. We consider the *critical set*

$$C = \{w \in W : \dim \ker \begin{pmatrix} \nabla_w t \\ Jf(w) \end{pmatrix} \geq p+1\},$$

where $\nabla_w t$ is the gradient of t at w . We have that $\nabla_w t = e_1$, the first element of the standard basis of the \mathbb{C} -vector space \mathbb{C}^{d+1} . As a result, to have the rank of $\begin{pmatrix} e_1 \\ Jf(w) \end{pmatrix}$ less than or equal to $(d+1) - (p-1)$ we must have all the minors of size $(d-p+1) \times (d-p+1)$ equal to zero, which are the defining equations for C . The set Zariski-closed set C contains all the points such that either

- $w \in W$ is regular and the intersection $W \cap \pi_1^{-1}(\pi_1(w))$ is not transversal. Note that in this case, $\text{Image}(Jf(w)) \subseteq \ker(e_1)$.
- $w \in W$ is singular, and therefore $\ker(Jf(w)) > p+1$.

Since π_1 is a closed map, if we observe that $\pi_1(C) \subsetneq \mathbb{C}^*$, then $\pi_1(C)$ is a finite set and we conclude. Note that $\pi_1(W_{\text{reg}} \cap C) = \mathbb{C}^*$, implies that W is contained in $\{t = t_0\}$ for some $t_0 \in \mathbb{C}^*$, which is not the case as $\pi_1 : W \rightarrow \mathbb{C}^*$ is surjective. Further, $\pi(W_{\text{sing}}) \subsetneq \mathbb{C}^*$ by assumption of the lemma, and we conclude. \square

Proof of Theorem 7.8. By the preceding lemma, and the fact that Φ_m preserves transversal intersection, we have

$$\frac{1}{m^{d-p}} \Phi_m^*[W_{e^m}] = \frac{1}{m^{d-(p+1)}} \Phi_m^*[W] \wedge \frac{1}{m} \Phi_m^*[\{t = e^m\}],$$

for a large m . Since $\text{Trop}_0(W)$ is a fan, it is transversal to the plane $\{u = -1\} \subset \mathbb{R}^{d+1}$. In consequence, we can use Proposition 6.5 to write

$$\lim_{m \rightarrow \infty} \frac{1}{m^{d-p}} \Phi_m^*[W_{e^m}] = \left(\lim_{m \rightarrow \infty} \frac{1}{m^{d-(p+1)}} \Phi_m^*[W] \right) \wedge \left(\lim_{m \rightarrow \infty} \frac{1}{m} \Phi_m^*[\{t = e^m\}] \right)$$

By Theorem 7.6, restricted to $(\mathbb{C}^*)^{n+1}$, and the fact that we used $\text{Log} = (-\log |\cdot|, \dots, -\log |\cdot|)$ in the definition of tropical currents, the above limits yield

$$\lim_{m \rightarrow \infty} \frac{1}{m^{d-p}} \Phi_m^*[W_{e^m}] = \mathcal{T}_{\text{Trop}_0(W)} \wedge \mathcal{T}_{\{u = -1\}}.$$

Applying Theorem 5.13 and Lemma 7.2, we obtain the equality.

For the assertion (b), note that the limit $\mathcal{T}_{\text{Trop}_\nu(I)}$ is a closed current and Theorem 4.3 implies that $\text{Trop}_\nu(I)$ is naturally balanced. To observe (c), note that (a) implies

$$\limsup \left(\frac{1}{m^{d-p}} \Phi_m^*[W_{e^m}] \right) \supseteq \text{supp}(\mathcal{T}_{\text{Trop}_\nu(I)}).$$

However, because of transversality, $\text{supp}(\mathcal{T}_{\text{Trop}_\nu(I)}) = \text{supp}(\mathcal{T}_{\text{Trop}_0(W)}) \cap \text{supp}(\mathcal{T}_{\{u = -1\}})$. At the same time,

$$\limsup(\Phi_m^*[W_{e^m}]) = \limsup(\Phi_m^*[W]) \cap \text{supp}(\Phi_m^*[\{t = e^m\}]).$$

Moreover, for the Hausdorff limit of sets $\lim(A_i \cap B_i) \subseteq (\lim A_i) \cap (\lim B_i)$. This implies

$$\lim \operatorname{supp}(\Phi_m^*[W_{e^m}]) \subseteq \operatorname{supp}(\mathcal{T}_{\operatorname{Trop}_0(W)}) \cap \operatorname{supp}(\mathcal{T}_{\{u=-1\}}),$$

which implies (c). Now, (d) is implied by Theorem 6.6. \square

In the setting of the previous theorem, a generalisation of Bergman's theorem (see Remark 7.4) asserts that

$$\operatorname{Log}_t(W_t) \longrightarrow \operatorname{Trop}(I), \quad \text{as } t \longrightarrow \infty,$$

where Log_t is the logarithm with base t . This theorem can be understood as a counterpart of Lemma 7.2 for tropicalisation with Log . This generalisation was finally proved by Jonsson in [Jon16], and we can now deduce a sequential analogue:

Corollary 7.10. In the setting of the previous theorem

$$\frac{1}{m} \operatorname{Log}(W_{e^m}) \longrightarrow \operatorname{Trop}_\nu(I), \quad \text{as } m \rightarrow \infty,$$

in the Hausdorff metric in compact subsets of \mathbb{R}^d .

Proof. Note that for any variety $Z \subseteq (\mathbb{C}^*)^{d+1}$, as a set $\operatorname{Log}(\Phi_m^{-1}(Z)) = \frac{1}{m} \operatorname{Log}(Z)$. Therefore,

$$\operatorname{Log} \operatorname{supp}\left(\frac{1}{m^{d-p}} \Phi_m^*[W_{e^m}]\right) = \frac{1}{m} \operatorname{Log}(W_{e^m}).$$

The assertion now follows from Theorem 7.8(c) and continuity of Log with respect to the Hausdorff metric on compact sets. \square

Let us first prove the analogous to [BJS⁺07, Lemma 3.2] and [OP13, Theorem 1.2].

Theorem 7.11. Assume that W and V are two algebraic subvarieties of $(\mathbb{C}^*)^d$ with respective dimensions p and q , with $p + q \geq d$. Assume that $\operatorname{Trop}_0(V)$ and $\operatorname{Trop}_0(W)$ intersect transversely, then

$$\frac{1}{m^{2d-(p+q)}} \Phi_m^*([W] \wedge [V]) \longrightarrow \mathcal{T}_{\operatorname{Trop}_0(W) \cdot \operatorname{Trop}_0(V)}, \quad \text{as } m \rightarrow \infty.$$

Moreover, $\mathcal{T}_{\operatorname{Trop}_0(W \cap V)} \leq \mathcal{T}_{\operatorname{Trop}_0(W) \cdot \operatorname{Trop}_0(V)}$. In particular, the corresponding induced weights for $\tau \in \operatorname{Trop}_0(W \cap V)$, is less than or equal to the weight of τ induced by $\operatorname{Trop}_0(W) \cdot \operatorname{Trop}_0(V)$.

Example 7.12. To see that the inequality in the previous theorem can be strict, let us consider the subvarieties of $(\mathbb{C}^*)^2$, $W = \mathbb{V}(z_2 - 1)$ and $V = \mathbb{V}(z_2 - z_1^2 - 1)$. We have that

$$\begin{aligned} \operatorname{Trop}_0(W) &= \operatorname{Trop}(\max\{x_2, 2x_1, 0\}), \\ \operatorname{Trop}_0(V) &= \operatorname{Trop}(\max\{x_2, 0\}). \end{aligned}$$

In Example 3.6, we discussed that the stable intersection of these two cycles is the origin $(0, 0) \in \mathbb{R}^2$ with multiplicity 2. Note that the (set-theoretic) intersection $W \cap V = \{(0, 1)\}$, thus $\operatorname{Trop}_0(W \cap V) = (0, 0)$ with multiplicity 1, whereas $\operatorname{Trop}_0(W) \cdot \operatorname{Trop}_0(V) = (0, 0)$ with multiplicity 2. We obtain

$$2 \mathcal{T}_{\operatorname{Trop}_0(W \cap V)} = \mathcal{T}_{\operatorname{Trop}_0(W) \cdot \operatorname{Trop}_0(V)}.$$

Note that this inequality, gladly, does not contradict [OP13, 3.3.1] (a consequence of [Ful98, 8.2]), which the scheme-theoretic intersection $W \cap V$ is considered.

Proof of Theorem 7.11. Assume that X_Σ is a toric variety compatible with $\text{Trop}_0(W) + \text{Trop}_0(V)$. We need to show that the hypothesis (a), ..., (d) of Theorem 6.9 for $\overline{W}_m = m^{p-n}\Phi_m^*[\overline{W}]$ and $\overline{V}_m = m^{q-n}\Phi_m^*[\overline{V}]$ is satisfied. Note that hypotheses (a) and (b) are implied by Theorem 7.6. The hypotheses (c) and (d) for \overline{W}_1 and \overline{V}_1 are a result of [OP13, Proposition 3.3.2]. The hypotheses (c) and (d) for any m are implied by the fact that $\Phi_m^*([\overline{W}] \wedge [\overline{V}]) = \Phi_m^*[\overline{W}] \wedge \Phi_m^*[\overline{V}]$. Therefore,

$$\overline{W}_m \wedge \overline{V}_m \longrightarrow \overline{\mathcal{T}}_{\text{Trop}_0(W)} \wedge \overline{\mathcal{T}}_{\text{Trop}_0(V)}, \quad \text{as } m \rightarrow \infty,$$

and the first assertion is obtained by restricting to $T_N \subseteq X_\Sigma$ and Theorem 5.13. To see the second assertion, note that by Theorem 7.6,

$$\frac{1}{m^{2d-(p+q)}} \Phi_m^*[W \cap V] \longrightarrow \mathcal{T}_{\text{Trop}_0(W \cap V)}, \quad \text{as } m \rightarrow \infty.$$

However, as currents $[W \cap V] \leq [W] \wedge [V]$ since the right-hand side might produce multiplicities on the intersection. Applying Φ_m^* to both sides preserves this inequality. Therefore, for every m ,

$$\Phi_m^*[W \cap V] \leq \Phi_m^*[W] \wedge \Phi_m^*[V].$$

We can conclude by taking the limit $m \rightarrow \infty$. □

The tropical version of the following was observed in various places [OP13, MS15], which does not need to assume the proper intersection of tropicalisations.

Theorem 7.13. Let W and V be two smooth algebraic subvarieties of $(\mathbb{C}^*)^d$, and for $0 < \epsilon < 1$, let $U_\epsilon((S^1)^d)$ be an ϵ -neighbourhood of $(S^1)^d$. Then,

$$\frac{1}{m^{2d-(p+q)}} \int_{(t_1, t_2) \in U_\epsilon((S^1)^{2d})} \Phi_m^*([t_1 V] \wedge [t_2 W]) d\nu(t_1) \otimes d\nu(t_2) \longrightarrow \mathcal{T}_{\text{Trop}_0(V)} \wedge \mathcal{T}_{\text{Trop}_0(W)},$$

as $m \rightarrow \infty$. Here, the $d\nu$ is the normalised Lebesgue measures on $U_\epsilon((S^1)^d)$. Moreover, when V and W are smooth the integrand in the left-hand-side can be replaced by $\Phi_m^*[t_1 V \cap t_2 W]$.

Proof. By a classical result of Kleiman [Kle74], the intersection of generic translations $t_1 V \cap t_2 W$ is proper and transversal in case V and W are smooth. Since Φ_m^* preserves proper intersections and transversality, we can separate the above integrand into $\Phi_m^*[t_1 V] \wedge \Phi_m^*[t_2 W]$. Using polar coordinates (r_i, θ_i) for t_i , we have

$$\int_{t_1 \in U_\epsilon((S^1)^d)} \Phi_m^*[t_1 V] d\nu(t_1) = \int_{t_1 \in U_\epsilon((S^1)^d)} t_1^{1/m} \Phi_m^*[V] d\nu(t_1).$$

Here, with abuse of notation, we choose the m -th root $t_1^{1/m}$ to be the first root of $t_1^{1/m}$. We can also obtain a similar equation for $t_2 W$. Further,

$$(\Phi_m^*[t_1 V] \otimes \Phi_m^*[t_2 W]) \wedge [\Delta] = (\Phi_m^*[V] \otimes \Phi_m^*[W]) \wedge (t_1)^{-1/m} (t_2)^{-1/m} [\Delta].$$

$$\int_{U_\epsilon((S^1)^{2d})} (t_1)^{-1/m} (t_2)^{-1/m} [\Delta] = \int_{[(1-\epsilon), 1+\epsilon]} \mathcal{T}_\Delta^{R_m} dR,$$

where $R_m = (|t_1|^{-1/m}, |t_2|^{-1/m})$, and $\mathcal{T}_\Delta^{R_m}$ is the tropical current associated with the diagonal $\Delta \subseteq \mathbb{R}^d \times \mathbb{R}^d$, where the compact torus is rescaled to $R_m(S^1)^{2d}$. In a toric

variety compatible with $\Delta \subseteq \mathbb{R}^d \times \mathbb{R}^d$, for any R , $\overline{\mathcal{T}_{\Delta}^{R_m}}$ has a continuous superpotential, and $\overline{\mathcal{T}_{\Delta}^{R_m}}$ is SP-convergent to $\overline{\mathcal{T}_{\Delta}}$. Using Proposition 2.6, Theorem 2.10, and restricting yields:

$$\frac{1}{m^{2d-(p+q)}} (\Phi_m^*[t_1 V] \otimes \Phi_m^*[t_2 W]) \wedge \int_{[(1-\epsilon), 1+\epsilon]} \mathcal{T}_{\Delta}^{R_m} dR \longrightarrow (\mathcal{T}_{\text{Trop}_0(V)} \otimes \mathcal{T}_{\text{Trop}_0(W)}) \wedge \mathcal{T}_{\Delta},$$

as $m \rightarrow \infty$. The latter equals the tropical current

$$\mathcal{T}_{(\text{Trop}_0(V) \times \text{Trop}_0(W)) \cdot \Delta},$$

which can be identified with $\mathcal{T}_{\text{Trop}_0(V)} \wedge \mathcal{T}_{\text{Trop}_0(W)}$. \square

REFERENCES

- [AESK⁺21] Mats Andersson, Dennis Eriksson, Håkan Samuelsson Kalm, Elizabeth Wulcan, and Alain Yger, *Global representation of Segre numbers by Monge-Ampère products*, Math. Ann. **380** (2021), no. 1-2, 349–391. MR4263687
- [AR10] Lars Allermann and Johannes Rau, *First steps in tropical intersection theory*, Math. Z. **264** (2010), no. 3, 633–670.
- [ASKW22] Mats Andersson, Håkan Samuelsson Kalm, and Elizabeth Wulcan, *On non-proper intersections and local intersection numbers*, Math. Z. **300** (2022), no. 1, 1019–1039. MR4359551
- [Bab14] Farhad Babaee, *Complex tropical currents*, Ph.D. Thesis, 2014.
- [Bab23] Farhad Babaee, *Dynamical tropicalisation*, J. Geom. Anal. **33** (2023), no. 3, Paper No. 74, 38. MR4531051
- [BD20] François Berteloot and Tien-Cuong Dinh, *The Mandelbrot set is the shadow of a Julia set*, Discrete Contin. Dyn. Syst. **40** (2020), no. 12, 6611–6633. MR4160083
- [BDR24] Fabrizio Bianchi, Tien-Cuong Dinh, and Karim Rakhimov, *Monotonicity of dynamical degrees for Hénon-like and polynomial-like maps*, Trans. Amer. Math. Soc. **377** (2024), no. 9, 6595–6618. MR4855319
- [Ber71] George M. Bergman, *The logarithmic limit-set of an algebraic variety*, Trans. Amer. Math. Soc. **157** (1971), 459–469. MR280489
- [BG84] Robert Bieri and J. R. J. Groves, *The geometry of the set of characters induced by valuations*, J. Reine Angew. Math. **347** (1984), 168–195. MR733052 (86c:14001)
- [BGPS14] José Ignacio Burgos Gil, Patrice Philippon, and Martín Sombra, *Arithmetic geometry of toric varieties. Metrics, measures and heights*, Astérisque **360** (2014), vi+222. MR3222615
- [BH17] Farhad Babaee and June Huh, *A tropical approach to a generalized Hodge conjecture for positive currents*, Duke Math. J. **166** (2017), no. 14, 2749–2813. MR3707289
- [BJS⁺07] T. Bogart, A. N. Jensen, D. Speyer, B. Sturmfels, and R. R. Thomas, *Computing tropical varieties*, J. Symbolic Comput. **42** (2007), no. 1-2, 54–73. MR2284285
- [BS11] José Ignacio Burgos Gil and Martín Sombra, *When do the recession cones of a polyhedral complex form a fan?*, Discrete Comput. Geom. **46** (2011), no. 4, 789–798. MR2846179 (2012j:14070)
- [BS14] Erwan Brugallé and Kristin M. Shaw, *A bit of tropical geometry*, 2014. arXiv:1311.2360v3.
- [BT82] Eric Bedford and B. A. Taylor, *A new capacity for plurisubharmonic functions*, Acta Math. **149** (1982), no. 1-2, 1–40. MR674165
- [CLS11] David A. Cox, John B. Little, and Henry K. Schenck, *Toric varieties*, Graduate Studies in Mathematics, vol. 124, American Mathematical Society, Providence, RI, 2011. MR2810322 (2012g:14094)
- [Dem12] Jean-Pierre Demailly, *Analytic methods in algebraic geometry*, Surveys of Modern Mathematics, vol. 1, International Press, Somerville, MA; Higher Education Press, Beijing, 2012. MR2978333
- [Dem92] Jean-Pierre Demailly, *Courants positifs et théorie de l’intersection*, Gaz. Math. **53** (1992), 131–159.

- [Dem] Jean-Pierre Demailly, *Complex analytic and differential geometry*, Open Content Book, Version of June 21, 2012. <https://www-fourier.ujf-grenoble.fr/~demailly/manuscripts/agbook.pdf>.
- [DNV18] Tien-Cuong Dinh, Viêt-Anh Nguyễn, and Duc-Viet Vu, *Super-potentials, densities of currents and number of periodic points for holomorphic maps*, Adv. Math. **331** (2018), 874–907. MR3804691
- [DR25] Jeffrey Diller and Roland Roeder, *Equidistribution without stability for toric surface maps*, Commentarii Mathematici Helvetici (2025).
- [DS04] Tien-Cuong Dinh and Nessim Sibony, *Regularization of currents and entropy*, Ann. Sci. École Norm. Sup. (4) **37** (2004), no. 6, 959–971. MR2119243
- [DS09] Tien-Cuong Dinh and Nessim Sibony, *Super-potentials of positive closed currents, intersection theory and dynamics*, Acta Math. **203** (2009), no. 1, 1–82. MR2545825 (2011b:32052)
- [DS10] Tien-Cuong Dinh and Nessim Sibony, *Super-potentials for currents on compact Kähler manifolds and dynamics of automorphisms*, J. Algebraic Geom. **19** (2010), no. 3, 473–529. MR2629598
- [Fed69] Herbert Federer, *Geometric measure theory*, Die Grundlehren der mathematischen Wissenschaften, Band 153, Springer-Verlag New York Inc., New York, 1969. MR0257325
- [FS95] John Erik Fornaess and Nessim Sibony, *Oka’s inequality for currents and applications*, Math. Ann. **301** (1995), no. 3, 399–419.
- [FS97] William Fulton and Bernd Sturmfels, *Intersection theory on toric varieties*, Topology **36** (1997), no. 2, 335–353. MR1415592 (97h:14070)
- [Ful98] William Fulton, *Intersection theory*, Second, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], vol. 2, Springer-Verlag, Berlin, 1998. MR1644323
- [Gub16] Walter Gubler, *Forms and current on the analytification of an algebraic variety (after Chambert-Loir and Ducros)*, Nonarchimedean and tropical geometry, 2016, pp. 1–30. MR3700066
- [Jon16] Mattias Jonsson, *Degenerations of amoebae and berkovich spaces*, Mathematische Annalen **364** (2016), no. 1-2, 293–311.
- [JY16] Anders Jensen and Josephine Yu, *Stable intersections of tropical varieties*, J. Algebraic Combin. **43** (2016), no. 1, 101–128. MR3439302
- [Kat09] Eric Katz, *A tropical toolkit*, Expo. Math. **27** (2009), no. 1, 1–36. MR2503041
- [Kaz03] B. Ya. Kazarnovskiĭ, *c-fans and Newton polyhedra of algebraic varieties*, Izv. Ross. Akad. Nauk Ser. Mat. **67** (2003), no. 3, 23–44. MR1992192
- [Kle74] Steven L. Kleiman, *The transversality of a general translate*, Compositio Math. **28** (1974), 287–297. MR360616
- [Lag12] Aron Lagerberg, *Super currents and tropical geometry*, Math. Z. **270** (2012), no. 3-4, 1011–1050. MR2892935
- [McM89] Peter McMullen, *The polytope algebra*, Adv. Math. **78** (1989), no. 1, 76–130. MR1021549
- [Mih21] Andreas Mihatsch, *On tropical intersection theory*, arXiv preprint arXiv:2107.12067 (2021).
- [Mik06] Grigory Mikhalkin, *Tropical geometry and its applications*, International Congress of Mathematicians. Vol. II, 2006, pp. 827–852. MR2275625 (2008c:14077)
- [MR20] Thomas Markwig and Yue Ren, *Computing tropical varieties over fields with valuation*, Found. Comput. Math. **20** (2020), no. 4, 783–800. MR4130539
- [MS15] Diane Maclagan and Bernd Sturmfels, *Introduction to tropical geometry*, Graduate Studies in Mathematics, vol. 161, American Mathematical Society, Providence, RI, 2015. MR3287221
- [OP13] Brian Osserman and Sam Payne, *Lifting tropical intersections*, Doc. Math. **18** (2013), 121–175. MR3064984
- [Ras01] Alexander Rashkovskii, *Indicators for plurisubharmonic functions of logarithmic growth*, Indiana Univ. Math. J. **50** (2001), no. 3, 1433–1446. MR1871763 (2002i:32027)

- [RGST05] Jürgen Richter-Gebert, Bernd Sturmfels, and Thorsten Theobald, *First steps in tropical geometry*, Idempotent mathematics and mathematical physics, 2005, pp. 289–317. MR2149011 (2006d:14073)
- [Sha13] Kristin M. Shaw, *A tropical intersection product in matroidal fans*, SIAM J. Discrete Math. **27** (2013), no. 1, 459–491. MR3032930
- [ST08] Bernd Sturmfels and Jenia Tevelev, *Elimination theory for tropical varieties*, Math. Res. Lett. **15** (2008), no. 3, 543–562. MR2407231
- [Tev07] Jenia Tevelev, *Compactifications of subvarieties of tori*, Amer. J. Math. **129** (2007), no. 4, 1087–1104. MR2343384
- [Yge13] Alain Yger, *Tropical geometry and amoebas*, 2013. Lecture notes, <http://cel.archives-ouvertes.fr/cel-00728880>.

SCHOOL OF MATHEMATICS, UNIVERSITY OF BRISTOL

Email address: farhad.babae@bristol.ac.uk

DEPARTMENT OF MATHEMATICS, NATIONAL UNIVERSITY OF SINGAPORE

Email address: matdte@nus.edu.sg