Contents

1	Convert Binary to Decimal
2	Prime Number Pyramid
3	Print 1-10 Numbers in Pyramid fashion
4	Right angled triangle using *
5	Sum of an AP series
6	Sum of a GP series
7	HCF or GCD of two numbers
8	LCM of two numbers
9	Display the following pyramids

1 Convert Binary to Decimal

Accept Number from User Divide Number by 10 and Store Remainder in variable rem Divide Original Number by 10. sum = sum + rem * pow(2,power);

/*pow is a function to find the power of a number.

Inside the First Iteration power = 0. Power is Incremented in each Iteration.

Calculate sum using the above formula, calculated sum is nothing but the decimal representation of the given binary number.

2 Prime Number Pyramid

3 Print 1-10 Numbers in Pyramid fashion

4 Right angled triangle using *

5 Sum of an AP series

Sample output:

```
Enter the first number of the A.P. series: 1
Enter the total numbers in the A.P. series: 5
Enter the common difference of A.P. series: 3
Sum of the series: 1 + 4 + 7 + 10 + 13 = 35
```

Type man pow on terminal for help*/

Definition of arithmetic progression (A.P.):

A series of numbers in which difference of any two consecutive numbers is always a same number that is constant. This constant is called as common difference.

Example of A.P. series: 5 10 15 20 25

Here common difference is 5 since difference of any two consecutive numbers for example 20 15 or 25-20 is 5.

Sum of A.P. series: Sn = n/2(2a + (n-1) d)

Tn term of A.P. series:

Tn = a + (n-1) d

6 Sum of a GP series

Sample output:

Enter the first number of the G.P. series: 1 Enter the total numbers in the G.P. series: 5 Enter the common ratio of G.P. series: 2 tn term of G.P.: 16.000000 Sum of the G.P.: 63.000000

Definition of geometric progression (G.P.):

A series of numbers in which ratio of any two consecutive numbers is always a same number that is constant. This constant is called as common ratio.

Example of G.P. series:

2 4 8 16 32 64

Here common difference is 2 since ratio any two consecutive numbers for example 32 / 16 or 64/32 is 2.

Sum of G.P. series: $Sn = a(1-(r^(n+1))/(1-r))$

Tn term of G.P. series:

 $\operatorname{Tn} = \operatorname{a}(\operatorname{r}(n-1))$

Sum of infinite G.P. series:

$$Sn = a/(1-r)$$
 if $1 > r$
= $a/(r-1)$ if $r > 1$

7 HCF or GCD of two numbers

This is the best way to find HCF of two numbers. In this method, smaller number is subtracted from larger number and that number is stored in place of larger number. This process is continued **until**, two numbers become equal which will be HCF.

or

The smallest of two integers entered by user is stored in variable min. Then i is initialized to min and for loop is executed. In each looping iteration, whether i is factor of these two numbers is checked. If i is a factor of these two numbers then, i will be the highest common divisor and loop is terminated using break statement.

8 LCM of two numbers

LCM=(num1*num2)/GCD

or

The largest of two integers is assigned to variable max. Then, while loop is executed and in each iteration it is checked whether max is perfectly divisible by two numbers entered by user or not. If max is not perfectly divisible, max is increased by 1 and this process goes not until max is perfectly divisible by both numbers.

9 Display the following pyramids

```
2)
Α
ВВ
C C C
D\ D\ D\ D
E E E E E
3)
4)
1 2 3 4 5
1 2 3 4
1 2 3
1 2
1
5)
6)
         1
       2 3 2
     3\ 4\ 5\ 4\ 3
  4 \ 5 \ 6 \ 7 \ 6 \ 5 \ 4
5 6 7 8 9 8 7 6 5
```

8)

1 1 1
1 1 1
1 3 3 1 1
1 4 6 4 1
1 5 10 10 5 1