Ref. [32] — Cubic Regularization of Newton Method

Yu. Nesterov and B. T. Polyak | Mathematical Programming 108 (2006)

Source blocker. The original full article/proofs were not available for full inspection in this session. Bibliography, abstract-level contribution and the canonical cubic model are source-backed; the explanatory derivations below are independently checked mathematical background, *not* a reconstruction of the paper's exact proof or all its constants. Do not mark the complete paper verified.

1. Why ordinary Newton steps are troublesome at saddles

Let $g = \nabla f(x)$ and $H = \nabla^2 f(x)$. Newton's second-order model is

$$m_N(s) = f(x) + g^T s + \frac{1}{2} s^T H s.$$

If H is indefinite, then $s^T H s$ can be negative along a direction with negative eigenvalue, making the unconstrained quadratic model unbounded below. A standard Newton step $-H^{-1}g$ is therefore not automatically a descent or saddle-escape method.

2. The proposed mechanism: regularize by a cubic norm

Under a Hessian-Lipschitz condition

$$\|\nabla^2 f(x+s) - \nabla^2 f(x)\| \le M\|s\|,$$

Taylor's theorem bounds the remainder by $M\|s\|^3/6$. This suggests the *cubic model*

$$\boxed{m_x(s) = f(x) + g^T s + \tfrac{1}{2} s^T H s + \tfrac{M}{6}\|s\|^3.}$$

Unlike an indefinite quadratic, the cubic term grows positively as $\|s\| \to \infty$; the model has a global minimizer. A globally minimizing step naturally takes account of negative curvature rather than treating it as a numerical error.

3. From the model to an update

For $s \neq 0$, the first-order condition for the model is

$$0 = \nabla_s m_x(s) = g + Hs + \frac{M}{2}\|s\| s.$$

Thus the step solves a shifted Newton-like system

$$\left(H + \frac{M\|s\|}{2} I\right) s = -g.$$

The shift depends on the resulting step length. This is why the step is *not* simply Newton's update with a fixed regularizer. Finding a global solution of the subproblem also requires more than merely solving the first-order equation.

Documented result. The original article establishes global worst-case complexity results, across classes including non-convex objectives, and studies local convergence of cubic-regularized Newton steps. Standard modern interpretations connect this framework to approximate second-order stationarity, with rates depending on the exact tolerance definition and oracle model.

4. What can and cannot be concluded

The core analytical premise is regularity of the Hessian and the ability to solve or accurately approximate cubic subproblems. Counting *outer iterations* is different from counting Hessian calculations, linear algebra operations or wall-clock time. The method does not say that arbitrary nonsmooth ReLU networks are globally optimized or that a second-order stationary point is necessarily the global optimum.

Survey relevance (RQ3). The method is a foundational second-order escape framework: modify the local quadratic so negative curvature does not destroy the model and analyze global complexity. Any theorem-numbered claims or exact constants should be checked after acquiring the full source.

5. Our notes

The cubic term prevents the indefinite Newton model from running off to minus infinity and makes curvature-aware steps mathematically manageable. The trade-off is a harder subproblem and Hessian-related cost per iteration. Source access is incomplete, so detailed original proofs are not asserted here. **Original bibliographic record:** https://doi.org/10.1007/s10107-006-0706-8.