

Convex Optimization

Lecture 1: Unconstrained Optimization for Differentiable Functions

Lecturer: *Dr.* Wan-Lei Zhao
Autumn Semester 2025

Outline

- 1 Overview about the Course
- 2 Optimization for Unconstrained Differentiable Problems
- 3 Maximum/Minimum by Gradient Descent Method
- 4 Solving Equation by Newton's Method-I
- 5 Minimization by Newton's Method-II

Syllabus

- 1 Optimization for Unconstrained Differentiable Functions (2)
- 2 Linear Programming
 - a Introduction (2)
 - b Simplex (4)
 - c Degenerated and Two-phase Simplex (2)
 - d Duality (2)
- 3 Quadratic Optimization
 - a Introduction (2)
 - b Convex Set (2)
 - c Convex Function and Convex Problem (3)
 - d Lagrange Multiplier and KKT (2)
 - e Dual of Lagrangian (2)
- 4 Portfolio Problem (2)
- 5 Support Vector Machine (2)
- 6 Integer Programming (2)

Assessment

① Assignment (10%)

- Coding
- Manual calc. exercises

② Attendance (10%)

③ Middle exam (20%)


④ Final exam (60%)

$$Score = 0.1 \times \sum_i Assignment_i + 0.1 \times Attend. + 0.2 \times mid + 0.6 \times final$$

- No cheating!
- No bargaining!
- No chatGPT!

Covered Topics

- **Optimization:** seek for an optimal solution for a defined objective function
- Under/without constraints





Nothing happens in the universe that does not have a sense of either certain maximum or minimum.

—Leonhard Euler (1707 – 1783)

Outline

- 1 Overview about the Course
- 2 Optimization for Unconstrained Differentiable Problems**
- 3 Maximum/Minimum by Gradient Descent Method
- 4 Solving Equation by Newton's Method-I
- 5 Minimization by Newton's Method-II

Derivatives on Function of Vector Variable

- Given $f(x) = a \cdot x_1 + b \cdot x_2$, $x = [x_1, x_2]^T \in \mathbb{R}^2$
- Take partial derivative on $f(x)$, we have

$$\frac{\partial f(x)}{\partial x_1} = a$$


$$\frac{\partial f(x)}{\partial x_2} = b$$

- The gradient of $f(x)$ is $[a, b]^T$ or $[a, b]$ depending on the context

Extreme Values of Univariate Function (1-1)

- Find out the inscribed rectangular of an ellipse that holds maximum area

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$



Extreme Values of Univariate Function (1-2)

- Find out the maximum point of the area function $A(x)$

- $A(x) = 4 * x * b * \sqrt{(1 - x^2/a^2)}$

- $A'(x) = 0$

```

1 syms a, b, x;
2 A = 4*b*x*(1 - x^2/a^2)^(1/2)
3 dA = diff(A,x)
4 solve(4*b*(1 - x^2/a^2)^0.5 - (4*b*x^2)/(a^2*(1 - x^2/a^2)^0.5)==0,x)


```

- Answer: $x = \frac{\sqrt{2}a}{2}$, viz $\arg \max_x A(x) = \frac{\sqrt{2}a}{2}$
- $\text{Max}(A(x)) = 2*a*b$

Extreme Values of multi-variable Function without Constraint (2-1)

- Given function:

$$f(x, y) = x^3 - y^3 + 3x^2 + 3y^2 - 9x \quad (1)$$



- Calculate its Hessian matrix

$$H = \begin{bmatrix} \frac{\partial^2 f(x,y)}{\partial x^2} & \frac{\partial^2 f(x,y)}{\partial x \partial y} \\ \frac{\partial^2 f(x,y)}{\partial y \partial x} & \frac{\partial^2 f(x,y)}{\partial y^2} \end{bmatrix} \quad (2)$$

Extreme Values of multi-variable Function without Constraint (2-1)

$$H = \begin{bmatrix} \frac{\partial^2 f(x,y)}{\partial x^2} & \frac{\partial^2 f(x,y)}{\partial x \partial y} \\ \frac{\partial^2 f(x,y)}{\partial y \partial x} & \frac{\partial^2 f(x,y)}{\partial y^2} \end{bmatrix}$$

$$\Downarrow$$

$$H = \begin{bmatrix} 6x + 6 & 0 \\ 0 & 6 - 6y \end{bmatrix}$$

$$\Downarrow$$

$$|H| = (6x + 6)(-6y + 6) - 0 = -36xy - 36y + 36x + 36 \quad (3)$$

Extreme Values of multi-variable Function without Constraint (2-2)

- Given function:

$$f(x, y) = x^3 - y^3 + 3x^2 + 3y^2 - 9x \quad (4)$$

$$\begin{cases} \frac{\partial f(x, y)}{\partial x} = 3x^2 + 6x - 9 = 0, \Rightarrow x = -3, 1 \\ \frac{\partial f(x, y)}{\partial y} = -y^2 + 6y = 0, \Rightarrow y = 0, 2 \end{cases}$$


x	y	$a = f_{xx}$	$ H(x, y) = a \cdot c - b^2$	f	Ext.
-3	0	-12	-72	27	uncertain
-3	2	-12	72	31	Ext. large
1	0	6	72	-5	Ext. small
1	2	6	-72	-1	uncertain

Outline

- 1 Overview about the Course
- 2 Optimization for Unconstrained Differentiable Problems
- 3 Maximum/Minimum by Gradient Descent Method
- 4 Solving Equation by Newton's Method-I
- 5 Minimization by Newton's Method-II


Why Gradient Descent?

- In practice, it is very hard to solve out a complex function




Why Gradient Descent?

- Given $f(x, y)$
- We want to find its minimum value



(a)



(b)

Derivative of a function and its tangent

$$g(x) = 2x^3 + 3x^2 - 12x + 7$$

- Given $g(x)$, tangent at x_0 is defined as:

$$f(x) = g'(x_0)(x - x_0) + g(x_0)$$





Figure: $g(x)$ is curve in blue, tangent of $g(x)$ at $x=-1$ is curve in red.

The gradient descent procedure

- Given we are going to find the minimum of $f(x)$, $x \in R^d$
- The learning rate is α
- ① *Initialize*(x)
- ② Repeat
 - a. $x^+ := x - \alpha f'(x)$
 - b. $x = x^+$
- Once we reaches the extreme, $f'(x) = 0$
- The procedure converges

Why Gradient Descent? (1)




- Given $f(x)$, $x \in R^d$, $f'(x_0) = \frac{\Delta y}{\Delta x}$
- We want to find its extreme value
- According to Taylor expansion $f(x) \approx f(x_0) + (x - x_0)f'(x_0)$

$$f(x) \approx f(x_0) + (x - x_0)f'(x_0) \quad (5)$$

- According to the above figure, we have

$$f(x) = f(x_0) - f'(x_0)\Delta x \quad (6)$$




$$f(x) = f(x_0) - f'(x_0)\Delta x$$

- $$\begin{aligned} f(x_0) + (x - x_0)f'(x_0) &= f(x_0) - f'(x_0)\Delta x \\ \Rightarrow x &= x_0 - \Delta x \end{aligned} \quad (6)$$

- $$x = x_0 - \alpha \cdot f'(x_0) \quad (7)$$


Why Gradient Descent? (3)



- Moreover, $\theta \rightarrow 0$, x drops the steepest

$$\begin{aligned}
 f(x_0) + (x - x_0)f'(x_0) &= f(x_0) - f'(x_0)\Delta x \\
 \implies x &= x_0 - \Delta x \\
 \implies x &= x_0 - \alpha f'(x_0)
 \end{aligned} \tag{6}$$

The General Steps in Gradient Descent



$$x = x_0 - \alpha f'(x_0) \quad (6)$$

1 *Initialize*(x)


2 Repeat

a. $x^+ := x - \alpha f'(x)$

b. $x = x^+$

Why Gradient Ascent? (1)


- Given we want to find the maximum of $f(x)$, $x \in R^d$



$$f(x) = f(x_0) + (x - x_0)f'(x_0)$$

$$f(x) = f(x_0) + f'(x_0)\Delta x$$

Why Gradient Ascent? (2)




$$f(x) = f(x_0) + (x - x_0)f'(x_0)$$

$$f(x) = f(x_0) + f'(x_0)\Delta x$$

\Downarrow

$$x = x_0 + \alpha f'(x_0) \quad (6)$$


The General Steps in Gradient Descent



$$x = x_0 + \alpha f'(x_0) \quad (6)$$


- 1 Initialize(x)
- 2 Repeat
 - a. $x^+ := x + \alpha f'(x)$
 - b. $x = x^+$

Minimum found by Gradient Descent (1)



$$f(x, y) = x^2 * y^3 + 3 * y^2 + 2 * x * y + x + 3; \quad (7)$$

Minimum found by Gradient Descent (2)



$$f(x, y) = x^2 * y^3 + 3 * y^2 + 2 * x * y + x + 3$$

$$\text{Grad}(f) = [2 * x * y^3 + 2 * y + 1, 3 * x^2 * y^2 + 6 * y + 2 * x]^T$$

Minimum found by Gradient Descent (3)

- Given learning rate $\alpha = 0.00005$


```
1 lr      = 0.00005;
2 xy      = rand(2,1);
3 iter = 0;
4 while iter < 500
5     xy = xy + lr*df(xy)';
6     iter = iter+1;
7 end
8
9 function [fval]=f(xy)
10     x = xy(1);  y = xy(2);
11     fval = x.^2*y.^3+3*y.^2+2*x.*y+x+3;
12 end
13
14 function [dx, dy]=df(xy)
15     x = xy(1);  y = xy(2);
16     dx = 2*x.*y.^3+2*y+1;
17     dy = 3*x.^2.*y.^2+6*y+2*x;
18 end
```

Outline

- 1 Overview about the Course
- 2 Optimization for Unconstrained Differentiable Problems
- 3 Maximum/Minimum by Gradient Descent Method
- 4 Solving Equation by Newton's Method-I**
- 5 Minimization by Newton's Method-II

The motivation


- We want to solve $f(x) = 0$, for instance $e^x - x^2 + 3x + 4 = 0$
- It is a little bit complicated



- We can solve it by an iterative procedure
- Which is known as Newton method

How it works? (1)

- We want to solve $f(x) = 0$, for instance $e^x - x^2 + 3x + 4 = 0$
- It is a little bit complicated



- Given we are at $(x_n, f(x_n))$, we want to move to $(x_{n+1}, f(x_{n+1}))$,
- which is closer to $f(x) = 0$

How it works? (2)


- The tangent line at $(x_n, f(x_n))$ is given as $y = f'(x_n)x + b$
- Since this line passes through $(x_n, f(x_n))$
- We have $y = f'(x_n) \cdot (x - x_n) + f(x_n)$
- We can easily find out its intersect point with x-axis

$$y = f'(x_n) \cdot (x - x_n) + f(x_n)$$

$$\xrightarrow{y=0} f'(x_n) \cdot (x - x_n) + f(x_n)$$

$$\Rightarrow x = \frac{x_n \cdot f'(x_n) - f(x_n)}{f'(x_n)}$$

$$\Rightarrow x = x_n - \frac{f(x_n)}{f'(x_n)}$$



How it works? (3)


$$y = f'(x_n) \cdot (x - x_n) + f(x_n)$$

$$\xrightarrow{y=0} f'(x_n) \cdot (x - x_n) + f(x_n)$$

$$\implies x = \frac{x_n \cdot f'(x_n) - f(x_n)}{f'(x_n)}$$


$$\implies x = x_n - \frac{f(x_n)}{f'(x_n)}$$

- $x = x_n - \frac{f(x_n)}{f'(x_n)}$ is the next iteration step
- The iteration continues until $f(x_n)$ reaches to 0




The Newton's method procedure

- 1 $x_n = x_0$
- 2 Repeat
 - a. $x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$
 - b. $x_n = x_{n+1}$
- 3 Until $f(x_n)$ close to 0



Practice with Newton's method (1)

- Solve $e^x - x^2 + 3x + 4 = 0$



- $f'(x) = e^x - 2x + 3$
- Notice that $f(x)$ is defined by ourselves

Practice with Newton's method (2)

- $f(x) = e^x - x^2 + 3x + 4$
- $f'(x) = e^x - 2x + 3$

```

1 function [x] = newtonsolve()
2     xn    = 6;
3     fval  = 8;
4     while fval > 0.001 do
5         fval    = f(xn);
6         dfval   = df(xn);
7         xp      = xn - fval/dfval;
8         xn      = xp;
9     end
10    x = xn;
11 end
12
13 function [fval]=f(x)
14     fval=exp(x) - x.^2+3*x + 4;
15 end
16
17 function [dfval]=df(x)
18     dfval=exp(x)-2*x+3;
19 end

```

Outline

- 1 Overview about the Course
- 2 Optimization for Unconstrained Differentiable Problems
- 3 Maximum/Minimum by Gradient Descent Method
- 4 Solving Equation by Newton's Method-I
- 5 Minimization by Newton's Method-II**

The Motivation

- Given a twice differentiable function: $f : R \rightarrow R$
- We have following minimization problem

$$\min_{x \in R} f(x) \quad (8)$$

- We want to construct a sequence $\{x_k\}$ based on initial guess x_0
- They allow $f(x_0) > f(x_1) > \dots > f(x_k) > \dots > f(x^*)$
- Each time, we are going to search the minimum value within the neighborhood of x_k
- So that we jump to the local minimum, namely x_{k+1}

How it works? (1)

- Given we are at x_k , t is a small value
- Then we expand $f(x_k)$ by *Taylor expansion*

$$f(x_k + t) \approx f(x_k) + f'(x_k)t + \frac{1}{2}f''(x_k)t^2 \quad (9)$$

- The next location x_{t+1} is defined to minimize $f(\cdot)$, $x_{k+1} = x_k + t$
- Given $f''(x_k)$ is positive, the minimum exists, and we can find it by

$$\begin{aligned} 0 &= \frac{d}{dt}(f(x_k) + f'(x_k)t + \frac{1}{2}f''(x_k)t^2) = f'(x_k) + f''(x_k)t \\ &\Rightarrow t = -\frac{f'(x)}{f''(x)} \end{aligned} \quad (10)$$

How it works? (2)

$$f(x_k + t) \approx f(x_k) + f'(x_k)t + \frac{1}{2}f''(x_k)t^2 \quad (11)$$

- The next location x_{k+1} is defined to minimize $f(\cdot)$, $x_{k+1} = x_k + t$
- Given $f''(x_k)$ is positive, the minimum exists, and we can find it by


$$\begin{aligned} 0 &= \frac{d}{dt}(f(x_k) + f'(x_k)t + \frac{1}{2}f''(x_k)t^2) = f'(x_k) + f''(x_k)t \\ &\Rightarrow t = -\frac{f'(x)}{f''(x)} \end{aligned} \quad (12)$$

- Since $x_{k+1} = x_k + t$, we have

$$x_{k+1} = x_k - \frac{f'(x)}{f''(x)} \quad (13)$$

The Newton's method procedure for minimization

- 1 $x_k = x_0$
- 2 Repeat
 - a. $x_{k+1} = x_k - \frac{f'(x_k)}{f''(x_k)}$
 - b. $x_k = x_{k+1}$
- 3 Until $|f(x_k) - f(x_{k+1})|$ is close to 0



Extend Newton's method to Multiple Dimension Cases

- Given a twice differentiable function: $f : R^n \rightarrow R$
- We have following minimization problem

$$\min_{x \in R^n} f(x) \quad (14)$$

- The Hessian matrix for $f(x)$ is defined as $H_f(x) \in R^{d \times d}$

$$x_{k+1} = x_k - H_f(x)^{-1} f'(x) \quad (15)$$

- For stableness, we also introduce the step size α for the iteration

$$x_{k+1} = x_k - \alpha H_f(x)^{-1} f'(x) \quad (16)$$