

Class 12 (Online)

Farnam (PID: A17628539)

2024-02-18

Section 1. proportion of G/G in a population

Downloaded a CSV file from ensemble < https://useast.ensembl.org/Homo_sapiens/Variation/Sample?db=core;r=17:39840102-39950103;v=rs8067378;vdb=variation;vf=959672880#373531_tablePanel

Here we read this CSV file

```
mxl <- read.csv("373531-SampleGenotypes-Homo_sapiens_Variation_Sample_rs8067378.csv")
head(mxl)
```

```
## Sample..Male.Female.Unknown. Genotype..forward.strand. Population.s. Father
## 1 NA19648 (F) A|A ALL, AMR, MXL -
## 2 NA19649 (M) G|G ALL, AMR, MXL -
## 3 NA19651 (F) A|A ALL, AMR, MXL -
## 4 NA19652 (M) G|G ALL, AMR, MXL -
## 5 NA19654 (F) G|G ALL, AMR, MXL -
## 6 NA19655 (M) A|G ALL, AMR, MXL -
## Mother
## 1 -
## 2 -
## 3 -
## 4 -
## 5 -
## 6 -
```

```
table(mxl$Genotype..forward.strand.)
```

```
##
## A|A A|G G|A G|G
## 22 21 12 9
```

```
table(mxl$Genotype..forward.strand.) / nrow(mxl) * 100
```

```
##
## A|A A|G G|A G|G
## 34.3750 32.8125 18.7500 14.0625
```

Section 2 and 3 would be done in Galaxy Portal

Now let's dig into this further:

One sample is obviously not enough to know what is happening in a population. You are interested in assessing genetic differences on a population scale.

How many samples do we have?

```
expr <- read.table("Last section.txt")
head(expr)
```

```
##      sample geno      exp
## 1 HG00367  A/G 28.96038
## 2 NA20768  A/G 20.24449
## 3 HG00361  A/A 31.32628
## 4 HG00135  A/A 34.11169
## 5 NA18870  G/G 18.25141
## 6 NA11993  A/A 32.89721
```

```
nrow(expr)
```

```
## [1] 462
```

```
table(expr$geno)
```

```
##
## A/A A/G G/G
## 108 233 121
```

```
library(ggplot2)
```

Let's make a boxplot:

```
ggplot(expr) + aes(geno, exp, fill=geno) +
geom_boxplot(notch = TRUE)
```

