

Run	Train data (#sample)	CG R@1	WIT R@1	Bloom R@1	Mammoth R@1	ImageNet	R@1-Avg
SEA-CLIP-Tiny-c1 (SEA blend)	CG-OE-filt + WIT-hf-base + Bloom (<i>1.21M</i>)	63.7	54.6	24.9	0.4	5.6	9.4
SEA-CLIP-Tiny-c2 (CC12M)	CC12M (10.97M)	0.1	1.9	1.2	0.5	40.0	6.5
SEA-CLIP-Tiny-c3 (CC12M + SEA)	CC12M + CG-OE-filt + WIT-hf + Bloom (<i>12.18M</i>)	33.3	25.2	16.1	0.8	37.4	14.2
SEA-CLIP-Tiny-c4	CC12M + CG-OE-filt + WIT-hf + Bloom + Mammoth-VL-SEA (<i>12.87M</i>)	29.3	21.8	23.3	15.3	36.4	17.4
SEA-CLIP-Tiny-WIT	WIT-hf-base (487K)	0.2	27.7	2.2	0.1	1.3	6.6
SEA-CLIP-Tiny-Bloom	Bloom-only (21K)	0.0	0.0	1.0	0.0	0.1	4.7
SEA-CLIP-Tiny-CG	CG-only (703K)	46.5	0.2	0.6	0.0	0.2	5.1
SEA-CLIP-Tiny w/o KD losses	CC12M + CG-OE-filt + WIT-hf + ML (<i>12.33M</i>)	47.4	41.5	28.8	—	—	—
SEA-CLIP-Tiny (CKD only)	CC12M + CG-OE-filt + WIT-hf + Bloom + Mammoth-VL-SEA (<i>12.87M</i>)	35.6	28.0	23.2	—	—	—
Teacher	—	<i>12.9</i>	<i>32.2</i>	<i>9.6</i>	<i>16.6</i>	<i>71.1</i>	<i>46.4</i>

Table 1: The dataset ablation study. All models are trained on the same training pipeline and hyperparameters. We ran the ablation on SEA-CLIP-Tiny using 3 choices: (c1) SEA blend, (c2) CC12M, and (c3) CC12M+SEA. We also tested on a single source of the SEA dataset, i.e., only WIT, Bloom, or CulturalGround (CG). We use **R@1-Avg** from Total Average@1 in Table ??.