
A
KEGG: higher in adult
Cadherin signaling
Cytokine−cytokine receptor interaction
NF−kappa B signaling pathway
Cell adhesion molecule (CAM)
Viral protein interaction with cytokine and cytokine receptor
ECM−receptor interaction
Rheumatoid arthritis
Pancreatic secretion
Complement and coagulation cascades
Salivary secretion
Staphylococcus aureus infection
Intestinal immune network for IgA production
0.02
0.04
0.06
0.08
GeneRatio
p.adjust
1e−03
3e−03
1e−02
Count
10
15
20
B
Hallmark: higher in adult
HALLMARK TNFA SIGNALING VIA NFKB
HALLMARK INFLAMMATORY RESPONSE
HALLMARK IL2 STAT5 SIGNALING
HALLMARK KRAS SIGNALING UP
HALLMARK IL6 JAK STAT3 SIGNALING
0.050
0.075
0.100
0.125
0.150
GeneRatio
p.adjust
1e−06
1e−04
1e−02
Count
10
15
20
25
30
C
KEGG: higher in infant
Bile secretion
Carbohydrate digestion and absorption
PPAR signaling pathway
Protein digestion and absorption
Fat digestion and absorption
Vitamin digestion and absorption
0.03
0.04
GeneRatio
p.adjust
1e−03
3e−03
1e−02
Count
6
8
10
12