

CropSage Texas Crop Catalog

Twenty-two sourced crop profiles for statewide Texas screening. Version 1.1.0 adds mature-crop effective root-zone depth ranges for crop-specific SSURGO available-water calculations.

Scoring guardrail: root-depth ranges determine the SSURGO integration depth; they do not prove current soil moisture or seasonal water sufficiency.

Catalog Scope

Catalog version	Jurisdiction	Review date	Crops	References
1.1.0	Texas, United States	2026-08-28	22	30

Texas Regions

Region ID	Display name	Catalog interpretation	Sources
plains	Texas Plains	High Plains, Rolling Plains, and adjoining north-central production areas; generally semi-arid with important irrigated and dryland field-crop systems.	SRC-002, SRC-003
eastern	Eastern Texas and Gulf Coast	Humid eastern counties and the Gulf Coast production belt, including the principal Texas rice area and substantial forage and vegetable production.	SRC-002, SRC-003
central_south_winter_garden	Central and South Texas / Winter Garden	Central and southern production areas outside the Lower Rio Grande Valley, including the irrigated Winter Garden vegetable district.	SRC-002, SRC-003
lower_rio_grande_valley	Lower Rio Grande Valley	The subtropical southern tip of Texas, with mild winters, hot summers, and predominantly irrigated commercial agriculture.	SRC-002, SRC-003
far_west	Far West Texas	Arid Trans-Pecos and El Paso production areas where commercial cropping is strongly constrained by irrigation-water availability.	SRC-002, SRC-003

Region labels remain screening zones; coordinate-to-region polygons are a separate unresolved geographic dataset.

Crop Profiles 1-8

Crop	Use	Regional ratings	Planting windows	Maturity	Temperature	Soil and pH	Root zone	Drainage	Water and irrigation	Evidence
Upland cotton <i>Gossypium hirsutum</i>	lint and cottonseed	Texas Plains: primary; Eastern Texas and Gulf Coast: suitable; Central and South Texas / Winter Garden: primary; Lower Rio Grande Valley: primary; Far West Texas: conditional	Texas Plains: May-Jun; Eastern Texas and Gulf Coast: Apr-May; Central and South Texas / Winter Garden: Mar-May; Lower Rio Grande Valley: Feb-Apr; Far West Texas: Apr-May	150-180 days	Optimum 26.7-32.2 C Heat > 37.8 C	loam, sandy loam, clay loam pH 5.7-7.0	100-170 cm regional_transfer SRC-028, SRC-030	well drained	Demand: high (700-1300 mm) Drought: high Irrigation: conditional	medium confidence provisional SRC-001, SRC-003, SRC-004, SRC-005, SRC-006, SRC-028, SRC-030
Corn grown for grain <i>Zea mays</i>	dry grain	Texas Plains: primary; Eastern Texas and Gulf Coast: suitable; Central and South Texas / Winter Garden: primary; Lower Rio Grande Valley: suitable; Far West Texas: conditional	Texas Plains: Apr-May; Eastern Texas and Gulf Coast: Feb-Apr; Central and South Texas / Winter Garden: Feb-Apr; Lower Rio Grande Valley: Jan-Mar; Far West Texas: Mar-Apr	110-140 days	Optimum 24.0-30.0 C Heat > 35.0 C	loam, silt loam, clay loam, sandy loam pH 5.5-7.0	100-170 cm regional_transfer SRC-028, SRC-030	well drained	Demand: moderate (500-800 mm) Drought: low Irrigation: conditional	medium confidence provisional SRC-001, SRC-003, SRC-004, SRC-005, SRC-007, SRC-028, SRC-030
Hard red winter wheat <i>Triticum aestivum</i>	food grain	Texas Plains: primary; Eastern Texas and Gulf Coast: suitable; Central and South Texas / Winter Garden: suitable; Lower Rio Grande Valley: limited; Far West Texas: conditional	Texas Plains: Sep-Nov; Eastern Texas and Gulf Coast: Oct-Nov; Central and South Texas / Winter Garden: Oct-Dec; Lower Rio Grande Valley: Nov-Dec; Far West Texas: Sep-Nov	210-270 days	Optimum 23.9-25.0 C Heat > 30.0 C	loam, clay loam, silt loam, sandy loam pH 5.5-7.0	150-180 cm generalized SRC-028	well drained	Demand: moderate (450-650 mm) Drought: moderate Irrigation: conditional	medium confidence provisional SRC-001, SRC-003, SRC-004, SRC-005, SRC-008, SRC-009, SRC-028
Grain sorghum <i>Sorghum bicolor</i>	dry grain	Texas Plains: primary; Eastern Texas and Gulf Coast: suitable; Central and South Texas / Winter Garden: primary; Lower Rio Grande Valley: primary; Far West Texas: conditional	Texas Plains: Apr-Jun; Eastern Texas and Gulf Coast: Mar-May; Central and South Texas / Winter Garden: Feb-Jun; Lower Rio Grande Valley: Feb-Mar, Jul-Aug; Far West Texas: Apr-May	90-120 days	Optimum 26.0-32.0 C Heat > 37.2 C	loam, clay loam, sandy loam, silty clay loam pH 5.5-7.0	100-200 cm regional_transfer SRC-028, SRC-030	well drained	Demand: moderate (450-650 mm) Drought: high Irrigation: often rainfed	medium confidence provisional SRC-001, SRC-003, SRC-004, SRC-005, SRC-009, SRC-028, SRC-030
Runner-type peanut <i>Arachis hypogaea</i>	edible seed and oil	Texas Plains: primary; Eastern Texas and Gulf Coast: suitable; Central and South Texas / Winter Garden: suitable; Lower Rio Grande Valley: conditional; Far West Texas: limited	Texas Plains: May-Jun; Eastern Texas and Gulf Coast: Apr-May; Central and South Texas / Winter Garden: Apr-May; Lower Rio Grande Valley: Mar-Apr	130-160 days	Optimum 25.0-30.0 C Heat not numeric	loamy sand, sandy loam, fine sandy loam pH 5.8-7.0	50-100 cm generalized SRC-028	very well drained	Demand: moderate (500-700 mm) Drought: moderate Irrigation: conditional	medium confidence provisional SRC-001, SRC-003, SRC-004, SRC-005, SRC-011, SRC-028
Long-grain rice <i>Oryza sativa</i>	milled long-grain rice	Texas Plains: not_supported; Eastern Texas and Gulf Coast: primary; Central and South Texas / Winter Garden: limited; Lower Rio Grande Valley: limited; Far West Texas: not_supported	Eastern Texas and Gulf Coast: Mar-May	100-150 days	Optimum 25.0-30.0 C Heat not numeric	clay, clay loam, silty clay loam pH 5.5-7.0	50-100 cm regional_transfer SRC-028, SRC-030	poorly drained tolerant	Demand: very_high (450-700 mm) Drought: very_low Irrigation: usually required	medium confidence provisional SRC-001, SRC-003, SRC-004, SRC-014, SRC-028, SRC-030
Soybean <i>Glycine max</i>	oilseed and protein meal	Texas Plains: conditional; Eastern Texas and Gulf Coast: primary; Central and South Texas / Winter Garden: suitable; Lower Rio Grande Valley: conditional; Far West Texas: limited	Texas Plains: May-Jun; Eastern Texas and Gulf Coast: Mar-Jun; Central and South Texas / Winter Garden: Mar-May; Lower Rio Grande Valley: Feb-Apr	100-150 days	Optimum 24.0-29.4 C Heat > 29.4 C	loam, silt loam, clay loam, sandy loam pH 5.8-7.0	60-130 cm regional_transfer SRC-028, SRC-030	well drained	Demand: moderate (450-700 mm) Drought: moderate Irrigation: conditional	medium confidence provisional SRC-001, SRC-003, SRC-004, SRC-005, SRC-010, SRC-028, SRC-030
Grain oats <i>Avena sativa</i>	grain	Texas Plains: suitable; Eastern Texas and Gulf Coast: primary; Central and South Texas / Winter Garden: suitable; Lower Rio Grande Valley: limited; Far West Texas: conditional	Texas Plains: Sep-Oct; Eastern Texas and Gulf Coast: Sep-Nov; Central and South Texas / Winter Garden: Oct-Nov	180-240 days	Optimum 20.0-21.1 C Heat > 27.8 C	loam, clay loam, silt loam, sandy loam pH 5.5-7.0	100-150 cm regional_transfer SRC-028, SRC-030	well drained	Demand: moderate (450-650 mm) Drought: moderate Irrigation: conditional	medium confidence provisional SRC-001, SRC-003, SRC-004, SRC-005, SRC-008, SRC-028, SRC-030

Crop Profiles 9-16

Crop	Use	Regional ratings	Planting windows	Maturity	Temperature	Soil and pH	Root zone	Drainage	Water and irrigation	Evidence
Oilseed sunflower <i>Helianthus annuus</i>	oilseed	Texas Plains: primary; Eastern Texas and Gulf Coast: suitable; Central and South Texas / Winter Garden: suitable; Lower Rio Grande Valley: conditional; Far West Texas: conditional	Texas Plains: Apr-Jul; Eastern Texas and Gulf Coast: Mar-May; Central and South Texas / Winter Garden: Mar-Jun; Lower Rio Grande Valley: Feb-Apr, Aug-Sep; Far West Texas: Apr-May	90-120 days	Optimum 15.0-25.0 C Heat > 35.0 C	loam, silt loam, clay loam, sandy loam pH 5.5-7.5	80-150 cm generalized SRC-028	well drained	Demand: high (600-1000 mm) Drought: high Irrigation: often rainfed	medium confidence provisional SRC-001, SRC-003, SRC-004, SRC-012, SRC-028
Sesame <i>Sesamum indicum</i>	oilseed and edible seed	Texas Plains: primary; Eastern Texas and Gulf Coast: conditional; Central and South Texas / Winter Garden: primary; Lower Rio Grande Valley: suitable; Far West Texas: conditional	Texas Plains: May-Jun; Eastern Texas and Gulf Coast: May-Jun; Central and South Texas / Winter Garden: Apr-Jun; Lower Rio Grande Valley: Mar-May; Far West Texas: May-Jun	90-120 days	Optimum 25.0-32.0 C Heat not numeric	sandy loam, loam, silt loam pH 5.5-8.0	100-150 cm generalized SRC-028	very well drained	Demand: low (not established) Drought: very_high Irrigation: often rainfed	low confidence provisional SRC-002, SRC-003, SRC-013, SRC-028
Corn grown for silage <i>Zea mays</i>	whole-plant silage	Texas Plains: primary; Eastern Texas and Gulf Coast: suitable; Central and South Texas / Winter Garden: suitable; Lower Rio Grande Valley: conditional; Far West Texas: conditional	Texas Plains: Apr-May; Eastern Texas and Gulf Coast: Mar-Apr; Central and South Texas / Winter Garden: Feb-Apr; Lower Rio Grande Valley: Jan-Mar; Far West Texas: Mar-Apr	80-110 days	Optimum 24.0-30.0 C Heat > 35.0 C	loam, silt loam, clay loam, sandy loam pH 5.5-7.0	100-170 cm regional_transfer SRC-028, SRC-030	well drained	Demand: moderate (500-800 mm) Drought: low Irrigation: recommended	medium confidence provisional SRC-001, SRC-003, SRC-004, SRC-005, SRC-007, SRC-028, SRC-030
Forage sorghum <i>Sorghum bicolor</i>	hay, greenchop, or silage forage	Texas Plains: primary; Eastern Texas and Gulf Coast: suitable; Central and South Texas / Winter Garden: primary; Lower Rio Grande Valley: suitable; Far West Texas: conditional	Texas Plains: May-Jun; Eastern Texas and Gulf Coast: Apr-Jul; Central and South Texas / Winter Garden: Mar-Jul; Lower Rio Grande Valley: Feb-Aug; Far West Texas: May-Jun	60-90 days	Optimum 26.0-32.0 C Heat not numeric	loam, clay loam, sandy loam, silty clay loam pH 5.5-7.0	100-200 cm regional_transfer SRC-028, SRC-030	well drained	Demand: moderate (450-650 mm) Drought: high Irrigation: often rainfed	medium confidence provisional SRC-003, SRC-004, SRC-005, SRC-009, SRC-026, SRC-028, SRC-030
Sorghum-sudangrass <i>Sorghum bicolor x Sorghum bicolor var. sudanense</i>	grazing, hay, greenchop, or silage forage	Texas Plains: primary; Eastern Texas and Gulf Coast: suitable; Central and South Texas / Winter Garden: primary; Lower Rio Grande Valley: suitable; Far West Texas: conditional	Texas Plains: May-Jun; Eastern Texas and Gulf Coast: Apr-Jul; Central and South Texas / Winter Garden: Mar-Jul; Lower Rio Grande Valley: Feb-Aug; Far West Texas: May-Jun	50-80 days	Optimum 26.0-32.0 C Heat not numeric	loam, clay loam, sandy loam, silt loam pH 5.5-7.0	100-150 cm generalized SRC-028	well drained	Demand: moderate (450-650 mm) Drought: high Irrigation: often rainfed	medium confidence provisional SRC-003, SRC-004, SRC-005, SRC-009, SRC-026, SRC-028
Alfalfa grown for hay <i>Medicago sativa</i>	perennial hay	Texas Plains: primary; Eastern Texas and Gulf Coast: limited; Central and South Texas / Winter Garden: suitable; Lower Rio Grande Valley: conditional; Far West Texas: primary	Texas Plains: Sep-Oct, Mar-Apr; Central and South Texas / Winter Garden: Sep-Nov, Feb-Mar; Far West Texas: Sep-Oct, Feb-Mar	60-90 days	Optimum 18.0-30.0 C Heat not numeric	deep loam, silt loam, clay loam, well-drained sandy loam pH 6.3-7.5	100-200 cm regional_transfer SRC-028, SRC-030	very well drained	Demand: very_high (800-1600 mm) Drought: moderate Irrigation: usually required	medium confidence provisional SRC-003, SRC-004, SRC-005, SRC-024, SRC-028, SRC-030
Bermudagrass grown for hay <i>Cynodon dactylon</i>	perennial hay	Texas Plains: conditional; Eastern Texas and Gulf Coast: primary; Central and South Texas / Winter Garden: primary; Lower Rio Grande Valley: primary; Far West Texas: conditional	Texas Plains: May-Jun; Eastern Texas and Gulf Coast: Mar-Jun; Central and South Texas / Winter Garden: Mar-Jun; Lower Rio Grande Valley: Feb-Jun; Far West Texas: Apr-Jun	60-90 days	Optimum 21.1-35.0 C Heat not numeric	sandy loam, loam, clay loam, silt loam pH 5.7-7.0	100-150 cm regional_transfer SRC-028, SRC-030	well drained	Demand: high (not established) Drought: very_high Irrigation: conditional	medium confidence provisional SRC-003, SRC-005, SRC-025, SRC-028, SRC-030
Annual ryegrass grown for forage <i>Lolium multiflorum</i>	cool-season grazing, baleage, or hay	Texas Plains: conditional; Eastern Texas and Gulf Coast: primary; Central and South Texas / Winter Garden: suitable; Lower Rio Grande Valley: conditional; Far West Texas: limited	Texas Plains: Sep-Oct; Eastern Texas and Gulf Coast: Oct-Nov; Central and South Texas / Winter Garden: Oct-Nov; Lower Rio Grande Valley: Nov-Dec	60-90 days	Optimum 10.0-21.0 C Heat not numeric	loam, silt loam, clay loam, sandy loam pH 5.5-7.0	60-100 cm generalized SRC-028	moderate	Demand: moderate (not established) Drought: low Irrigation: conditional	medium confidence provisional SRC-003, SRC-005, SRC-027, SRC-028

Crop Profiles 17-22

Crop	Use	Regional ratings	Planting windows	Maturity	Temperature	Soil and pH	Root zone	Drainage	Water and irrigation	Evidence
White potato <i>Solanum tuberosum</i>	fresh-market and processing tubers	Texas Plains: suitable; Eastern Texas and Gulf Coast: conditional; Central and South Texas / Winter Garden: primary; Lower Rio Grande Valley: suitable; Far West Texas: conditional	Texas Plains: Mar-Apr, Jul-Aug; Eastern Texas and Gulf Coast: Feb-Mar, Aug-Sep; Central and South Texas / Winter Garden: Jan-Mar, Aug-Sep; Lower Rio Grande Valley: Nov-Feb; Far West Texas: Feb-Mar, Jul-Aug	90-120 days	Optimum 15.0-24.0 C Heat not numeric	sandy loam, loam, fine sandy loam pH 5.0-7.8	40-60 cm regional_transfer SRC-028, SRC-030	very well drained	Demand: high (508-1016 mm) Drought: very_low Irrigation: recommended	medium confidence provisional SRC-002, SRC-003, SRC-015, SRC-020, SRC-023, SRC-028, SRC-030
Sweet potato <i>Ipomoea batatas</i>	fresh-market and processing storage roots	Texas Plains: conditional; Eastern Texas and Gulf Coast: primary; Central and South Texas / Winter Garden: suitable; Lower Rio Grande Valley: suitable; Far West Texas: limited	Texas Plains: May-Jun; Eastern Texas and Gulf Coast: Apr-Jun; Central and South Texas / Winter Garden: Mar-May; Lower Rio Grande Valley: Feb-May	90-150 days	Optimum 24.0-32.0 C Heat not numeric	fine sandy loam, sandy loam, loam pH 5.0-7.5	100-150 cm generalized SRC-028	very well drained	Demand: moderate (254-508 mm) Drought: moderate Irrigation: conditional	medium confidence provisional SRC-002, SRC-003, SRC-015, SRC-019, SRC-023, SRC-028
Seedless watermelon <i>Citrullus lanatus</i>	fresh-market fruit	Texas Plains: suitable; Eastern Texas and Gulf Coast: suitable; Central and South Texas / Winter Garden: primary; Lower Rio Grande Valley: primary; Far West Texas: conditional	Texas Plains: May-Jun; Eastern Texas and Gulf Coast: Mar-Jun; Central and South Texas / Winter Garden: Feb-May; Lower Rio Grande Valley: Jan-Apr; Far West Texas: Apr-May	65-75 days	Optimum 26.7-35.0 C Heat not numeric	loamy sand, sandy loam, light-textured loam pH 5.5-8.0	80-150 cm regional_transfer SRC-028, SRC-030	very well drained	Demand: moderate (254-381 mm) Drought: low Irrigation: recommended	high confidence provisional SRC-002, SRC-003, SRC-015, SRC-016, SRC-023, SRC-028, SRC-030
Dry-bulb onion <i>Allium cepa</i>	cured dry bulbs	Texas Plains: suitable; Eastern Texas and Gulf Coast: conditional; Central and South Texas / Winter Garden: primary; Lower Rio Grande Valley: primary; Far West Texas: suitable	Texas Plains: Feb-Mar; Eastern Texas and Gulf Coast: Sep-Nov, Jan-Feb; Central and South Texas / Winter Garden: Sep-Nov; Lower Rio Grande Valley: Sep-Nov; Far West Texas: Jan-Mar	150-210 days	Optimum 15.6-26.7 C Heat > 35.0 C	sandy loam, loam, silt loam pH 6.0-8.4	30-60 cm generalized SRC-028	very well drained	Demand: high (635-762 mm) Drought: very_low Irrigation: usually required	high confidence provisional SRC-002, SRC-003, SRC-015, SRC-017, SRC-023, SRC-028
Fresh-market cabbage <i>Brassica oleracea</i> var. <i>capitata</i>	fresh-market heads	Texas Plains: suitable; Eastern Texas and Gulf Coast: suitable; Central and South Texas / Winter Garden: primary; Lower Rio Grande Valley: primary; Far West Texas: conditional	Texas Plains: Feb-Mar, Jul-Aug; Eastern Texas and Gulf Coast: Jan-Mar, Aug-Oct; Central and South Texas / Winter Garden: Sep-Nov, Jan-Feb; Lower Rio Grande Valley: Sep-Dec; Far West Texas: Feb-Mar, Aug-Sep	70-120 days	Optimum 15.6-21.1 C Heat not numeric	loam, silt loam, clay loam, sandy loam pH 6.0-7.5	50-80 cm generalized SRC-028	well drained	Demand: high (508-762 mm) Drought: very_low Irrigation: recommended	high confidence provisional SRC-002, SRC-003, SRC-015, SRC-021, SRC-022, SRC-023, SRC-028
Fresh-market spinach <i>Spinacia oleracea</i>	fresh-market leaves	Texas Plains: suitable; Eastern Texas and Gulf Coast: conditional; Central and South Texas / Winter Garden: primary; Lower Rio Grande Valley: primary; Far West Texas: suitable	Texas Plains: Feb-Mar, Aug-Oct; Eastern Texas and Gulf Coast: Jan-Mar, Sep-Nov; Central and South Texas / Winter Garden: Sep-Feb; Lower Rio Grande Valley: Oct-Feb; Far West Texas: Feb-Mar, Sep-Oct	35-50 days	Optimum 18.3-23.9 C Heat > 23.9 C	clay loam, loam, silt loam pH 6.5-8.0	30-50 cm generalized SRC-028	well drained	Demand: moderate (254-381 mm) Drought: very_low Irrigation: recommended	high confidence provisional SRC-002, SRC-003, SRC-015, SRC-018, SRC-023, SRC-028

References 1-15

ID	Title	Publisher	Supports	URL
SRC-001	Field Crops: Usual Planting and Harvesting Dates	USDA National Agricultural Statistics Service	Texas statewide field-crop planting windows; harvest timing	https://www.nass.usda.gov/Publications/Todays_Reports/reports/fcdate10.pdf
SRC-002	Descriptions of Geographic Regions in Texas	Texas A&M AgriLife Extension	Texas region vocabulary; regional climate and production context	https://aggie-horticulture.tamu.edu/vegetable/guides/the-crops-of-texas/appendix-1-descriptions-of-geographic-regions-in-texas/
SRC-003	The Crops of Texas: Highlights and Overviews	Texas A&M AgriLife Extension	regional crop distribution; Texas production context	https://aggie-horticulture.tamu.edu/vegetable/guides/the-crops-of-texas/highlights-and-overviews/
SRC-004	Crop Water Needs and Sensitivity to Drought	Food and Agriculture Organization of the United Nations	indicative seasonal crop-water ranges; relative drought sensitivity; critical growth stages	https://www.fao.org/4/S2022E/s2022e07.htm
SRC-005	OSU Soil Test Interpretations	Oklahoma State University Extension	Southern Plains crop soil-pH preferences	https://extension.okstate.edu/fact-sheets/osu-soil-test-interpretations
SRC-006	When Is the Best Time to Plant Cotton in the Oklahoma Panhandle?	Oklahoma State University Extension	cotton thermal basis; cotton heat warning; cotton maturity heat units	https://extension.okstate.edu/fact-sheets/when-is-the-best-time-to-plant-cotton-in-the-oklahoma-panhandle
SRC-007	Dry Conditions During Corn Pollination	University of Minnesota Extension	corn heat thresholds; pollination sensitivity; interaction between heat and soil moisture	https://extension.umn.edu/agriculture/crop-production/corn/dry-conditions-during-corn-pollination
SRC-008	Heat Stress and Drought Stress in Small Grains	University of Minnesota Extension	wheat and oat temperature ranges; small-grain heat-sensitive stages; small-grain heat thresholds	https://extension.umn.edu/growing-small-grains/heat-stress-and-drought-stress-small-grains
SRC-009	Agriculture, Climate, and Change: Temperature Effects on Wheat and Sorghum	Kansas State University Extension	wheat and sorghum thermal response; flowering heat sensitivity	https://eupdate.agronomy.ksu.edu/article/new-eupdate-series-on-agriculture-climate-and-change-part-1-298
SRC-010	How Do High Temperatures Impact My Soybeans?	North Carolina State Extension	soybean optimum temperature; soybean heat stress; R5 sensitivity	https://soybeans.ces.ncsu.edu/news/how-do-high-temperatures-impact-my-soybeans/
SRC-011	Peanut Production Field Guide	University of Georgia Cooperative Extension	runner peanut production requirements; soil and irrigation management; maturity	https://extension.uga.edu/publications/detail.html?number=B1146
SRC-012	Sunflower Planting Depth and Population in the Oklahoma Panhandle	Oklahoma State University Extension	oilseed sunflower soil-temperature range; germination heat warning; drought tolerance	https://extension.okstate.edu/fact-sheets/sunflower-planting-depth-and-population-in-the-oklahoma-panhandle
SRC-013	Sesame Seed Oil Properties	Oklahoma State University Extension	commercial sesame scope; Texas and Oklahoma production context; drought tolerance	https://extension.okstate.edu/fact-sheets/sesame-seed-oil-properties
SRC-014	Louisiana Rice Production Handbook	LSU AgCenter	US Gulf Coast rice production requirements; soil, water, and crop calendar context	https://www.lsuagcenter.com/~media/system/9/0/e/9/90e93160aba5daccea90c6d955299f74/pub2331riceproductionhandbook2014completebook.pdf
SRC-015	Appendix of Useful Tables: Organic Vegetable Production Guide	Texas A&M AgriLife Extension	Texas vegetable water use; critical water stages; vegetable pH ranges; variety context	https://aggie-horticulture.tamu.edu/vegetable/guides/organic-vegetable-production-guide/appendix-of-useful-tables/

References 16-30

ID	Title	Publisher	Supports	URL
SRC-016	Seedless Watermelon Crop Guide	Texas A&M AgriLife Extension	seedless watermelon temperature; soil; pH; water; maturity	https://aggie-horticulture.tamu.edu/vegetable/watermelon/seedless-watermelon-crop-guide/
SRC-017	Onion Commercial Crop Guide	Texas A&M AgriLife Extension	dry-bulb onion temperature; day-length scope; soil; pH; planting conditions	https://aggie-horticulture.tamu.edu/vegetable/wp-content/uploads/sites/10/2011/10/onion1.pdf
SRC-018	Spinach Commercial Crop Guide	Texas A&M AgriLife Extension	fresh-market spinach temperature; soil; pH; frost tolerance	https://aggie-horticulture.tamu.edu/vegetable/wp-content/uploads/sites/10/2011/10/spinach.pdf
SRC-019	Easy Gardening: Sweet Potatoes	Texas A&M AgriLife Extension	sweet-potato temperature; soil; pH; frost response	https://aggie-horticulture.tamu.edu/vegetable/files/2010/10/EHT-026-Easy-Gardening-Sweet-Potatoes.pdf
SRC-020	Easy Gardening: Irish Potatoes	Texas A&M AgriLife Extension	white-potato planting temperature; Texas planting windows; soil and crop duration	https://aggie-horticulture.tamu.edu/wp-content/uploads/sites/12/2011/03/E-511_irish_potato.pdf
SRC-021	Easy Gardening: Cole Crops	Texas A&M AgriLife Extension	cabbage cold tolerance; Texas seasonal context	https://aggie-horticulture.tamu.edu/vegetable/wp-content/uploads/sites/10/2010/10/E-279_cole_crops.pdf
SRC-022	Timely Tips on Starting Seedlings at Home	Texas A&M AgriLife Extension	vegetable day and night growing temperature ranges	https://aggie-horticulture.tamu.edu/earthkind/landscape/starting-seedlings-at-home/
SRC-023	Fall Vegetable Gardening Guide	Texas A&M AgriLife Extension	Texas vegetable planting zones; fall planting timing; frost susceptibility; maturity classes	https://aggie-horticulture.tamu.edu/wp-content/uploads/sites/10/2013/09/EHT-056.pdf
SRC-024	Texas Alfalfa Production	Texas A&M AgriLife Extension	alfalfa Texas adaptation; soil depth; water and drainage requirements; cutting interval	https://agrilifeextension.tamu.edu/wp-content/uploads/2026/03/texasalfalaproduction.pdf
SRC-025	Bermudagrass Varieties, Hybrids, and Blends for Texas	Texas A&M AgriLife Extension	bermudagrass Texas adaptation; soil drainage; summer growth; hay use	https://agrilifeextension.tamu.edu/wp-content/uploads/2025/07/Bermudagrass-Varieties-and-Blends-for-Texas.pdf
SRC-026	Warm Season Annual Forages	Texas A&M AgriLife Extension Forage Fax	forage sorghum and sorghum-sudangrass Texas adaptation; soil moisture and system fit	https://foragefax.tamu.edu/warm-season-annual-forages/
SRC-027	Cool Season Annual Forages: To Plant or Not To Plant	Texas A&M AgriLife Extension Forage Fax	annual ryegrass Texas planting timing; soil-moisture requirement; forage season	https://foragefax.tamu.edu/cool-season-annual-forages-to-plant-or-not-to-plantnow-or-later/
SRC-028	Crop Evapotranspiration: Guidelines for Computing Crop Water Requirements	Food and Agriculture Organization of the United Nations	crop coefficient context; water-demand comparison for field crops and forages; maximum effective rooting-depth ranges for mature crops; soil-water depletion fractions	https://www.fao.org/4/x0490e/x0490e0e.htm
SRC-029	SSURGO Web Soil Survey and Soil Data Access	USDA Natural Resources Conservation Service	runtime comparison of mapped soil texture; pH; drainage; available water capacity	https://www.nrcs.usda.gov/resources/data-and-reports/soil-survey-geographic-database-ssurgo
SRC-030	Irrigation Monitoring with Soil Water Sensors	Texas A&M AgriLife Extension	Texas-context root-depth ranges for selected field crops, forages, and vegetables; soil and management limits on effective root-zone depth; soil-water monitoring context	https://moore.agrilife.org/files/2024/05/E-618_-_Irrigation-Monitoring-with-Soil-Water-Sensors-resized.pdf