

Accounts and clinical sources
Patient
patient_id : UUID
physician_id : UUID
owns
1
0..N
Physician
physician_id : UUID
1
has
0..N
1
ClinicalInput
input_id : UUID
encounter_id : UUID
contains
1
0..N
Encounter
encounter_id : UUID
patient_id : UUID
1
1
versions
1
1..N
InputRevision
input_id : UUID
revision_no : integer
1
analysed by
0..N
SpecialistFinding
finding_id : UUID
0..1
1
produces
provides
performs
Drafts and evidence
0..N
Proposal
proposal_id : UUID
encounter_id : UUID
1
versions
1..N
EvidenceItem
evidence_id : UUID
ProposalRevision
proposal_id : UUID
revision_no : integer
supplies
1
1
1
1
1
linked by
uses
supports
assessed by
0..N
0..N
0..N
0..N
0..N
ProposalInput
proposal_id : UUID
proposal_revision : integer
ProposalEvidence
proposal_id : UUID
proposal_revision : integer
CriticReview
critic_id : UUID
represented by
Research evaluation
0..N
EvaluationCase
case_id : UUID
encounter_id : UUID
EvaluationRun
run_id : UUID
1
1
evaluated in
records
0..N
0..N
EvaluationOutput
output_id : UUID
reviewed in
accepted as
Review and memory
0..N
0..N
PhysicianAction
action_id : UUID
physician_id : UUID
1
authorises
0..1
0..1
ClinicalDecision
decision_id : UUID
0..1
represented by
0..N
0..N
SemanticMemory
memory_id : UUID
decision_id : UUID