

# Number Theory

2018A7PS0193P



## Contents

Chapter 1. Fundamentals	5
1. Notation	5
2. Induction	5
3. Well Ordering Principle	6
4. Binomial Theorem	7
5. Pigeonhole Principle	7
Chapter 2. Division	9
1. Division Algorithm	9
2. Base b representations	10
Chapter 3. Properties of Numbers	13
1. Prime and Composite Numbers	13
2. Prime Counting function	15
3. Gaps between Primes	16
4. Fermat Numbers	17
5. Fibonacci Numbers	18
6. Lucas Numbers	19
Chapter 4. Greatest Common Divisor and Least Common Multiple	21
1. Greatest Common Divisor	21
2. Euclidean Algorithm	24
3. Least Common Multiple	25
4. Fundamental Theorem of Arithmetic	26
Chapter 5. Linear Diophantine Equations	27
1. Linear Diophantine Equations in Two Variables	27
2. Fibonacci Linear Diophantine Equation	27
3. Linear Diophantine Equations in Many Variables	28
Chapter 6. Congruence	31
1. Introduction	31
2. Square and Multiply Algorithm	35

3. Cancellation in Congruences	35
4. Combining Congruences with different moduli	35
5. Linear Congruences	36
6. System of Linear Congruences	37
7. Modular Inverse	39

## CHAPTER 1

### Fundamentals

#### 1. Notation

For the rest of this course, the following notation will be followed:

- (1)  $\mathbb{N}$  is the set of natural numbers
- (2)  $\mathbb{Z}$  is the set of integers
- (3)  $\mathbb{W}$  is the set of whole numbers, i.e.  $\mathbb{W} = \mathbb{N} \cup \{0\}$

#### 2. Induction

Often in number theory, we use inductive proofs to prove our arguments. Induction consists of the following steps:

- (1) Define an induction hypothesis  $P(k)$
- (2) Verify it works for some base case  $k = b$ . It is possible multiple base cases need to be verified.
- (3) Assuming  $P(k)$  is true, show that it implies that  $P(k + 1)$  is true

Remember that  $P(k)$  is a statement, not a function. You cannot multiply it by some constant or perform any operations on it.

In weak induction (like in the steps given above), we only assume that  $P(k)$  is true. However in strong induction, we assume that  $P(i)$  is true  $\forall i \in [b, k]$ , and use this to prove that  $P(k+1)$  is true.

EXERCISE. Prove that the principle of strong induction is true given that the principle of weak induction is true.

SOLUTION. Let us assume that  $P(1), \dots, P(b)$  is true. If  $P(1), \dots, P(k)$  are true for some  $k \geq b$ , then  $P(k + 1)$  is true. Then, we must show that  $P(n)$  is true for all  $n \geq 1$ .

Let  $Q(n)$  be the statement that  $P(1), \dots, P(n)$  are true. Of course, in the base case,  $Q(1)$  is true. Let  $Q(k)$  be true, where  $K \geq 1$ . This means that  $P(1), \dots, P(k)$  is true, so  $P(k + 1)$  must be true. Hence,  $Q(k + 1)$  is true.

So, by Weak induction,  $Q(n)$  is true  $\forall n \geq 1$ , which implies that  $P(n)$  is true  $\forall n \geq 1$ . ■

### 3. Well Ordering Principle

**THEOREM 1.1** (Well Ordering Principle). Every non empty set of non-negative integers has a least element.

This is not true about negative integers - consider the case of infinite sets, like the set of all integers. There is no well defined least element.

**LEMMA 1.2.** The well ordering principle is equivalent to the principle of mathematical induction.

**PROOF.** First, let us prove that WOP  $\Rightarrow$  PMI. Let  $P(n)$  be a statement that depends on  $n \in \mathbb{N}$ . Suppose that:

- $P(1)$  is true
- $P(k)$  is true implies  $P(k+1)$  is true for all  $k \in \mathbb{N}$ .

We have to show that  $P(n)$  is true for all  $n \in \mathbb{N}$ . Let :

$$S = \{n \in \mathbb{N} : P(n) \text{ is true}\}$$

This means we must show that  $S = \mathbb{N}$ . Let  $T := \mathbb{N} \setminus S$ , i.e.  $T$  is the complement. Let us assume that  $S \neq \mathbb{N}$ .

By WOP,  $T$  has a least element, say  $m$ . Note that  $m \geq 2$  since  $1 \in S$ . Then,  $m-1 \notin T$  and  $m-1 \in S$ . As such,  $P(m-1)$  must be true! However, by our initial assumptions, that would mean  $P(m)$  is true as well, so  $m \in S$ . This creates a contradiction, since  $m \in T$ . Hence,  $S = \mathbb{N}$ .

Now, let us prove that PMI  $\Rightarrow$  WOP.

Consider the statement  $P(n)$  that every non empty set of non-negative integers of size  $n$  has a least element. It is clear that the base case  $P(1)$  is true. Now, let us assume that  $P(k)$  is true - what can we say about  $P(k+1)$ . When we insert an element, we have two cases:

- (1) The inserted element is less than the least element. In this case, there is a new least element, and  $P(k+1)$  is true.

- (2) The inserted element is not less than the least element. In this case, the least element is the same, and  $P(k+1)$  is true.

Hence, by PMI, we can say that  $P(n)$  is true  $\forall n \in \mathbb{N}$ , i.e., WOP is true.

Since  $\text{PMI} \Rightarrow \text{WOP}$  and  $\text{WOP} \Rightarrow \text{PMI}$ ,  $\text{PMI} \Leftrightarrow \text{WOP}$ . □

#### 4. Binomial Theorem

**THEOREM 1.3** (Binomial Theorem). Let  $x, y \in \mathbb{C}$  and let  $n \in \mathbb{N}$ , then

$$(x + y)^n = \sum_{k=0}^n \binom{n}{k} x^k y^{n-k}$$

**COROLLARY 1.3.1.**

$$\sum_{k=0}^n \binom{n}{k} = 2^n$$

**LEMMA 1.4** (Pascal's Identity).

$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$$

**LEMMA 1.5.**

$$\sum_{k=0}^{\lfloor n/2 \rfloor} \binom{n-k}{k} = F_n$$

#### 5. Pigeonhole Principle

**THEOREM 1.6.** If  $n$  items are put into  $m$  containers, with  $n > m$ , then at least one container must contain more than one item.





## CHAPTER 2

### Division

#### 1. Division Algorithm

**THEOREM 2.1.** Let  $a, b \in \mathbb{Z}$  with  $b > 0$ . Then, there exist unique integers  $q$  and  $r$  such that  $a = bq + r$ ,  $r \in [0, b)$ .

**PROOF.** Let  $S = \{a - bn : n \in \mathbb{Z}, a - bn \geq 0\}$ . This set is always non-empty:

- If  $a \geq 0$ , then  $a \in S$
- If  $a < 0$ , then if  $n = a$ , we have  $a - ab \in S$  since  $b \geq 1$ .

By WOP,  $S$  has a least element, say  $r$ . So, there exists  $q \in \mathbb{Z}$  such that  $r = a - bq$ . Since  $r \in S$ , we have  $r \geq 0$ .

Suppose  $r \geq b$ . Then:

$$\begin{aligned} a - b(q + 1) &= a - bq - b = r - b \geq 0 \\ \Rightarrow a - b(q + 1) &\in S \\ \Rightarrow r - b &\in S \end{aligned}$$

However,  $r - b < r$ , and  $r$  is the least element! This gives us a contradiction. So,  $r < b$ .

As such, we have proved the existence of this solution. Now we must prove it's uniqueness.

Suppose there exists  $p, r, q', r'$ , such that:

$$\begin{aligned} a &= bq + r, 0 \leq r < b \\ a &= bq' + r', 0 \leq r' < b \end{aligned}$$

Assume WLOG  $q \geq q'$ . Now,

$$r' - r = b(q - q')$$

If  $q > q'$ , then  $r' - r \geq b$ . However,  $r' - r < b$ . So, this is a contradiction, and  $q' = q$ . The solution must be unique.

□

DEFINITION 1. If  $a, b \in \mathbb{Z}$ , we say that  $a$  divides  $b$  if  $b = ak$  for some  $k \in \mathbb{Z}$ . This is denoted by  $a|b$

Some properties of division are:

- If  $a|b$ , then  $\pm a|\pm b$
- If  $a|b$  and  $b|c$  then  $a|c$  (Transitivity)
- If  $a|b$  and  $a|c$  then  $a|bx + cy$  (Linear Combination)
- If  $a|b$  and  $b \neq 0$ , then  $|a| \leq |b|$  (Bounds by divisibility)
- $a|b$  and  $b|a$ , then  $b = \pm a$ .

EXERCISE. Prove that  $x^a - 1|x^b - 1 \Leftrightarrow a|b$ .

SOLUTION. First, let us prove that if  $a|b$ , then  $x^a - 1|x^b - 1$ . Let  $b = qa$ . Then,

$$x^b - 1 = (x^a)^q - 1^q = (x^a - 1)((x^a)^{q-1} + \dots + x^a + 1)$$

So,  $x^b - 1|x^a - 1$ . Now to prove the converse. Let  $b = aq + r$ . Assume  $a \nmid b$ , then  $0 < r < a$ . Then,

$$x^b - 1 = x^b - x^r + x^r - 1 = x^r(x^{aq} - 1) + x^r - 1$$

$x^a - 1|x^{qa} - 1$ , so  $x^r - 1$  is the remainder. Since  $r < a$ ,  $x^a - 1 \nmid x^r - 1$ . This would mean that  $x^a - 1 \nmid x^b - 1$ , which is a contradiction. So,  $r = 0$ . Hence proved. ■

## 2. Base b representations

THEOREM 2.2. Let  $b \in \mathbb{N}$  with  $b \geq 2$ . Then every positive integer can be expressed uniquely as

$$N = a_k b^k + \dots + a_1 b + a_0$$

where  $k \geq 0$ ,  $a_k \neq 0$  and  $0 \leq a_i < b$  for  $i = 0, \dots, k$ . This is denoted by  $N = (a_k, \dots, a_1 a_0)_b$

PROOF. By the division algorithm, there exist unique integers  $q_0$  and  $a_0$  such that:

$$N = q_0 b + a_0, a_0 \in [0, b)$$

Note that  $q_0 < N$ . If  $q_0 \neq 0$  we apply the division algorithm again to find unique integers  $q_1$  and  $a_q$  such that:

$$q_0 = q_1b + a_1, a_1 \in [0, b)$$

Then,

$$N = (q_1b + a_1)b + a_0 = q_1b^2 + a_1b + a_0$$

We continue till we get a quotient  $q_k = 0$ . This will terminate since  $q_k < \dots < q_2 < q_1 < q_0 < N$ , forming a decreasing sequence of non-negative integers and eventually reaching zero. From this, we get:

$$N = a_kb^k + \dots + a_1b + a_0$$

Hence, the solution always exists.

Suppose  $N$  has two distinct expansions. We can write it as:

$$\begin{aligned} N &= a_kb^k + \dots + a_1b + a_0 \\ &= c_kb^k + \dots + c_1b + c_0 \end{aligned}$$

where  $0 \leq a_i, c_j < b$  for all  $i, j$ . Let  $d_i = a_i - c_i$ . Then,  $\sum_{i=0}^k d_ib^i = 0$ . The  $d_i$  cannot all be zero as the two expansions are assumed distinct. Let  $j$  be the least integer,  $0 \leq j \leq k$ , such that  $d_j \neq 0$ . Then,  $\sum_{i=j}^k d_ib^i = 0$ . Dividing by  $b^j$ , we find that  $\sum_{i=j}^k d_ib^{i-j} = 0$ . Thus,

$$d_j + b \left( \sum_{i=j+1}^k d_ib^{i-j-1} \right) = 0$$

This implies that the  $b|d_j$  and since  $d_j \neq 0$ , we get that  $b = |b| \leq |d_j|$ . However,  $|d_j| < b$ . Hence, we have a contradiction, and the two expansions cannot be distinct. Hence, the solution is also always unique.  $\square$

LEMMA 2.3. If  $N = (a_k \dots a_1 a_0)_b$ , then:

$$\begin{aligned} bN &= (a_k \dots a_1 a_0 0)_b \\ \left\lfloor \frac{N}{b} \right\rfloor &= (a_k \dots a_1)_b \end{aligned}$$

This is a trivial result, which can be thought of as a left or right bitwise shift.

LEMMA 2.4 (Particular case of Legendre's formula). Let  $n \in \mathbb{N}$  and let  $e$  denote the highest power of 2 dividing  $n!$ . Then

$$e = \sum_{k=1}^{\infty} \left\lfloor \frac{n}{2^k} \right\rfloor$$

This is always a finite sum. This can alternatively be expressed as, if  $n = (a_k \dots a_1 a_0)_2$ , then:

$$e = n - (a_k + \dots + a_1 + a_0)$$

PROOF. It is clear that  $e$  is the sum of the no. of positive multiples of  $2^i$  which are  $\leq n$ , for all  $i$ . So, this can be calculated by:

$$e = \sum_{k=1}^{\infty} \left\lfloor \frac{n}{2^k} \right\rfloor$$

□

Thus, if  $r$  denotes the number of ones in the binary expansion of  $n$ , then  $2^{n-r}$  is the highest power of 2 dividing  $n!$ . Further,

- $2^n \nmid n!$  for  $n \in \mathbb{N}$
- $2^{n-1} \mid n!$  if and only if  $n$  is a power of 2.

## CHAPTER 3

### Properties of Numbers

#### 1. Prime and Composite Numbers

DEFINITION 2. A positive integer  $p > 1$  is called prime if its only positive divisors are 1 and  $p$ . A positive integer which is not prime is called composite.

The number 1 is neither prime nor composite.

LEMMA 3.1. Every integer  $n \geq 2$  has a prime factor.

PROOF. Let  $P(n)$  be the statement that  $n$  has a prime factor. Then  $P(2)$  is true, since 2 is a prime factor of 2. Let  $k \geq 2$ . Assume  $P(2) \dots P(k)$  are true.

If  $k + 1$  is prime, then  $k + 1$  is a prime factor of itself. So  $P(k + 1)$  is true.

If  $k + 1$  is composite, then there exists  $d \in [2, k]$  such that  $d | k + 1$ . By the induction hypothesis,  $d$  has a prime factor  $p$ . Since  $p | d$  and  $d | k + 1$ ,  $p | k + 1$ . So  $p$  is a prime factor of  $k + 1$ , and  $P(k + 1)$  is true. By PSI,  $P(n)$  is true for all  $n \geq 2$ .  $\square$

THEOREM 3.2 (Euclid). There are infinitely many primes.

PROOF. Suppose there are finitely many primes  $p_1, \dots, p_k$ . Let

$$N = p_1 \dots p_k + 1$$

Since  $N \geq 2$ , it must have a prime factor. Hence, there exists  $i \in [1, k]$  such that  $p_i | N$ . Since  $p_i | N$  and  $p_i | p_1 p_2 \dots p_k$ , we get that  $p_i | N - p_1 p_2 \dots p_k$ , i.e.,  $p_i | 1$ . However,  $p_i \geq 2$ , which gives us a contradiction. So, there must be infinitely many primes.  $\square$

EXERCISE. For  $n \geq 1$ , let  $p_n$  be the  $n$ th prime. Prove that

$$p_n \leq 2^{2^{n-1}}$$

SOLUTION. Let  $P(n)$  be the statement that  $p_n \leq 2^{2^{n-1}}$ . It is clear that this is true for the base case  $P(1)$ . Let us assume that  $P(1), \dots, P(k)$  is true for  $k \geq 1$ . We observed in

Euclid's proof that  $p_1 \dots p_k + 1$  is not divisible by any of  $p_1 \dots p_k$ . Hence if  $p_i$  denotes a prime factor of  $p_1 \dots p_k + 1$ , then  $i \geq k + 1$ .

$$p_{k+1} \leq p_i \leq p_1 \dots p_k + 1$$

Using the inductive hypothesis, we find that

$$\begin{aligned} p_{k+1} &\leq p_1 \dots p_k + 1 \leq 2 \cdot 2^2 \cdot 2^{2^2} \dots 2^{2^{k-1}} + 1 \\ &= 2^{\sum_{j=0}^{k-1} 2^j} + 1 = 2^{2^k - 1} + 1 \leq 2^{2^k} \end{aligned}$$

So,  $P(k + 1)$  is true. So, by PSI, the result has been proven. ■

EXERCISE. Prove that there are infinitely many primes of the form  $6n + 5$ .

SOLUTION. Suppose that there are finitely many primes of the form  $6n + 5$ , called  $p_1, \dots, p_n$ . Consider  $q = 6p_1 \dots p_n + 5$ .

For our first case, consider that  $q$  is a prime. This would mean there is a new prime of the form  $6n + 5$ , which would be a contradiction.

Now consider that  $q$  is a composite number. All odd primes besides 3 are of the form  $6n + 1$  or  $6n + 5$ . We can prove (by checking) that the product of two numbers of the form  $6n + 1$  will only be of the form  $6n + 1$ . Hence, at least one of the prime factors of  $q$  must be of the form  $6n + 5$ . However, the primes  $p_i$  cannot divide  $q$ . Hence, this is a contradiction, since there must be a new prime. ■

DEFINITION 3. The product of the first  $n$  prime numbers is called the  $n^{\text{th}}$  primorial and is denoted by  $p_n\#$ .

DEFINITION 4. Euclid numbers are integers of the form  $E_n = p_n\# + 1$ .

All Euclid numbers are not primes -  $E_6$  is not a prime!

THEOREM 3.3. Every composite number  $n$  has a prime factor  $\leq \lfloor \sqrt{n} \rfloor$

PROOF. Since  $n$  is composite, there exists integers  $k, l \in (1, n)$  such that

$$n = kl$$

If  $k > \sqrt{n}$  and  $l > \sqrt{n}$  then  $kl > n$ , which is false. So, one of them must be less than or equal to  $\sqrt{n}$ . □

So, if  $n > 1$  has no prime factors  $\leq \lfloor \sqrt{n} \rfloor$ , then  $n$  is prime. We can use this as a test of primality.

It is faster to do this using the Sieve of Eratosthenes. Using this, we can test primality of the first  $n$  integers in  $O(n \log \log n)$  instead of  $O(n\sqrt{n})$ . This is a pretty well known algorithm so it's left to the reader to see it on cp-algorithms.

**THEOREM 3.4.** There is no non-constant polynomial  $f(x)$  with integer coefficients such that  $f(n)$  is prime for all integer  $n$ .

**PROOF.** Suppose such a polynomial  $f(x)$  exists:

$$f(x) = a_k x^k + \dots + a_1 x + a_0, k \geq 1, a_k \neq 0$$

Let  $b \in \mathbb{Z}$ . Then  $f(b)$  is a prime number, say  $p$ . Let  $t \in \mathbb{Z}$ . We have:

$$\begin{aligned} f(b+tp) &= a_k(b+tp)^k + \dots + a_1(b+tp) + a_0 \\ &= (a_k b^k + \dots a_1 b + a_0) + p \cdot g(t) \\ &= f(b) + p \cdot g(t) = p(1 + g(t)) \end{aligned}$$

where  $g(t)$  is a polynomial in  $t$ . Since  $p|f(b+tp)$  and it must be prime, so  $f(b+tp) = p$ . This implies that  $f$  assumes the value  $p$  infinitely many times. This is a contradiction, since a polynomial of degree  $k$  cannot assume the same value  $k$  times.  $\square$

## 2. Prime Counting function

Let  $x$  be a positive real number. We define :

$$\pi(x) = \sum_{p \leq x} 1$$

where  $p$  denotes a prime. So  $\pi(x)$  counts the number of primes  $\leq x$ . This is called the prime counting function.

**THEOREM 3.5 (Prime Number Theorem).**

$$\lim_{x \rightarrow \infty} \frac{\pi(x)}{x / \log x} = 1$$

This essentially states that  $\pi(x) \sim \frac{x}{\log x}$ . The proof is too complicated to be covered in this course.

### 3. Gaps between Primes

The following lemma states that we can find a gap between primes of any arbitrary length.

LEMMA 3.6. For every  $n \in \mathbb{N}$ , there are  $n$  consecutive integers that are all composite.

PROOF. Consider the numbers:

$$(n+1)! + 2, (n+1)! + 3, \dots, (n+1)! + (n+1)$$

It is clear that for  $n \geq 1$ ,  $2|(n+1)! + 2$ . However,  $(n+1)! + 2 \neq 2$ . So,  $(n+1)! + 2$  cannot be prime, and must be composite. We can extend this to each of the given numbers, and prove that they are all composite.  $\square$

DEFINITION 5. A pair  $(p, q)$  of primes with  $p < q$  is called a twin prime pair if  $q - p = 2$ .

It is unknown how many twin primes exist. It is conjectured that there are infinitely many twin primes, but this has not yet been proved.

THEOREM 3.7 (Bertrand's Postulate). For every integer  $n \geq 2$ , there is always at least one prime between  $n$  and  $2n$ .

This was verified by Bertrand but proved by Chebyshev. It is sometimes called Chebyshev's theorem. The proof of this result goes beyond the scope of this course.

REMARK. Do not use this result unless mentioned that we can, in the exam .

EXERCISE. Using Bertrand's postulate, prove that for  $n \geq 2$ :

$$p_n < 2^n$$

SOLUTION. Consider the statement  $P(n)$  that  $p_n < 2^n$ . This is true for the base case that  $P(2)$ . Now assume that  $P(k)$  is true. This means:

$$p_k < 2^k$$

From Bertrand's postulate,

$$k < p_k < 2k$$

$$k + 1 \leq p_k$$

We also know from Bertrand's postulate that:

$$k + 1 < p_{k+1} < 2(k + 1)$$



$$p_{k+1} < 2p_k$$

So, from the induction hypothesis,

$$p_{k+1} < 2 \cdot 2^k$$

$$p_{k+1} < 2^{k+1}$$

Hence,  $P(k) \Rightarrow P(k+1)$ . From PMI,  $P(n)$  is true  $\forall n \geq 2$ . ■

EXERCISE. Prove that if  $2^m + 1$  is prime, then  $m = 2^n$  for some  $n$ .

SOLUTION. Here, we use the following lemma - if  $k$  is odd, then  $x^k + 1$  is divisible by  $x + 1$ . Suppose that  $m$  has an odd factor  $k$ . Then, we can express  $m$  as  $kp$ . So,

$$2^{kp} + 1 = (2^p)^k + 1$$

From our lemma, this is divisible by  $2^p + 1$  which is a number other than 1 and itself. This means  $2^m + 1$  cannot be prime if it has an odd factor, and hence  $m$  must be a power of 2. ■

#### 4. Fermat Numbers

DEFINITION 6. Fermat numbers are  $f_n$  such that:

$$f_n = 2^{2^n} + 1$$

LEMMA 3.8 (Recursive definition).

$$f_n = f_{n-1}^2 - 2f_{n-1} + 2$$

This result is obvious from expanding the RHS, so the proof is not given here.

EXERCISE. Prove that  $f_n$ ,  $n \geq 2$ , all end in 7.

SOLUTION. Let  $P(n)$  be the statement that  $f_n$  ends in 7. This is true for our base case  $P(2)$ . Let us assume that  $P(k-1)$  is true, i.e.  $f_k \mod 10 = 7$ . So, by the recursive definition:

$$\begin{aligned} f_k &= f_{k-1}^2 - 2f_{k-1} + 2 \mod 10 \\ &= 7^2 - 2 \cdot 7 + 2 \mod 10 \\ &= 7 \mod 10 \end{aligned}$$

So, by PMI,  $P(n)$  is true for all  $n \geq 2$ . ■

LEMMA 3.9 (Duncan's Identity).

$$f_0 f_1 \dots f_{n-1} = f_n - 2$$

PROOF. Let  $P(n)$  be the statement that this is true for  $f_n$ . This is clearly true for the base case  $P(1)$ . Let us assume  $P(k)$  is true, i.e.

$$f_0 f_1 \dots f_{k-1} = f_k - 2$$

$$f_0 f_1 \dots f_k = f_k(f_k - 2) = f_k^2 - 2f_k = f_{k+1} - 2$$

The above result comes from the recursive definition. Since  $P(k+1)$  follows from  $P(k)$ , by PMI, Duncan's identity is true.  $\square$

THEOREM 3.10. Every prime factor of  $f_n$ ,  $n \geq 2$ , is of the form  $k \cdot 2^{n+2} + 1$ .

PROOF. To be discussed later in the course.  $\square$

This theorem can be helpful to quickly find the primality of  $f_n$ . For instance,  $f_4$  is prime - we can see this by checking all the numbers of the form  $2^6 k + 1$  which are less than  $\sqrt{f_4}$ . This cuts down the search space and makes primality checking faster.

## 5. Fibonacci Numbers

DEFINITION 7. Fibonacci numbers are numbers of the form:

$$F_n = F_{n-1} + F_{n-2}$$

where  $F_1 = 1, F_2 = 1$

LEMMA 3.11.

$$\sum_{i=1}^k F_i = F_{k+2} - 1$$

LEMMA 3.12 (Cassini's Formula).

$$F_{n-1} F_{n+1} - F_n^2 = (-1)^n, n \geq 2$$

The proofs of lemmas 3.11 and 3.12 come directly from induction, so I am not discussing the proof here.

LEMMA 3.13.

$$F_{n+m} = F_m F_{n+1} + F_{m-1} F_n, m \geq 2, n \geq 1$$

PROOF. This is a non-trivial case of induction. Let us fix  $m \in \mathbb{N}$ , and do induction on  $n$ . For  $n = 1$ , the RHS is:

$$F_{m-1} F_1 + F_m F_2 = F_{m-1} + F_m = F_{m+1}$$

This is also true for  $n = 2$ . Assume that the result is true for  $k = 3, 4, \dots, n$ . We want to show that the result is true for  $k = n + 1$ . For  $k = n - 1$ ,

$$F_{m+n-1} = F_{m-1} F_{n-1} + F_m F_n$$

For  $k = n$ ,

$$F_{m+n} = F_{m-1} F_n + F_m F_{n+1}$$

Add both sides, we get:

$$F_{m+n-1} + F_{m+n} = F_{m+n+1} = F_{m-1} F_{n+1} + F_m F_{n+2}$$

Hence Proved. □

## 6. Lucas Numbers

DEFINITION 8. Lucas numbers are numbers  $L_n$  such that:

$$L_n = L_{n-1} + L_{n-2}$$

where  $L_1 = 1, L_2 = 3$ .

THEOREM 3.14 (Binet's formulas). Let  $\alpha = \frac{1+\sqrt{5}}{2}$  and  $\beta = \frac{1-\sqrt{5}}{2}$ .

$$F_n = \frac{\alpha^n - \beta^n}{\alpha - \beta}$$

$$L_n = \alpha^n + \beta^n$$

PROOF. TODO □



## CHAPTER 4

# Greatest Common Divisor and Least Common Multiple

### 1. Greatest Common Divisor

DEFINITION 9. Let  $a, b \in \mathbb{Z}$ , not both zero. The greatest common divisor of  $a$  and  $b$  is the positive integer  $d$  such that:

- $d|a$  and  $d|b$
- If  $c$  is a positive integer such that  $c|a$  and  $c|b$ , then  $c \leq d$ .

This is generally denoted by  $(a, b)$ .

The GCD of two non-zero numbers always exists and is unique. Observe that  $(a, b) = (a, -b) = (-a, b) = (-a, -b)$ . If  $a \neq 0$ , then  $(a, 0) = |a|$ .

DEFINITION 10.  $a, b \in \mathbb{Z}$  are relatively prime (coprime) if  $(a, b) = 1$ .

EXERCISE. Prove that  $(F_n, F_{n+1}) = 1$  for  $n \geq 1$ .

SOLUTION. From Cassini's formula, we know that:

$$F_{n-1}F_{n+1} - F_n^2 = (-1)^n$$

Let  $d = (F_n, F_{n+1})$ . Since  $d|F_n$  and  $d|F_{n+1}$ , we have  $d|F_{n-1}F_{n+1} - F_n^2$ . So  $d|(-1)^n$ . This means that  $|d| \leq |(-1)^n|$ . Since  $d$  is a positive number,  $d = 1$ . ■

EXERCISE. Prove that  $(f_m, f_n) = 1$  for distinct non-negative  $m, n$ .

SOLUTION. Suppose  $m < n$ . Let  $d = (f_m, f_n)$ . Since  $d|f_m$  and  $d|f_n$ ,  $d|f_n - (f_0 \dots f_m \dots f_{n-1})$ . From Duncan's identity, this implies that  $d|2$ . Since  $d > 0$ ,  $d = 1$  or  $d = 2$ . However, Fermat numbers are always odd - so  $d \neq 2$ . This implies that  $d = 1$ . ■

THEOREM 4.1. Let  $a, b \in \mathbb{Z}$ , not both zero. Then there exist integers  $x_0, y_0$  such that:

$$(a, b) = ax_0 + by_0$$

PROOF. Consider the set  $S = \{ax + by > 0 : a, b \in \mathbb{Z}\}$ . Let  $d = \min S$ . Suppose that  $d$  does not divide  $a$ . Then by division algorithm:

$$\begin{aligned} a &= qd + r \\ qd &= a - r \\ q(ax + by) &= a - r \\ r &= a(1 - qx) - bqy \end{aligned}$$

So,  $r$  is a linear combination of  $a$  and  $b$ , and since  $r > 0$ ,  $r \in S$ . From division algorithm,  $r < d$ , which contradicts the fact that  $d = \min S$ . So, by contradiction,  $d$  divides  $a$  (and by similar argument,  $d$  divides  $b$ ). We also know that any common divisor of  $a$  and  $b$  must divide  $d$ . This is obvious since if  $a = uc$  and  $b = vc$ , then  $d = ax + by = c(ux + vy)$ , so  $c|d$ . From these two facts, it is clear that  $d$  is the GCD, and is of the form  $ax + by$ .  $\square$

EXERCISE. Let  $a, b \in \mathbb{N}$ . If  $b = aq + r$ , then  $(a, b) = (a, r)$ .

SOLUTION. Let  $d = (a, b)$  and  $e = (b, r)$ . We need to show that  $d = e$ . Since  $d|a$  and  $d|b$ ,  $d|a - bq = r$ . So  $d$  is a common divisor of  $b$  and  $r$ . Hence  $d \leq e$ . Similarly, as  $e|b$  and  $e|r$ ,  $e$  divides  $bq + r = a$ . Thus  $e$  is a common divisor of  $a$  and  $b$ , so  $e \leq d$ . Thus,  $e = d$ .  $\blacksquare$

EXERCISE. Let  $a, b, c \in \mathbb{N}$ . Prove that  $(ac, bc) = c(a, b)$ .

SOLUTION. Let  $d = (a, b)$ . Then  $d|a$  and  $d|b$ , so  $d|ca$  and  $d|cb$ . There exist integers  $x, y$  such that:

$$\begin{aligned} d &= ax + by \\ cd &= (ac)x + (bc)y \end{aligned}$$

If  $e$  is a positive integer such that  $e|ac$  and  $e|bc$ , then  $e|(ac)x + (bc)y$ , i.e.  $e|cd$ . So,  $cd$  is the GCD of  $ac$  and  $bc$ .  $\blacksquare$

THEOREM 4.2. Let  $a, b \in \mathbb{Z}$ , not both zero. Then  $(a, b) = 1$  if and only if there exist integers  $x_0, y_0$  such that  $ax_0 + by_0 = 1$ .

PROOF. If  $(a, b) = 1$ , there must exist  $x_0, y_0$  such that  $ax_0 + by_0 = 1$ , from Theorem 4.1. Conversely, suppose there exists  $x_0, y_0 \in \mathbb{Z}$ , such that

$$ax_0 + by_0 = 1$$

Let  $d = (a, b)$ . Then  $d|ax_0 + by_0$ , which means that  $d|1$ . Since  $d \in \mathbb{N}$ ,  $d = 1$ .  $\square$

COROLLARY 4.2.1. Let  $d = (a, b)$ . Then,  $(\frac{a}{d}, \frac{b}{d}) = 1$

COROLLARY 4.2.2. If  $(a, b) = 1$ , and  $a$  and  $b$  both divide  $c$ , then  $ab|c$ .

THEOREM 4.3 (Euclid's Lemma). If  $a|bc$  and  $(a, b) = 1$ , then  $a|c$ .

PROOF.

$$ax + by = 1$$

$$acx + bcy = c$$

Since  $a|acx$  and  $a|bcy$ , so  $a|c$ . □

EXERCISE. Let  $m, n \in \mathbb{N}, m > 2$ . If  $F_m|F_n$ , prove that  $m|n$ .

SOLUTION. We know that:

$$F_n = F_{n-m}F_{m-1} + F_{n-m+1}F_m$$

Since  $F_m|F_n$  and  $F_m|F_{n-m+1}F_m$ , we get  $F_m|F_n - F_{n-m+1}F_m$  and hence  $F_m|F_{n-m}F_{m-1}$ . But, we also know that  $(F_m, F_{m-1}) = 1$ . By Euclid's Lemma,  $F_m|F_{n-m}$ .

From the division algorithm, let  $n = mq + r$ . Suppose  $r > 0$ . From our previous result,  $F_m|F_{n-m}$ , so  $F_m|F_{n-2m} \dots F_m|F_r$ . This means that  $F_m \leq F_r$ . But  $r < m$ , so  $F_r < F_m$ . This is a contradiction. So  $r = 0$ . Hence proved. ■

DEFINITION 11. Let  $n \geq 2$  and let  $a_1, \dots, a_n \in \mathbb{Z}$ , not all zero. The GCD of  $a_1, \dots, a_n$  is the largest positive integer that divides each  $a_i$ . This is denoted by  $a_1, \dots, a_n$ .

This has the following properties:

- $(a_1, a_2, \dots, a_n)$  is the least positive integer that is a linear combination of  $a_1, \dots, a_n$ .
- $(a_1, \dots, a_n) = ((a_1, \dots, a_{n-1}), a_n)$
- If  $d|a_1, \dots, a_n$  and  $(d, a_i) = 1$  for all  $i \in [1, n-1]$ , then  $d|a_n$ .
- If  $a_1, \dots, a_n$  are pairwise relatively prime, then  $(a_1, \dots, a_n) = 1$ .

## 2. Euclidean Algorithm

We are given  $a, b \in \mathbb{Z}$ , not both zero, and want to compute  $(a, b)$ . The algorithm to do this works as follows:

- (1) If  $a$  or  $b$  are negative, replace with their absolute value.
- (2) If  $a > b$ , then swap  $a$  and  $b$ .
- (3) If  $a = 0$ , then  $(a, b) = b$ .
- (4) If  $a > 0$ , write  $b = aq + r$ . Then,

$$(a, b) = (r, a)$$

Go to step 3 with  $a = r$  and  $b = a$  respectively.

This is called the Euclidean Algorithm.

To express  $(a, b)$  as a linear combination of  $a$  and  $b$  where  $0 \leq a \leq b$ , we create a table with four columns with headings  $x, y, r, q$ . We denote the rows as  $R_{-1}, R_0, R_1, \dots, R_{i-1}$  and the entries in  $R_i$  as  $x_i, y_i, r_i, q_i$ .  $R_{-1} = (0, 1, b, 0)$  and  $R_0 = (1, 0, a, 0)$ . Suppose we have filled till  $R_{i-1}$  for some  $i \geq 1$ . To fill  $R_i$ , we first compute  $q_i$ , which is the quotient obtained on dividing  $r_{i-2}$  by  $r_{i-1}$ . Next,  $R_i = R_{i-2} - q_i R_{i-1}$ . We continue this until  $r = 0$ . Then, the GCD is the value of  $r$  in the row before, and the coefficients are the values of  $x$  and  $y$  in the row before. This is known as the Extended Euclidean Algorithm.

**THEOREM 4.4** (Lame's theorem). Let  $b \geq a \geq 2$ . The number of divisions required to compute  $(a, b)$  by the Euclidean algorithm is at most 5 times the number of decimal digits in  $a$ .

**PROOF.** Suppose that  $a$  contains  $k$  decimal digits and takes  $n$  divisions to compute  $(a, b)$ . We need to show that  $n \leq 5k$ . Let  $r_0 = b, r_1 = a$ . Applying Division algorithm repeatedly, we have:

$$r_0 = r_1 q_1 + r_2, 0 < r_2 < r_1$$

$$r_1 = r_2 q_2 + r_3, 0 < r_3 < r_2$$

$\dots$

$$r_{n-1} = r_n \cdot q_n + 0$$

We can prove that  $q_i \geq 1$  for  $1 \leq i \leq n-1$  and  $q_n \geq 2$ . This is because if  $q_n$  is 1, then  $r_{n-1} = r_n$ , which is a contradiction. If  $q_i = 0$ , then  $r_{i-1} = r_{i+1}$ , which is also a contradiction.



We claim that  $r_{n-i} \geq F_{i+2}$  for  $1 \leq i \leq n-1$ . This is true for the base case, where  $r_{n-1} \geq F_3 = 2$ .

$$\begin{aligned} r_{n-j} &= r_{n-(j-1)}q_{n-(j-1)} + r_{n-(j-2)} \\ &\geq r_{n-(j-1)} + r_{n-(j-2)} \\ &\geq F_{j+1} + F_j = F_j + 2 \end{aligned}$$

In particular,  $a = r_1 \geq F_{n+1}$ . Now  $10^k > a \geq F_{n+1}\alpha^{n-1}$ , where  $\alpha = \frac{1+\sqrt{5}}{2}$  and  $n \geq 3$ . Taking logarithms and using the fact that  $\log \alpha > 1/5$ , we get that  $n \leq 5k$ .  $\square$

EXERCISE. Prove that for all  $a, m, n \in \mathbb{N}$ ,

$$(a^n - 1, a^m - 1) = a^{(n,m)} - 1$$

SOLUTION. Let  $f_n = a^n - 1$ . Our goal is to prove that  $(f_n, f_m) = f_{(n,m)}$ .

$$\begin{aligned} f_n &= a^n - 1 \\ &= a^{n-m}(a^m - 1) + a^{n-m} - 1 \\ &= f_{n-m} + a^{n-m} - 1 \end{aligned}$$

So now, when performing Euclid's algorithm to find the GCD,  $(f_n, f_m) = (f_{n-m}, f_m)$ . This occurs recursively, and as we can see it is also performing Euclid's algorithm in the subscript. Hence,  $(f_n, f_m) = f_{(n,m)}$ .  $\blacksquare$

### 3. Least Common Multiple

DEFINITION 12. Let  $a, b \in \mathbb{N}$ . We say that a positive integer  $l$  is the least common multiple of  $a$  and  $b$  if

- $a$  and  $b$  both divide  $l$
- $c$  is a positive integer divisible by both  $a$  and  $b$ , then  $l \leq c$ .

It is generally denoted by  $[a, b]$ .

EXERCISE. Prove that  $(a, b)[a, b] = ab$ .

SOLUTION. Let  $d = (a, b)$  and let  $k = ab/d$ . Obviously,  $k \in \mathbb{N}$ . We have to show that  $k = [a, b]$ . We first prove that  $a$  and  $b$  both divide  $k$ . Write

$$a = da_1, b = db_1, a_1, b_1 \in \mathbb{Z}$$

Then,  $k = \frac{ab}{d} = ab_1 = ba_1$ . Hence  $a$  and  $b$  both divide  $k$ . Suppose  $c$  is a positive integer divisible by both  $a$  and  $b$ . Write

$$c = aa', c = bb', a', b' \in \mathbb{Z}$$

Since  $d = (a, b)$ , there exists integers  $x$  and  $y$  such that:

$$d = ax + by$$

Then

$$\frac{c}{k} = \frac{cd}{ab} = \frac{c(ax + by)}{ab} = \frac{c}{b}x + \frac{c}{a}y = b'x + a'y \in \mathbb{Z}$$

Hence  $k$  divides  $c$ . Since  $c \neq 0$ , this implies  $k \leq c$ . Therefore, by definition,  $k = [a, b]$  and  $(a, b)[a, b] = ab$ . ■

#### 4. Fundamental Theorem of Arithmetic

**THEOREM 4.5** (Fundamental Theorem of Arithmetic). Every integer  $n \geq 2$  can be written as a product of primes and the factorization into primes is unique up to the order of the factors.

**PROOF.** Let  $P(n)$  be the statement that  $n$  can be written as a product of primes.  $P(2)$  is true, since it is a prime. Let  $k \geq 2$  and  $P(2), \dots, P(k)$  is true. If  $k+1$  is prime, then  $P(k+1)$  is immediately true. If it is composite, then  $k+1$  is of the form  $a \cdot b$  where  $2 \leq a, b \leq k$ . By induction hypothesis, since  $P(a)$  and  $P(b)$  is true,  $P(k+1)$  is also true.

Suppose that

$$n = p_1 p_2 \dots p_r = q_1 \dots q_s$$

Assume  $r \leq s$ . Since if  $p_1 | q_1 \dots q_s$ ,  $p_1 = q_i$  for some  $i$ ,  $1 \leq i \leq s$ . We can cancel this out and keep doing this till we get

$$1 = q_{i_1} q_{i_2} \dots q_{i_{s-r}}$$

However,  $q_i$  is a prime. This is a contradiction unless  $s = r$ . Hence uniqueness is proved. □

The largest power  $x$  such that  $p^x | a$  for a prime  $p$  is called the multiplicity of  $p$  in  $a$ .

## CHAPTER 5

### Linear Diophantine Equations

#### 1. Linear Diophantine Equations in Two Variables

Linear Diophantine equations in two variables are of the form:

$$ax + by = c$$

An integer solution exists if and only if  $(a, b) | c$ . In this case, for any initial solution  $x_0, y_0$ , the general solution is given by:

$$x = x_0 + \frac{b}{d}n, y = y_0 - \frac{a}{d}n$$

where  $d = (a, b)$  and  $n \in \mathbb{Z}$ . The initial solution  $x_0, y_0$  can be found using the Extended Euclidean Algorithm.

#### 2. Fibonacci Linear Diophantine Equation

THEOREM 5.1. Let  $k \in \mathbb{N}, c \in \mathbb{Z}$ . The linear Diophantine equation:

$$F_{k+1}x + F_k y = c$$

is always solvable. If  $k$  is even, the complete solution is given by:

$$x = F_{k-1}c + F_k n, y = -F_k c - F_{k+1}n$$

If  $k$  is odd, the complete solution is given by:

$$x = -F_{k-1}c + F_k n, y = F_k c - F_{k+1}n$$

PROOF.  $(F_k, F_{k+1}) = 1$ , so the solution always exists. If  $k$  is even, then:

$$x_0 = F_{k-1}c, y_0 = -F_k c$$

always solves it (prove it via Cassini's Formula). If  $k$  is odd, then:

$$x_0 = -F_{k-1}c, y_0 = F_k c$$

always solves it. □

### 3. Linear Diophantine Equations in Many Variables

THEOREM 5.2. Let  $a_1, \dots, a_k$  be integers, not all zero. The Diophantine equation:

$$a_1x_1 + \dots + a_kx_k = x$$

has an integer solution if and only if  $(a_1, \dots, a_k) | c$ . In that case, there are infinitely many solutions.

EXERCISE. Find the complete solution to:

$$6x + 8y + 12z = 10$$

SOLUTION.  $(6, 8, 12) = 2$ , and  $2 | 10$ , so this equation is solvable. Let us consider the “subequation”  $8y + 12z$ . Since it is a linear combination of 8 and 12, it must be a multiple of  $(8, 12) = 4$ . So,

$$8y + 12z = 4u$$

This gives us a new equation:

$$6x + 4u = 10$$

The general solution of this is given by:

$$x = 1 + 2n, u = 1 - 3n$$

Substituting  $u$  in the first equation,

$$8x + 12y = 4(1 - 3n)$$

This equation is also solvable, and we can find its general solution as well. Since  $4 = 2 \cdot 8 + (-1) \cdot 12$ , we get  $4(1 - 3n) = (2 - 6n)8 + (-1 - 3n)12$ . Finally, we can write the complete solution:

$$x = 1 + 2n$$

$$y = 2 - 6n + 3m$$

$$z = -1 + 3n - 2m$$

■

EXERCISE. Suppose we have a coin system. Let's say we use 4 rupee and 9 rupee coins only. Then which amounts can be exchanged? For instance, 1 rupee cannot be exchanged; 17 rupees can be exchanged.

SOLUTION. This problem can be expressed as - what numbers can be expressed as a linear combination of 4 and 9 with positive coefficients.

In fact, only finitely many integers cannot be exchanged. Generally, if  $p, q \in \mathbb{N}$  with  $(p, q) = 1$ , then the largest integer which cannot be represented as  $ap + bq$  with  $a, b \in \mathbb{Z}, a, b \geq 0$  is  $pq - p - q$ .

Let  $x = ap + bq$ . Since  $(p, q) = 1$ , each multiple of  $q$  leaves a different remainder when divided by  $p$  which does not change when one adds multiples of  $p$  to it. This means that if  $x$  leaves a remainder  $r$  when divided by  $q$ , there has been some  $ap$  that leaves the same remainder. The last remainder to be taken is  $q$  with  $p(q - 1)$ .

*TO FIX*





## CHAPTER 6

# Congruence

### 1. Introduction

DEFINITION 13. Let  $m \in \mathbb{N}$ . Two integers  $a, b$  are said to be congruent modulo  $m$  if  $m|a-b$ . This is denoted by  $a \equiv b \pmod{m}$

Congruence has the following properties:

- $a \equiv a \pmod{m}$
- $a \equiv b \pmod{m}$ , then  $b \equiv a \pmod{m}$
- $a \equiv b \pmod{m}$  and  $b \equiv c \pmod{m}$ , then  $a \equiv c \pmod{m}$

As such, it is an equivalence relation. The set of all integers congruent to  $a$  modulo  $m$  is denoted by  $[a]$ , and is called the congruence class of  $a \pmod{m}$

LEMMA 6.1. Two integers are congruent modulo  $m$  if and only if  $a, b$  have the same remainder upon division by  $m$

PROOF. Suppose  $a$  and  $b$  have the same remainder upon division by  $m$ . Then:

$$a = mq_1 + r$$

$$b = mq_2 + r$$

$$a - b = m(q_1 - q_2) + 0$$

$$m|a - b \Rightarrow a \equiv b \pmod{m}$$

Suppose  $a \equiv b \pmod{m}$ . This means that  $a - b = mk, k \in \mathbb{Z}$ . By division algorithm, there exist unique integers  $q, r$  such that:

$$b = mq + r, 0 \leq r < m$$

$$a = m(q + k) + r$$

Since  $0 \leq r < m$ ,  $r$  is the remainder of  $a$  upon division by  $m$ . So, they have the same remainder. Hence proved.  $\square$

So, every integer  $a$  is congruent to its remainder  $r$  modulo  $m$ . We call  $r$  the least residue of  $a$  modulo  $m$ . Every integer is congruent to exactly one of the least residues  $0, 1, \dots, m-1$  modulo  $m$ . Hence,

$$\mathbb{Z} = \bigcup_{i=0}^{m-1} [i]_m$$

LEMMA 6.2. Suppose  $a \equiv b \pmod{m}$  and  $c \equiv d \pmod{m}$ . Then:

- $a \pm c \equiv b \pm d \pmod{m}$
- $ac \equiv bd \pmod{m}$
- $a^n \equiv b^n \pmod{m}$

LEMMA 6.3. Let  $p$  be a prime and let  $a \in \mathbb{Z}$  with  $(a, p) = 1$ . Then the least residues of:

$$a, 2a, \dots, (p-1)a$$

modulo  $p$  are a permutation of the integers  $1, 2, \dots, p-1$ .

PROOF. Let  $i \in \{1, 2, \dots, p-1\}$ . Suppose  $ia \equiv 0 \pmod{p}$ . Then,  $i \equiv 0 \pmod{p}$ , which is false. So,  $ia$  cannot be congruent to 0 modulo  $p$ .

Let  $i, j \in \{1, 2, \dots, p-1\}, i \neq j$ . Suppose that  $ia \equiv ja \pmod{p}$ . Since  $(a, p) = 1$ ,  $i \equiv j \pmod{p}$ . However,  $|i-j| \leq p-1$ . Then,  $p \mid |i-j|$  if and only if  $i = j$ , but this is a contradiction. Hence,  $ia$  for  $i \in \{1, 2, \dots, p-1\}$  are all incongruent to one another. Hence, they will all have different least residues, none of which are equal to 0. So, the least residues must be a permutation of  $\{1, 2, \dots, p-1\}$ .  $\square$

EXERCISE. Find the remainder when  $1! + 2! + \dots + 100!$  is divided by 15.

SOLUTION. If  $n \geq 5$ , then  $n!$  is divisible by 15. So,  $n! \equiv 0 \pmod{15}$ . Hence, this sum modulo 15 is equivalent to  $1! + 2! + 3! + 4!$  modulo 15. So, the remainder is 3.  $\blacksquare$

EXERCISE. Find the positive integers  $n$  for which  $\sum_{k=1}^n k!$  is a square.

SOLUTION. If  $k \geq 5$ , then  $k! \equiv 0 \pmod{10}$ . So,

$$\sum_{k=1}^n k! \equiv 1! + 2! + 3! + 4! \pmod{10}$$



Squares are congruent to 0,1,4,5,6 or 9 modulo 10. However, for  $n \geq 5$ , this sum is 3 modulo 10. So,  $n$  must be less than 5. We can now try each of these cases by hand. From this, we find  $n = 1$  or  $n = 3$ . ■

DEFINITION 14. A set  $a_1, \dots, a_m$  is said to be a complete set of residues modulo  $m$  if every integer is congruent modulo  $m$  to exactly one of them.

REMARK. A set of  $m$  integers form a complete set of residues modulo  $m$  if and only if no two of the integers are congruent modulo  $m$ .

EXERCISE. Let  $m \in \mathbb{N}$  with  $m \geq 3$ . Prove that the set  $\{1^2, 2^2, \dots, m^2\}$  doesn't form a complete set of residues modulo  $m$ .

SOLUTION. If they do not form a complete set of residues modulo  $m$ , then two of the integers must be congruent modulo  $m$ . Since  $m - 1 \equiv -1 \pmod{m}$ , we have:

$$(m - 1)^2 \equiv 1 \pmod{m}$$

Thus, this set cannot form a complete set of residues modulo  $m$ . ■

EXERCISE. Find all positive integers  $x, y, z$  such that:

$$3^x + 4^y + 5^z$$

SOLUTION. Let us first consider this modulo 4. Then,

$$3^x \equiv 1 \pmod{4}$$

We can prove that this is only possible if  $x$  is even.  $3 \equiv 3 \pmod{4}$  and  $3^2 \equiv 1 \pmod{4}$ . This repeats again and again, and can be proved by induction.

Now let us consider this modulo 4. Then,

$$1 \equiv 2^z \pmod{3}$$

From the same argument above, we get that  $z$  must be even too.

From this, we get a new equation:

$$3^{2x_1} + 2^{2y} = 5^{2z_1}$$

$$2^{2y} = 5^{2z_1} - 3^{2x_1} = (5^{z_1} + 3^{x_1})(5^{z_1} - 3^{x_1})$$

From the fundamental theorem of arithmetic, we know that:

$$5^{z_1} + 3^{x_1} = 2^s, s \geq 0$$

$$5^{z_1} - 3^{x_1} = 2^t, t \geq 0$$

From this set of equations,

$$5^{z_1} = \frac{2^t + 2^s}{2} = 2^{t-1}(2^{s-t} + 1)$$

Since the LHS is odd,  $t = 1$ . So,

$$5^{z_1} = 2^{s-1} + 1 \text{ and } 3^{x_1} = 2^{s-1} - 1$$

By considering the  $3^{x_1}$  modulo 3, we get that:

$$2^{s-1} \equiv 1 \pmod{3}$$

This once again implies that  $s - 1$  is even.

$$3^{x_1} = 2^{2s_1} - 1 = (2^{s_1} - 1)(2^{s_1} + 1)$$

Once again we apply the fundamental theorem of arithmetic, and get the equations

$$2^{s_1} - 1 = 3^a$$

$$2^{s_1} + 1 = 3^b$$

$$3^b - 3^a = 2$$

This implies that  $b = 1, a = 0$ . From this we know that  $s_1 = 1$  and  $s = 3$ . These results also tell us that  $z = 2$  and  $x = 2$ .

$$3^2 + 4^y = 5^2$$

$$4^y = 5^2 - 3^2 = 16$$

Hence, the only positive integers satisfying the equation are  $x = y = z = 2$ . ■

EXERCISE. Show that  $x^2 + 2y^2 = 8z + 5$  has no integral solution.

SOLUTION. This equation is solvable if and only if there  $\exists x, y$  such that

$$x^2 + 2y^2 \equiv 5 \pmod{8}$$

If  $a$  is an integer, we can show that  $a^2 \equiv 0, 1, 4 \pmod{8}$ . So if  $x$  and  $y$  are integers,  $x^2$  is equivalent to 0, 1, 4, and  $2y^2$  is equivalent to 0, 2. The sum cannot be equivalent to 5. Hence, no integral solution exists. ■

EXERCISE. Show that the sequence 71, 771, 7771, ... has no perfect squares.

**SOLUTION.** Remember that a perfect square is equivalent to either 0 or 1 modulo 4. We now prove that the  $i^{\text{th}}$  term  $r_i$  of the given sequence is such that  $r_i \equiv 3 \pmod{4}$ .

Let  $P(n)$  be a statement that the term  $r_i$  is such that  $r_i \equiv 3 \pmod{4}$ . This is true for a  $i = 1$ , since  $71 \equiv 3 \pmod{4}$ . Assume  $P(k)$  is true. We can express  $r_{i+1}$  by the recurrence:

$$\begin{aligned} r_{i+1} &= 10r_i + 61 \\ r_{i+1} &\equiv 10 \cdot 3 + 1 \pmod{4} = 3 \pmod{4} \end{aligned}$$

By PMI,  $P(n)$  is true  $\forall n \geq 1$ . Hence, no member of this sequence may be a perfect square.

As an aside, remember that there is no perfect square that can end with two odd digits. ■

## 2. Square and Multiply Algorithm

Let  $a, l, m$  be integers  $\geq 2$ . We want to compute  $a^l \pmod{m}$ . To do this, we could use the following steps:

- (1) Write the base-2 expansion on  $l = (a_l \cdots a_1 a_0)_2$ , where  $k \geq 0, a_k \neq 0, 0 \leq a_i < 2$ .
- (2) Compute  $a^{2^i} \pmod{m}$  for  $0 \leq i \leq k$ . This can be done in  $O(k)$  via squaring.
- (3) Multiply the  $a^{2^i} \pmod{m}$  for all  $i$  with  $a_i = 1$  to get the result.

## 3. Cancellation in Congruences

**THEOREM 6.4.** Let  $a, b, c, m \in \mathbb{Z}$  with  $m \geq 2$ . Then,

- If  $ac \equiv bc \pmod{m}$  and  $(c, m) = 1$ , then  $a \equiv b \pmod{m}$ .
- Let  $d = (c, m)$ . If  $ac \equiv bc \pmod{m}$  and  $(c, m) = 1$ , then  $a \equiv b \pmod{\frac{m}{d}}$ .

**PROOF.** We have  $m|(a - b)c$ , which implies that  $\frac{m}{d}|(a - b)\frac{c}{d}$ . Since  $(\frac{m}{d}, \frac{c}{d}) = 1$ , we get that  $\frac{m}{d}|(a - b)$ . This is the second result. The first result is a consequence of the second. □

## 4. Combining Congruences with different moduli

THEOREM 6.5. Let  $a, b, m_1, \dots, m_k \in \mathbb{Z}$  with  $m_1, m_2, \dots, m_k \geq 2$ . If

$$a \equiv b \pmod{m_1}$$

$$a \equiv b \pmod{m_2}$$

...

$$a \equiv b \pmod{m_k}$$

then  $a \equiv b \pmod{[m_1, m_2, \dots, m_k]}$

### 5. Linear Congruences

Let  $a, b, m \in \mathbb{Z}, m \geq 2$ . Then  $ax \equiv b \pmod{m}$  is called a linear congruence. Our goal is to find possible  $x$  that satisfy the given congruence.

THEOREM 6.6. Let  $a, b, m \in \mathbb{Z}$  and  $d = (a, m)$ . The linear congruence  $ax \equiv b \pmod{m}$  has a solution if and only if  $d|b$ . If  $d|b$ , then there are  $d$  mutually incongruent solutions modulo  $m$ .

PROOF. The linear congruence  $ax \equiv b \pmod{m}$  can be expressed as a linear Diophantine equation:

$$ax = b + my$$

$$ax - my = b$$

Hence, it has a solution if and only if that LDE is solvable. This means that the linear congruence is also only solvable if and only if  $(a, m)|b$ .

Suppose  $d|b$ . Then, there are infinitely many solutions to the LDE. If  $(x_0, y_0)$  is a particular solution, then the complete solution is given by

$$x = x_0 + \frac{m}{d}n, y = y_0 + \frac{a}{d}n$$

Then, we will get the solutions:

$$x_0, x_0 + \frac{m}{d}, x_0 + 2\frac{m}{d}, \dots, x_0 + d\frac{m}{d}, \dots$$

Here we can see that after that last term, all the solutions will be congruent to one of the first  $d$  terms. Hence, if the first  $d$  solutions are incongruent, we will have proved our theorem.

Assume

$$\begin{aligned} x_0 + \frac{m}{d}n_1 &\equiv x_0 + \frac{m}{d}n_2 \pmod{m} \\ \frac{m}{d}n_1 &\equiv \frac{m}{d}n_2 \pmod{m} \\ n_1 &\equiv n_2 \pmod{d} \text{ (From Theorem 6.3)} \end{aligned}$$

This means that two solutions will only be congruent if  $n_1$  and  $n_2$  are congruent modulo  $d$ . So,  $ax \equiv b \pmod{m}$  has  $d$  mutually incongruent solutions:

$$x_0, x_0 + \frac{m}{d}, x_0 + 2\frac{m}{d}, \dots, x_0 + (d-1)\frac{m}{d}$$

□

## 6. System of Linear Congruences

Let  $a_1, \dots, a_k \in Z$ . Let  $n_1, \dots, n_k$  be integers  $\geq 2$ . We want to find integers  $x$  that satisfy the system of linear congruences

$$x \equiv a_1 \pmod{n_1}$$

...

$$x \equiv a_k \pmod{n_k}$$

**THEOREM 6.7** (Chinese Remainder Theorem). If  $n_1, \dots, n_k$  are pairwise coprime, then the system:

$$x \equiv a_1 \pmod{n_1}$$

...

$$x \equiv a_k \pmod{n_k}$$

has a unique solution modulo  $n_1n_2 \cdots n_k$

**PROOF.** Let  $n = n_1n_2 \cdots n_k$ . For  $i = 1, \dots, k$  let  $N_i = \frac{n}{n_i}$ . Note that  $(N_i, n_i) = 1$ . Let the unique solution of the congruence

$$N_i x \equiv 1 \pmod{n_i}$$

be  $x_i$ .

Let us prove that  $x = a_1N_1x_1 + \cdots + a_kN_kx_k$  satisfies the given system of congruences. First, let us prove that it satisfies the first congruence. Since  $N_i \equiv 0 \pmod{n_1}$  for  $i = 2, \dots, k$ , we get:

$$x \equiv a_1N_1x_1 + 0 \pmod{n_1}$$

We have already assumed that  $N_1x_1 \equiv 1 \pmod{n_1}$ . So,

$$x \equiv a_1 \pmod{n_1}$$

Hence proved. The same argument can be made for all the congruences.

Now let us prove that the solution is unique. Suppose  $x, y$  satisfy the system of congruences. Then

$$x \equiv a_1 \equiv y \pmod{n_1}$$

$$\dots$$

$$x \equiv a_1 \equiv y \pmod{n_k}$$

$$x \equiv y \pmod{[n_1, n_2, \dots, n_k]}$$

Since  $(n_i, n_j) = 1$ ,  $x \equiv y \pmod{n_1n_2\dots n_k}$ . Hence the solution is unique.  $\square$

EXERCISE. Solve the following system of congruences:

$$x \equiv 11 \pmod{12}$$

$$x \equiv 1 \pmod{5}$$

$$x \equiv 0 \pmod{7}$$

SOLUTION.

$$n = 12 \times 5 \times 7 = 420$$

$$N_1 = \frac{n}{12} = 35$$

$$N_2 = 84$$

$$N_3 = 60$$

First, let us solve  $N_1x \equiv 1 \pmod{n_1}$ . Then,

$$35x \equiv 1 \pmod{12}$$

$$x \equiv -1 \pmod{12}$$

Let  $x_1 = -1$ . The same way, we find  $x_2 = -1$  and  $x_3 = 0$ . So,

$$x = 11 \times 35 \times (-1) + 1 \times 84 \times (-1) + 0$$

$$x \equiv 371 \pmod{420}$$

■

## 7. Modular Inverse

DEFINITION 15. Let  $p$  be a prime and let  $a \in \mathbb{Z}$  with  $(a, p) = 1$ . Then  $a$  is invertible modulo  $p$  and its inverse is a number  $x$  such that:

$$a \cdot x \equiv 1 \pmod{p}$$

LEMMA 6.8.  $a$  is self-invertible modulo  $p$  if and only if  $a \equiv \pm 1 \pmod{p}$ .

PROOF. By definition of being “self invertible”,

$$a^2 \equiv 1 \pmod{p}$$

$$(a - 1)(a + 1) \equiv 0 \pmod{p}$$

So,  $(a - 1) \equiv 0 \pmod{p}$  or  $(a + 1) \equiv 0 \pmod{p}$ . From this,  $a \equiv \pm 1 \pmod{p}$ . □

THEOREM 6.9 (Wilson’s Theorem). If  $p$  is a prime number, then  $(p - 1)! \equiv -1 \pmod{p}$

PROOF. If  $p = 2$ , then  $(p - 1)! = 1$ , which is  $-1 \pmod{p}$ .

Suppose  $p > 2$ . We group the  $p - 3$  integers, namely  $2, \dots, p - 2$  into pairs of inverses  $(a, b)$  such that  $ab \equiv 1 \pmod{p}$ . Hence,

$$(p - 1)! \equiv 1 \cdot 2 \cdot 3 \cdots (p - 2)(p - 1) \equiv (p - 1) \equiv -1 \pmod{p}$$

□

The following are some consequences of Wilson’s theorem:

- The product of  $p - 1$  integers between any two consecutive multiples of  $p$  is congruent to  $-1$  modulo  $p$ .
- Let  $p$  be a prime and  $0 \leq r \leq p - 1$ . Then :

$$r!(p - 1 - r)! + (-1)^r \equiv 0 \pmod{p}$$

EXERCISE. Let  $p$  be a prime and let  $n \in \mathbb{N}$ . Prove that

$$\frac{(np)!}{n!p^n} \equiv (-1)^n \pmod{p}$$

SOLUTION. TODO ■

THEOREM 6.10 (Fermat's Little Theorem). Let  $p$  be a prime and let  $a \in \mathbb{Z}$  with  $(a, p) = 1$ . Then,

$$a^{p-1} \equiv 1 \pmod{p}$$

PROOF. From Lemma 6.3,

$$\begin{aligned} a \cdot 2a \cdots (p-1)a &\equiv 1 \cdot 2 \cdots (p-1) \pmod{p} \\ (p-1)!a^{p-1} &\equiv (p-1)! \pmod{p} \\ a^{p-1} &\equiv 1 \pmod{p} \end{aligned}$$

The last step follows from the fact that  $(p, (p-1)!) = 1$ . □

An important consequence of this result is that the inverse of any number  $a$  modulo  $p$  is  $a^{p-2}$ . By using this result, we can solve any congruence of the form:

$$ax \equiv b \pmod{p}$$

by multiplying both sides by the inverse of  $a$ , getting the equation:

$$x \equiv a^{p-2}b \pmod{p}$$

THEOREM 6.11. Let  $p$  be a prime and let  $a \in \mathbb{Z}$ . Then:

$$a^p \equiv a \pmod{p}$$

PROOF. If  $(a, p) = 1$ , then by Fermat's little theorem,

$$a^{p-1} \equiv 1 \pmod{p}$$

$$a^p \equiv a \pmod{p}$$

If  $(a, p) \neq 1$ , then  $p|a$ . Hence  $a \equiv a^p \equiv 0 \pmod{p}$ . □

Theorem 6.11 gives us the base for Fermat's test for non primality. Let  $n \in \mathbb{N}$ . If  $a^n \not\equiv a \pmod{n}$ , then  $n$  must be composite. Clearly, the converse does not hold - if  $a^n \equiv a \pmod{n}$ , this is not proof that  $n$  is prime! Composite numbers that follow the result in Fermat's little theorem are called Carmichael numbers.



THEOREM 6.12. Let  $p_1, p_2, \dots, p_k$  be distinct prime numbers. Let  $a \in \mathbb{N}$  and let

$$l = [p_1 - 1, p_2 - 1, \dots, p_k - 1]$$

Then,

$$a^{l+1} \equiv a \pmod{p_1 p_2 \cdots p_k}$$

PROOF. Let  $i \in \{1, 2, \dots, k\}$ . By Fermat's little theorem,

$$a^{p_i-1} \equiv 1 \pmod{p_i}$$

$$a^l = a^{(p_i-1) \times \frac{l}{p_i-1}} \equiv 1 \pmod{p_i}$$

$$a^{l+1} \equiv a \pmod{p_i}$$

The above result is also true if  $(a, p) \neq 1$  since then  $p|a$ , so  $a^{l+1} \equiv a \equiv 0 \pmod{p_i}$ . The result now follows from Theorem 6.5.  $\square$

EXERCISE. Let  $p$  be a prime number with  $p \geq 5$ . Let  $a \in \mathbb{Z}$ . Prove that:

$$a^p \equiv a \pmod{6p}$$

SOLUTION. TODO, let  $k = 3, p_1 = 2, p_2 = 3, p_3 = p$ . ■