

# Tracking San Francisco Bay water quality using generalized additive models in an R Shiny framework

Marcus W. Beck<sup>1</sup> Ian Wren<sup>2</sup> Rebecca Murphy<sup>3</sup>  
Perry de Valpine<sup>4</sup> David Senn<sup>5</sup>

<sup>1</sup>Tampa Bay Estuary Program


<sup>2</sup>San Francisco Baykeeper

<sup>3</sup>University of Maryland at Chesapeake Bay Program

<sup>4</sup>University of California Berkeley

<sup>5</sup>San Francisco Estuary Institute

Nov. 4, 2019



# Why do we care about trends?



- Provide information on natural variation of water quality parameters
  - ▶ What are the 1<sup>st</sup> order principles that describe a system?

# Why do we care about trends?



- Provide information on natural variation of water quality parameters
  - ▶ What are the 1<sup>st</sup> order principles that describe a system?
- Document historical changes in response to management actions
  - ▶ Did investments make a difference?

# Why do we care about trends?




- Provide information on natural variation of water quality parameters
  - ▶ What are the 1<sup>st</sup> order principles that describe a system?
- Document historical changes in response to management actions
  - ▶ Did investments make a difference?
- Anticipate future changes with proposed restoration or management
  - ▶ Can we understand the past to predict the future?

# Trends vary in space and time



*Observed data represent effects of many processes*



## Climate

precipitation  
temperature  
wind events  
ENSO effects

## Local

light/turbidity  
residence time  
invasive species  
trophic effects


## Regional/historical

watershed inputs  
point sources  
management actions  
flow changes

# Must translate data into information



*Observed data represents effects of many processes*




*Models should describe components to evaluate effects*

# Must translate data into information




*Observed data represents effects of many processes*




*Models should describe components to evaluate effects*

Annual




Seasonal



# Must translate data into information




*Observed data represents effects of many processes*




*Models should describe components to evaluate effects*


Annual




Residual



Seasonal



Flow effects



# South San Francisco Bay




- A high-nutrient,  
high-turbidity,  
low-productivity system

[Cole and Cloern, 1984,

Alpine and Cloern, 1988]

## South San Francisco Bay Long-term monitoring stations



# South San Francisco Bay



- A high-nutrient,  
high-turbidity,  
low-productivity system


[Cole and Cloern, 1984,

Alpine and Cloern, 1988]

- Recent increases observed in  
summer/fall chl-a

[Cloern et al., 2007, Cloern and Jassby, 2012]

South San Francisco Bay  
Long-term monitoring stations




# South San Francisco Bay



- A high-nutrient,  
high-turbidity,  
low-productivity system  
[Cole and Cloern, 1984,  
Alpine and Cloern, 1988]
- Recent increases observed in  
summer/fall chl-a  
[Cloern et al., 2007, Cloern and Jassby, 2012]
- Nutrient Management  
Strategy (NMS) to  
characterize status/trends and  
management needs

South San Francisco Bay  
Long-term monitoring stations




# South San Francisco Bay



Questions of concern:

South San Francisco Bay  
Long-term monitoring stations




# South San Francisco Bay



## Questions of concern:

- Since changes are visually apparent, which are significant?

## South San Francisco Bay Long-term monitoring stations




# South San Francisco Bay

## Questions of concern:

- Since changes are visually apparent, which are significant?
- What has been the estimated rate and direction of any linear or non-monotonic change?

South San Francisco Bay  
Long-term monitoring stations




# South San Francisco Bay



## Questions of concern:

- Since changes are visually apparent, which are significant?
- What has been the estimated rate and direction of any linear or non-monotonic change?
- Do any of these changes coincide with changes in other water quality parameters?

## South San Francisco Bay Long-term monitoring stations



# Application of additive models



- The Chesapeake Bay Program (CBP) has been wrestling with similar issues [Beck and Murphy, 2017, Murphy et al., 2019b]

# Application of additive models



- The Chesapeake Bay Program (CBP) has been wrestling with similar issues [Beck and Murphy, 2017, Murphy et al., 2019b]
- We applied Generalized Additive Models (GAMs) as implemented by CBP to characterize long-term trends at nine stations over thirty years in South SF Bay

# Application of additive models



- The Chesapeake Bay Program (CBP) has been wrestling with similar issues [Beck and Murphy, 2017, Murphy et al., 2019b]
- We applied Generalized Additive Models (GAMs) as implemented by CBP to characterize long-term trends at nine stations over thirty years in South SF Bay
- For each station, chlorophyll was modelled as a function of annual and seasonal changes over time using four model structures  
[Murphy et al., 2019a, baytrends R package]

# Application of additive models



- The Chesapeake Bay Program (CBP) has been wrestling with similar issues [Beck and Murphy, 2017, Murphy et al., 2019b]
- We applied Generalized Additive Models (GAMs) as implemented by CBP to characterize long-term trends at nine stations over thirty years in South SF Bay
- For each station, chlorophyll was modelled as a function of annual and seasonal changes over time using four model structures  
[Murphy et al., 2019a, baytrends R package]
  - ▶  $\text{gam0: chl} \sim \text{year} + \text{s(doy)}$

# Application of additive models



- The Chesapeake Bay Program (CBP) has been wrestling with similar issues [Beck and Murphy, 2017, Murphy et al., 2019b]
- We applied Generalized Additive Models (GAMs) as implemented by CBP to characterize long-term trends at nine stations over thirty years in South SF Bay
- For each station, chlorophyll was modelled as a function of annual and seasonal changes over time using four model structures  
[Murphy et al., 2019a, baytrends R package]
  - ▶ `gam0: chl ~ year + s(doy)`
  - ▶ `gam1: chl ~ year + s(doy) + s(year)`

# Application of additive models



- The Chesapeake Bay Program (CBP) has been wrestling with similar issues [Beck and Murphy, 2017, Murphy et al., 2019b]
- We applied Generalized Additive Models (GAMs) as implemented by CBP to characterize long-term trends at nine stations over thirty years in South SF Bay
- For each station, chlorophyll was modelled as a function of annual and seasonal changes over time using four model structures  
[Murphy et al., 2019a, baytrends R package]
  - ▶ `gam0: chl ~ year + s(doy)`
  - ▶ `gam1: chl ~ year + s(doy) + s(year)`
  - ▶ `gam2: chl ~ year + s(doy) + s(year) + ti(doy, year)`



# Application of additive models



- The Chesapeake Bay Program (CBP) has been wrestling with similar issues [Beck and Murphy, 2017, Murphy et al., 2019b]
- We applied Generalized Additive Models (GAMs) as implemented by CBP to characterize long-term trends at nine stations over thirty years in South SF Bay
- For each station, chlorophyll was modelled as a function of annual and seasonal changes over time using four model structures  
[Murphy et al., 2019a, baytrends R package]
  - ▶ `gam0: chl ~ year + s(doy)`
  - ▶ `gam1: chl ~ year + s(doy) + s(year)`
  - ▶ `gam2: chl ~ year + s(doy) + s(year) + ti(doy, year)`
  - ▶ `gam6: chl ~ year + s(doy) + s(year, k = large)`



# Application of additive models

gam0: chl ~ year + s(doy)





# Application of additive models

gam1:  $\text{chl} \sim \text{year} + \text{s(doy)} + \text{s(year)}$





# Application of additive models

gam2:  $\text{chl} \sim \text{year} + \text{s(doy)} + \text{s(year)} + \text{ti(doy, year)}$



# Application of additive models

gam6:  $\text{chl} \sim \text{year} + \text{s(doy)} + \text{s(year, k = large)}$




# Application of additive models



# Application of additive models

All years




# Application of additive models



# Application of additive models

All years



# Application of additive models



Table: Generalized cross-validation (GCV) summary statistics by station and model

Model	Stations (north to south)								
	18	21	22	24	27	30	32	34	36
gam0	-117.5	-30.6	-20.6	38.9	174.4	205.5	232.9	256.6	238.7
gam1	-138.8	-89.3	-70.3	-18.7	104.1	111.5	162.6	196	182
gam2	<b>-141.8</b>	-147.3	<b>-116.4</b>	-98.5	4	49.6	108.1	189.9	147.2
gam6	-139.5	<b>-235.2</b>	-116.3	<b>-176.8</b>	<b>-92.1</b>	<b>-115.9</b>	<b>-149</b>	<b>-3.3</b>	<b>-65.5</b>

# Application of additive models



Table: Generalized cross-validation (GCV) summary statistics by station and model

Model	Stations (north to south)								
	18	21	22	24	27	30	32	34	36
gam0	-117.5	-30.6	-20.6	38.9	174.4	205.5	232.9	256.6	238.7
gam1	-138.8	-89.3	-70.3	-18.7	104.1	111.5	162.6	196	182
gam2	<b>-141.8</b>	-147.3	<b>-116.4</b>	-98.5	4	49.6	108.1	189.9	147.2
gam6	-139.5	<b>-235.2</b>	-116.3	<b>-176.8</b>	<b>-92.1</b>	<b>-115.9</b>	<b>-149</b>	<b>-3.3</b>	<b>-65.5</b>

Table: R-squared summary statistics by station and model

Model	Stations (north to south)								
	18	21	22	24	27	30	32	34	36
gam0	0.47	0.41	0.37	0.37	0.33	0.36	0.32	0.31	0.32
gam1	0.51	0.48	0.43	0.44	0.43	0.48	0.41	0.41	0.43
gam2	<b>0.53</b>	0.54	0.48	0.53	0.54	0.54	0.49	0.41	0.48
gam6	0.51	<b>0.68</b>	<b>0.54</b>	<b>0.66</b>	<b>0.68</b>	<b>0.72</b>	<b>0.75</b>	<b>0.69</b>	<b>0.75</b>

# Conclusions from additive models



- More complex models provided more “parsimonious” fit
  - ▶ Non-monotonic trends were obvious
  - ▶ Interactions between year/season were common

# Conclusions from additive models



- More complex models provided more “parsimonious” fit
  - ▶ Non-monotonic trends were obvious
  - ▶ Interactions between year/season were common
- Clear demarcation of annual and seasonal trends allowed a description of the separate components
  - ▶ Annual trends showed increase, with some decreases in recent period of record
  - ▶ Change in dominance between spring/fall peaks were observed

# Conclusions from additive models



- More complex models provided more “parsimonious” fit
  - ▶ Non-monotonic trends were obvious
  - ▶ Interactions between year/season were common
- Clear demarcation of annual and seasonal trends allowed a description of the separate components
  - ▶ Annual trends showed increase, with some decreases in recent period of record
  - ▶ Change in dominance between spring/fall peaks were observed

***How can this information support decision-making??***

<https://sccwrp.shinyapps.io/sfbaytrends>

## GAM evaluation - SF South Bay

### Exploratory plots

The following plots show the raw data for all monitoring stations and parameters in South Bay, 1990 - 2017. Select the parameter, plot type (total time series, by year, or by month), and variable transformation. The year and month plots are aggregated boxplots of all observations at a station for each selected time period. The variable transformation can be used to show the observations in arithmetic or logarithmic space.

Choose station:



32

Choose plot type:

tot

Log-space:

TRUE




# Shiny interactive web platform




*Explore results for each station, by model*

model	smoother	edf	Ref.df	F	p.value
gam0	s(doy)	7.10	8.00	31.64	0.00
gam1	s(dec_time)	2.45	27.00	2.74	0.00
gam1	s(doy)	7.17	8.00	37.97	0.00
gam2	s(dec_time)	21.40	27.00	5.05	0.00
gam2	s(doy)	7.11	8.00	42.39	0.00
gam2	ti(dec_time,doy)	8.98	12.00	3.98	0.00
gam6	s(dec_time)	218.84	334.00	3.06	0.00
gam6	s(doy)	5.07	8.00	2.24	0.00

model	k	AIC	GCV	R2
gam0	NA	232.88	0.09	0.32
gam1	28	162.57	0.08	0.41
gam2	28	108.06	0.07	0.49
gam6	336	-148.96	0.06	0.75




## *Perform trend tests with selected years*



# Shiny interactive web platform




*Perform trend tests with selected years*



# Shiny interactive web platform




*Evaluate trends between years, by season*



# Shiny interactive web platform



*Evaluate trends between years, by season*



# Summary and next steps



- This analysis was a proof of concept
  - ▶ Does the GAM approach developed by the CBP transfer to SF Bay?
  - ▶ Do additional considerations need to be made?

# Summary and next steps



- This analysis was a proof of concept
  - ▶ Does the GAM approach developed by the CBP transfer to SF Bay?
  - ▶ Do additional considerations need to be made?
- Shiny platform helps communicate results to stakeholder
  - ▶ Which station, model, and time period do I care about?
  - ▶ How can I understand limitations of the different models?

# Summary and next steps



- This analysis was a proof of concept
  - ▶ Does the GAM approach developed by the CBP transfer to SF Bay?
  - ▶ Do additional considerations need to be made?
- Shiny platform helps communicate results to stakeholder
  - ▶ Which station, model, and time period do I care about?
  - ▶ How can I understand limitations of the different models?
- Follow-up work:
  - ▶ Extend to other locations in the Bay
  - ▶ Explore trend analysis of aggregated stations
  - ▶ Incorporate additional variables - as response or as explanatory

## Acknowledgments and contact info:

Research staff and employees at the San Francisco Estuary Institute, Delta Regional Monitoring Program, Southern California Coastal Water Research Project, and the Tampa Bay Estuary Program



mbeck@tbep.org

Phone (TBEP): (727) 893-2765



## Links:

This presentation: [https://github.com/fawda123/CERF\\_2019](https://github.com/fawda123/CERF_2019)

Shiny app: <https://sccwrp.shinyapps.io/sfbaytrends/>

Detailed results: <http://fawda123.github.io/SFbaytrends/README>

# References



- Alpine AE, Cloern JE. 1988.  
Phytoplankton growth rates in a light-limited environment, San Francisco Bay.  
*Marine Ecology Progress Series*, 44(2):167–173.
- Beck MW, Murphy RR. 2017.  
Numerical and qualitative contrasts of two statistical models for water quality change in tidal waters.  
*Journal of the American Water Resources Association*, 53(1):197–219.
- Cloern JE, Jassby AD. 2012.  
Drivers of change in estuarine-coastal ecosystems: Discoveries from four decades of study in San Francisco Bay.  
*Reviews of Geophysics*, 50(4):1–33.
- Cloern JE, Jassby AD, Thompson JK, Hieb KA. 2007.  
A cold phase of the East Pacific triggers new phytoplankton blooms in San Francisco Bay.  
*Proceedings of the National Academy of Sciences of the United States of America*, 104(47):18561–18565.
- Cole BE, Cloern JE. 1984.  
Significance of biomass and light availability to phytoplankton productivity in San Francisco Bay.  
*Marine Ecology Progress Series*, 17(1):15–24.
- Murphy R, Perry E, Keisman J, Harcum J, Leppo EW. 2019a.  
baytrends: Long Term Water Quality Trend Analysis.  
R package version 1.1.0.
- Murphy RR, Perry E, Harcum J, Keisman J. 2019b.  
A Generalized Additive Model Approach to evaluating water quality: Chesapeake Bay case study.  
*Environmenal Modelling & Software*, 118:1–13.