Important declarations

Please remove this info from manuscript text if it is also present there.

Associated Data

Data supplied by the author:

All source materials for reproducing the paper are available in GitHub: https://github.com/fawda123/bioassess opensci

Required Statements

Competing Interest statement:

The authors declare that they have no competing interests. M.W. Beck is employed by the Tampa Bay Estuary Program, but completed the manuscript as an employee of the Southern California Coastal Water Research Project (SCCWRP). Other non-academic affiliations include R.D. Mazor, S. Theroux, D.J. Gillett employed by SCCWRP and G. Gearheart employed by the California State Water Resources Control Board.

Funding statement:

The authors received no funding for this work.



The importance of open science for biological assessment of aquatic environments

Corresponding Author: Marcus W Beck Email address: mbeck@tbep.org

Open science principles that seek to improve science can effectively bridge the gap between researchers and environmental managers. However, widespread adoption has yet to gain traction for the development and application of bioassessment products. At the core of this philosophy is the concept that research should be reproducible and transparent, in addition to having long-term value through effective data preservation and sharing. In this paper, we review core open science concepts that have recently been adopted in the ecological sciences and emphasize how adoption can benefit the field of bioassessment for both prescriptive condition assessments and proactive applications that inform environmental management. An example from the state of California demonstrates effective adoption of open science principles through data stewardship, reproducible research, and engagement of stakeholders with multimedia applications. We also discuss technical, sociocultural, and institutional challenges for adopting open science, including practical approaches for overcoming these hurdles in bioassessment applications.

¹ Southern California Coastal Water Research Project, Costa Mesa, California, USA

² Tampa Bay Estuary Program, Saint Petersburg, Florida, USA

Bren School of Environmental Sciences & Management, University of California, Santa Barbara, Santa Barbara, California, United States

⁴ National Center for Ecological Analysis and Synthesis, Santa Barbara, California, USA

⁵ Department of Civil and Environmental Engineering, Utah Water Research Laboratory, Utah State University, Logan, Utah, United States

⁶ California Water Resources Control Board, Sacramento, California, USA



The importance of open science for biologicalassessment of aquatic environments

3

- 4 Marcus W. Beck^{Corresp., 1, 2}, Casey O'Hara³, Julia S. Stewart Lowndes⁴, Raphael D. Mazor¹,
- 5 Susanna Theroux¹, David J. Gillett¹, Belize Lane⁵, Greg Gearheart⁶

6

- 7 ¹ Southern California Coastal Water Research Project, 3535 Harbor Blvd, Costa Mesa, CA,
- 8 92626, USA
- 9 ² Tampa Bay Estuary Program (current address), 263 13th Ave S Ste 350, St. Petersburg, FL,
- 10 33701, USA
- 11 ³ Bren School of Environmental Sciences & Management, University of California Santa
- 12 Barbara, 2400 Bren Hall, Santa Barbara, CA, 93117, USA
- ⁴ National Center for Ecological Analysis and Synthesis, 735 State St, Santa Barbara, CA,
- 14 93101, USA
- 15 Department of Civil and Environmental Engineering, Utah Water Research Laboratory, Utah
- 16 State University, 1600 Canyon Rd, Logan, UT, 84321, USA
- 17 6 California State Water Resources Control Board, 1001 I Street, Sacramenta, CA, 95814, USA

18

- 19 Corresponding Author: Marcus W. Beck
- 20 Email address: mbeck@tbep.org

21 Abstract

- 22 Open science principles that seek to improve science can effectively bridge the gap between
- 23 researchers and environmental managers. However, widespread adoption has yet to gain traction
- 24 for the development and application of bioassessment products. At the core of this philosophy is
- 25 the concept that research should be reproducible and transparent, in addition to having long-term
- 26 value through effective data preservation and sharing. In this paper, we review core open science
- 27 concepts that have recently been adopted in the ecological sciences and emphasize how adoption
- 28 can benefit the field of bioassessment for both prescriptive condition assessments and proactive
- 29 applications that inform environmental management. An example from the state of California
- 30 demonstrates effective adoption of open science principles through data stewardship,
- 31 reproducible research, and engagement of stakeholders with multimedia applications. We also
- 32 discuss technical, sociocultural, and institutional challenges for adopting open science, including
- 33 practical approaches for overcoming these hurdles in bioassessment applications.



34 Introduction

- 35 Bioassessment is an essential element of aquatic monitoring programs that helps guide decisions
- 36 for managing the ecological integrity of environmental resources. Legal mandates to assess
- 37 biological condition have stimulated the development of bioassessment programs and tools in the
- 38 United States (Clean Water Act, CWA), Canada (Canada Waters Act), Europe (Water
- 39 Framework Directive), China (Environmental Quality Standards for Surface Water), South
- 40 Africa (National Water Act), and elsewhere (Borja et al. 2008). Decades of research have
- 41 supported the development of assessment indices for multiple assemblages with regional
- 42 applications in streams, rivers, lakes, and marine environments (Karr et al. 1986, Kerans and
- 43 Karr 1994, Fore and Grafe 2002, Beck and Hatch 2009, Borja et al. 2009, 2016). Substantial
- 44 technical advances have been made in measuring biological responses to environmental change
- 45 (Hawkins et al. 2000a, 2000b), how these responses can be distinguished from natural
- 46 environmental variation (Stoddard et al. 2006, Hawkins et al. 2010), and interpreting the impacts
- 47 of these changes (Davies and Jackson 2006).
- 48 Integrating bioassessment products (e.g., scoring indices, causal assessment protocols) into
- 49 management or regulatory frameworks can be challenging, despite the technological advances
- 50 (Kuehne et al. 2017). How a bioassessment product is used in practice to inform decisions and
- 51 prioritize management actions can differ from why it may have been originally developed.
- 52 Numerous assessment products have been developed for specific regional applications (Birk et
- al. 2012) and concerns about redundancy, comparability, duplicated effort, and lack of
- 54 coordinated monitoring have recently been highlighted (Cao and Hawkins 2011, Poikane et al.
- 55 2014, Kelly et al. 2016, Nichols et al. 2016). Kuehne et al. (2019) recently highlighted a lack of
- 56 institutional connectivity among actors with expertise in freshwater assessment as a hallmark of
- 57 the status quo in which applied science is conducted. Moreover, existing indices may not be
- 58 easily calculated by others beyond initial research applications (Hering et al. 2010, Nichols et al.
- 59 2016) or may be incorrectly applied based on differences between goals for developing an index
- and the needs of management programs (Dale and Beyeler 2001, Stein et al. 2009). The
- abundance of available products can be a point of frustration for managers given a lack of
- 62 guidance for choosing among alternatives, particularly as to how different assessment products
- relate to specific management, monitoring, or policy objectives (Dale and Beyeler 2001, Stein et
- 64 al. 2009).
- To address these challenges, a new mode of operation is needed where method development is
- open and transparent, developed products are discoverable and reproducible, and most
- 67 importantly, implementation in the management community is intuitive and purposeful. Open
- science principles that improve all aspects of the scientific method can help meet these needs and
- 69 there is a unique opportunity in bioassessment to leverage openness to support public resources.
- 70 Open science and its ideals originated partly due to failures of reproducibility and biases in the
- 71 primary literature that were revealed as systematic concerns in research fields with immediate
- 72 implications for human health (Makel et al. 2012, Franco et al. 2014). These ideas and the
- 73 failures that they address have slowly permeated the ecological and environmental sciences
- 74 (Hampton et al. 2015, 2016, Lowndes et al. 2017). Open science has also influenced how
- 75 research workflows are conceptualized in other disciplines (e.g., archaeology, Marwick et al.
- 76 2016, behavioral ecology, Ihle et al. 2017, hydrology, Slater et al. 2019, vegetation sciences,
- 77 Collins 2016) and has enabled a shift towards publishing structures that are more fair and



- 78 transparent through open access (Oudenhoven et al. 2016, Essl et al. 2020). Limited examples
- 79 have suggested that open access databases can be leveraged to develop bioassessment products
- 80 that increase transparency among stakeholders (Borja et al. 2019). Adopting an open science
- 81 paradigm in bioassessment is particularly relevant compared with other fields given the explicit
- 82 need to develop products that are accessible to the management community. Legal and ethical
- 83 precedents in bioassessment may also necessitate open data sharing given that environmental
- 84 monitoring programs are often established to protect and maintain publicly-owned natural
- 85 resources.

Survey Methodology and Objectives

- 87 This review draws on previous literature to describe approaches for open science that can
- 88 empower the research and management community to embrace a new mode of thinking for
- 89 bioassessment applications. These approaches are expected to benefit the bioassessment research
- 90 community by providing new tools that augment existing workflows for developing assessment
- 91 products and improving their ability to address environmental issues by bridging the gap
- between the scientific, management, and regulatory communities. The intended audience for this
- 93 review is primarily the research team that develops bioassessment products, but we also write for
- 94 the funders and users (e.g., regulators and managers) of these products to emphasize the value of
- 95 investing in open science for the protection of public resources.
- 96 This traditional review covers literature published in recent years advocating for open science in
- 97 different fields of study. Because no similar efforts have yet been made to apply these principles
- 98 to bioassessment, we draw on examples from the previous literature that demonstrate successful
- 99 applications in other fields to motivate researchers and practitioners to embrace these new ideas
- in bioassessment. Comprehensive and unbiased coverage of the previous literature was
- accomplished by querying online search engines, primarily Google Scholar, with search terms as
- they relate to open science (e.g., "reproducibility", "data science", "open source") and with
- Boolean operators to find applications to bioassessment (e.g., "reproducibility AND
- bioassessment"). Studies were included if they provided general overview of open science
- 105 concepts that were relevant to bioassessment or if they directly described open science
- applications to bioassessment, although the latter were scarce. Emphasis was given to the breadth
- of research that has supported the development of open source software applications that can aid
- 108 bioassessment, both as general tools and more specific programs tailored for indicator
- development. We excluded studies that described applications with citizen science components.
- Although citizen science can be a valuable tool for researchers and managers, methods for
- effective implementation are beyond the scope of this review.
- Our objectives are to 1) provide a general overview of principles of open science and 2)
- empower the research community by providing examples of how these principles can be applied
- to bioassessment. For the second objective, we also provide a case study of stream bioassessment
- in the urban landscape of southern California to demonstrate a successful proof of concept.
- Herein, open science 'tools' describe best practices and specific applications that use an open
- philosophy to support applied science. We structure the review by first introducing open science
- principles, then describing how these principles could be applied to bioassessment (i.e.,
- developing goals, curating data, and applying open-source software) including a case study



- example, and lastly providing a discussion of limitations and opportunities to better contextualize
- real world applications of open science.

Principles of open science

- 123 Conventional modes of creating scientific products and more contemporary approaches that align
- with open science principles share the same goals. Both are motivated by principles of the
- scientific method that make the process of discovery transparent and repeatable. Where the
- 126 conventional and open science approaches diverge is the extent to which technological advances
- facilitate the entire research process. Distinction between the two approaches can be
- 128 conceptualized as the "research paper as the only and final product" for the conventional
- approach, whereas the open science approach is inherently linked to advances in communication
- and analysis that have been facilitated by the Internet and computer sciences (Table 1). As a
- result, the open science approach can enhance all aspects of the scientific process from initial
- 132 conception of a research idea to the delivery and longevity of a research product (Figure 1). The
- process is iterative where products are improved by the individual and/or others, facilitated by
- open science tools that enhance access and reproducibility of data.
- 135 The paradigm of the research paper as a final scientific product can inhibit the uptake of research
- methods and findings by environmental managers. The research paper is conventionally viewed
- as a communication tool for scientists to report and share results among peers. Researchers
- access periodicals to stay informed of scientific advances and use the information to replicate
- and improve on methods for follow-up analysis. Although the primary literature continues to
- serve this critical role, this workflow is problematic when scientific products are needed to serve
- interests outside of the research community. For example, the paper as an endpoint for
- environmental managers fails to deliver products that are easily accessible from the practitioner's
- perspective, both in application and interpretation. A research paper is less likely to effect
- environmental change because it does not provide a mechanism to transfer actionable
- information to those that require scientific guidance for decision-making, such as sharing
- analysis code or results that describe output from assessment products. Numerous studies have
- documented implementation failures as a result of siloing among research communities where
- the flow of information does not extend beyond institutional walls (e.g., Mitchell 2005, Liu et al.
- 2008). Information loss over time is another concern associated with the paradigm of research
- paper as final product (Michener et al. 1997), particularly as intimate knowledge of study details
- is lost as new projects are initiated or individuals leave institutions.

Open data as a component of the open science process

- 153 Open data is a fundamental component of the broader open science process described in Figure
- 1. Under this mode of thinking, the research team becomes stewards of its data. For
- bioassessment data, government institutions may be the primary stewards of information that
- supports product development within a broader research team. Stewardship allows the data to be
- treated as a dynamic product with a traceable and replicable provenance (i.e., origin), rather than
- proprietary and serving only the internal needs of an immediate research goal. Metadata that
- describe the structure and history of a dataset ensure the data have an identity. Metadata also
- 160 encourage adoption of core data structures that allow integration across different sources, which
- is critical for collaboration across institutional boundaries (Horsburgh et al. 2016, Hsu et al.



- 162 2017). Other open science practices, such as integration of data with dynamic reporting tools or
- submitting data to a federated repository (i.e., a decentralized database system for coordination
- and sharing), can facilitate communication for researchers and those for which the research was
- developed (Bond-Lamberty et al. 2016). Prominent examples that can benefit next-generation
- bioassessment methods, such as molecular-based techniques for species identification, include
- the BarCode of Life Data Systems (BOLD) and GenBank repositories.
- Open data can benefit research by contributing to an increase in novel products created through
- 169 collaboration. Collaborative publications have increased in the environmental sciences as
- 170 research teams leverage open data to create synthesis products that allow novel insights from
- 171 comparisons across multiple datasets. Quantitative meta-analyses and systematic reviews are
- increasingly used to extract information from the primary literature (Carpenter et al. 2009, Lortie
- 173 2014). In addition, open data products can increase efficiency of the individual researcher and a
- 174 collective research team by encouraging collaborators to adopt an open science workflow. Many
- tools developed within the software and computer science community to facilitate open process
- and the creation of open data are now easily accessible to environmental scientists (Yenni et al.
- and the creation of open data are now easily accessible to environmental scientists (Yenni et al.
- 177 2019). Version control software (e.g., Git, GitHub), open source programming languages (e.g.,
- 178 R, Python), and integrated development environments (IDEs, e.g., RStudio, Spyder) can all be
- leveraged to dynamically create and share open data products that can build institutional
- 180 memory. These tools promote deliberate and shared workflows among researchers that can lead
- to better science in less time (Lowndes et al. 2017) and have proven useful in recent applications
- in the hydrologic sciences (Idaszak et al. 2017, Slater et al. 2019).
- Open access to data can also benefit management and regulatory communities. Openness can
- improve the value of data from monitoring programs by facilitating data discovery and synthesis,
- often through the adoption of a common metadata structure and integration of data within
- 186 federated data networks (e.g., DataONE, iRODS). Research institutions can also use open data
- maintained by management or regulatory communities to develop products that directly support
- the mission of the latter, e.g., assessment methods developed from long-term monitoring datasets
- that identify priority areas to focus management actions or fulfill regulatory obligations. Open
- data can also improve public trust in scientific findings by exposing the underlying information
- used to develop a research product (Grand et al. 2012). Similar concepts are used in
- 192 "blockchain" technologies that allow public financial transactions in an open, distributed format,
- as for trading in cryptocurrencies (Pilkington 2016). Increased trust could facilitate eventual
- adoption of proposed rules or regulations that are based on research products created from open
- data. More efficient and effective implementation of potential regulations may also be possible if
- supporting data are openly available.

Applying open science principles to bioassessment

- Here we provide a detailed description of open science processes that the bioassessment
- 199 community could leverage to create reproducible, transparent, and discoverable research
- 200 products for environmental managers. The below examples require understanding the distinction
- between the general open science process in Figure 1, open data as an individual component of
- the open science process, and the technology-based tools that can be used to achieve these ends.
- 203 Both the tools and open data are critical components that facilitate the broader process to achieve
- 204 the principles outlined in Table 1. "Openness" of process, tools, and data exists on a continuum,



- and incremental improvements can transform an individual's and research group's practice over
- 206 time. We encourage awareness that an open process adopts the open science tools that are
- appropriate for a research question and the creation of open data can be a fundamental
- 208 component of the process. Acceptance by the research team and collaborators of the concepts
- 209 described in Table 1 is critical to achieving openness.
- 210 The overall process is shown in Figure 2 as an expansion of general concepts in Figure 1. This
- iterative flow of information is facilitated by 1) openly sharing planning documents, 2) using
- established metadata standards to document synthesized data products, 3) hosting data products
- 213 on open repositories, 4) creating reproducible summary documents that integrate the data and
- 214 research products, and 5) incorporating the developed product into interactive applications that
- 215 deliver the results to the managers and stakeholders. The technical phase of defining research
- 216 goals, collecting and synthesizing data, and developing the bioassessment product are primary
- 217 tasks of the research team. However, the open science process is distinguished by the flow of
- 218 information to and from the research phase that can benefit the specific project and the science of
- 219 bioassessment as a whole.

Developing bioassessment goals

- 221 In an open science process, the goals identified by the research team for developing a
- bioassessment product should occur through direct, two-way interaction with the management or
- regulatory institution that requires the product. Although such an approach has historically been
- used to develop bioassessment products, the interaction in an open science workflow differs in
- 225 how information is exchanged. This exchange can be accomplished through direct
- communication and sharing of planning documents to ensure all decisions are transparent, i.e.,
- open planning. In person meetings are ideal, but planning documents are dynamic and will
- require remote sharing and revision as ideas progress. Online tools such as Google documents,
- 229 Slack discussion channels, and open lab notebooks can be instrumental for collaboration. More
- 230 informal approaches, such as blogging and sharing ideas on social media, can expose new
- concepts to the broader community for guidance (Woelfle et al. 2011, Darling et al. 2013).
- Overall, the research team should use these tools to identify stakeholder needs while also
- considering the balance between the research goals and limitations of the data to meet these
- 234 goals. This approach will ensure that the needs of the management and stakeholder communities
- will be consistent with the services provided by the research product.
- 236 An important practice that is not often used in bioassessment for project planning is study pre-
- registration. This is a relatively new addition to the philosophy of open science that allows a
- research team to define their study procedures, expected outcomes, and statistical analysis plans
- 239 in advance of the actual study (Munafó et al. 2017). Although the standard scientific method may
- seem to support such proactive practices, pre-registration is an explicit declaration to make the
- intentions of a study design clear to avoid publication bias where only positive outcomes are
- reported and to prevent an interpretive result where the researcher retrospectively defines study
- objectives after initial results are obtained if they do not agree with expectations. This latter issue
- is a serious concern where scientists use postdiction with significant hindsight bias in place of
- 245 prediction and conventional hypothesis testing to inform scientific discovery (Nosek et al. 2018).
- Registered reports can also be used as a publishing format where an initial study design is peer-
- reviewed and the article is provisionally accepted by a journal if the results are created with
- 248 methods that do not deviate from the accepted design. This promotes greater coverage in the



- 249 primary literature of null conclusions that otherwise may not have been published, reducing bias
- 250 for publishing positive results.
- 251 Pre-registration has been used extensively in clinical research (Dickersin and Rennie 2003),
- 252 where outcomes often have immediate implications for human health and well-being. In contrast,
- 253 bioassessment studies often focus on developing applied products, where conventional
- 254 hypothesis testing is less a concern. Studies are typically methods-focused where a research
- 255 product is developed to address a management or regulatory need, rather than a specific research
- 256 question with a testable hypothesis. However, pre-registration could be an important tool for the
- 257 environmental sciences where an explicit declaration of study intent as being applied or
- 258 methods-focused could prevent postdiction or an otherwise misuse of study results after a project
- 259 is completed. Existing venues that support pre-registration of studies across multiple disciplines
- 260 could be used in bioassessment study planning (e.g., Open Science Framework, AsPredicted).

Curating bioassessment data

- 262 After project goals are established, the research team identifies requirements and sources of data
- that need to be synthesized to meet the research needs. Bioassessment data, or more generally,
- biological data obtained from field sampling have a unique set of challenges that require added
- vigilance in data stewardship (Cao and Hawkins 2011). Species identification requires a tradeoff
- between taxonomic specificity and cost (Lenat and Resh 2001, Chessman et al. 2007). Species
- 267 names also change regularly requiring updates to standard taxonomic effort (STE) tables that are
- 268 critical for many biological indices, although some standardized databases have facilitated broad-
- scale comparisons (e.g., World Register of Marine Species, Costello et al. 2013). Unidentified or
- ambiguous taxa must also be explicitly treated in analysis workflows (Cuffney et al. 2007), e.g.,
- are they treated as missing values or are they substituted with coarser taxonomic designations? In
- 272 contrast, molecular techniques based on DNA barcoding eliminate the need for direct species
- 273 collection and morphological identification (Deiner et al. 2017, Hering et al. 2018). These next-
- 274 generation approaches have capitalized on advances in database development that allow open
- 275 access by diverse researchers across disciplines and are well-suited for the development of
- additional open science tools. Despite these advances, molecular-based approaches have also
- 277 suffered from challenges related to standardization of workflows and coverage of reference
- databases (White et al. 2014, Elbrecht et al. 2017).
- Open science tools can facilitate the curation of bioassessment data by addressing the above
- 280 challenges. For example, a multimetric index may require taxonomic data collected at multiple
- sites by different institutions, whereas the output data may include summary scores, individual
- metrics, and any additional supporting information to assess the quality of the output. In an open
- 283 science workflow, these data products can be documented using a standardized metadata
- language (e.g., Ecological Metadata Language Standard, or EML) which describes the who,
- 285 what, and why to ensure the data have an identity. Adoption of a metadata standard also ensures
- that a machine-readable file is produced to allow integration into a data repository. This will
- allow a synthesized data product to be discoverable beyond the specific research application and
- will provide metadata to help others understand the context of the data (e.g., Idaszak et al. 2017).
- Finally, the dataset can be assigned a unique digital object identifier (DOI, e.g., through Zenodo)
- 290 that provides a permanent address and is also citable to allow researchers to track usage of a
- 291 bioassessment data product.



- In an open paradigm, the data itself is a product to achieve the research goals and also becomes
- 293 available to the research and management community as a fully documented source of
- information that has value beyond the specific project. The openness of the synthesized data
- 295 product is one of the primary means of facilitating the application of a bioassessment product.
- 296 The synthesized data product can be used by the research team to create interactive applications
- 297 for stakeholders to share and explore the data and is also fully integrated into summary reports
- using software for generating dynamic documents (e.g., using knitr, Xie 2015, RMarkdown,
- 299 Allaire et al. 2018, Jupyter notebooks, Kluyver et al. 2016). Continuous integration services can
- 300 automate quality control and regularly update data products as new information is collected
- 301 (Yenni et al. 2019). The data product also becomes available on an open data repository that is
- discoverable by other researchers and can contribute to alternative scientific advances beyond the
- immediate goals (e.g., Hydroshare for the hydrologic sciences, Idaszak et al. 2017).

Using R for bioassessment application

- 305 The R statistical programming language (RDCT 2018) is one of the most commonly used
- analysis platforms in the environmental sciences (Lai et al. 2019, Slater et al. 2019) and many
- 307 existing R packages have value for the bioassessment community (Table 2). For managing the
- 308 day to day tasks of working with multiple datasets, the tidyverse suite of packages provides the
- necessary tools to import, wrangle, explore, and plot almost any data type (Wickham 2017). The
- 310 tidyverse also includes the powerful ggplot2 package that is based on a syntactical grammar
- of graphics for plotting (Wilkinson 2005, Wickham 2009). This package provides a set of
- 312 independent plotting instructions that can be built piecewise and is a departure from other
- 313 graphics packages that represent a collection of special cases that limit the freedom of the
- analyst. In bioassessment, ggplot2 can be used both in an exploratory role during the
- development phase and also to create publication quality graphics.
- 316 Bioassessment data are inherently spatial and recent package development has greatly improved
- 317 the ability to analyze and map geospatial data in R. The raster package can used to read/write,
- 318 manipulate, analyze, and model grid-based spatial data (Hijmans 2019), which are often common
- 319 supporting layers for bioassessment (e.g., elevation or climate data). For vector data (i.e., points,
- 320 lines, and polygons), the sf package ("simple features", Pebesma 2018) uses principles of data
- 321 storage that parallel those from the tidyverse by representing spatial objects in a tidy and
- 322 tabular format. This facilitates analysis by presenting complex spatial structures in a readable
- format that can be integrated in workflows with existing packages, including other mapping
- packages (e.g., leaflet, Cheng et al. 2018, or mapview, Appelhans et al. 2018). This allows the
- research team to use a workflow that is focused in a single environment, rather than using
- 326 separate software for statistical and geospatial analysis.
- R is fundamentally a statistical language and several existing R packages can be used to evaluate
- and support bioassessment data. Random forest models have been used to develop predictive
- 329 bioassessment indices that compare observed taxa to modeled expectations (i.e., O/E indices).
- The randomForest package (Liaw and Wiener 2002) uses an ensemble learning approach that is
- robust to complex, non-linear relationships and interactions between variables. These models are
- particularly useful with large, regional datasets that describe natural and anthropogenic gradients
- in condition (Laan and Hawkins 2014, Mazor et al. 2016). The nlme package can be used to
- create non-linear mixed effect models that are more flexible than standard regression approaches
- 335 (Pinheiro et al. 2018). The nlme package can develop models for nested sampling designs, such



- as repeat visits to sample sites or otherwise confounding variables that contribute information but
- are not unique observations (Mazor et al. 2014). The mgcv package provides similar functionality
- as nlme, but uses an additive modeling approach where individual effects can be evaluated as the
- sum of smoothed terms (Wood 2017). The mgcv package is often applied to model biological
- response to stressor gradients (Yuan 2004, Taylor et al. 2014).
- Other R packages have been developed specifically for bioassessment. For example, the TITAN2
- package can be used to develop quantitative evidence of taxon-specific changes in abundance
- and occurrence across environmental gradients (Baker et al. 2015). Results from this package
- 344 can support exploratory analysis for developing bioassessment products, such as identifying
- indicator species or establishing numeric criteria (Taylor et al. 2018). The results can be also be
- 346 used post hoc to evaluate potential response of a biological index with changing environmental
- conditions, such as proposed management actions for rehabilitation (King et al. 2011).
- 348 Alternatively, the indicspecies package provides similar functionality but is based only on
- 349 species occurrence or abundance matrices across sites (De Caceres and Legendre 2009). This
- package can be used to identify species at sites if continuous environmental data are unavailable,
- such as those that are representative of reference conditions (Bried et al. 2014). Finally, the
- vegan package has been a staple among community ecologists for multivariate analyses in R
- 353 (Oksanen et al. 2018).
- 354 Although the R network includes over 15000 user contributed packages, only a handful of these
- packages are specific to bioassessment. Community practices have allowed R to reach new
- audiences where new packages build on the work of others and are transportable between users
- and operating systems. Formalized communities, such as rOpenSci, encourage standardization
- and review of contributed packages within the ecological sciences to make scientific data
- retrieval reproducible. Several tools have also been developed and published in the last five years
- that greatly simplify the process of creating new packages in R (Wickham 2015, Wickham et al.
- 361 2018). The advantages of creating and sharing R packages that are specific to bioassessment
- applications are important for several reasons. First, an R package compartmentalizes technical
- instructions developed during the research phase that can be executed by anyone with access to
- 364 the software. R packages also require explicit documentation of the functions and data
- requirements. As such, package users will not only have access to underlying code but also
- understand the why and what for different package functions.
- Finally, R can be used to create interactive applications that deliver bioassessment products to
- 368 stakeholders and managers in entirely novel contexts. In particular, the shiny package provides
- programming tools built around concepts of reactivity, where data inputs and outputs can be
- modified in real time (Chang et al. 2018). A shiny application is an interactive user interface
- that is developed with R code, but is a standalone product that can be used without any
- 372 programming experience. These applications are deployed online and can extend the reach of
- bioassessment products to those that require the information for decision-making but otherwise
- do not have the time or resources to learn R. Applications built in shiny can also be easily linked
- 375 to other R packages. For example, a shiny website could be created to allow users to upload raw
- data and estimate and report bioassessment scores using an R package developed externally.
- Moreover, shiny applications are completely customizable and can be tailored by the developer
- 378 to the specific needs of any user. This distinction separates shiny from other web-based analysis
- 379 platforms.



Open science in practice: The SCAPE project

- 381 Although bioassessment products have been sufficiently developed in California (USA), there
- are no narrative or numeric criteria in place to support designated aquatic life uses in wadeable
- 383 streams, nor are bioassessment data actively used to support conservation or watershed
- 384 management. Indices using benthic macroinvertebrates and algae have been developed that
- provide consistent indications of biological condition across the diverse geography and climates
- in the state (Fetscher et al. 2013, Mazor et al. 2016, Ode et al. 2016). A physical habitat index
- has also been developed that provides complementary information supporting bioassessment data
- 388 (Rehn et al. 2018). Combined, these indices represent significant achievements in overcoming
- 389 technical challenges for developing accurate and interpretable bioassessment products. However,
- 390 these products are not used at a statewide scale to inform decisions and past efforts for stream
- 391 management have only used a fraction of available products. A synthesis of condition
- 392 assessments is needed to effectively implement bioassessment products in California and data
- must be presented in a context that is relevant to the needs of decision makers.
- 394 Recent regulatory initiatives in California have established a foundation for openness that could
- 395 greatly improve the application of bioassessment products to support decision-making. In
- 396 particular, these initiatives have set a precedent for openly sharing data collected with public
- funds. The Open and Transparent Water Data Act passed by the state legislature in 2016 requires
- 398 water quality institutions to "create, operate, and maintain a statewide integrated water data
- 399 platform that, among other things, would integrate existing water and ecological data information
- 400 from multiple databases and provide data on completed water transfers and exchanges" (AB
- 401 1755, Dodd, 2015-2016). This legislation also calls for state agencies to "develop protocols for
- data sharing, documentation, quality control, public access, and promotion of open-source
- 403 platforms and decision support tools related to water data". These aspirations were further
- 404 supported by a resolution on July 10, 2018 that formally committed the State Water Resources
- 405 Control Board to "provide broader access to data used to make local, regional, and statewide
- 406 water management and regulatory decisions in California". These recent initiatives in California
- 407 have similarly been observed at the national level. For example, the Data Coalition is an
- 408 advocacy group that operates on behalf of the private and public sector for the publication of
- 409 government data in a standardized and open format. The Internet of Water also operates at the
- ational-level by focusing on strengthening connections between data producers and users
- 411 through centralized data hubs and data standards.
- 412 Open science tools have recently been used in California to address bioassessment
- 413 implementation challenges in developed landscapes. The Stream Classification and Priority
- Explorer, or SCAPE (Beck 2018a, Beck et al. 2019), was developed using an open science
- 415 framework to help identify reasonable management goals for wadeable streams using existing
- bioassessment and watershed data. The SCAPE tool represents both a modeling approach to help
- 417 prioritize management goals (Figure 3) and a set of open science products for direct application
- 418 to environmental managers. The modeling component addresses a practical problem of achieving
- 419 reference conditions in developed landscapes, where channel modification is common. Using the
- 420 National Hydrography Dataset (NHD-Plus; McKay et al. 2012) and watershed predictors
- 421 (StreamCat; Hill et al. 2016), the model classifies stream segments as biologically "constrained"
- or "unconstrained" by landscape alteration. This classification system can be used to set
- 423 management priorities based on the constraint class. For example, a monitoring site with an



- observed biological index score that is above a predicted range could be assigned a higher
- 425 management priority relative to a site that is scoring within the range that is expected based on
- 426 landscape development.
- 427 Open science tools were critically important for translating and delivering SCAPE products to
- 428 decision-makers. Local stakeholder engagement to identify research goals guided the technical
- 429 development process of SCAPE. All analyses, including model development and validation,
- were conducted using R. A version control system (Git) and online hosting (GitHub) also
- allowed full transparency of decisions that were made to create the SCAPE model. A permanent
- 432 DOI was assigned through Zenodo to track downloads and portability of source code (Beck
- 433 2018a). Importantly, an online, interactive web page (https://sccwrp.shinyapps.io/SCAPE)
- greatly increased the impact and relevance of SCAPE by improving stakeholder understanding
- 435 through direct interaction with key decision points that influenced model output. A manuscript
- describing the technical components of the model was created using knitr and RMarkdown (Xie
- 437 2015, Allaire et al. 2018). This increased efficiency of the writing process also minimized the
- 438 potential of introducing errors into tables or figures by eliminating the need to copy results
- between different writing platforms. Finally, a geospatial data file from the model was also made
- public on a federated data repository, which included metadata and plain language
- documentation to track provenance of the original information (Beck 2018b).

Limitations and opportunities

- 443 Although the case for open science in bioassessment is appealing, the widespread adoption of
- 444 these principles in practice is inhibited by inertia of existing practices, disciplinary culture, and
- institutional barriers. Conventional and closed workflows used by many scientists are adopted
- and entrenched because of ease of use, precedence, and familiarity, yet they can be inefficient,
- 447 inflexible, and difficult to communicate or replicate. Open science tools can improve analysis.
- documentation, and implementation through greater flexibility, but they expose research teams to
- entirely new concepts and skillsets in which they may never have been trained (e.g., Idaszak et
- al. 2017). Not only are the required skillsets demanding, but the open science toolbox continues
- 451 to expand as new methods are developed and old methods become obsolete. This requires a
- research team to stay abreast of new technologies as they are developed and weigh the tradeoffs
- of adopting different workflows for different research tasks.
- 454 Advocates for open science are well aware of the technical challenges faced by individuals and
- research teams that have never been exposed to the core concepts. Most importantly, education
- and training (e.g., through The Carpentries) remain key components for developing skillsets
- among researchers where the focus is both on learning new skills for transferability and realizing
- 458 their value for improving science as a whole (Hampton et al. 2017). A goal of many training
- 459 curricula is to instill confidence in new users by developing comfort with new workflows, such
- as replacing a point-and-click style of analysis with one focused on using a command line
- through a computer terminal. Other approaches to demonstrate the value of new techniques use a
- side by side approach of closed vs open workflows to show the increased efficiency and power
- of the latter. Adoption becomes much more reasonable once users realize the value of investing
- 464 in learning a new skill.



488

489

490

491

492

493

494

495

496

497

498

499

500

501

505

507

508

465 Advocates of open science also recognize the limitations of teaching in that not all audiences can 466 be reached and not all materials are retained or even used after training. A strategy of empowering trainees to become trainers and teach others at their home institutions (e.g., train-467 468 the-trainer workshops and programs) enables open science to reach more individuals, and 469 benefits science more broadly as they develop technical and communication skills, and build local communities. Those that also adopt new workflows through training can also direct their 470 471 research products to facilitate collaboration with non-adopters rather than the latter synthesizing 472 and analyzing their data in potentially suboptimal ways (Touchon and McCoy 2016). These 473 "champions" can be a voice of encouragement for others by demonstrating how new tools can be 474 introduced and learned over time through shared experiences (Lowndes et al. 2017). This also 475 encourages the development of a community of practice that shares and learns together to navigate the collection of existing and developing open science tools (Stevens et al. 2018). 476 477 Champions of open science should also be vocal proponents that spread awareness of the value 478 of open science tools, particularly to those that make decisions on project resources. Department 479 heads or administrative leaders may not be aware of the value of investing in open science. 480 particularly if the consequences of not doing so are externalized in indirect costs that are not 481 budgeted. A change in mindset may be needed where open science is viewed as a core tool that 482 is critical to maintaining relevance of a research program in the future (Hampton et al. 2017). 483 Many scientists feel they cannot prioritize learning new skills given existing demands on their 484 time, particularly if the benefits of these approaches, such as the value for the research team of 485

sharing their data, are not apparent or immediate. Short-term funding and even political cycles can disincentivize scientists from spending time on anything but contractually obligated deliverables, which as noted above, may not effectively apply science in decision-making. This is an acute concern for early career scientists that have higher demands on establishing reputation and credentials, where investments in open science may be seen as detracting from progress (Allen and Mehler 2019). As an alternative, a practical solution is to actively encourage the investment in open science within the research team or lab, as opposed to placing the burden on the individual as an isolated researcher (i.e., team science, Cheruvelil and Soranno 2018). Laboratory or department heads should allow and encourage research staff to invest time in learning new skills and exploring new ideas, even if this does not immediately benefit the latest project. Over time, small investments in developing new skills will have long-term payoffs, particularly if a growing skillset among the research team encourages networking and peer instruction (Lowndes et al. 2017, Allen and Mehler 2019). Developing an environment where open science tools are highly valued and encouraged may also increase job satisfaction and benefit recruitment and retention if researchers are allowed the space and time to develop skills beyond the current project.

The scientific culture within a discipline or institution may inhibit the adoption of open science 502 methods. A common argument against open science is the protection of data that an individual 503 research team may view as proprietary or sensitive. There are reasonable arguments to treat data 504 as personal property, particularly if exceptional effort was spent to secure funding for a project and if the data were hard-earned or sensitive, e.g., detailed location data on endangered species 506 or medical/socioeconomic data (Zipper et al. 2019). These issues are less of a concern for bioassessment where many datasets are collected by institutions that are publicly funded and data accessibility may be mandated by law. However, an open science process dictates that both

interim and completed research products derived from public data should be available to the 509



- 510 broader bioassessment community. This raises an additional concern that research teams using
- transparent workflows could expose themselves to increased criticism by their peers and the
- 512 public (Lewandowsky and Bishop 2016, Allen and Mehler 2019), particularly where the
- 513 developed products can have important regulatory implications.
- Feedback and criticism are fundamental and natural parts of the scientific process. Scientists
- receive feedback at many stages in the conventional scientific workflow (e.g., internal review,
- 516 peer-review, presentations at conferences). Potentially new and challenging avenues for feedback
- are created in an open workflow. A concern is that openness can provide a platform for
- antagonistic or even hostile views, which could alter or degrade the scientific product (Landman
- and Glantz 2009, Lewandowsky and Bishop 2016). However, opportunities for addressing
- alternative viewpoints are critical to the open process of creating applied products, even if some
- voices are politically charged. This is especially true in bioassessment where finished products
- 522 that could be adopted in regulation are often heavily scrutinized. It is in the interest of applied
- scientists to hear the concerns of all parties during the development phase. This is not to provide
- an avenue to erode the integrity or objectives of the science, but to enable full knowledge of the
- very real barriers to adoption that exist when science is applied in regulation. Openness that
- 526 invites all voices to participate is a much more agreeable path to consensus than producing the
- science in isolation of those that it affects (Pohjola and Tuomisto 2011). Ultimately, these
- 528 products are developed to improve the environment as a public resource and the ideals promoted
- 529 by an open science process directly align with these goals.
- Institutional barriers can inhibit open science given the scale of change that must occur for
- adoption. Bureaucratic hurdles can disincentivize initiatives that promote change, particularly if
- that change originates from researchers not in administrative roles. Regulatory institutions may
- also prefer some level of opacity for how research products that influence policy are made
- available during development. The level of transparency advocated by open science could be
- viewed as opening the floodgates to increased legal scrutiny that can unintentionally hinder
- forward progress. Despite these reservations, many public institutions now advocate for
- 537 increased openness because of the benefits that facilitate and engender public trust. Open data
- 538 initiatives are now fairly common and represent a form of advocacy by public institutions for
- broader adoption of open science principles. Many national-level data products already exist that
- embrace openness to invest in the quality and availability of data (e.g., National Water Quality
- Monitoring Council initiatives, US Geological Survey products through NWIS and BioData, US
- 542 Environmental Protection Agency through STORET/WQX). Internationally, institutions in
- Europe and Canada that have projects supported by public funds are obligated to publish data
- and papers as open access (Horizon 2020, Tri-Agency Open Access Policy). Although past
- efforts and recent changes represent progress, many institutions have yet to strictly define open
- science and how it is applied internally and externally. As open science continues to build
- recognition, means of integrating toolsets that promote openness and transparency beyond
- 548 publicly shared data will have to be adopted by regulatory and management institutions.

Conclusions

549

- 550 The relevance of bioassessment applications can be improved with open science by using
- reproducible, transparent, and effective tools that bridge the gap between research and
- management. Many open science tools can improve communication between researchers and



- managers to expose all aspects of the research process and facilitate implementation to support
- 554 policy, regulation, or monitoring efforts. Communication ensures that the developed product is
- created through an exchange of ideas to balance the potentially competing needs of different
- sectors and institutions. The documentation and archiving of data used to create a bioassessment
- 557 product also ensures that other researchers can discover and build on past efforts, rather than
- constantly rebuilding the wheel. Incremental improvements of existing products can reduce the
- 559 proliferation of site- and taxon-specific methods with limited regional applications by exploring
- new ways to integrate biological indicators across space and time.
- Efforts to formally recognize and integrate open science in bioassessment are needed now more
- than ever. The transition of bioassessment from taxonomic-based indices to molecular
- approaches presents novel challenges that will only increase in severity as researchers continue
- to refine methods for molecular applications (Baird and Hajibabaei 2012). Although molecular-
- based indices share similar assessment objectives as conventional indices, the data requirements
- and taxonomic resolution are substantially more complex. Bioassessment researchers developing
- molecular methods are and will continue to be inundated with data from high-throughput DNA
- sequencers. Systematic approaches to document, catalog, and share this information will be
- required to advance and standardize the science. Molecular approaches are also dependent on
- 570 existing reference libraries for matching DNA samples for taxonomic identification. The
- 571 integrity of reference libraries depends greatly on the quality of metadata and documentation for
- 572 contributed samples. Open science principles should be leveraged in this emerging arena to
- ensure that new bioassessment methods continue to have relevance for determining the condition
- of aquatic resources.

Acknowledgments

- 576 The authors acknowledge support from the San Gabriel River Regional Monitoring Program and
- 577 the Sanitation Districts of Los Angeles County. We thank Eric Stein for reviewing an earlier
- draft of the manuscript. We thank Mike McManus for assistance with the conceptual diagrams.

579 **References**

- 580 Allaire, J., Y. Xie, J. McPherson, J. Luraschi, K. Ushey, A. Atkins, H. Wickham, J. Cheng, and
- 581 W. Chang. 2018. Rmarkdown: Dynamic documents for R. https://CRAN.R-
- 582 project.org/package=rmarkdown.
- Allen, C., and D. M. A. Mehler. 2019. Open science challenges, benefits and tips in early career
- and beyond. PLoS Biology 17:e3000246.
- Appelhans, T., F. Detsch, C. Reudenbach, and S. Woellauer. 2018. Mapview: Interactive
- viewing of spatial data in R. https://github.com/r-spatial/mapview.
- Baird, D. J., and M. Hajibabaei. 2012. Biomonitoring 2.0: A new paradigm in ecosystem
- assessment made possible by next-generation dna sequencing. Molecular Ecology 21:2039–
- 589 2044.



- Baker, M. E., R. S. King, and D. Kahle. 2015. TITAN2: Threshold indicator taxa analysis.
- 591 https://CRAN.R-project.org/package=TITAN2.
- 592 Beck, M. W. 2018a. SCCWRP/SCAPE: v1.0 (Version 1.0). Zenodo,
- 593 http://doi.org/10.5281/zenodo.1218121.
- Beck, M. W. 2018b. Constrained streams for biological integrity in California. Knowledge
- Network for Biocomplexity. urn:uuid:75411f50-32ed-42a5-bbfd-26833c7a441f.
- Beck, M. W., and L. K. Hatch. 2009. A review of research on the development of lake indices of
- 597 biotic integrity. Environmental Reviews 17:21–44.
- 598 Beck, M. W., R. D. Mazor, S. Johnson, K. Wisenbaker, J. Westfall, P. R. Ode, R. Hill, C. Loflen,
- 599 M. Sutula, and E. D. Stein. 2019. Prioritizing management goals for stream biological integrity
- within the developed landscape context. Freshwater Science 38:883–898.
- Birk, S., W. Bonne, A. Borja, S. Brucet, A. Courrat, S. Poikane, A. Solimini, W. van de Bund, N.
- Zampoukas, and D. Hering. 2012. Three hundred ways to assess Europe's surface waters: An
- almost complete overview of biological methods to implement the Water Framework Directive.
- 604 Ecological Indicators 18:31–41.
- Bond-Lamberty, B., A. P. Smith, and V. Bailey. 2016. Running an open experiment:
- Transparency and reproducibility in soil and ecosystem science. Environmental Research Letters
- 607 11:1–7.
- Borja, A., S. B. Bricker, D. M. Dauer, N. T. Demetriades, J. G. Ferreira, A. T. Forbes, P.
- Hutchings, X. Jia, R. Kenchington, J. C. Marques, and C. Zhu. 2008. Overview of integrative
- 610 tools and methods in assessing ecological integrity in estuarine and coastal systems worldwide.
- Marine Pollution Bulletin 56:1519–1537.
- Borja, A., M. Elliot, J. H. Anderson, T. Berg, J. Carstensen, B. S. Halpern, A. S. Heiskanen, S.
- 613 Korpinen, J. S. S. Lowndes, G. Martin, and N. Rodriguez-Ezpeleta. (2016). Overview of
- 614 integrative assessment of marine systems: The ecosystem approach in practice. Frontiers in
- 615 Marine Science 3:20.
- Borja, A., J. M. Garmendia, I. Menchaca, A. Uriarte, and Y. Sagarmínaga. 2019. Yes, we can!
- 617 Large-scale integrative assessment of european regional seas, using open access databases.
- 618 Frontiers in Marine Science 6.
- Borja, A., A. Ranasinghe, and S. B. Weisberg. 2009. Assessing ecological integrity in marine
- waters, using multiple indices and ecosystem components: Challenges for the future. Marine
- 621 Pollution Bulletin 59:1–4.
- 622 Bried, J. T., S. K. Jog, A. R. Dzialowski, and C. A. Davis. 2014. Potential vegetation criteria for
- 623 identifying reference-quality wetlands in the south-central United States. Wetlands 34:1159–
- 624 1169.
- 625 Cao, Y., and C. Hawkins. 2011. The comparability of bioassessments: A review of conceptual
- and methodological issues. Journal of the North American Benthological Society 30:680–701.



- 627 Carpenter, S. R., E. V. Armbrust, P. W. Arzberger, F. S. C. III, J. J. Elser, E. J. Hackett, A. R.
- 628 Ives, P. M. Kareiva, M. A. Leibold, P. Lundberg, M. Mangel, N. Merchant, W. W. Murdoch, M.
- A. Palmer, D. P. C. Peters, S. T. A. Pickett, K. K. Smith, D. H. Wall, and A. S. Zimmerman.
- 630 2009. Accelerate synthesis in ecology and environmental sciences. BioScience 59:699–701.
- 631 Chang, W., J. Cheng, J. Allaire, Y. Xie, and J. McPherson. 2018. Shiny: Web application
- framework for R. https://CRAN.R-project.org/package=shiny.

- 634 Cheng, J., B. Karambelkar, and Y. Xie. 2018. Leaflet: Create interactive web maps with the
- javascript 'leaflet' library. https://CRAN.R-project.org/package=leaflet.
- 636 Cheruvelil, K. S., and P. A. Soranno. 2018. Data-intensive ecological research is catalyzed by
- open science and team science. BioScience 68:813–822.
- 638 Chessman, B., S. Williams, and C. Besley. 2007. Bioassessment of streams with
- 639 macroinvertebrates: Effect of sampled habitat and taxonomic resolution. Journal of the North
- American Benthological Society 26:546–565.
- Collins, S. L. 2016. Vegetation science in the age of big data. Journal of Vegetation Science
- 642 27:865–867.
- 643 Costello, M. J., P. Bouchet, G. Boxshall, K. Fauchald, D. Gordon, B. W. Hoeksema, G. C. B.
- Poore, R. W. M. van Soest, S. Stöhr, T. C. Walter, B. Vanhoorne, W. Decock, and W. Appeltans.
- 645 2013. Global coordination and standardisation in marine biodiversity through the World Register
- of Marine Species (WoRMS) and related databases. PLoS One 8:e51629.
- 647 Cuffney, T. F., M. D. Bilger, and A. M. Haigler. 2007. Ambiguous taxa: Effects on the
- characterization and interpretation of invertebrate assemblages. Journal of the North American
- 649 Benthological Society 26:286–307.
- 650 Dale, V. H., and S. C. Beyeler. 2001. Challenges in the development and use of ecological
- 651 indicators. Ecological Indicators 1:3–10.
- Darling, E. S., D. Shiffman, I. M. Côte, and J. A. Drew. 2013. The role of Twitter in the life
- 653 cycle of a scientific publication. Ideas in Ecology and Evolution 6:32–43.
- Davies, S. P., and S. K. Jackson. 2006. The biological condition gradient: A descriptive model
- 655 for interpreting change in aquatic ecosystems. Ecological Applications 16:1251–1266.
- De Caceres, M., and P. Legendre. 2009. Associations between species and groups of sites:
- 657 Indices and statistical inference. Ecology. http://sites.google.com/site/miqueldecaceres/.
- Deiner, K., H. M. Bik, E. Mächler, M. Seymour, A. Lacoursiére-Roussel, F. Altermatt, S. Creer,
- 659 I. Bista, D. M. Lodge, N. de Vere, M. E. Pfrender, and L. Bernatchez. 2017. Environmental
- DNA metabarcoding: Transforming how we survey animal and plant communities. Molecular
- 661 Ecology 26:5872–5895.
- 662 Dickersin, K., and D. Rennie. 2003. Registering clinical trials. 290 4:516–523.



- Elbrecht, V., E. E. Vamos, K. Meissner, J. Aroviita, and F. Leese. 2017. Assessing strengths and
- weaknesses of dna metabarcoding-based macroinvertebrate identification for routine stream
- monitoring. Methods in Ecology and Evolution 8:1265–1275.
- Essl, F., F. Courchamp, S. Dullinger, J. M. Jeschke, and S. Schindler. 2020. Make open access
- publishing fair and transparent! BioScience 70:201–204.
- Fetscher, A. E., R. Stancheva, J. P. Kociolek, R. G. Sheath, E. D. Stein, R. D. Mazor, P. R. Ode,
- and L. B. Busse. 2013. Development and comparison of stream indices of biotic integrity using
- diatoms vs. Non-diatom algae vs. A combination. Journal of Applied Phycology 26:433–450.
- Fore, L. S., and C. Grafe. 2002. Using diatoms to assess the biological condition of large rivers
- 672 in Idaho (U.S.A.). Freshwater Biology 47:2015–2037.
- 673 Franco, A., N. Malhotra, and G. Simonovits. 2014. Publication bias in the social sciences:
- Unlocking the file drawer. Science 345:1502–1505.
- 675 Grand, A., C. Wilkinson, K. Bultitude, and A. F. T. Winfield. 2012. Open science: A new "trust
- 676 technology"? Science Communication 34:679–689.
- Hampton, S. E., S. S. Anderson, S. C. Bagby, C. Gries, X. Han, E. M. Hart, M. B. Jones, W. C.
- Lenhardt, A. MacDonald, W. K. Michener, J. Mudge, A. Pourmokhtarian, M. P. Schildhauer, K.
- H. Woo, and N. Zimmerman. 2015. The tao of open science for ecology. Ecosphere 6:1–13.
- Hampton, S. E., M. B. Jones, L. A. Wasser, M. P. Schildhauer, S. R. Supp, J. Brun, R. R.
- Hernandez, C. Boettiger, S. L. Collins, L. J. Gross, D. S. Fernández, A. Budden, E. P. White, T.
- 682 K. Teal, S. G. Labou, and J. E. Aukema. 2017. Skills and knowledge for data-intensive
- environmental research. Bioscience 67:546–557.
- Hampton, S. E., C. A. Strasser, J. J. Tewksbury, W. K. Gram, A. E. Budden, A. L. Batcheller, C.
- S. Duke, and J. H. Porter. 2016. Big data and the future of ecology. Frontiers in Ecology and the
- 686 Environment 11:156–162.
- Hawkins, C. P., R. H. Norris, J. Gerritsen, R. M. Hughes, S. K. Jackson, R. K. Johnson, and R. J.
- Stevenson. 2000a. Evaluation of the use of landscape classifications for the prediction of
- 689 freshwater biota: Synthesis and recommendations. Journal of the North American Benthological
- 690 Society 19:541-556.
- 691 Hawkins, C. P., R. H. Norris, J. N. Hogue, and J. W. Feminella. 2000b. Development and
- 692 evaluation of predictive models for measuring the biological integrity of streams. Ecological
- 693 Applications 10:1456–1477.
- Hawkins, C. P., J. R. Olson, and R. A. Hill. 2010. The reference condition: Predicting
- benchmarks for ecological and water-quality assessments. Journal of the North American
- 696 Benthological Society 29:312–343.
- 697 Hering, D., A. Borja, J. Carstensen, L. Carvalho, M. Elliott, C. K. Field, A. S. Heiskanen, R. K.
- Johnson, J. Moe, D. Pont, A. L. Solheim, and W. de Bund. 2010. The European Water
- 699 Framework Directive at the age of 10: A critical review of the achievements with
- recommendations for the future. Science of the Total Environment 408:4007–4019.



- Hering, D., A. Borja, J. I. Jones, D. Pont, P. Boets, A. Bouchez, K. Bruce, S. Drakare, B.
- Hänfling, M. Kahlert, F. Leese, K. Meissner, P. Mergen, Y. Reyjol, P. Segurado, A. Vogler, and
- 703 M. Kelly. 2018. Implementation options for DNA-based identification into ecological status
- assessment under the European Water Framework Directive. Water Research 138:192–205.
- Hijmans, R. J. 2019. Raster: Geographic data analysis and modeling. https://CRAN.R-
- 706 project.org/package=raster.
- Hill, R. A., M. H. Weber, S. G. Leibowitz, A. R. Olsen, and D. J. Thornbrugh. 2016. The
- 708 Stream-Catchment (StreamCat) dataset: A database of watershed metrics for the conterminous
- 709 United States. Journal of the American Water Resources Association 52:120–128.
- Horsburgh, J. S., M. M. Morsy, A. M. Castronova, J. L. Goodall, T. Gan, H. Yi, M. J. Stealey,
- and D. G. Tarboton. 2016. Hydroshare: Sharing diverse environmental data types and models as
- social objects with application to the hydrology domain. Journal of the American Water
- 713 Resources Association 52:873–889.
- Hsu, L., E. Mayorga, J. S. Horsburgh, M. R. Carter, K. A. Lehnert, and s. L. Brantley. 2017.
- 715 Enhancing interoperability and capabilities of earth science data using the Observations Data
- 716 Model 2 (ODM2). Data Science Journal 16:1–16.
- 717 Idaszak, R., D. G. Tarboton, H. Yi, L. Chistopherson, M. J. Stealey, B. Miles, P. Dash, A.
- 718 Couch, C. Spealman, D. P. Ames, and J. S. Horsburgh. 2017. HydroShare a case study of the
- 719 application of modern software engineering to a large distributed federally-funded scientific
- software development project. Pages 217–233 in J. Carver, N. P. C. Hong, and G. K.
- 721 Thiruvathukal, editors. Chapter 10 in software engineering for science. Taylor & Francis CRC
- 722 Press. Boca Raton, Florida, USA.
- 723 Ihle, M., I. S. Winney, A. Krystalli, and M. Croucher, 2017. Striving for transparent and credible
- research: Practical guidelines for behavioral ecologists. Behavioral Ecology 28:348–354.
- 725 Karr, J. R., K. D. Fausch, P. L. Angermeier, P. R. Yant, and I. J. Schlosser. 1986. Assessing
- biological integrity in running waters: A method and its rationale. Illinois Natural History
- 727 Survey, Champaign, Illinois.
- 728 Kelly, M. G., S. Birk, N. J. Willby, L. Denys, S.Drakare, M. Kahlert, S. M. Karjalainen, A.
- 729 Marchetto, J. A. Pitt, G. Urbanič, and S. Poikane, 2016. Redundancy in the ecological
- 730 assessment of lakes: Are phytoplankton, macrophytes and phytobenthos all necessary? Science
- 731 of the Total Environment 15:594–602.
- Kerans, B. L., and J. R. Karr. 1994. A benthic index of biotic integrity (B-IBI) for rivers of the
- 733 Tennessee Valley. Ecological Applications 4:768–785.
- King, R. S., M. E. Baker, P. F. Kazyak, and D. E. Weller. 2011. How novel is too novel? Stream
- community thresholds at exceptionally low levels of catchment urbanization. Ecological
- 736 Applications 21:1659–1678.



- Kluyver, T., B. Ragan-Kelley, F. Pérez, B. Granger, M. Bussonnier, J. Frederic, K. Kelley, J.
- Hamrick, J. Grout, S. Corlay, P. Ivanov, D. Avila, S. Abdalla, and C. Willing. 2016. Jupyter
- 739 notebooks a publishing format for reproducible computational workflows. IOS Press.
- Kuehne, L. M., and A. L. Strecker, and J. D. Olden. 2019. Knowledge exchange and social
- capital for freshwater ecosystem assessments. BioScience 70:174–183.
- Kuehne, L. M., J. D. Olden, A. L. Strecker, J. J. Lawler, and D. M. Theobald. 2017. Past,
- present, and future of ecological integrity assessment for fresh waters. Frontiers in Ecology and
- 744 the Environment 15:197–205.
- Laan, J. J. V., and C. P. Hawkins. 2014. Enhancing the performance and interpretation of
- 746 freshwater biological indices: An application in arid zone streams. Ecological Indicators 36:470–
- 747 482.
- Lai, J., C. J. Lortie, R. A. Muenchen, J. Yang, and K. Ma. 2019. Evaluating the popularity of R
- 749 in ecology. Ecosphere 10:e02567.
- Landman, A., and S. A. Glantz. 2009. Tobacco industry efforts to undermine policy-relevant
- research. American Journal of Public Health 99:45–58.
- Lenat, D. R., and V. H. Resh. 2001. Taxonomy and stream ecology the benefits of genus- and
- species-level identifications. Journal of the North American Benthological Society 20:287–298.
- Lewandowsky, S., and D. Bishop. 2016. Research integrity: Don't let transparency damage
- 755 science. Nature 529:459–461.
- 756 Liaw, A., and M. Wiener. 2002. Classification and regression by randomForest. R News 2:18–
- 757 22
- Liu, Y., H. Gupta, E. Springer, and T. Wagener. 2008. Linking science with environmental
- decision making: Experiences from an integrated modeling approach to supporting sustainable
- 760 water resources management. Environmental Modelling & Software 23:846–858.
- Lortie, C. J. 2014. Formalized synthesis opportunities for ecology: Systematic reviews and meta-
- 762 analyses. OIKOS 123:897–902.
- Lowndes, J. S. S., B. D. Best, C. Scarborough, J. C. Afflerbach, M. R. Frazier, C. C. O'Hara, N.
- Jiang, and B. S. Halpern. 2017. Our path to better science in less time using open data science
- 765 tools. Nature Ecology & Evolution 1:1–7.
- Makel, M. C., J. A. Plucker, and B. Hegarty. 2012. Replications in psychology research: How
- often do they really occur? Perspectives on Psychological Science 7:537–542.
- 768 Marwick, B., J. d'Alpoim Guedes, C. M. Barton, L. A. Bates, M. Baxter, A. Bevan, E. A.
- 769 Bollwerk, R. K. Bocinsky, T. Brughmans, A. K. Carter, C. Conrad, D. A. Contreras, S. Costa, E.
- 770 R. Crema, A. Daggett, B. Davies, B. L. Drake, T. S. Dye, P. France, R. Fullagar, D. Giusti, S.
- 771 Graham, M. D. Harris, J. Hawks, S. Heath, D. Huffer, E. C. Kansa, S. W. Kansa, M. E. Madsen,
- J. Melcher, J. Negre, F. D. Neiman, R. Opitz, D. C. Orton, P. Przystupa, M. Raviele, J. Riel-
- Salvatore, P. Riris, I. Romanowska, N. Strupler, I. I. Ullah, H. G. V. Vlack, E. C. Watrall, C.



- Webster, J. Wells, J. Winters, and C. D. Wren. 2016. Open science in archaeology. SAA
- 775 Archaeological Record 17:8–14.
- 776 Mazor, R. D., A. C. Rehn, P. R. Ode, M. Engeln, K. C. Schiff, E. D. Stein, D. J. Gillett, D. B.
- Herbst, and C. P. Hawkins. 2016. Bioassessment in complex environments: Designing an index
- for consistent meaning in different settings. Freshwater Science 35:249–271.
- Mazor, R. D., E. D. Stein, P. R. Ode, and K. Schiff. 2014. Integrating intermittent streams into
- 780 watershed assessments: Applicability of an index of biotic integrity. Freshwater Science 35:459–
- 781 474.
- 782 McKay, L., T. Bondelid, T. Dewald, J. Johnston, R. Moore, and A. Reah. 2012. NHDPlus
- 783 Version 2: User Guide.
- Michener, W. K., J. W. Brunt, J. J. Helly, T. B. Kirchner, and S. G. Stafford. 1997.
- Nongeospatial metadata for the ecological sciences. Ecological Applications 7:330–342.
- 786 Mitchell, B. 2005. Integrated water resource management, institutional arrangements, and land
- use planning. Environmental Planning A: Economy and Space 37:1335–1352.
- Munafó, M. R., B. A. Nosek, D. V. M. Bishop, K. S. Button, C. D. Chambers, N. P. du Sert, U.
- 789 Simonsohn, E. J. Wagenmakers, J. J. Ware, and J. P. A. Loannidis. 2017. A manifesto for
- 790 reproducible science. Nature Human Behavior 1.
- Nichols, S. J., L. A. Barmuta, B. C. Chessman, P. E. Davies, F. J. Dyer, E. T. Harrison, C. P.
- Hawkins, I. Jones, B. J. Kefford, S. Linke, R. Marchant, L. Metzeling, K. Moon, R. Ogden, M.
- 793 Peat, T. B. Reynoldson, and R. M. Thompson. 2016. The imperative need for nationally
- 794 coordinated bioassessment of rivers and streams. Marine and Freshwater Research 68:599–613.
- Nosek, B. A., C. R. Ebersole, A. C. DeHaven, and D. T. Mellor. 2018. The preregistration
- revolution. Proceedings of the National Academy of Sciences 115:2600–2606.
- 797 Ode, P. R., A. C. Rehn, R. D. Mazor, K. C. Schiff, E. D. Stein, J. T. May, L. R. Brown, D. B.
- Herbst, D. Gillett, K. Lunde, and C. P. Hawkins. 2016. Evaluating the adequacy of a reference-
- 799 site pool for ecological assessments in environmentally complex regions. Freshwater Science
- 800 35:237–248.
- Oksanen, J., F. G. Blanchet, M. Friendly, R. Kindt, P. Legendre, D. McGlinn, P. R. Minchin, R.
- 802 B. O'Hara, G. L. Simpson, P. Solymos, M. H. H. Stevens, E. Szoecs, and H. Wagner. 2018.
- 803 Vegan: Community ecology package. https://CRAN.R-project.org/package=vegan.
- Oudenhoven, A. P. E. van, M. Schröter, and R. de Groot. 2016. Open access to science on
- 805 ecosystem services and biodiversity. International Journal of Biodiversity Science, Ecosystem
- 806 Services & Management 12:239–241.
- Pebesma, E. 2018. Sf: Simple features for r. https://CRAN.R-project.org/package=sf.
- 808 Pilkington, M. 2016. Blockchain technology: Principles and applications. Pages 225–253 in F. X.
- 809 Olleros and M. Zhegu, editors. Research handbook on digital transformations. Edward Elgar
- 810 Publishing, Cheltenham, United Kingdom.



- Pinheiro, J., D. Bates, S. DebRoy, D. Sarkar, and R Core Team. 2018. nlme: Linear and
- 812 nonlinear mixed effects models. https://CRAN.R-project.org/package=nlme.
- Pohjola, M. V., and J. T. Tuomisto. 2011. Openness in participation, assessment, and policy
- making upon issues of environment and environmental health: A review of literature and recent
- 815 project results. Environmental Health 10:1–13.
- Poikane, S., N. Zampoukas, A. Borja, S. P. Davies, W. van de Bund, and S. Birk. 2014.
- 817 Intercalibration of aquatic ecological assessment methods in the European Union. Environmental
- 818 Science & Policy 44:237–246.
- Powers, S. M., and S. E. Hampton. 2019. Open science, reproducibility, and transparency in
- 820 ecology. Ecological Applications 29:e01822.
- 821 RDCT (R Development Core Team). 2018. R: A language and environment for statistical
- computing, v3.5.1. R Foundation for Statistical Computing, Vienna, Austria.
- Rehn, A. C., R. D. Mazor, and P. R. Ode. 2018. An index to measure the quality of physical
- 824 habitat in California wadeable streams. California Water Boards, Surface Water Ambient
- 825 Monitoring Program, California Department of Fish; Wildlife, Southern California Coastal
- 826 Water Research Project, Sacramento, California.
- 827 Slater, L. J., G. Thirel, s. Harrigan, O. Delaigue, A. Hurley, A. Khouakhi, I. Prosdocimi, C.
- Vitolo, and K. smith. 2019. Using R in hydrology: A review of recent developments and future
- directions. Hydrology and Earth System Sciences 23:2939–2963.
- 830 Stein, E. D., M. Brinson, M. C. Rains, W. Kleindl, and F. R. Hauer. 2009. Wetland assessment
- alphabet soup: How to choose (or not choose) the right assessment method. Society of Wetland
- 832 Scientists Bulletin 26:20–24.
- Stevens, S. L. R., M. Kuzak, C. Martinez, A. Moser, P. Bleeker, and M. Galland. 2018. Building
- a local community of practice in scientific programming for life scientists. PLoS Biology
- 835 16:e2005561.
- 836 Stoddard, J. L., D. P. Larsen, C. P. Hawkins, R. K. Johnson, and R. H. Norris. 2006. Setting
- 837 expectations for the ecological condition of streams: The concept of reference condition.
- 838 Ecological Applications 16:1267–1276.
- 839 Taylor, J. M., J. A. Back, B. W. Brooks, and R. S. King. 2018. Spatial, temporal and
- 840 experimental: Three study design cornerstones for establishing defensible numeric criteria in
- freshwater ecosystems. Journal of Applied Ecology 55:2114–2123.
- Taylor, J. M., R. S. King, A. A. Pease, and K. O. Winemiller. 2014. Nonlinear response of
- stream ecosystem structure to low-level phosphorus enrichment. Freshwater Biology 59:969–
- 844 984
- Touchon, J. C., and M. W. McCoy. 2016. The mismatch between current statistical practice and
- 846 doctoral training in ecology. Ecosphere 7:e01394.



- White, B. P., E. M. Pilgrim, L. M. Boykin, E. D. Stein, and R. D. Mazor. 2014. Comparison of
- four species-delimitation methods applied to a DNA barcode data set of insect larvae for use in
- routine bioassessment. Freshwater Science 33:338–348.
- Wickham, H. 2009. Ggplot2: Elegant graphics for data analysis. Springer-Verlag. New York.
- 851 http://had.co.nz/ggplot2/book
- Wickham, H. 2015. R packages. Page 182. O'Reilly, Sebastopol, California.
- Wickham, H. 2017. Tidyverse: Easily install and load the 'tidyverse'. https://CRAN.R-
- 854 project.org/package=tidyverse
- Wickham, H., J. Hester, and W. Chang. 2018. Devtools: Tools to make developing r packages
- easier. https://CRAN.R-project.org/package=devtools
- Wilkinson, L. 2005. The grammar of graphics. Page 691. Second ed. Statistics; Computing,
- 858 Springer, New York.
- Woelfle, M., P. Olliaro, and M. H. Todd. 2011. Open science is a research accelerator. Nature
- 860 Chemistry 3:745–748.
- Wood, S. N. 2017. Generalized additive models: An introduction with r. Page 476. 2nd editions.
- Chapman; Hall, CRC Press, London, United Kingdom.
- Xie, Y. 2015. Dynamic documents with R and knitr. 2nd editions. Chapman; Hall/CRC, Boca
- 864 Raton, Florida.
- Yenni, G. M., E. M. Christensen, E. K. Bledsoe, S. R. Supp, R. M. Diaz, E. P. White, and s. K.
- M. Ernest. 2019. Developing a modern data workflow for regularly updated data. PLoS Biology
- 867 17:e3000125.
- Yuan, L. L. 2004. Assigning macroinvertebrate tolerance classifications using generalised
- additive models. Freshwater Biology 49:662–677.
- 270 Zipper, S. C., K. S. S. Whitney, J. M. Deines, K. M. Befus, U. Bhatia, S. J. Albers, J. Beecher, C.
- Brelsford, M. Garcia, T. Gleeson, F. O'Donnell, D. Resnik, and E. Schlager. 2019. Balancing
- open science and data privacy in the water sciences. Water Resources Research 55:1–10.

Figure 1

A simplified workflow of the open science paradigm (adapted from Hampton et al. 2015).

All aspects of the research process, from the conception of an idea to publishing a product, can be enhanced using open science tools. The workflow is iterative where products are continually improved through collaborations facilitated through discovery and reproducibility of open data.






Figure 2

An idealized open approach for bioassessment applications.

The green box represents the technical steps of the individual research team for developing the product, the manager and stakeholder box are those that require or motivate the creation of bioassessment products, the gray boxes indicate sources of external information (data and guidance documents) as input into the technical process, and the open text indicates open components of the planning, application, or implementation phase of a bioassessment product. Figures were adapted from Hampton et al. (2015). NGO: non-government organization, RMP: regional monitoring program.






Figure 3

Schematic demonstrating how the Stream Classification and Community Explorer (SCAPE) can be used to identify potential management actions for stream sites.

Stream segment classifications are defined as biologically constrained or unconstrained based on landscape characteristics (A) and sites with bioassessment scores are evaluated relative to the classifications. Sites can be under-scoring, as expected, or over-scoring relative to the segment classification and expected range of scores (middle plot). Unconstrained sites are those where present landscape conditions do not limit biological potential and constrained sites are those where landscape conditions limit biological potential (B). Management actions and priorities can be defined based on site scores relative to segment classifications. TMDL: Total Maximum Daily Load. Photo credit: Raphael Mazor.








Table 1(on next page)

Core definitions and principles of open science.

Content adapted from Open Knowledge International, http://opendefiniion.org/, Creative Commons, http://openscience.org/, and Powers and Hampton (2019).



Concepts and principles	Description	
Open	Anyone can freely access, use, modify, and share for any purpose	
Open Science	Practicing science in such a way that others can collaborate and contribute, where research data, lab notes and other research processes are freely available, under terms that enable reuse, redistribution and reproduction of the research and its underlying data and methods	
Reproducible	Producing equivalent outcomes from the same data set, or in the case of computational reproducibility, producing equivalent outcomes from the same data set using the same code and software as the original study	
Principle 1	Transparency in experimental methods, observations, and collection of data	
Principle 2	Public availability and reusability of scientific data	
Principle 3	Public accessibility and transparency of scientific communication	
Principle 4	The use of web-based tools to facilitate scientific collaboration and reproducibility	



Table 2(on next page)

R packages that can be used in the development and application of bioassessment products.



Task	Package	Description
General	tidyverse (Wickham 2017)	A suite of packages to import, wrangle, explore, and plot data. Includes the popular ggplot2 and dplyr packages.
Mapping, geospatial	sf (Pebesma 2018)	A simple features architecture for working with vectorized spatial data, including common geospatial analysis functions
	raster (Hijmans 2019)	Reading, writing, manipulating, analyzing, and modeling gridded spatial data
	leaflet (Cheng et al. 2018)	Integration of R with the popular JavaScript leaflet library for interactive maps
	mapview (Appelhans et al. 2018)	Creates interactive maps to quickly examine and visually investigate spatial data, built off leaflet and integrated with sf
Statistical modeling	randomForest (Liaw and Wiener 2002)	Create classification and regression trees for predictive modeling
	nlme (Pinheiro et al. 2018)	Non-linear, mixed effects modeling
	mgcv (Wood 2017)	Generalized additive modeling
Community analysis	TITAN2 (Baker et al. 2015)	Ecological community threshold analysis using indicator species scores
	indicspecies (De Caceres and Legendre 2009)	Indicator species analysis
	vegan (Oksanen et al. 2018)	Multivariate analysis for community ecology
Science communication	shiny (Chang et al. 2018)	Reactive programming tools to create interactive and customizable web applications
	rmarkdown (Allaire et al. 2018)	Tools for working with markdown markup languages in .Rmd files
	knitr (Xie 2015)	Automated tools for markdown files that process integrated R code chunks