

# Estuaries and Coasts

## Assessment of the cumulative effects of restoration activities on water quality in Tampa Bay, Florida --Manuscript Draft--

Manuscript Number:	ESCO-D-19-00135R1	
Full Title:	Assessment of the cumulative effects of restoration activities on water quality in Tampa Bay, Florida	
Article Type:	Management Applications	
Keywords:	chlorophyll; Gulf of Mexico; long-term monitoring; restoration; Tampa Bay; trends	
Corresponding Author:	Marcus William Beck Southern California Coastal Water Research Project UNITED STATES	
Corresponding Author Secondary Information:		
Corresponding Author's Institution:	Southern California Coastal Water Research Project	
Corresponding Author's Secondary Institution:		
First Author:	Marcus William Beck	
First Author Secondary Information:		
Order of Authors:	Marcus William Beck	
	Edward T Sherwood	
	Jessica Renee Henkel	
	Kirsten Dorans	
	Kathryn Ireland	
	Patricia Varela	
Order of Authors Secondary Information:		
Funding Information:	Gulf Research Program	Not applicable
Abstract:	<p>Habitat and water quality restoration projects are commonly used to enhance coastal resources or mitigate negative impacts of water quality stressors. Significant resources have been expended for restoration projects, yet much less attention has focused on evaluating broad regional outcomes beyond site-specific assessments. This study presents an empirical framework to evaluate multiple datasets in the Tampa Bay area (Florida, USA) to identify 1) the types of restoration projects that have produced the greatest improvements in water quality, and 2) over which time frames different projects may produce water quality benefits. Information on the location and date of completion of 887 restoration projects from 1971 to 2017 were spatially and temporally matched with water quality records at each of 45 long-term monitoring stations in Tampa Bay. The underlying assumption was that the developed framework could identify differences in water quality changes between types of restoration projects based on aggregate estimates of chlorophyll-a concentrations before and after the completion of one to many projects. Water infrastructure projects to control point source nutrient loading into the Bay were associated with the highest likelihood of chlorophyll-a reduction, particularly for projects occurring prior to 1995. Habitat restoration projects were also associated with reductions in chlorophyll-a, although the likelihood of reductions from the cumulative effects of these projects were less than those from infrastructure improvements alone. The framework is sufficiently flexible for application to different spatiotemporal contexts and could be used to develop reasonable expectations for implementation of future water quality restoration activities throughout the Gulf of Mexico.</p>	



**SOUTHERN CALIFORNIA COASTAL WATER RESEARCH PROJECT**  
*A Public Agency for Environmental Research*

June 13<sup>th</sup>, 2019

Dr. Paul Montagna  
Dr. Charles A. "Si" Simenstad  
Co-Editors-In-Chief  
Estuaries and Coasts

I am pleased to resubmit our manuscript, "Assessment of the cumulative effects of restoration activities on water quality in Tampa Bay, Florida," to be considered as a management applications paper in Estuaries and Coasts.

We greatly appreciate the comments provided by our reviewers and have taken all efforts to address the concerns in our revision. In short, we have made editorial changes in all sections to provide clarity on our objectives, the methods used to address our objectives, primary results, and implications of the results in the discussion. Additional references and context were also added where needed. Please see our point by point response to all reviewer comments for details.

We are confident that readers of Estuaries and Coasts will find this information useful and appreciate the opportunity to publish our work in this venue. Please feel free to reach out should there be any questions about our submission.

Marcus W. Beck ([marcusb@sccwrp.org](mailto:marcusb@sccwrp.org)), 714-755-3217  
S. California Coastal Water Research Project  
3535 Harbor Blvd. Suite 110  
Costa Mesa, CA 92626

*Response to reviewer comments on manuscript “Assessment of the cumulative effects of restoration activities on water quality in Tampa Bay, Florida”, by M. W. Beck, E. T. Sherwood, J. R. Henkel, K. Dorans, K. Ireland, and P. Varela.*

*We sincerely thank the reviewers for providing helpful comments on our manuscript. Our responses to each of the comments are below. Line numbers provided by the reviewers are those in the original draft, whereas those in our responses are those in the revised draft.*

## **Reviewer 1 comments:**

Overall comments.

It was enjoyable to consider the substantial contributions made by this original and creative research effort. The spatial analysis appears to have been done carefully and documented well. Mostly, the assumptions and limitations of the study were made clear, which I appreciated; a few exceptions are noted in my detailed comments. The figures are exceptionally clear and communicative throughout, and even highly original and informative in some cases such as Fig. 7. The tables are clean and well-designed.

*We appreciate the substantial comments provided herein and have addressed each concern on a point-by-point basis below.*

Editing is a problem easily rectified by engaging an experienced editor, which will be very important for this paper to communicate the strength of its contributions, as it should. A very short list of examples is provided here but there was not time in a peer review to document the many editorial exceptions:

39-40 "widespread chronic" is awkward  
*Changed to “widespread”.*

42 lack of Oxford comma  
*Added.*

46 starting sentence with And is pretty informal for a journal publication  
*Changed to “Further,”.*

50-52 rigorous used twice in one sentence  
*Revised to “...can further constrain evaluations of restoration success”*

50-52 tense problem; difficulty and constrain. (Difficulty constrains. Difficulties constrain.)  
*Changed to “difficulties”.*

150-154. Awkward. 2 sentences.  
*Revised: “We developed and applied a two-level classification scheme that described each restoration project as 1) a habitat or water infrastructure improvement and 2) more specifically as enhancement, establishment, or protection for habitat or as nonpoint or point source controls for water infrastructure.”*

168 included, not include  
*Changed.*

176 "matching between" is awkward

Revised to *"The matchings began with..."*

Overall: Inconsistent use of bay versus Bay when the word stands alone from Tampa

*All uses were checked and changed to "Bay".*

Detailed Comments by Line Number and Figure Number:

Introduction. The rationale developed in the opening sections of the Introduction is solid and well cited with a combination of peer-reviewed papers and a small number of necessary reports in the gray literature: both natural and social/institutional challenges to long-term monitoring of restoration projects and to large-scale restoration success; so the unprecedented current opportunity in Gulf following Deepwater Horizon requires careful examination of what has worked before in the region.

Line 76 Given the previously well-developed argument about the challenges of long-term, intensive monitoring, it would be interesting for the reader if the "secrets" to doing so in Tampa Bay were described at this point, even just a single sentence on what made Tampa Bay long-term intensive monitoring "work." eg was it citizen science? Volunteers? A charismatic individual? Etc. (I am guessing it may come down to Holly Greening herself, but even that would be interesting if it has to do with continuity of leadership, for example.) Or maybe it comes down to regional partnerships (Sherwood et al. 2016, Regional Studies in Marine Science)

*The long-term monitoring program in Tampa Bay has been the foundation for assessing and tracking the recovery of water quality since the 1970s. We agree with the reviewer that it's worthwhile pointing out why this program has been so successful. We have added a paragraph to the study area section to describe the history of the monitoring program: "A long-term monitoring program in Tampa Bay has been instrumental in assessing and tracking restoration efforts. In the early 1970s, the initial baywide ambient monitoring program was established by a local environmental leader (Roger Stewart), which was subsequently institutionalized through State legislation by the creation of the Environmental Protection Commission of Hillsborough County. This occurred largely in response to citizen out-cry of the Bay's deteriorating ecology (Greening et al. 2014). Ongoing local support for this program has remained since 1972, and other local, municipal governments have created complementary water quality monitoring programs, all of which now support Tampa Bay water quality assessments and management efforts spear-headed by the Tampa Bay Estuary Program (Sherwood et al. 2016). Some of the key attributes supporting the maintenance of this long-term monitoring program are summarized in Schiff et al. (2016) and Gross and Hagy (2017)."*

*Also note that this paragraph in the introduction has been revised in response to the following comment.*

78-79 It is difficult to believe that "no previous efforts have been made to directly quantify potential associations between these activities and estuarine water quality." I think the authors owe readers to follow this statement with a concise lit review, a full paragraph, of eg Sherwood 2016,2017; Greening 2014; and others' published work; stating what they did do, so as to clarify how the authors' work here is unique and different. It may also be worth consulting gray literature because surely there was some "method to the madness" as to which remediation and restoration projects were funded and most importantly why they were prioritized, which was very likely based on past experience of what worked even if it was never published. I note that lines 78-88 are citation free. Deeper consideration of what has

come before is warranted here to assure the reader that what's being proposed by the authors is really needed. I believe that it is needed, but that history needs to be documented here in brief.

*We agree that our general statement that “no previous efforts have been made” was imprecise and have revised this paragraph to describe the context more clearly. We have moved some examples from the text on lines 124-131 in the previous draft to this paragraph. The revised paragraph is as follows:*

*“Tampa Bay (Florida, USA) is the second largest estuarine embayment in the GOM and improvement in condition over the last four decades is one of the most exceptional success stories for coastal water quality management (Greening et al. 2006; Greening et al. 2014). Most notably, seagrass coverage in 2016 was reported as 16,857 hectares baywide, surpassing the goal of restoring coverage to 95% that occurred in 1950 (Sherwood et al. 2017). Reductions in nutrient loading (Poe et al. 2005; Greening et al. 2014), chlorophyll-a concentrations (Wang et al. 1999; Beck and Hagy 2015), and improvements in water clarity (Morrison et al. 2006; Beck et al. 2018) have also preceded the seagrass recovery. Most of these positive changes have resulted from management efforts to reduce point source controls on nutrient pollution in the highly developed areas of Hillsborough Bay (Johansson 1991; Johansson and Lewis III 1992). These controls allowed nutrient and chlorophyll-a targets to be met by the early 1990s. However, numerous smaller projects, including watershed-focused efforts (Lewis et al. 1998), may have had a supporting role in maintaining water quality improvements through contemporary periods. The cumulative effects of over 900 restoration projects, relative to broad watershed-scale management efforts, are not well understood. Understanding how implementation of these projects is associated with adjacent estuarine water quality at various spatio-temporal scales will provide an improved understanding of the link between overall estuary improvements and specific restoration activities.”*

89-90 The argument seems thin. Could probably add several other reasons to this sentence.

*The beginning of this paragraph was revised: “Demonstrating success for restoration activities is challenging for several reasons (Ruiz-Jaen and Aide 2005; Wortley et al. 2013). Success may be vaguely or even subjectively defined because the effects of restoration could be described in different ways depending on project goals (Zedler 2007). For example, site-specific measures of before/after condition are commonly used to measure of success, whereas downstream effects may be more important to consider for baywide conditions (Diefenderfer et al. 2011). More importantly, quantifying success as a measure of environmental improvements is challenged by the variety of factors that affect water quality across space and time. New tools are needed...”*

95-97 Excellent objectives. Challenging and worthwhile.

Methods: Overall, there is a lack of documentation of other published work particularly in regard to the spatial methods which informed those used in the study, which should be added throughout Methods section. It is clear that the team is experienced in spatial modeling and I presume familiar with the literature in GIS methods. That said, the essential work appears to be an important contribution.

*The following was added to the first paragraph of the data synthesis and analysis framework: “Spatial and temporal matching can be accomplished using several methods that vary in complexity. For example, hydrologic distances or other non-Euclidean distance weightings by watershed topology can be used to link measurements to modelled locations in space (Curriero 2006; Gardner et al. 2011).*

*However, we adopted a relatively simple approach with limited data requirements to maximize potential applications in other regions (e.g., no hydrology data are needed, only spatial location).”*

*Additional info was added to line 228: “A weighted average by distance (or parametric distance weights, Sickles and Johnson 2008) was used based on the assumption that restoration projects farther from a water quality station will have a weaker association with potential changes in chlorophyll-a.”*

The software name and version number should be mentioned somewhere in Methods.

*The software and version were already mentioned on lines 272-273. No other software was used. We also note that source materials were made available through a DOI link in the supplement. We have expanded material in the supplement to also include a web application to explore results: “Supplement: Data analysis frameworks, analysis code, and manuscript source materials are available at [DOI:10.5281/zenodo.2539623](https://doi.org/10.5281/zenodo.2539623). An interactive application exploring results for Tampa Bay and individual Bay segments is available at [https://shiny.tbepitech.org/restorebayes/ind\\_eval.Rmd](https://shiny.tbepitech.org/restorebayes/ind_eval.Rmd).”*

115 “purportedly” implies you disagree with this conclusion. Instead, suggest removing that word and adding a citation to the fact.

*Removed “purportedly” and added two citations (Comp 1985; Lombardo and Lewis 1985).*

121 Actually, according to Sherwood, the goal was 95% of 1950s level, not 100%. Suggest re-reading Sherwood and citing original literature for the goal if available.

*Text was changed.*

124-131. It would have helped to make this nuance clear at the end of the introduction (see comments lines 78-79). Tampa Bay is such a well known success story that it was not credible to state no cause-and-effect research had been reported. Clarifying that it is the other smaller projects that haven't been evaluated in the Intro would help make the authors believable.

*Please see our response above to the comment on the paragraph starting on line 76 in the original draft.*

129-131 “have likely” suggests bias. Instead saying that reports indicate they may have had an effect, which you sought to evaluate, would be a cleaner approach.

*Changed to “may have had”. We are unaware of any reports that explicitly evaluated the hundreds of restoration projects that have occurred in the Bay or its watershed.*

135-148 Please address whether there was overlap in the projects in these datasets.

*There was no overlap between the datasets. A sentence was added to line 150 to clarify: “Each database was unique and no overlap in documented projects was observed.”*

156-158 move down as the topic sentence of the next section on Data Analysis, and remove “methods described below”

*Moved.*

168-170 Why list data you didn't use? Also, previously, it was stated that chlorophyll was the metric. Now two others have been added. Confusing.

*These lines were revised: “Chlorophyll-a ( $\mu\text{g/L}$ ) and total nitrogen ( $\text{mg/L}$ ) measurements at each site were used for analysis, totaling up to 515 discrete observations for each station. Total nitrogen concentrations were included in initial data assessments, as this nutrient is considered limiting in Tampa Bay (Greening et al. 2014).”*

The assumption seems to be that the nearest sites have the most effect, which seems to disregard flow direction. However, the time binning does help to ameliorate this concern.

*Please see our response above regarding the citations added for geospatial methods and our clarifications below on the methods for time binning.*

In the absence of any other explanation, I assume that even projects up in the watershed were digitized for hydrologic distance "as the crow flies" and it is rather interesting that despite this apparent weakness, the results were significant. Implications should be treated in the Discussion. Potentially an avenue of future research to build on this?

*We have added a final subsection in the discussion on "future directions": "Our approach is not without limitations and future research could build on the methods to provide an improved assessment of restoration effectiveness. Our geospatial analyses were relatively simple, in that spatial matchings were accomplished through Euclidean distances. Alternative distance measures could be used that consider hydrologic distances following flow networks in the watershed. The importance of these approaches could provide insight into pollutant dispersal pathways in environments with low elevation gradients, such as Florida. Weighting restoration projects by relative effort could also facilitate an improved assessment of effectiveness, such as considering total restoration area as an important variable to consider for water quality improvements. Some of these data were available in our compiled dataset, although coverage was insufficient for a complete analysis. Finally, the social and human dimensions of different restoration projects were not considered herein but are important factors that can be equally or even more relevant determinants of success that should be considered when weighing restoration options. Future restoration and monitoring activities should adopt a more comprehensive evaluation of success measures that includes and extends beyond water quality changes."*

188-190 Explanations of time windows are not complete enough to understand.

\* Also raises a question of how these were determined. A priori, as a hypothesis? Or after trial and error and discovering which ones yielded results. Important to state this.

*By design, the time windows are flexible to allow exploration of potential periods of time within which water quality improvements may be expected after project completion. Text was added to this paragraph to make this clear: "...where the latter could be manually changed to characterize a potential duration of time within which water quality could improve after project completion dates."*

\* Also need to state how long projects take to complete, and if this length of time is different for the different categories of projects.

*This information was not available and we have indicated as such to make this limitation of our analysis more clear: "Lastly, no data were available on project duration and we assumed that the year associated with each project was generally inclusive of project implementation and completion."*

\* Fig 5b seems to indicate much longer time slices than are described in the text.

*This was not well-described in the text. Lines 205-211 were revised: "The before estimate was the average of observations for the year preceding the completion of a project and the after estimate was the average of observations for a selected window of time (e.g., five years) that occurred after completion of a project. The before estimate for the year prior established the basis of comparison for the water quality estimates in the selected window of time after project completion."*

\* Also raises a question of how much interannual variability there is in hydrologic connectivity in this system. Is one year of after data sufficient to capture effects? I realize the assumption may be that after one year, the likelihood of other changes affecting the results increases.

*We agree that there can be many factors influencing our analysis (many of which are described more thoroughly in the discussion). We hope our additions to this paragraph have clarified our methods and their intent.*

198-200 I think this should be summations (plural) were (not was) correct? The before and after data were separately divided by the number of projects, correct?

*Yes, changed.*

Data synthesis and analysis framework Section: Lack of citations to related methods. Need to contextualize the work with that of others in spatial data processing and water quality analysis. *References were added in response to the above general comments for the methods section.*

212-213 Did these hypotheses have a direction or were they as general as is described here?

*The sentence was revised: "We make no assumptions about the expected magnitude of an association given that the model does not describe a specific mechanism of change, nor do we make any explicit assumption about the direction of change (i.e., two-tailed hypothesis tests were used), although a general assumption was that chlorophyll-a would decrease over time in agreement with known changes in water quality."*

213-214 Why wouldn't the statements be quantitative, given that this is a quantitative analysis by distance and time? Qualitative may undersell the analytical effort made here.

*This statement was revised: "An expected outcome is that explicit, quantitative conclusions can be made...". However, the interpretation of the results can be nuanced, as noted in the discussion.*

217-221 This is a promising application.

228-230 A strength of this paper is this multi-scale approach. Others have similarly looked at eg drift cell reaches along shorelines versus watershed-scale effects on coasts, references which could be incorporated in the methods. I think the point here is larger than an "alternative context." Rather each scale provides information about potential stressors and processes controlling the impacts. For stream restoration examples see eg Roni et al., N. Am J of Fisheries Management 2002; Roni et al. 2012 book chapter, available online, "Prioritization of Watersheds and Restoration Projects"; for coastal restoration examples see eg multi-scale analysis of restoration priorities papers by Thom et al., Ecol Rest 2011; Diefenderfer et al. Env Man 2009.

*We agree that the value of this spatial flexibility was understated. We have added a sentence to line 259 that includes the suggested citations and better describes the value of the analysis: "Evaluating the results at different scales can also provide insights into potential (or lack of) stressors and processes controlling the impacts, which can help prioritize management actions by location (Diefenderfer et al. 2009; Thom et al. 2011)."*

247-249 good idea/verification of assumptions. This is a little bit unclear, "but with random date and location assignments for each restoration project". I assume that both randomized date/location, and actual date/location were analyzed and then compared, correct?

*Yes, this was clarified: "...but with random date and location assignments for each restoration project that were then compared to the actual results."*

253-58 a bit cryptic

*This section was revised for clarity: "In other words, the randomization creates a null model where the estimated effects of restoration projects would not differ from a simple evaluation of trends in the raw data - slicing the observed time series by random dates and evaluating before/after averages with random projects is expected to reflect the known decline of chlorophyll-a in the raw data. Alternatively, evidence that our framework provides meaningful results would be supported by differences in chlorophyll-a changes between project types and the timing associated with the changes."*

Results. Overall comment. Consider organizing with the key findings and figures first, and lesser/background findings second.



*We appreciate the concern that presenting background (i.e., observed) findings first somewhat diminishes the importance of our key results. However, these findings are limited only to the first page and we feel it provides a succinct and necessary background to the primary results that follow. We have retained the original structure.*

Compare 264-266 with 277-279. These are not in fact "coincident," as stated. The monotonic rate of decline of pollution slowed after 20 years of the 40-year period; but the rate of increase in projects on the ground increased after the year 2000! These are only coincident over the entire 40-year period, not within the 40-year period. In fact, the patterns of rate of pollution reduction and rate of new projects (a project count, not an effect magnitude) appear to be the opposite.

*We were careful in our language not to overstate the associations, but we agree these trends are not simply "coincident" and have removed this from the text.*

307-320 Very important points that may belong in the Discussion. It's worth adding what these analyses don't say about the value of Protection, as well, because Protection in an area such as Tampa Bay with severe development pressure is very important. (338-341)

*Much of this content was moved to the "analysis limitations" section. We have also added a note about the challenges associated with protection in developed areas where land is in high demand (line 494).*

368-370 I don't much mind this here but many would call it Discussion material and it might receive more attention there if highlighted as part of the contribution of these Methods.

*This statement was removed. The importance of this flexibility is highlighted in the discussion (line 551).*

377-378 not a complete thought (or sentence)

*Sentence revised: "Results for habitat protection projects varied for different Bay segments and year window/closest project combinations, with no clear patterns."*

381-382 Very important and unique finding.

383 Effects of random restoration dates and locations Section: Strengthens the paper.

Discussion. Overall: The opening of the Discussion is rather boring in comparison to the strength of what came before. I think the fact that prior findings were supported belongs much further down. In fact, all of that material seems to belong in a "Verification of Approach" type of subsection of the Discussion. *We agree this is a soft start to the discussion. We have moved the conclusions paragraph to the beginning of the discussion to emphasize our key findings and value of the approach.*

440-442 Sounds a bit naïve relative to cost-benefit. Land acquisition can be extremely expensive in coastal regions with high demand and difficult to obtain grant money to achieve. Still, I support the conclusion in 445-448 based on experience with other conservation/restoration programs.

*This sentence was revised for clarity: "Categorization by effort combined with the associated estimates of water quality improvements provides a coarse evaluation of the tradeoffs associated with each project type" We also added a sentence about the difficulties of protection in response to a previous comment.*

Adequate consideration should be given to circling back to the objectives of the paper developed in the Introduction, and verbally checking off the boxes you've completed.

*The first paragraph of the discussion (formerly the conclusions paragraph) now explicitly acknowledges our objectives: "Consistent with our objectives, this new tool provides 1) a unique process..."*

#### Figures

1. Nice figure. Need to credit data sources in caption. "Have been monitored" implies a role by the authors without crediting others.

*Data sources were added to the figure caption.*

2. Very useful, especially the maps in b but also a.

3. Also very useful because this was hard to visualize from the text. However, the projects shown are the "easy" ones in terms of digitizing hydrologic distance, because they are in the bay so it was done "as the crow flies." How was hydrologic distance calculated for projects higher in the watershed? Hydrologic distance is usually digitized from the streamlines in a dataset but this was not described. It would be useful to add a second set of panels for the projects in the watershed that reveals this.

*The methods were clarified in the revisions, i.e., see our response to the general comment about the methods section. Based on this clarification, it is not appropriate to include such a figure and we have explicitly identified this approach as a future research objective.*

4. Reasonable effort. A little hard to understand.

*We hope our clarifications on the temporal slices in the methods have made this figure clearer.*

5. clear and useful but Fig 5b seems to indicate much longer time slices than are described in the text.

*Please see our revisions to the methods.*

6. This took a while to understand and is rather original, but ultimately worth the time for the information communicated.

7. Very interesting and informative! This type of information is gold for monitoring program design. May wish to highlight this more in the Discussion.

#### **Reviewer 2 comments:**

*We appreciate the comments provided below. Many of the comments about organization and language have been addressed by our revisions in response to the first reviewer. We have noted specific instances below.*

1) The organization might be improved.

P23 Assumptions and limitations section (Line 475) should be separated and the Assumptions should be moved to the section "Data synthesis and analysis framework",

*All of the assumptions of our approach were previously included in the methods section (e.g., lines 228, 234, 235). We retained this section in the discussion in its current form but have changed the title to more clearly emphasize the discussion points on the limitations, i.e., subsection title was changed to "Analysis limitations".*

2) The paragraphs (Line 307-320) are suggested to be moved to the Discussion section,  
*This content was moved to the discussion.*

3) Discussion section should be re-arranged with several sub-sections by the major findings, for example, the effectiveness of the proposed framework, limitation of this approach, etc.

*Please see our response to comments on the discussion from the first reviewer. Also note that we have moved the conclusions paragraph to the beginning of the introduction to highlight the major findings and value of this work.*

2. The language should be concise, particularly (1)the section Introduction is too tedious and the readers cannot catch the key point easily,

*Please see our responses to the first reviewer regarding our revisions to the introduction. The first reviewer notes several editorial changes that could improve the language in the introduction, similar to the comment herein. A majority of our responses to comments that cover lines 35 – 100 in the original draft address these concerns.*

# **Assessment of the cumulative effects of restoration activities on water quality in Tampa Bay, Florida**

Marcus W. Beck\* ([marcusb@sccwrp.org](mailto:marcusb@sccwrp.org)), Southern California Coastal Water Research Project,  
Costa Mesa, CA

Edward T. Sherwood ([esherwood@tbep.org](mailto:esherwood@tbep.org)), Tampa Bay Estuary Program, St. Petersburg, FL

Jessica Renee Henkel ([jessica.henkel@restorethegulf.gov](mailto:jessica.henkel@restorethegulf.gov)), Gulf Coast Ecosystem Restoration  
Council, New Orleans, LA

Kirsten Dorans ([kdorans@tulane.edu](mailto:kdorans@tulane.edu)), Tulane University School of Public Health and Tropical  
Medicine, New Orleans, LA

Kathryn Ireland ([kathryn.b.ireland@gmail.com](mailto:kathryn.b.ireland@gmail.com)), Montana State University, Bozeman, MT

Patricia Varela ([pvarela@geosyntec.com](mailto:pvarela@geosyntec.com)), Geosyntec Consultants Inc., Houston, TX

\*Corresponding author

## Abstract

Habitat and water quality restoration projects are commonly used to enhance coastal resources or mitigate negative impacts of water quality stressors. Significant resources have been expended for restoration projects, yet much less attention has focused on evaluating broad regional outcomes beyond site-specific assessments. This study presents an empirical framework to evaluate multiple datasets in the Tampa Bay area (Florida, USA) to identify 1) the types of restoration projects that have produced the greatest improvements in water quality, and 2) over which time frames different projects may produce water quality benefits. Information on the location and date of completion of 887 restoration projects from 1971 to 2017 were spatially and temporally matched with water quality records at each of 45 long-term monitoring stations in Tampa Bay. The underlying assumption was that the developed framework could identify differences in water quality changes between types of restoration projects based on aggregate estimates of chlorophyll-a concentrations before and after the completion of one to many projects. Water infrastructure projects to control point source nutrient loading into the Bay were associated with the highest likelihood of chlorophyll-a reduction, particularly for projects occurring prior to 1995. Habitat restoration projects were also associated with reductions in chlorophyll-a, although the likelihood of reductions from the cumulative effects of these projects were less than those from infrastructure improvements alone. The framework is sufficiently flexible for application to different spatiotemporal contexts and could be used to develop reasonable expectations for implementation of future water quality restoration activities throughout the Gulf of Mexico.

Key words: chlorophyll, long-term monitoring, restoration, Tampa Bay, trends

## Introduction

Despite considerable investments over the last four decades in coastal and estuarine ecosystem restoration (Diefenderfer et al. 2016), numerous challenges still impede comprehensive success. In the Gulf of Mexico (GOM), chronic and discrete drivers contribute to the difficulty in restoring and managing coastal ecosystems. For example, the synergistic effects of widespread coastal urbanization and climate change impacts will likely limit future habitat management effectiveness in the southeast United States (Enwright et al. 2016). Competing management and policy directives for flood protection, national commerce, and energy development complicate and prolong efforts to abate coastal hypoxia and other coastal water quality issues (Rabotyagov et al. 2014; Alfredo and Russo 2017). Disputes surrounding fair and equitable natural resource allocation often result in contentious implementation plans for the long-term sustainability of coastal resources (GMFMC 2017). Further, discrete tropical storm (Greening et al. 2006) and large-scale pollution events (Beyer et al. 2016) often reset, reverse or delay progress in restoring coastal ecosystems. These factors contribute to a complex setting for successful implementation of ecosystem restoration activities within the GOM.

In addition to these challenges, the difficulties of rigorously monitoring and understanding an ecosystem's condition and restoration trajectory at various spatial and temporal scales can further constrain evaluations of restoration success (Hobbs and Harris 2001; Liang et al. 2019). The lack of long-term environmental monitoring is a primary impediment to understanding pre- versus post- restoration change (Schiff et al. 2016) and also impedes recognition of any coastal ecosystem improvements derived from prolonged management, policy and restoration activities. Long-term coastal monitoring programs can facilitate a broader sense of how management,

policy and restoration activities affect coastal ecosystem quality (Borja et al. 2016). Utilizing lessons-learned from environmental monitoring programs, new frameworks are starting to emerge to better understand and facilitate coastal restoration ecology (Bayraktarov et al. 2016; Diefenderfer et al. 2016).

A very large, comprehensive and concerted effort to restore Gulf of Mexico coastal ecosystems is currently underway (GCERC 2013; 2016). Primary funding for this effort is derived from the legal settlements resulting from the 2010 Deepwater Horizon oil spill. Funding sources include: early restoration investments that were made immediately following the spill, natural resource damage assessments resulting from the spill's impacts (NRDA 2016), a record legal settlement of civil and criminal penalties negotiated between the responsible parties and the US government with strict US congressional oversight (United States vs. BPXP, et. al.), and matching funds from research, monitoring and restoration practitioners worldwide. These funds, equating to >\$20B US, present the Gulf of Mexico community an unprecedented opportunity to revitalize regional restoration efforts that will span multiple generations (GCERC 2013; 2016). Consequently, the restoration investments being made with these funds will be highly scrutinized. Better understanding the environmental outcomes of past restoration investments will help identify how, where and when future resources should be invested so that the Gulf Coast community can achieve the highest degree of restoration success.

Tampa Bay (Florida, USA) is the second largest estuarine embayment in the GOM and improvement in condition over the last four decades is one of the most exceptional success stories for coastal water quality management (Greening and Janicki 2006; Greening et al. 2014). Most notably, seagrass coverage in 2016 was reported as 16,857 hectares baywide, surpassing the goal of restoring coverage to 95% that occurred in 1950 (Sherwood et al. 2017). Reductions

in nutrient loading (Poe et al. 2005; Greening et al. 2014), chlorophyll-a concentrations (Wang et al. 1999; Beck and Hagy 2015), and improvements in water clarity (Morrison et al. 2006; Beck et al. 2018) have also preceded the seagrass recovery. Most of these positive changes have resulted from management efforts to reduce point source controls on nutrient pollution in the highly developed areas of Hillsborough Bay (Johansson 1991; Johansson and Lewis 1992). These controls allowed nutrient and chlorophyll-a targets to be met by the early 1990s. However, numerous smaller projects, including watershed-focused efforts (Lewis et al. 1998), may have had a supporting role in maintaining water quality improvements through contemporary periods. The cumulative effects of over 900 restoration projects, relative to broad watershed-scale management efforts, are not well understood. Understanding how implementation of these projects is associated with adjacent estuarine water quality at various spatio-temporal scales will provide an improved understanding of the link between overall estuary improvements and specific restoration activities.

Demonstrating success for restoration activities is challenging for several reasons (Ruiz-Jaen and Aide 2005; Wortley et al. 2013). Success may be vaguely or even subjectively defined because the effects of restoration could be described in different ways depending on project goals (Zedler 2007). For example, site-specific measures of before/after condition are commonly used measures of success, whereas downstream effects may be more important to consider for baywide conditions (Diefenderfer et al. 2011). More importantly, quantifying success as a measure of environmental improvements is challenged by the variety of factors that affect water quality across space and time. New tools are needed that can address these challenges to help guide and support GOM restoration. Here, we present an empirical framework for evaluating the influence of restoration projects on water quality improvements within Tampa Bay. The



framework helps synthesize routine, ambient monitoring data across various spatio-temporal scales to demonstrate how coastal restoration activities cumulatively affect estuarine water quality improvement. Data on water quality and restoration projects in the Tampa Bay area were used to demonstrate application of the analysis framework. Water quality and restoration datasets were evaluated to identify: 1) the types of restoration activities that most improve water quality, and 2) the time frames over which water quality benefits resulting from restoration may be resolved. Changes in chlorophyll-a concentrations, a proxy for negative eutrophication effects within Tampa Bay (Greening et al. 2014), were used as the success metric to evaluate estuarine restoration activities.

## Methods

### Study area

Tampa Bay is located on the west-central GOM coast of the Florida peninsula. Its watershed is among the most highly developed regions in Florida (Figure 1). More than 60 percent of land-use within 15 km of the Bay shoreline is urban or suburban (SWFWMD 2018). The Bay has been a focal point of economic activity since the 1950s and currently supports a mix of industrial, private, and recreational activities. The watershed includes one of the largest phosphate production regions in the country, which is supported by port operations primarily in the northeast portion of the Bay (Greening et al. 2014).

Current water quality in Tampa Bay is dramatically improved from the degraded historical condition. Nitrogen loads into the Bay in the mid-1970s have been estimated as  $8.9 \times 10^6$  kg/y, largely from wastewater effluent (Greening et al. 2014). In addition to reduced aesthetics,

124 hypereutrophic environmental conditions were common and included elevated chlorophyll-a and  
125 harmful algae, and reduced bottom water dissolved oxygen, water clarity, and seagrass coverage.  
126 Yields for some commercial and recreational species were also depressed, although emergent  
127 tidal wetland loss and fisheries practices (i.e., widespread Bay trawling and gill-netting) likely  
128 also contributed to declines (Comp 1985; Lombardo and Lewis 1985).

129 A long-term monitoring program in Tampa Bay has been instrumental in assessing and tracking  
130 restoration efforts. In the early 1970s, the initial baywide ambient monitoring program was  
131 established by a local environmental leader (Roger Stewart), which was subsequently  
132 institutionalized through State legislation by the creation of the Environmental Protection  
133 Commission of Hillsborough County. This occurred largely in response to citizen out-cry of the  
134 Bay's deteriorating ecology (Greening et al. 2014). Ongoing local support for this program has  
135 remained since 1972, and other local, municipal governments have created complementary water  
136 quality monitoring programs, all of which now support water quality assessments and  
137 management efforts spear-headed by the Tampa Bay Estuary Program (Sherwood et al. 2016).  
138 Some of the key attributes supporting the maintenance of this long-term monitoring program are  
139 summarized in Schiff et al. (2016) and Gross and Hagy (2017).

140 Nearly 900 public and private projects to improve water quality have also been completed in  
141 Tampa Bay and its watershed over the past 4 decades. These projects represent numerous  
142 voluntary (e.g., coastal habitat acquisition, restoration, preservation, etc.) and compliance-driven  
143 (e.g., stormwater retrofits, process water treatment upgrades, site-level permitting, power plant  
144 scrubber upgrades, improved agricultural practices, residential fertilizer use ordinances, etc.)  
145 activities. Linking data from the long-term monitoring program with data on these projects will

provide an understanding of how the cumulative effects of small-scale restoration have contributed to water quality relative to the historical water infrastructure upgrades.

## **Data sources**

Several databases were combined to document restoration projects in Tampa Bay and its watershed. Each database was unique and no overlap in documented projects was observed. Data from the Tampa Bay Water Atlas (version 2.3, <http://maps.wateratlas.usf.edu/tampabay/>; TBEP 2017) documented 253 projects from 1971 to 2007 that were primarily focused on habitat establishment, enhancement, or protection along the Bay's immediate shoreline or within the larger watershed area. Examples include restoration of salt marshes and mangroves, exotic vegetation control, and conversion of agricultural lands to natural habitats. Information on an additional 265 recent (2008-2017) projects was acquired from the US EPA's National Estuary Program Mapper (<https://gispub2.epa.gov/NEPmap/>). This database provides only basic information, such as year of completion, geographic coordinates, general activities, and areal coverage. Data from the TBEP Action Plan Database Portal (<https://apdb.tbeptech.org/index.php>) documented locations of infrastructure improvement projects, structural best management practices, and policy-driven stormwater or wastewater management actions. This database included 368 projects from 1992 to 2016 for county, municipal or industrial activities, such as implementation of best management practices at treatment plants, creation of stormwater retention or treatment controls, or site-specific controls of industrial and municipal point sources.

For all restoration datasets, shared information included the project location, year of completion, and classification of the restoration activity. We developed and applied a two-level classification

scheme that described each restoration project as 1) a habitat or water infrastructure improvement and 2) more specifically as enhancement, establishment, or protection for habitat or as nonpoint or point source controls for water infrastructure. These categories were used to provide a broad characterization of restoration activities that were considered to contribute to improvements in water quality over time. The final combined dataset included 887 projects from 1971 to 2017 (Figure 2). Projects with incomplete information were not included in the final dataset.

Water quality data in Tampa Bay have been collected consistently since 1974 by the Environmental Protection Commission of Hillsborough County (Sherwood et al. 2016; TBEP 2017). Data were collected monthly at forty-five stations using a water sample or monitoring sonde at bottom, mid- or surface depths, depending on parameter. The locations of monitoring stations were fixed and cover the entire Bay from the uppermost mesohaline sections to the lowermost euhaline portions that have direct interaction with the GOM (Figure 1). Mid-depth water samples at each station are laboratory processed immediately after collection. Chlorophyll-*a* ( $\mu\text{g/L}$ ) and total nitrogen ( $\text{mg/L}$ ) measurements at each site were used for analysis, totaling up to 515 discrete observations for each station. Total nitrogen concentrations were included in initial data assessments, as this nutrient is considered limiting in Tampa Bay (Greening et al. 2014).

## **Data synthesis and analysis framework**

The five sub-categories for each project (habitat enhancement, establishment, and protection; nonpoint and point source controls) were separately evaluated to describe the likelihood of changes in water quality associated with each type. Water quality monitoring sites were matched

to the closest selected restoration projects and changes in the water quality data were evaluated relative to the completion dates of the selected projects. Spatial and temporal matching can be accomplished using several methods that vary in complexity. For example, hydrologic distances or other non-Euclidean distance weightings by watershed topology can be used to link measurements to modelled locations in space (Curriero 2006; Gardner et al. 2011). However, we adopted a relatively simple approach with limited data requirements to maximize potential applications in other regions (e.g., no hydrology data are needed, only spatial location). The matchings began with a spatial join wherein the Euclidean distances between each water quality station and each restoration project were quantified. The restoration projects closest to each water quality station were identified using the distances between projects and water quality stations. The distances were also grouped by the five restoration project types (i.e., habitat protection, nonpoint source control, etc.) such that the closest  $n$  sites of a given project type could be identified for any water quality station (Figure 3).

For each spatial match, temporal matching between water quality stations and restoration projects was obtained by subsetting the water quality data within a time window before and after the completion date of each restoration project (Figure 4). For the closest  $n$  restoration sites for each of five project types, two summarized water quality estimates were obtained to quantify a before and after estimate of chlorophyll-a associated with each project. The before estimate was the average of observations for the year preceding the completion of a project and the after estimate was the average of observations for a selected window of time (e.g., five years) that occurred after completion of a project. The before estimate for the year prior established the basis of comparison for the water quality estimates in the selected window of time after project completion, where the latter could be manually changed to characterize a potential duration of

time within which water quality could improve after project completion. The final two estimates of the before and after values of the five types of restoration projects at each water quality station were based on an average of the  $n$  closest restoration sites, weighted inversely by distance from the monitoring station. Lastly, no data were available on project duration and we assumed that the year associated with each project was generally inclusive of project implementation and completion. Time windows that overlapped the start and end date of the water quality time series were discarded.

Change in water quality relative to each type of restoration project was estimated as:

$$\Delta WQ = \frac{\sum_{i=1}^n \widehat{wq} \in win + proj_{i,dt}}{n \cdot dist_{i \in n}} - \frac{\sum_{i=1}^n \widehat{wq} \in proj_{i,dt} - win}{n \cdot dist_{i \in n}} \quad (1)$$

where  $\Delta WQ$  was the difference between the after and before averages for each of  $n$  spatially matched restoration projects. For each  $i$  of  $n$  projects ( $proj$ ), the average water quality ( $\widehat{wq}$ ) within the window ( $win$ ) either before ( $proj_{i,dt} - win$ ) or after ( $win + proj_{i,dt}$ ) the completion date ( $dt$ ) for project  $i$  was summed. The summations of water quality before and after each project were then divided by the total number of  $n$  matched projects, multiplied by the distance of the projects from a water quality station ( $dist_{i \in n}$ ). This created a weighted average of the before-after estimates for each project that was inversely related to the distance from a water quality station. A weighted average by distance (or parametric distance weights; Sickie and Johnson 2008) was used based on the assumption that restoration projects farther from a water quality station will have a weaker association with potential changes in chlorophyll-a. The total change in water quality for a project type was simply the difference in weighted averages. This process was repeated for every station (Figure 5). Overall differences between project types were

evaluated by ANOVA F-tests, whereas pairwise differences between project types were evaluated by t-tests with corrected probability values for multiple comparisons.

One of the key assumptions of our approach is that restoration projects will benefit water quality through a reduction in chlorophyll-a. We make no assumptions about the expected magnitude of an association given that the model does not describe a specific mechanism of change, nor do we make any explicit assumption about the direction of change (i.e., two-tailed hypothesis tests were used), although a general assumption was that chlorophyll-a would decrease over time in agreement with known changes in water quality. However, we hypothesized that the magnitude of chlorophyll-a changes varies by project type, and number of projects or length of time window evaluated. An expected outcome is that explicit, quantitative conclusions can be made about the relative differences between projects types, particularly regarding how additional projects of a particular type could benefit water quality and within what general time windows a change might be expected (Diefenderfer et al. [2011](#)).

The model was also designed to quantify cumulative relationships of restoration projects with water quality at different spatial scales. In eqn. (1), the association of a restoration type with chlorophyll-a is estimated for one water quality station, whereas estimates from several water quality stations can be combined to develop an overall description of a particular restoration type as it applies to an areal unit of interest, potentially over broad regional scales. For example, estimated associations of point source control projects with each water quality station in the Bay can be combined to develop an overall narrative of how these projects could (assuming a causal relationship) influence environmental change across the entire Bay. Estimates across stations were evaluated to describe associations in baywide improvements from various restoration project types throughout the watershed. Estimates were also evaluated by individual Bay

segments that have specific management targets for chlorophyll-a concentration (Florida Statute 62-302.532; Janicki et al. 1999). Stratification by Bay segments provided an alternative context for interpreting the results based on areal differences between segments and how restoration projects varied in space and time. Evaluating the results at different scales can also provide insights into potential (or lack of) stressors and processes controlling the impacts, which can help prioritize management actions by location (Diefenderfer et al. 2009; Thom et al. 2011).

The analysis of each project type was bounded by two key parameters in eqn. (1). These include  $n$ , the number of spatially-matched restoration projects used to average the cumulative estimate of each project type, and  $win$ , the time windows before and after a project completion date that were used to subset a station's water quality time series. These boundaries affected our ability to characterize each restoration project type with water quality changes. Identifying values that maximized the difference between before and after water quality measurements was necessary to quantify how many projects were most strongly associated with a change in water quality, the time within which a change is expected, and the magnitude of an expected change between project types. For simplicity, we evaluated different combinations of five or ten year time windows from the date of each project completion and the five or ten closest projects to each water quality station. All analyses were conducted with customized scripts created for the R statistical programming language (RDCT 2018).

## **Testing effects of restoration dates and location**

Because of the documented improvements in water quality in Tampa Bay, a concern with our approach is that any association between restoration projects and chlorophyll-a may result from correlations between the two parameters, confounding a true demonstration of water quality



improvements in relation to restoration activities. To address this challenge, estimated changes in chlorophyll-a were evaluated in response to temporal and spatial matching with restoration projects, as above, but with random date and location assignments for each restoration project that were then compared to the actual results. An expected outcome of randomization is that no differences are observed between project types and that all associations between projects and chlorophyll-a changes should reflect the continuous decline of chlorophyll-a over time, as observed in the independent water quality record. In other words, the randomization creates a null model where the estimated effects of restoration projects would not differ from a simple evaluation of trends in the raw data - slicing the observed time series by random dates and evaluating before/after averages with random projects is expected to reflect the known decline of chlorophyll-a in the raw data. Alternatively, evidence that our framework provides meaningful results would be supported by differences in chlorophyll-a changes between project types and the timing associated with the changes.

## **Results**

### **Water quality observations**

Chlorophyll-a in Tampa Bay decreased over the forty-year record consistent with documented changes (Wang et al. 1999; Greening et al. 2014; Beck and Hagy 2015) (Table 1). Median concentrations were highest from 1977 to 1987 (median 13.40  $\mu\text{g/L}$  at low salinity stations  $<26.5$  psu, 7.30  $\mu\text{g/L}$  at high salinity stations  $>26.5$  psu). Declines were monotonic throughout the period of record with the largest reductions occurring during the first twenty years (34% decrease), followed by consistent but smaller reductions in concentrations later in the time series.

300 A 34% decrease at low salinity stations and a 30% decrease at high salinity stations was  
301 observed between the periods of 1977-1987 to 1987-1997. Seasonally, chlorophyll-a  
302 concentrations were highest in the late summer/early fall periods (median 13.80  $\mu\text{g/L}$  at low  
303 salinity stations, 7.23  $\mu\text{g/L}$  at high salinity stations, across all years). Total nitrogen  
304 concentrations had similar trends as chlorophyll-a, although a steady decline was observed  
305 across the entire time series rather than primarily in the first two decades in contrast to  
306 chlorophyll-a (Poe et al. 2005; Greening et al. 2014). An exception for total nitrogen was  
307 observed at high salinity stations where concentrations were relatively constant at approximately  
308 0.55 mg/L from 1987 to 2007. Seasonally, total nitrogen concentrations peaked in the late  
309 summer/early fall period.

310 The monotonic decline in chlorophyll-a concentrations was mirrored by increases in the number  
311 and types of restoration projects in the watershed, where the number of documented projects  
312 increased after 2000 (Figure 2). For the entire record, 275 (31% of total) habitat enhancement,  
313 259 (29%) habitat establishment, 45 (5%) habitat protection, 248 (28%) nonpoint source, and 60  
314 (7%) point source control projects were documented. Individual point source controls early in the  
315 record were those that occurred in the historically polluted upper Hillsborough Bay and adjacent  
316 to the city of St. Petersburg (Johansson 1991; Johansson and Lewis 1992; Lewis et al. 1998).  
317 Prior to 1995, only 11 water infrastructure projects (three nonpoint control, eight point source  
318 controls) were documented in the database, whereas 70 habitat projects were recorded (50 habitat  
319 establishment, 20 habitat enhancement). Nearly ten times as many restoration projects were  
320 completed in 1995 to present (806 total), with notable increases in the number of nonpoint source  
321 controls (245) and habitat protection projects (45).

## Associations between restoration projects and water quality change

Before employing our analytical approach, we evaluated temporal trends in water quality and possible drivers of water quality change to develop an analytical baseline for comparison. A simple analysis of water quality measurements versus the cumulative number of restoration projects over time showed a decrease in both total nitrogen and chlorophyll-a with additional restoration effort. Analysis of median water quality estimates across all monitoring stations for a given year versus the cumulative number of restoration projects as of that year showed that water quality was related to the number of projects for all project types (based on linear models,  $\alpha = 0.05$ ; Figure 6). Associations with number of projects were relatively strong for total nitrogen and relatively weaker for chlorophyll-a across project types. Decreases in total nitrogen were most strongly associated with water infrastructure projects for nonpoint source ( $F = 65.5, df = 1,23, p < 0.005$ ) and point source controls ( $F = 60.8, df = 1,21, p < 0.005$ ), as expected. Habitat protection projects were also strongly associated with decreases in total nitrogen ( $F = 34.8, df = 1,14, p < 0.005$ ). For chlorophyll-a, the strongest associations were observed with habitat establishment ( $F = 20.8, df = 1,35, p < 0.005$ ) and point source control ( $F = 13.7, df = 1,22, p < 0.005$ ) projects. A marginally significant association was observed between chlorophyll-a and cumulative habitat protection projects ( $F = 4.6, df = 1,14, p = 0.049$ ).

In contrast to results in Figure 6, baywide estimates of the effects of restoration projects using spatial-temporal matching depended on the year window sizes and number of nearby restoration projects matched to each water quality station (Figure 7). Estimated associations of different projects types with chlorophyll-a at individual stations are shown in the left maps and the

baywide aggregate associations across all stations for a given project type in the right plots. Station points in the left maps correspond to the change estimate for the year window and closest project type selections for each project type that were obtained through the steps in Figure 5 and eqn. (1). Stations outlined in black have significant results based on t-tests of the mean estimates of chlorophyll-a relative to zero change. The plots on the right are based on the baywide distributions of the estimated water quality changes for all stations for the corresponding project types in the maps on the left. The plots on the right also include statistical summaries for 1) an analysis of variance (ANOVA) F-test to compare the distribution of water quality changes between project types, 2) individual t-tests for each project type to evaluate changes that were different from zero, and 3) a multiple comparison test denoted by letters to identify which project types had changes that were different from each other.

For site-specific estimates of water quality changes, longer time windows and more project matches with each monitoring station increased observations of significant associations (i.e., more black circles in the maps in Figure 7d, compared to Figure 7a). This was particularly true for habitat protection projects where no significant associations were observed for the 5 year window, 5 closest projects combination, but twelve stations had significant associations for the 10 year window, 10 closest projects combination. A similar trend was observed for point source control projects where more stations had more significant reductions in chlorophyll-a with the 10 year window, 10 closest projects. For nonpoint source projects, the greatest number of stations ( $n = 13$ ) with significant improvements in water quality was observed for the 5 year window, 10 closest projects combination. Associations of habitat enhancement and habitat establishment projects with water quality stations were inconsistent, with some sites showing an increase or decrease that varied by the year window, closest project combinations. Spatial patterns among

stations regarding associations with different project types were also not clear, although point source controls were more commonly associated with improvements in mid-Bay stations (Middle Tampa Bay segment, see Figure 1).

The estimated baywide effects for each project type showed that point source controls were more strongly associated with reductions in chlorophyll-a than the other project types (Figure 7, right plots). This association was particularly strong for the ten year window combinations (Figure 7c, d), where the results suggested an overall baywide reduction in chlorophyll-a of approximately 2  $\mu\text{g/L}$ , depending on the number of projects implemented (median change across all sites: reduction of 2.7  $\mu\text{g/L}$  for 10 years, 5 closest projects and 1.6  $\mu\text{g/L}$  for 10 years, 10 closest projects). Nonpoint source controls were also significantly associated with chlorophyll-a reductions, but only when a large number of projects were implemented (10 closest project combinations, Figure 7b, d). Additionally, the magnitude of nonpoint source control reductions were less than point source controls (reduction of 0.7  $\mu\text{g/L}$  for 5 years, 10 closest projects and 0.5  $\mu\text{g/L}$  for 10 years, 10 closest projects). Habitat protection projects were also significantly associated with baywide changes for all year window, closest project combinations, with the largest estimated reduction of 1  $\mu\text{g/L}$  for the 5 year window, 5 closest projects combination. Habitat enhancement and establishment projects were not strongly associated with baywide changes in chlorophyll-a, with the exception of habitat establishment for the 10 year window, 10 closest projects combination (0.9  $\mu\text{g/L}$  reduction).

The above analysis was repeated for individual Bay segments to identify spatial variation in associations of restoration projects with water quality changes. Table 2 provides similar information as the plots on the right side of Figure 7, where results are presented similarly but for each of four Bay segments (HB: Hillsborough Bay, LTB: Lower Tampa Bay, MTB: Middle

Tampa Bay, OTB: Old Tampa Bay, Figure 1) for each year window, closest projects combination. As for the baywide result, point source controls were most consistently associated with reductions in chlorophyll-a, particularly for ten year window combinations. Nonpoint source controls were also important, although significant associations with chlorophyll-a changes were limited to the Middle and Old Tampa Bay segments. Results for habitat protection projects varied for different Bay segments and year window/closest project combinations, with no clear patterns. Habitat establishment projects were most strongly associated with changes in each Bay segment for the 10 year window, 10 closest projects combination, with the exception of Hillsborough Bay where the relationship was not significant. Chlorophyll-a changes in Lower Tampa Bay were significantly associated with habitat enhancement and establishment projects for the 5 year window, 10 closest projects combination.

## **Effects of random restoration dates and locations**

A comparison of the baywide results (Figure 7, right side) to results from simulations where dates and locations were randomized for each restoration project suggested that the framework in eqn. (1) is robust. The same year windows and closest project combinations were evaluated as above (i.e., 5/10 year windows, 5/10 closest projects), but with 1000 simulations where the date and location of each restoration were randomized (i.e., random draw from uniform distribution of years from 1971 to 2017, random draw from uniform distribution of latitude and longitude based on the bounding box of the study area). Nearly all of the simulated results suggested that each project was associated with a decline in chlorophyll-a (Table 3, values in bold, mean < 0). This is consistent with our null hypothesis that randomization would simply reflect the long-term decline in chlorophyll-a that is apparent in the observed water quality records. Some differences

were observed in the 5 years, 5 projects combination where no change (mean = 0) was the most observed outcome from the simulations. These inconsistencies with our null hypothesis may be the result of using relatively small windows and project combinations, i.e., slicing the data too thin to detect the long-term decline in chlorophyll-a.

The “Actual” and “Agreement” columns in Table 3 indicate the estimated changes for each project using the actual restoration dates/locations and if the result agrees with those from the random simulations. In support of the alternative hypothesis, nine of the rows in the “Agreement” column indicate a result different than a consistent decline in chlorophyll-a expected under the null hypothesis. Compared to random simulations, different results were more often observed for habitat enhancement and habitat establishment projects, where the simulated results most often suggested a decrease and the actual results suggested no change in chlorophyll-a. Nonpoint and point source control projects were in agreement with simulated results, although this does not provide sufficient evidence that the results from the actual data are incorrect. Because the null hypothesis under randomization suggests projects will be associated with water quality improvements based on the independent chlorophyll-a time series, an observed decline in chlorophyll-a in relation to actual restoration projects could still suggest a signal rather than a false positive result. There is no way of identifying type I errors with the current dataset, although the differences from the null results for habitat enhancement and establishment projects do suggest the framework is robust.

## **Discussion**

A long-term record of restoration activities and water quality data in Tampa Bay provided the foundation to develop a novel decision support tool for coastal restoration practitioners and

managers. Consistent with our objectives, this new tool provides 1) a unique process to understand the associations between past restoration projects and known changes in water quality, and 2) establishes, under certain assumptions, an expectation of water quality improvements that could result from future restoration activities contingent upon the level of investments in different activities and the necessary time to monitor any observed downstream water quality benefits at local to watershed scales. Overall, we demonstrated a baywide association of water quality changes to different restoration activities that varied by project type, while refining parameters for estimating the results, including the spatial context of interpretation. The flexibility of our approach has potentially broad application and extension within the Gulf Coast restoration and management community.

The results support several conclusions that are consistent with recognized, long-term changes in water quality in Tampa Bay. Water infrastructure projects related to point and nonpoint source controls were consistently associated with improved water quality. The record of restoration projects included key point source nutrient controls that occurred primarily in upper Tampa Bay (Hillsborough Bay) and that were successful in reducing nutrient loads during the first two decades of observation (Johansson [1991](#); Johansson and Lewis [1992](#); Greening et al. [2014](#)).

These outcomes were expected and the ability of the results to clearly demonstrate these long-term associations provided a proof of concept for the overall approach. Moreover, efforts focused on mitigating effects of nonpoint sources of pollution were more common in the latter half of record after 1990 and our results provide evidence that these projects have been effective in improving water quality baywide, as well. Nonpoint source control efforts broadly described several activities that included, among others, street sweeping, education/outreach efforts, and various best management practices for stormwater, agricultural, and wetland management



programs. The ability to document effects of nonpoint control efforts on water quality relative to end-of-pipe controls is challenging (Hassett et al. 2005; Meals et al. 2010; Liang et al. 2019) and the results suggest that our approach is capable of detecting improvements in water quality from these projects when many are implemented.

Habitat restoration projects were also associated with reductions in chlorophyll-a, although to a lesser magnitude than water infrastructure projects. Our categorization of habitat projects as enhancement, establishment, and protection were developed to better understand potential effects on water quality related to the type and intensity of actions for each group. Specifically, the categories represented extremes from low to high intensity effort, where protection was low effort (e.g., direct land acquisition), establishment was highest effort (e.g., mangrove/seagrass plantings, creation of oyster reefs), and enhancement was moderate effort depending on the activity (e.g., hydrologic restoration for wetlands, exotic species control). Categorization by effort combined with the associated estimates of water quality improvements provides a coarse evaluation of the tradeoffs associated with each project type. For example, habitat protection was consistently linked to chlorophyll-a reductions independent of year windows and number of projects. The effort for land acquisition is minimal relative to the other habitat restoration projects. Conversely, habitat enhancement was not strongly associated with baywide improvements in water quality and such projects may require more intensive effort and monitoring assessments to understand and contribute towards downstream water quality benefits. Based on these results, habitat protection may be a more immediate and efficient approach than other types of habitat restoration projects, especially if the primary restoration objective is to quickly improve downstream water quality.

Our results also provide an approach to identify an expected range of time and number of projects that are associated with potential improvements in water quality (Diefenderfer et al. 2011). This information was included as an explicit component in eqn. (1) to quantify tradeoffs for different restoration activities based on how results varied by time and effort, similar to the categorization for habitat projects. Monitoring water quality improvements after a short period of time since project completion and with fewer projects (e.g., 5 year window, 5 closest projects) may be more efficient than those where improvements are observed after longer periods of time and with more projects (e.g., 10 year window, 10 closest projects), dependent on the project type. Given this logic, both habitat protection and point source controls could potentially provide the greatest measured water quality benefits for the least effort, whereas other projects provide lesser improvements, require more time to confer water quality benefits (e.g., full maturation of a habitat enhancement/establishment site), and require more projects to be implemented to contribute to significant water quality improvements (e.g., implementing many nonpoint source controls across a watershed). However, this approach assumes that immediate downstream water quality improvements with minimal effort are the primary restoration objectives and implicitly discounts the long-term effects that may or may not persist for a given project type or regional restoration challenges. Protection may also be costly in developed areas with competing land use interests, despite minimal restoration requirements. Likewise, other restoration objectives may be a primary driver for pursuing a particular project type (e.g. increasing biodiversity, improving fish and wildlife habitats, etc.).

Conclusions from our approach may also be sensitive to system hysteresis. An initial improvement of water quality associated with a particular project may be obfuscated by chronic degraded conditions, if additional restoration projects that directly address the underlying

problem are not pursued (i.e., Scheffer et al. 1998; Borja et al. 2010). As an example, habitat enhancement projects were associated with improvements in Lower Tampa Bay only during the five year window when ten projects were implemented; reductions in chlorophyll-a were not shown to persist at the ten year window.

## **Analysis limitations**

There are several limitations of our approach that affect the interpretation of the results. These assumptions and limitations are a reflection of 1) the inherent uncertainty in quantifying baywide effects of restoration projects that vary considerably in mechanisms affecting water quality (e.g., Baird 2005; Borja et al. 2010), and 2) explicit construction of the approach to best account for this uncertainty. Because we do not know the true effect of restoration projects, the results can be interpreted as worst- or best-case descriptions depending on how much certainty is reflected in the estimates. At worst, we provide an approach that can identify the closest restoration projects that have occurred near a water quality monitoring site and a means to compare water quality between project types. We consider this valuable information for managers even if there is considerable uncertainty in the estimates of change; there are currently no tools that match restoration projects to water quality records in Tampa Bay. At best, we provide a decision support tool that provides managers with an expectation of water quality benefits associated with restorations actions, and an estimate of time required for improvements to be observed (Diefenderfer et al. 2011). Both kinds of information are critical to inform and sustain environmental restoration programs.

The value of our approach to quantify cumulative effects of restoration activities on improving water quality is likely between the worst- and best-case scenarios outlined above given the

confidence that can be invested in the conclusions. The ability of our model to support previously and well-described changes in water quality in response to key management interventions (e.g., improvements from point source controls; Greening et al. 2014; Beck and Hagy 2015) provides additional assurance and weight of evidence that our approach improves our understanding of restoration effectiveness beyond basic relationships presented in Figure 6. The latter was specifically presented to demonstrate limitations of simpler analysis methods. For example, the separate effects of habitat establishment and point source projects on chlorophyll-a reductions cannot be separated through simple linear analyses because both increase over time, i.e., an association of chlorophyll-a with one project type could be an artifact of an association with another project type. Likewise, a weak association (e.g., habitat protection and chlorophyll-a) does not provide strong evidence that a particular project type is unimportant for water quality improvements, given that the simple analysis may lack sufficient power to detect an association. Operating under these constraints, an explanation of how the new approach can guide management is required to minimize extrapolation of conclusions beyond a reasonable level of confidence in what is provided by the results (i.e., a levels-of-evidence reasoning; Diefenderfer et al. 2011).

As an example, a practitioner could use the following logic derived from our approach to guide future restoration efforts. Figure 7 suggests that point source controls were responsible for an approximate baywide chlorophyll-a reduction of 2  $\mu\text{g/L}$ . These results were observed at the 10 year window and 5 (and 10) closest project combinations. What exactly does this information suggest and what expectations can be derived regarding the likely effects of future point source control efforts? First, an inaccurate conclusion is that baywide chlorophyll-a would be reduced by 2  $\mu\text{g/L}$  after ten years, if five projects are implemented in the future. A more correct

interpretation is that, historically, the aggregate effect of point source control projects for a “typical” station at any point in the record and location in the Bay has been a reduction of 2  $\mu\text{g/L}$  after about ten years in response to 5-10 projects, all of which were completed at different times. Importantly, this latter interpretation assumes causal relationships between the projects and changes in chlorophyll-a. This description is also understandably vague, but it provides context of an expectation relative to other types of projects, particularly in narrative terms. For example, the aggregate effects of habitat establishment projects were most apparent for the ten year window, ten closest projects combination. With this information, qualitative conclusions about the relative effects of point source control versus habitat establishment can be made. Point source control projects are “more effective” in improving downstream water quality because improvements are expected “quicker” and with “fewer” implemented projects, as applied to a “typical” water quality station that could be at any location in the Bay. Similar but alternative conclusions could be made in different spatial contexts (e.g., “typical” stations in Hillsborough Bay, Table 2).

The flexibility and simplicity of the approach to quantify associations between restoration projects and water quality was purposeful given the constraints of the data. Although the compiled restoration databases included additional information on effort (e.g., acreage restored), these data were not consistently collected and our approach was constrained to the most basic information about each project (i.e, type, completion date, and location). Explicit monitoring of project effectiveness pre- and post-completion is atypical (Neckles et al. 2002; NASEM 2017) and our minimal dataset describing the when and where for each project is a more available description of restoration effort across systems, especially in historic contexts. Within these constraints, the model was strictly associative and any conclusions do not provide a mechanistic

570 explanation. These limitations in our associative approach were apparent in some of the results.  
571 Specifically, significant increases in chlorophyll-a associated with particular projects and  
572 year/site combinations were observed (e.g., Figure 7d, three sites for habitat enhancement).  
573 These trends are contrary to expectations and highlight shortcomings where the simple design  
574 may not have adequately accounted for improvements in downstream water quality (e.g., full  
575 ecosystem maturation of a restoration site). Our aggregated estimates using all stations to  
576 describe a baywide association were partly meant to reduce the influence of some of these  
577 spurious results.

578 The simplicity of our approach also means that it is highly adaptable to novel contexts. A  
579 primary goal of this study was to develop a decision support tool that could be applied  
580 elsewhere. We used Tampa Bay as an example where the outcome was partially known and a  
581 rich dataset was available, affording us a prior expectation of the results. Application to  
582 additional systems would require, at minimum, water quality observations spanning multiple  
583 years and a similar dataset of completed restoration projects. Our categorization that described  
584 relevant water and habitat related restoration projects was specific to Tampa Bay, but our  
585 approach can include different project definitions and specificity depending on the types of  
586 activities that may have occurred and their expected benefits to water quality in a different  
587 system. Similarly, the flexibility of our approach to accommodate different year windows and  
588 number of projects provides a diagnostic that is sensitive to both the restoration effort expressed  
589 in a dataset and how the potential associations could be interpreted. Lastly, we demonstrated  
590 flexibility in the spatial context from estimated changes at discrete locations to entire system-  
591 wide responses. Although there is uncertainty associated with these interpretations (noted  
592 above), the ability to accommodate different spatial contexts means the approach can be readily

applied to different systems or management questions at various scales - from a single monitoring station to a regional monitoring network.

## **Future directions**

Our approach is not without limitations and future research could build on the methods to provide an improved assessment of restoration effectiveness. Our geospatial analyses were relatively simple, in that spatial matchings were accomplished through Euclidean distances. Alternative distance measures could be used that consider hydrologic distances following flow networks in the watershed. The importance of these approaches could provide insight into pollutant dispersal pathways in environments with low elevation gradients, such as Florida. Weighting restoration projects by relative effort could also facilitate an improved assessment of effectiveness, such as considering total restoration area as an important variable to consider for water quality improvements. Some of these data were available in our compiled dataset, although coverage was insufficient for a complete analysis. Finally, the social and human dimensions of different restoration projects were not considered herein but are important factors that can be equally or even more relevant determinants of success that should be considered when weighing restoration options. Future restoration and monitoring activities should adopt a more comprehensive evaluation of success measures that includes and extends beyond water quality changes.

## **Supplement**

Data analysis frameworks, analysis code, and manuscript source materials are available at <https://doi.org/10.5281/zenodo.3243530>. An interactive application exploring results for Tampa

Bay and individual Bay segments is available at  
[https://shiny.tbep.tech.org/restoreTB/ind\\_eval.Rmd](https://shiny.tbep.tech.org/restoreTB/ind_eval.Rmd).

## Acknowledgments

This manuscript was a direct result of the Open Science for Synthesis (OSS): Gulf Research Program workshop convened by the University of Santa Barbara, National Center for Ecological Assessment and Synthesis in July 2017. We are greatly indebted to the workshop instructors, particularly Matt Jones, Bryce Mecum, Julien Brun, Chris Lortie, Amber Budden, Leah Wasser, and Tracy Teal, for inspiring and motivating our use of open science tools in developing this manuscript. We would also like to thank the many TBEP partners and collaborators for their continuing efforts to restore and monitor Tampa Bay. The progress achieved in restoring the Tampa Bay ecosystem over recent decades would not be possible without the collaborative partnerships fostered in the region. Our partners' willingness to adapt and implement innovative monitoring and management actions in response to the ever evolving challenges threatening Tampa Bay is greatly appreciated. We also thank James Hagy, anonymous reviewers, and editors for their thoughtful comments in improving this paper. This project was partially funded through the OSS: Gulf Research Program workshop, EPA Section 320 Grant Funds, and TBEP's local government partners (Hillsborough, Manatee, Pasco and Pinellas Counties; the Cities of Clearwater, St. Petersburg, and Tampa; Tampa Bay Water; and the Southwest Florida Water Management District) through contributions to the TBEP's operating budget. K. S. Dorans was supported by the National Institute of General Medical Sciences (grant 1P20GM109036- 01A1).



## Figure captions

*Figure 1 Water quality stations and restoration projects in the Tampa Bay area. Water quality stations have been monitored monthly since 1974. Locations of restoration projects represent 887 records that are generally categorized as habitat or water infrastructure projects from 1971 to present. Bay segments as management units of interest are shown in the upper right inset. HB: Hillsborough Bay, LTB: Lower Tampa Bay, MTB: Middle Tampa Bay, OTB: Old Tampa Bay. Water quality data from the Hillsborough County Environmental Protection Commission (TBEP 2017), restoration project data from the Tampa Bay Water Atlas (<http://maps.wateratlas.usf.edu/tampabay/>; TBEP 2017), US EPA National Estuary Program Mapper (<https://gispub2.epa.gov/NEPmap/>), and the Tampa Bay Estuary Program Action Plan Database Portal (<https://apdb.tbep.tech.org/index.php>).*

*Figure 2 Counts (top) and locations (bottom) of restoration project types over time in the Tampa Bay watershed. Restorations were categorized as water infrastructure (blue; nonpoint source controls, point source controls) and habitat (green; enhancements, establishments, protection) projects. The compiled restoration database included records of 887 project types and locations from 1971 to 2017.*

*Figure 3 Spatial matching of water quality stations with restoration projects. Spatial matches of each water quality station (blue dots) with habitat (solid line to grey dots) and water infrastructure (dashed line to black dots) projects are shown as the closest single match by type on the left and the “n” closest matches on the right. The spatial matches were made for the five restoration project types within the broader habitat and water categories shown in the figure.*

Figure 4 Temporal matching of restoration project types with time series data at a water quality station. The restoration project locations that are spatially matched with a water quality station (a) are used to create a temporal slice of the water quality data within a window of time before and after the completion date of each restoration project (b). Slices are based on the closest “n” restoration projects by type ( $n = 2$  in this example) to a water quality station. The two broad categories of habitat and water infrastructure projects are shown in the figure as an example, whereas the analysis evaluated all five restoration categories.

Figure 5 Steps to estimate cumulative water quality changes at a single station associated with a selected number of projects and time windows. Subplot (a) shows a station in Middle Tampa Bay matched to the five nearest restoration projects for each of five types (some are co-located). The time slices of the water quality observations for one year prior and ten years after the completion of each project are shown in (b), ordered from near to far. The before/after water quality averages for the slices are shown in (c) and the differences between the two are shown in (d). Finally, the weighted averages for the five closest matches by project type are shown in (e) with 95% confidence intervals.

Figure 6 Relationships between cumulative number of restoration projects over time and water quality observations in Tampa Bay. The plot shows median total nitrogen (mg/L) and chlorophyll-a ( $\mu\text{g/L}$ ) across all monitoring stations for each year against the cumulative number of projects for all preceding years. Points are sized and shaded by year to show the progression of water quality and number of projects over time. Summary statistics are shown in the bottom left corner as the significance of the linear regression (stars) and R-squared value.  $p > 0.05$  ns,  $p < 0.05$  \*,  $p < 0.005$  \*\*

Figure 7 Associations of restoration projects with chlorophyll-a changes at all sites in Tampa Bay. Associations were evaluated based on different year windows (5, 10) since completion of restoration projects and number of closest restoration projects (5, 10) to each monitoring station (subfigures a-d). The left plots show the estimated changes at each site (green decreasing, red increasing) for each restoration project type, with significant changes at a site outlined in black. The right plots show the aggregated site changes for each project type. Overall differences between project types were evaluated by ANOVA F-tests (bottom left corner, right plots), whereas pairwise differences between project types were evaluated by t-tests with corrected p-values for multiple comparisons. Chlorophyll-a changes by project types within each subfigure that are not significantly different share a letter and significance of the within-group mean relative to zero is also shown.

## Tables

*Table 1: Summary of total nitrogen and chlorophyll-a observations from monitoring stations in Tampa Bay. Minimum, median, and maximum observed values for low and high salinity conditions are shown for seasonal and annual aggregations of water quality observations at all monitoring stations (See Figure 1). Low or high salinity is based on values below or above the long-term baywide median (26.5 psu). JFM: January, February, March; AMJ: April, May, June; JAS: July, August, September; OND: October, November, December.*

Salinity	Time period	Total Nitrogen (mg/L)			Chlorophyll-a (µg/L)		
		Min	Median	Max	Min	Median	Max
lo	JFM	0.00	0.46	2.69	0.12	5.30	114.40
	AMJ	0.03	0.59	3.03	0.20	8.40	183.40
	JAS	0.02	0.64	3.02	0.50	13.80	266.60
	OND	0.03	0.57	4.14	0.00	10.00	192.14
	1977-1987	0.02	0.88	3.03	0.10	13.40	266.60
	1987-1997	0.05	0.73	4.14	0.00	8.78	192.14
	1997-2007	0.00	0.54	2.89	0.12	7.86	261.90
	2007-2017	0.03	0.42	2.75	0.50	7.40	220.60
hi	JFM	0.03	0.43	1.65	0.00	3.20	55.80
	AMJ	0.02	0.48	1.95	0.10	5.40	74.90
	JAS	0.03	0.54	3.16	0.10	7.23	333.40
	OND	0.02	0.43	2.43	0.00	4.67	142.90
	1977-1987	0.02	0.57	1.92	0.30	7.30	136.80
	1987-1997	0.02	0.54	2.43	0.00	5.11	142.90
	1997-2007	0.02	0.56	3.16	0.00	4.80	72.30
	2007-2017	0.03	0.33	1.80	0.80	3.70	333.40

Table 2: Associations of restoration projects with chlorophyll-a changes for different segments of Tampa Bay (Figure 1). Associations were evaluated based on different year windows (5, 10) since completion of restoration projects and number of closest restoration projects (5, 10) to each monitoring station within each segment. Overall differences in chlorophyll-a changes between restoration project types by segment and year/project number combinations were evaluated by ANOVA F-tests, whereas pairwise differences (shown as letters) between project types were evaluated by t-tests with corrected p-values for multiple comparisons. Chlorophyll-a changes by project types that are not significantly different share a letter (comparisons are only valid within rows) and significance of the within-group mean relative to zero is also shown. HB: Hillsborough Bay, LTB: Lower Tampa Bay, MTB: Middle Tampa Bay, OTB: Old Tampa Bay.  $p > 0.05$  ns,  $p < 0.05$  \*,  $p < 0.005$  \*\*

Restoration project							
Combination	Bay segment	ANOVA	Habitat enhance	Habitat establish	Habitat protect	Nonpoint control	Point control
5 years, 5 projects	HB	F = 1.09, ns	a, 0	a, 0	a, < 0	a, 0	a, 0
	LTB	F = 17.37, **	a, 0	a, 0	a, 0	a, 0	b, < 0
	MTB	F = 13.5, **	b, 0	ab, 0	a, < 0	ab, < 0	c, < 0
	OTB	F = 1.5, ns	a, 0	a, 0	a, 0	a, 0	a, 0
5 years, 10 projects	HB	F = 0.66, ns	a, 0	a, 0	a, 0	a, 0	a, 0
	LTB	F = 2.64, ns	ab, < 0	ab, < 0	ab, < 0	a, 0	b, < 0
	MTB	F = 18.75, **	a, 0	a, 0	a, < 0	a, < 0	b, < 0
	OTB	F = 3.11, *	ab, 0	a, 0	ab, 0	b, < 0	a, 0
10 years, 5 projects	HB	F = 2.9, *	ab, 0	ab, 0	ab, < 0	a, 0	b, < 0
	LTB	F = 6.13, **	a, 0	a, 0	a, 0	ab, 0	b, < 0
	MTB	F = 14.11, **	a, 0	a, 0	a, 0	a, 0	b, < 0
	OTB	F = 3.15, *	b, 0	ab, 0	ab, 0	a, < 0	a, < 0

Restoration project

Combination	Bay segment	ANOVA	Habitat enhance	Habitat establish	Habitat protect	Nonpoint control	Point control
10 years, 10 projects	HB	F = 2.42, ns	a, 0	a, 0	a, 0	a, 0	a, < 0
	LTB	F = 1.78, ns	a, 0	a, < 0	a, < 0	a, 0	a, < 0
	MTB	F = 11.79, **	a, 0	a, < 0	a, 0	a, 0	b, < 0
	OTB	F = 2.35, ns	b, 0	ab, < 0	ab, 0	a, < 0	ab, < 0

710

711 *Table 3 Results comparing water quality associations for random date and location assignments*  
712 *of restoration projects to those from real data. The columns “mean < 0”, “mean = 0”, and*  
713 *“mean > 0” show the proportion of results for 1000 simulations of random dates and locations*  
714 *where the estimated effect had an overall decrease in observed chlorophyll-a (mean < 0), no*  
715 *change (mean = 0), or increase in chlorophyll-a (mean > 0). The actual estimate of the*  
716 *association with chlorophyll-a change from observed data for restoration projects is also shown.*  
717 *The agreement column shows whether the actual estimate is in agreement (y: yes, n: no) with the*  
718 *most frequently observed result from the random simulations (bold).*

Combination	Restoration project	mean < 0	mean = 0	mean > 0	Actual	Agreement
5 years, 5 projects	Habitat enhance	0.44	<b>0.51</b>	0.06	mean = 0	y
	Habitat establish	0.47	<b>0.49</b>	0.04	mean = 0	y
	Habitat protect	<b>0.49</b>	0.38	0.14	mean < 0	y
	Nonpoint control	0.45	<b>0.5</b>	0.05	mean = 0	y
	Point control	<b>0.49</b>	0.4	0.11	mean < 0	y
5 years, 10 projects	Habitat enhance	<b>0.59</b>	0.38	0.04	mean = 0	n
	Habitat establish	<b>0.61</b>	0.36	0.04	mean = 0	n
	Habitat protect	<b>0.58</b>	0.32	0.1	mean < 0	y
	Nonpoint control	<b>0.59</b>	0.37	0.04	mean < 0	y
	Point control	<b>0.57</b>	0.31	0.12	mean < 0	y
10 years, 5 projects	Habitat enhance	<b>0.82</b>	0.17	0	mean = 0	n
	Habitat establish	<b>0.82</b>	0.18	0	mean = 0	n
	Habitat protect	<b>0.79</b>	0.18	0.04	mean < 0	y
	Nonpoint control	<b>0.81</b>	0.18	0.01	mean = 0	n
	Point control	<b>0.75</b>	0.22	0.03	mean < 0	y
10 years, 10 projects	Habitat enhance	<b>0.92</b>	0.08	0	mean = 0	n
	Habitat establish	<b>0.94</b>	0.06	0	mean < 0	y
	Habitat protect	<b>0.83</b>	0.14	0.02	mean < 0	y
	Nonpoint control	<b>0.94</b>	0.06	0	mean < 0	y
	Point control	<b>0.87</b>	0.12	0.01	mean < 0	y

719

## References

- Alfredo, K. A., and T. A. Russo. 2017. Urban, agricultural, and environmental protection practices for sustainable water quality. *WIREs Water* 4: e1229. doi:[10.1002/wat2.1229](https://doi.org/10.1002/wat2.1229).
- Baird, R. C. 2005. On sustainability, estuaries, and ecosystem restoration: The art of the practical. *Restoration Ecology* 13: 154–158. doi:[10.1111/j.1526-100X.2005.00019.x](https://doi.org/10.1111/j.1526-100X.2005.00019.x).
- Bayraktarov, E., M. I. Saunders, S. Abdullah, M. Mills, J. Beher, H. P. Possingham, P. J. Mumby, and C. E. Lovelock. 2016. The cost and feasibility of marine coastal restoration. *Ecological Applications* 26: 1055–1074. doi:[10.1890/15-1077](https://doi.org/10.1890/15-1077).
- Beck, M. W., and J. D. Hagy III. 2015. Adaptation of a weighted regression approach to evaluate water quality trends in an estuary. *Environmental Modelling and Assessment* 20: 637–655. doi:[10.1007/s10666-015-9452-8](https://doi.org/10.1007/s10666-015-9452-8).
- Beck, M. W., J. D. Hagy III, and C. Le. 2018. Quantifying seagrass light requirements using an algorithm to spatially resolve depth of colonization. *Estuaries and Coasts* 41: 592–610. doi:[10.1007/s12237-017-0287-1](https://doi.org/10.1007/s12237-017-0287-1).
- Beyer, J., H. C. Trannum, T. Bakke, P. V. Hodson, and T. K. Collier. 2016. Environmental effects of the Deepwater Horizon oil spill: A review. *Marine Pollution Bulletin* 110: 28–51. doi:[10.1016/j.marpolbul.2016.06.027](https://doi.org/10.1016/j.marpolbul.2016.06.027).
- Borja, Á., D. M. Dauer, M. Elliott, and C. A. Simenstad. 2010. Median- and long-term recovery of estuarine and coastal ecosystems: Patterns, rates and restoration effectiveness. *Estuaries and Coasts* 33: 1249–1260. doi:[10.1007/s12237-010-9347-5](https://doi.org/10.1007/s12237-010-9347-5).



740 Borja, Á., G. Chust, J. G. Rodríguez, J. Bald, M. J. Belzunce-Segarra, J. Franco, J. M.  
 741 Garmendia, et al. 2016. ‘The past is the future of the present’: Learning from long-time series of  
 742 marine monitoring. *Science of the Total Environment* 566-567. Elsevier B.V.: 698–711.  
 743 doi:[10.1016/j.scitotenv.2016.05.111](https://doi.org/10.1016/j.scitotenv.2016.05.111).

744 Comp, G. S. 1985. A survey of the distribution and migration of the fishes in Tampa Bay. In  
 745 *Proceedings, tampa bay area scientific information symposium, may 1982*, ed. S. F. Treat, J. L.  
 746 Simon, R. R. Lewis III, and R. L. Whitman Jr., 393–425. Minneapolis, Minnesota: Burgess  
 747 Publishing Co., Inc. <https://tbep.tech.org/BASIS/BASIS1/BASIS1.pdf> (Accessed June, 2019)

748 Curriero, F. C. 2006. On the use of non-Euclidean distance measures in geostatistics.  
 749 *Mathematical Geology* 38: 907–926. doi:[10.1007/s11004-006-9055-7](https://doi.org/10.1007/s11004-006-9055-7).

750 Diefenderfer, H. L., G. E. Johnson, R. M. Thom, K. E. Buenau, L. A. Weitkamp, C. M.  
 751 Woodley, A. B. Borde, and R. K. Kropp. 2016. Evidence-based evaluation of the cumulative  
 752 effects of ecosystem restoration. *Ecosphere* 7: e01242. doi:[10.1002/ecs2.1242](https://doi.org/10.1002/ecs2.1242).

753 Diefenderfer, H. L., K. L. Sobocinski, R. M. Thom, C. W. May, A. B. Borde, S. L. Southard, J.  
 754 Vavrinec, and N. K. Sather. 2009. Multiscale analysis of restoration priorities for marine  
 755 shoreline planning. *Environmental Management* 44: 712–733. doi:[10.1007/s00267-009-9298-4](https://doi.org/10.1007/s00267-009-9298-4).

756 Diefenderfer, H. L., R. M. Thom, G. E. Johnson, J. R. Skalski, K. A. Vogt, B. D. Ebberts, G. C.  
 757 Roegner, and E. M. Dawley. 2011. A levels-of-evidence approach for assessing cumulative  
 758 ecosystem response to estuary and river restoration programs. *Ecological Restoration* 29: 111–  
 759 132. doi:[10.3368/er.29.1-2.111](https://doi.org/10.3368/er.29.1-2.111).

760 Enwright, N. M., K. T. Griffith, and M. J. Osland. 2016. Barriers to and opportunities for  
 761 landward migration of coastal wetlands with sea-level rise. *Frontiers in Ecology and the*  
 762 *Environment* 14: 307–316. doi:[10.1002/fee.1282](https://doi.org/10.1002/fee.1282).

763 Gardner, B., P. J. Sullivan, and A. J. Lembo. 2011. Predicting stream temperatures:  
 764 Geostatistical model comparison using alternative distance measures. *Canadian Journal of*  
 765 *Fisheries and Aquatic Sciences* 60: 344–351. doi:[10.1139/f03-025](https://doi.org/10.1139/f03-025).

766 GCERC (Gulf Coast Ecosystem Restoration Council). 2013. Comprehensive plan: Restoring the  
 767 Gulf Coast’s ecosystem and economy.  
 768 [https://www.restorethegulf.gov/sites/default/files/Initial%20Comprehensive%20Plan%20Aug%2](https://www.restorethegulf.gov/sites/default/files/Initial%20Comprehensive%20Plan%20Aug%202013.pdf)  
 769 [02013.pdf](https://www.restorethegulf.gov/sites/default/files/Initial%20Comprehensive%20Plan%20Aug%202013.pdf). (Accessed January, 2018)

770 GCERC (Gulf Coast Ecosystem Restoration Council). 2016. Comprehensive plan: Update 2016,  
 771 restoring the Gulf Coast’s ecosystem and economy.  
 772 [https://www.restorethegulf.gov/sites/default/files/COL\\_20161208\\_CompPlanUpdate\\_English.pd](https://www.restorethegulf.gov/sites/default/files/COL_20161208_CompPlanUpdate_English.pdf)  
 773 [f](https://www.restorethegulf.gov/sites/default/files/COL_20161208_CompPlanUpdate_English.pdf). (Accessed January, 2018)

774 GMFMC (Gulf of Mexico Fishery Management Council). 2017. State management program for  
 775 recreational red snapper summary: Draft amendment to the fishery management plan for the reef  
 776 fish resources of the Gulf of Mexico. [http://gulfcouncil.org/wp-content/uploads/B-7-a-](http://gulfcouncil.org/wp-content/uploads/B-7-a-Recreational-State-Management-for-Red-Snapper.pdf)  
 777 [Recreational-State-Management-for-Red-Snapper.pdf](http://gulfcouncil.org/wp-content/uploads/B-7-a-Recreational-State-Management-for-Red-Snapper.pdf). (Accessed January, 2018)

778 Greening, H. S., and A. Janicki. 2006. Toward reversal of eutrophic conditions in a subtropical  
 779 estuary: Water quality and seagrass response to nitrogen loading reductions in Tampa Bay,  
 780 Florida, USA. *Environmental Management* 38: 163–178.

781 Greening, H. S., P. Doering, and C. Corbett. 2006. Hurricane impacts on coastal ecosystems.  
 782 *Estuaries and Coasts* 29: 877–879. doi:[10.1007/BF02798646](https://doi.org/10.1007/BF02798646).

783 Greening, H. S., A. Janicki, E. T. Sherwood, R. Pribble, and J. O. R. Johansson. 2014.  
 784 Ecosystem responses to long-term nutrient management in an urban estuary: Tampa Bay,  
 785 Florida, USA. *Estuarine, Coastal and Shelf Science* 151: A1–A16.  
 786 doi:[10.1016/j.ecss.2014.10.003](https://doi.org/10.1016/j.ecss.2014.10.003).

787 Gross, C., and J. D. Hagy III. 2017. Attributes of successful actions to restore lakes and estuaries  
 788 degraded by nutrient pollution. *Journal of Environmental Management* 187: 122–136.  
 789 doi:[10.1016/j.jenvman.2016.11.018](https://doi.org/10.1016/j.jenvman.2016.11.018).

790 Hassett, B., M. Palmer, E. Bernhardt, S. Smith, J. Carr, and D. Hart. 2005. Restoring watersheds  
 791 project by project: Trends in Chesapeake Bay tributary restoration. *Frontiers in Ecology and the*  
 792 *Environment* 3: 259–267. doi:[10.1890/1540-9295\(2005\)003\[0259:RWPBPT\]2.0.CO;2](https://doi.org/10.1890/1540-9295(2005)003[0259:RWPBPT]2.0.CO;2).

793 Hobbs, R J, and J A Harris. 2001. Restoration Ecology: Repairing the Earth’s Ecosystems in the  
 794 New Millennium. *Restoration Ecology* 9: 239–246. doi:[10.1046/j.1526-100x.2001.009002239.x](https://doi.org/10.1046/j.1526-100x.2001.009002239.x).

795 Janicki, A., D. Wade, and J. R. Pribble. 1999. *Development of a process to track the status of*  
 796 *chlorophyll and light attenuation to support seagrass restoration goals in Tampa Bay*. 04-00. St.  
 797 Petersburg, Florida: Tampa Bay National Estuary Program.  
 798 [https://tbep.tech.org/TBEP\\_TECH\\_PUBS/2000/TBEP\\_04\\_00Chlor-A.pdf](https://tbep.tech.org/TBEP_TECH_PUBS/2000/TBEP_04_00Chlor-A.pdf) (Accessed June, 2019)

799 Johansson, J. O. R. 1991. Long-term trends of nitrogen loading, water quality and biological  
 800 indicators in Hillsborough Bay, Florida. Edited by S. F. Treat and P. A. Clark. Tampa, Florida,

801 USA: 2<sup>nd</sup> Tampa Bay BASIS Proceedings: 157–176.

802 <https://www.tbeptech.org/BASIS/BASIS2/BASIS2.pdf>. (Accessed June 2019)

803 Johansson, J. O. R., and R. R. Lewis III. 1992. Recent improvements in water quality and

804 biological indicators in Hillsborough Bay, a highly impacted subdivision of Tampa Bay, Florida,

805 USA. *Marine Coastal Eutrophication* Proceedings of an International Conference, Bologna,

806 Italy, 21-24 March 1990: 1199–1215.

807 Lewis, R. R., P. A. Clark, W. K. Fehring, H. S. Greening, R. O. Johansson, and R. T. Paul. 1998.

808 The rehabilitation of Tampa Bay Estuary, Florida, USA, as an example of successful integrated

809 coastal management. *Marine Pollution Bulletin* 37: 468–473. doi:[10.1016/S0025-](https://doi.org/10.1016/S0025-326X(99)00139-3)

810 [326X\(99\)00139-3](https://doi.org/10.1016/S0025-326X(99)00139-3).

811 Liang, D., L. A. Harris, J. M. Testa, V. Lyubchich, and S. Filoso. 2019. Detection of the effects

812 of stormwater control measures in streams using a Bayesian BACI power analysis. *Science of the*

813 *Total Environment* 661: 386–392. doi:[10.1016/j.scitotenv.2019.01.125](https://doi.org/10.1016/j.scitotenv.2019.01.125).

814 Lombardo, R., and R. R. Lewis III. 1985. A review of commercial fisheries data: Tampa Bay,

815 Florida. In *Proceedings, tampa bay area scientific information symposium, may 1982*, ed. S. F.

816 Treat, J. L. Simon, R. R. Lewis III, and R. L. Whitman Jr., 614–634. Minneapolis, Minnesota:

817 Burgess Publishing Co., Inc. <https://tbeptech.org/BASIS/BASIS1/BASIS1.pdf>. (Accessed June,

818 2019)

819 Meals, D. W., S. A. Dressing, and T. E. Davenport. 2010. Lag time in water quality response to

820 best management practices: A review. *Journal of Environmental Quality* 39: 85–96.

821 doi:[10.2134/jeq2009.0108](https://doi.org/10.2134/jeq2009.0108).

822 Morrison, G., E. T. Sherwood, R. Boler, and J. Barron. 2006. Variations in water clarity and  
 823 chlorophylla in Tampa Bay, Florida, in response to annual rainfall, 1985-2004. *Estuaries and*  
 824 *Coasts* 29: 926–931.

825 NASEM (National Academies of Sciences, Engineering, and Medicine). 2017. *Effective*  
 826 *monitoring to evaluate ecological restoration in the gulf of mexico*. Washington, DC: The  
 827 National Academies Press. doi:[10.17226/23476](https://doi.org/10.17226/23476).

828 Neckles, H. A., M. Dionne, D. M. Burdick, C. T. Roman, R. Buchsbaum, and E. Hutchins. 2002.  
 829 A monitoring protocol to assess tidal restoration of salt marshes on local and regional scales.  
 830 *Restoration Ecology* 10: 556–563. doi:[10.1046/j.1526-100X.2002.02033.x](https://doi.org/10.1046/j.1526-100X.2002.02033.x).

831 NRDA (Deepwater Horizon Natural Resource Damage Assessment Trustees). 2016. Deepwater  
 832 Horizon oil spill: Final programmatic damage assessment and restoration plan and final  
 833 programmatic environmental impact statement.  
 834 <http://www.gulfspillrestoration.noaa.gov/restoration-planning/gulf-plan>. (Accessed June, 2018)

835 Poe, A., K. Hackett, S. Janicki, R. Pribble, and A. Janicki. 2005. *Estimates of total nitrogen, total*  
 836 *phosphorus, total suspended solids, and biochemical oxygen demand loadings to Tampa Bay,*  
 837 *Florida: 1999-2003*. #02-05. St. Petersburg, Florida, USA: Tampa Bay Estuary Program.  
 838 [https://tbep.tech.org/TBEP\\_TECH\\_PUBS/2005/TBEP\\_02\\_05Loadings99\\_03.pdf](https://tbep.tech.org/TBEP_TECH_PUBS/2005/TBEP_02_05Loadings99_03.pdf). (Accessed  
 839 June, 2019)

840 Rabotyagov, S. S., C. L. Kling, P. W. Gassman, N. N. Rabalais, and R. E. Turner. 2014. The  
 841 Economics of Dead Zones: Causes, Impacts, Policy Challenges, and a Model of the Gulf of  
 842 Mexico Hypoxic Zone. *Review of Environmental Economics and Policy* 8: 58–79.  
 843 doi:[10.1093/reep/ret024](https://doi.org/10.1093/reep/ret024).

844 RDCT (R Development Core Team). 2018. R: A language and environment for statistical  
 845 computing, v3.5.1. R Foundation for Statistical Computing, Vienna, Austria. [http://www.R-](http://www.R-project.org)  
 846 [project.org](http://www.R-project.org)

847 Ruiz-Jaen, M. C., and T. M. Aide. 2005. Restoration success: How is it being measured?  
 848 *Restoration Ecology* 13: 569–577. doi:[10.1111/j.1526-100X.2005.00072.x](https://doi.org/10.1111/j.1526-100X.2005.00072.x).

849 Scheffer, M., S. Carpenter, J. A. Foley, C. Folkes, and B. Walker. 1998. Catastrophic shifts in  
 850 ecosystems. *Nature* 413: 591–596.

851 Schiff, K., P. R. Trowbridge, E. T. Sherwood, P. Tango, and R. A. Batiuk. 2016. Regional  
 852 monitoring programs in the United States: Synthesis of four case studies from Pacific, Atlantic,  
 853 and Gulf Coasts. *Regional Studies in Marine Science* 4: A1–A7. doi: [10.1016/j.rsma.2015.11.007](https://doi.org/10.1016/j.rsma.2015.11.007)

854 Sherwood, E. T., H. S. Greening, A. J. Janicki, and D. J. Karlen. 2016. Tampa Bay estuary:  
 855 Monitoring long-term recovery through regional partnerships. *Regional Studies in Marine*  
 856 *Science* 4: 1–11. doi:[10.1016/j.rsma.2015.05.005](https://doi.org/10.1016/j.rsma.2015.05.005).

857 Sherwood, E. T., H. S. Greening, J. O. R. Johansson, K. Kaufman, and G. Raulerson. 2017.  
 858 Tampa Bay (Florida, USA): Documenting seagrass recovery since the 1980s and reviewing the  
 859 benefits. *Southeastern Geographer* 57: 294–319. doi: [10.1353/sgo.2017.0026](https://doi.org/10.1353/sgo.2017.0026).

860 Sickle, J. Van, and C. B. Johnson. 2008. Parametric distance weighting of landscape influence on  
 861 streams. *Landscape Ecology* 23: 427–438. doi:[10.1007/s10980-008-9200-4](https://doi.org/10.1007/s10980-008-9200-4).

862 SWFWMD (Southwest Florida Water Management District). 2018. Southwest Florida Water  
 863 Management District Geospatial Open Data Portal. <https://data-swfwmd.opendata.arcgis.com/>.  
 864 (Accessed January, 2018)


865 TBEP (Tampa Bay Estuary Program). 2017. Tampa Bay Water Atlas.  
 866 <http://www.tampabay.wateratlas.usf.edu/>. (Accessed July, 2017)

867 Thom, R. M., G. D. Williams, N. R. Evans, and E. Haas. 2011. Lower Columbia River and  
 868 estuary habitat restoration prioritization framework. *Ecological Restoration* 29: 94–110.  
 869 doi:[10.3368/er.29.1-2.94](https://doi.org/10.3368/er.29.1-2.94).


870 Wang, P. F., J. Martin, and G. Morrison. 1999. Water quality and eutrophication in Tampa Bay,  
 871 Florida. *Estuarine, Coastal and Shelf Science* 49: 1–20.

872 Wortley, L., J. M. Hero, and M. Howes. 2013. Evaluating ecological restoration success: A  
 873 review of the literature. *Restoration Ecology* 21: 537–543. doi:[10.1111/rec.12028](https://doi.org/10.1111/rec.12028).

874 Zedler, J. B. 2007. Success: An unclear, subjective descriptor of restoration outcomes.  
 875 *Ecological Restoration* 25: 162–168. doi:[10.3368/er.25.3.162](https://doi.org/10.3368/er.25.3.162).









Water quality stations ● Restoration projects ● Habitat ● Water Infrastructure


Closest match by project type




Closest "n" matches



(a) Spatial match



(b) Temporal match








Figure6

[Click here to access/download;Figure;Fig6.png](#)




Figure7

