Quantitative approaches to understand nutrient pollution in estuaries: An example for the upper San Francisco Estuary

Marcus W. Beck, PhD

USEPA National Health and Environmental Effects Research Laboratory, Gulf Ecology Division, beck.marcus@epa.gov, Phone: 8509342480

Aug. 26, 2016

How do we collect and use data?

The foundation of environmental management is a strong monitoring network [National Research Council, 1990]

Monitoring provides information for decision-making based on apparent trends...

What are the changes in environmental condition over time?

Are these changes 'good' or 'bad' based on our management objectives?

What may have caused these changes?

How do we collect and use data?


The good news: We are getting better at monitoring - standardized, automated, increased coverage, real-time/continuous

The bad news: Our ability to use these data for decision-making has not kept pace with availability!

How do we collect and use data?

The good news: We are getting better at monitoring - standardized, automated, increased coverage, real-time/continuous

The bad news: Our ability to use these data for decision-making has not kept pace with availability!



How do we collect and use data?

Most of my research career has focused on using monitoring data to understand effects of eutrophication in one form or another

Eutrophication (noun) - an **increase** in the rate of supply of **organic matter** to an ecosystem

- [Nixon, 1995]

How do we collect and use data?

Most of my research career has focused on using monitoring data to understand effects of eutrophication in one form or another

Eutrophication (noun) - an increase in the rate of supply of organic matter to an ecosystem

- [Nixon, 1995]

Nutrient Loading


Adapted from [Cloern, 2001]

How do we collect and use data?

Most of my research career has focused on using monitoring data to understand effects of eutrophication in one form or another

Eutrophication (noun) - an **increase** in the rate of supply of **organic matter** to an ecosystem

- [Nixon, 1995]




Adapted from [Cloern, 2001]

How do we collect and use data?

Most of my research career has focused on using monitoring data to understand effects of eutrophication in one form or another

Eutrophication (noun) - an **increase** in the rate of supply of **organic matter** to an ecosystem

- [Nixon, 1995]



Adapted from [Cloern, 2001]

How do we collect and use data?

Today's talk: My experience evaluating monitoring data to inform our understanding of the eutrophication paradigm

Water quality trends in the Delta:

- Example 1: Model theory and application
- Example 2: Trends over time
- Example 3: Selected case studies


WRTDS adaptation for tidal waters

Increasing availability of records describing long-term changes

Observed data can provide a means to an end, potentially \boldsymbol{high} \boldsymbol{power} with large sample size


Can we develop and apply tools that leverage the descriptive capabilities of these large datasets?

Can we *link descriptions* to *causal events* to inform management or understanding?



WRTDS adaptation for tidal waters

Observed data represents effects of many processes



$\underline{Climate}$

precipitation temperature wind events ENSO effects

Local


light/turbidity residence time invasive species trophic effects

Regional/historical

watershed inputs point sources management actions flow changes

WRTDS adaptation for tidal waters


Observed data represents effects of many processes




Models should describe components to evaluate effects

WRTDS adaptation for tidal waters

Observed data represents effects of many processes



Models should describe components to evaluate effects



1990


2000

Time


2010

WRTDS adaptation for tidal waters


Observed data represents effects of many processes




Models should describe components to evaluate effects



Annual





WRTDS adaptation for tidal waters

Problem: Response endpoints of eutrophication vary naturally over time and with discharge or tidal patterns

Solution: Develop a model that accounts for changes in relationships between drivers of pollution over time

The weighted regression (WRTDS) model is being developed by USGS for pollutant modelling in rivers [Hirsch et al., 2010]

Models pollution concentration as a function of $\it time$, $\it discharge$, and $\it season$

Adaptation: Applied to Tampa Bay [Beck and Hagy III, 2015], further validated/compared in Patuxent Estuary [Beck and Murphy, In review]


WRTDS adaptation for tidal waters

How does weighted regression work?

WRTDS adaptation for tidal waters

Application to Delta:

- Nine stations (three Suisun, three middle, three delta)
- Three analytes (DIN, ammonium, nitrite/nitrate), two flow records
- Four decades of data



Acknowledgments:

Research staff and employees at USEPA Gulf Ecology Division

Funding sources and contact:



beck.marcus@epa.gov

Phone: 8509342480

References

Beck MW, Hagy III JD. 2015.

Adaptation of a weighted regression approach to evaluate water quality trends in an estuary.

Beck MW, Murphy RR. In review.

Numerical and qualitative contrasts of two statistical models for water quality change in tidal waters. Journal of the American Water Resources Assocation.

Cloern JE. 2001.

Our evolving conceptual model of the coastal eutrophication problem.

farine Ecology Progress Series, 210:223–253

Hirsch RM, Moyer DL, Archfield SA. 2010.

Weighted regressions on time, discharge, and season (WRTDS), with an application to Chesapeake Bay river inputs.

Journal of the American Water Resources Association, 46(5):857–880

National Research Council. 1990.

Managing Troubled Waters: The Role of Environmental Monitoring.

National Academy Press, Washington, DC

Nixon SW, 1995.

Coastal marine eutrophication: A definition, social causes, and future concerns.

Ophelia, 41:199-219.