

Four decades of water quality change in the upper San Francisco Estuary

Marcus W. Beck¹, David Senn², Phil Bresnahan², Emily Novick², James D. Hagy III¹, Thomas Jabusch²

¹*USEPA National Health and Environmental Effects Research Laboratory
Gulf Ecology Division, 1 Sabine Island Drive, Gulf Breeze, FL 32561
Phone: 850-934-2480, Fax: 850-934-2401
Emails: beck.marcus@epa.gov, hagy.jim@epa.gov*

²*San Francisco Estuary Institute
4911 Central Avenue, Richmond, CA 94804
Phone: 510-746-7334, Fax: 510-746-7300
Emails: davids@sfei.org, philb@sfei.org, emilyn@sfei.org, thomasj@sfei.org*

Version Date: Fri Sep 30 16:40:38 2016 -0500

Abstract

Recent methods for trend analysis have been developed that leverage the descriptive potential of long-term time series. Combined with these methods, multi-decadal datasets of water quality in coastal systems can provide valuable opportunities to gain insights into ecosystem properties and drivers of change. This study describes use of an estuarine adaptation of the Weighted Regressions on Time, Discharge, and Season (WRTDS) model to describe water quality trends over four decades in the Delta region of the San Francisco Estuary (SFE). This region is a complex mosaic of inflows that are primary sources of nutrients into the larger Bay. To date, a comprehensive evaluation of the long-term monitoring dataset at multiple stations in the Delta has not been conducted despite the importance of nutrient transport from the region for water quality in the entire bay. The WRTDS technique is data-driven where the parameterization of the functional model changes smoothly over time following dynamic patterns of season and flow. Water quality trends that have not been previously quantified can be described, including variation in flow-normalized concentrations, frequency occurrence of extreme events, and response to historical changes in the watershed, all of which are important needs for understanding changes in the SFE. Model results from multiple stations in the Delta provided novel descriptions of historical trends and relationships between key species of dissolved inorganic nitrogen (ammonium, nitrate/nitrite, total). This variation was described in the context of varying contributions of input flows from the Sacramento and San Joaquin rivers, as well as tidal exchange with the central SFE. Conceptual relationships between water quality and drivers of change were used to generate and test hypotheses of mechanistic relationships using selected examples from the trend descriptions. Overall, this analysis provides an ecological and management-based understanding of historical trends in the SFE as a means to interpret potential impacts of recent changes and expected trends in this dynamic system. An argument is also made for more comprehensive evaluations of long-term monitoring datasets to understand relationships between response endpoints and causal mechanisms in coastal waters.

1 Introduction

1. How and why are trends interpreted - assessment of raw data, surrogates, various methods (kendall, GAM, WRTDS), what have been implications of using different approaches, see Kratzer USGS report <http://pubs.usgs.gov/sir/2010/5228/pdf/sir20105228.pdf> and data

http://pubs.usgs.gov/sir/2010/5228/, need to interpret eutrophication trends in estuaries - it's confusing (Cloern and Jassby 2010)

2. WRTDS, original method (Hirsch et al. 2010, 2015)

3. WRTDS application to Tampa Bay as test set (Beck and Hagy III 2015), further validation in Patuxent and other tidal waters Beck et al. (2015)

4. SF estuary, unique and prominent location, full story is complex (historical context and recent changes) (Cloern and Jassby 2012), why is the delta important (a vigorous biogeochemical reactor) (Jassby and Cloern 2000, Jassby et al. 2002, Jassby 2008), no one has empirically described the data in the delta using data-intensive methods

San Francisco Bay on the Pacific Coast of the United States is one of the most prominent estuaries in the western hemisphere. Background nutrient concentrations in the Bay often exceed those associated with excessive primary production, although eutrophication events have historically been infrequent. Recent changes in response to additional stressors (e.g., variation in freshwater inputs/withdrawals, invasive species, climate change) suggests that Bay condition has not followed historical trajectories and more subtle spatial and temporal variation could provide clues that describe underlying properties of this system. The unique ecological and social context of the Bay, including a rich source of monitoring data from the last four decades, provides a valuable opportunity to gain insight into ecosystem properties of estuaries.

5. Study goal and objectives

- Provide a description of trends - annual, seasonal, spatial, response to flow, change by analytes
- Detailed description of selected sites in the context of conceptual relationships - 1) nonlinear or extreme quantile changes, site TBD, 2) P8 and WWTP improvements, 3) Suisun DIN, SiO₂, Chl_a, and clams
- What this means for understanding other systems

2 *Methods*

2.1 Study location and data

The San Francisco Estuary (SFE)... exchange with ocean, freshwater inputs (rivers and watershed drainage), runoff variation (drought years), climate, support of natural resources, subembayments, POTW (publicly owned treatment works), nutrient loading relative to other locations

2.2 Analysis method and application

$$\ln(N) = \beta_0 + \beta_1 t + \beta_2 \ln(Q) + \beta_3 \sin(2\pi t) + \beta_4 \cos(2\pi t) \quad (1)$$

nine discrete sampling stations with data from 1976 to 2012, dissolved inorganic nitrogen (DIN), ammonium, total nitrate, effects of different flow variables

Annual, seasonal trends of flow-normalized predictions

2.3 Case studies

These are science questions that are relevant outside of the region.

2.3.1 *Disaggregating observed nitrogen time series*

Hypothesis: Because multiple factors influence nutrient concentrations at different times, relationships between nutrients, time, and flow/salinity are non-linear and complex, so we expect 1) annual trend independent of seasonal trend, 2) changes in seasonal amplitudes and quantile trends over time, 3) varying flow contribution, either as difference between predicted/flow-normalized results or changes in nutrient v flow scatterplots at different annual periods.

2.3.2 *Effects of wastewater treatment*

Hypothesis: Modal response of nutrient concentrations at P8 over time is result of WWTP upgrades, so we expect 1) a shift in load contributions before/after upgrade, 2) a flow-normalized annual trend at P8 to show a change concurrent with WWTP upgrades, and 3) different nitrogen species will have different changes depending on change in load outputs. See [here](#)

2.3.3 *Effects of biological invasions*

Hypothesis: Biological invasions by benthic filter feeders have shifted abundance and composition of phytoplankton communities in Suisun Bay, so we expect 1) decline in annual, flow-normalized chlorophyll concentrations over time coincident with increase in abundance of invaders, 2) changes in ratios of limiting nutrients (nitrogen, SiO₂) suggesting different uptake rates with shift in community composition, and 3) seasonal shifts in limiting nutrients based on changes in community composition and relative abundances with seasonal succession.

3 *Results*

3.1 Trends

3.2 Selected examples

3.2.1 *Disaggregating observed nitrogen time series*

Fig. 3, Fig. 4

Emphasize the information the model provides relative to the observed time series. A distinct annual trend with a maximum in the middle of the time series is observed, with lower values at the beginning and end of the period. The seasonal patterns generally showed that DIN concentrations were highest in January with higher values at moderate to low flow rates depending on the year. Interestingly, summer and fall concentrations have showed a slight increase later in the time series (2004-2009). The confounding effect of flow is also very apparent such that higher flows were associated with lower concentration. Dynaplot showed that there was always a negative association between the two (i.e., no modal response). The quantile distributions showed similar trends over time in both predicted values and flow-normalized predictions, although some exceptions were observed. In particular, high flow (1984, 2008) reduced concentrations of all quantiles but the magnitude of the effect increased at higher quantiles (i.e., the effect was disproportionate). The opposite was observed for low flow, i.e., the ninetieth percent showed the greatest increase for low flow.

Emphasize the summer/fall change in the 2000s, why is this? Check (Cloern et al. 2007), showed seasonal changes in early 2000s in chlorophyll (NE Pacific shifted to cool phase), is there a mechanism here with DIN? Relate to conceptual diagram.

3.2.2 *Effects of wastewater treatment*

Overall reduction in total nitrogen load was observed as a result of reduction in ammonium (Fig. 5). Nitrate is the primary constituent of total nitrogen after 2007. Organic nitrogen is a larger percentage of the total after nitrification. What was reduction in ammonium starting in 2002?

Nitrogen trends at P8 shifted in response to upstream wastewater treatment plant (WWTP) upgrades (Fig. 6), with ammonium showing the largest reduction. Interestingly, nitrite/nitrate concentrations also showed a similar but less dramatic decrease. Percent changes are shown in Table 2, where both nitrogen species shows large percent increases prior to WWTP upgrades followed by decreases after upgrades with ammonium showing the largest percentage. Seasonally, increases prior to upgrades were most apparent in the July-August-September (JAS) months for both analytes. Seasonal reductions post-upgrades were also largest in JAS for nitrite/nitrate, whereas percent reductions were similar across all monthly groupings for ammonium.

Relationships of nitrogen with flow showed the typical inverse flow/concentration dynamic with flushing at high flow, although patterns differed by nitrogen species. Seasonal variation was more apparent for ammonium, although both typically had the highest concentrations in the winter. Additionally, strength of the flow/nutrient relationship changed throughout the time series the year where the strongest relationship differed by analyte. Nitrite/nitrate typically had the strongest relationship flow later in the time series, whereas ammonium had the strongest relationship with flow in the early 2000s.

3.2.3 *Effects of biological invasions*

Data from (Crauder et al. 2016), Jassby (2008) describes phytoplankton community changes in the upper estuary, including chlorophyll response to flow. Figure 10 in Jassby (2008) showed that chlorophyll generally decreased with flow in 1980 but increased with flow in 2000.

Note the decrease in *Potamocorbula* abundance in 2011, 2012. These are wet years where abundance/biomass of the clams is driven down by lower salinity. Contrasted with the annual chlorophyll trends in the same years, the predicted values are above the flow-normalized trend suggesting an increase in chlorophyll with higher flow. The potential mechanism is therefore a decrease in clam abundance with high flow that releases phytoplankton from filtration pressure. This also explains the positive association of chlorophyll with flow in recent years (bottom right

dynaplot).

Further, chlorophyll trends early in the time series generally show a decrease with high flow with a distinct maximum at moderate flow. This may suggest stratification events at moderate flow contributed to phytoplankton blooms early in the time series. Water withdrawals later in the time series could have also altered environmental conditions to reduce the frequency occurrence of stratification events. Look into this more...

What about biomass/density relationships for *Potamocorbula*? Although clam density increases throughout the period, What about initial decrease in chlorophyll prior to clam invasion? Is this related to water withdrawals (i.e., decrease in stratification events at moderate flow)?

Fig. 7, Fig. 8, Table 3, Table 4

4 Discussion

Trends as percent change depend on the mean value, lower values will have larger percent changes.

Second case study showed typical inverse relationships between nutrients and flow, more flow means greater flushing and dilution of nutrient concentrations. Conversely, low flow means less flushing and higher nutrient concentrations, although this may not always be observed if the available nutrients are biologically available. Low-flow events during warmer months show the lowest ammonium concentrations, which corresponds to seasonal maxima in chlorophyll concentration. A similar but weaker relationship was observed with nitrite/nitrate where increased flow was related to decreased concentration and lower concentrations overall were observed in the summer. However, low-flow events still had higher concentrations than high-flow events in July, as compared to ammonium which was low regardless of flow. This suggests that ammonium concentrations are driving phytoplankton production at P8. Annual trends in chlorophyll concentration (not shown) showed an overall decrease from the 1970s to present, although a slight peak is observed in the 2000s. This peak is likely related to the maximum ammonium concentration shown in Fig. 6. Moreover, flow/chlorophyll relationships have generally been constant throughout the period of record such that a change in flow has not been related to a change in phytoplankton production. This suggests that nutrient loads that contribute to production at P8 are primarily from point sources at WWTP outflows as a change in flow does not

172 affect the load output. But what are watershed loads?

173 What do nitrogen trends mean? Have to interpret relative to trends in other variables. A
174 decrease in nitrogen or constant nitrogen does not mean nitrogen inputs have stayed the same,
175 they might actually be increasing if nitrogen. A change in chlorophyll relative to change in
176 nitrogen could be informative, and even more so, a change in silica relative to change in
177 chlorophyll suggests diatom biomass has changed. However, there are mismatches in these trends
178 that suggest other processes are at play, e.g., residence times and flow inputs, etc. Trends in
179 Suisun relative to trends in Delta provide an example, e.g., Suisun is decrease in chlorophyll,
180 increase in silica, increase in nitrogen, delta is decrease in silica, increase/decrease in DIN
181 (depending on time period/season), decrease in chlorophyll, what's going on? See Senn slide 14
182 (from burial?). The WRTDS model lets us at least address trends in the context of season, time,
183 and flow. This allows for more improved interpretation relative to observing raw data. Also explain
184 more information by looking at ammonium, nitrate/nitrite, relative to DIN. What about other
185 variables (light level as suspended particulate matter, temperature)?

References

- Beck MW, Hagy III JD. 2015. Adaptation of a weighted regression approach to evaluate water quality trends in an estuary. *Environmental Modelling and Assessment*, 20(6):637–655.
- Beck MW, Hagy III JD, Murrell MC. 2015. Improving estimates of ecosystem metabolism by reducing effects of tidal advection on dissolved oxygen time series. *Limnology and Oceanography: Methods*, 13(12):731–745.
- Cloern JE, Jassby AD. 2010. Patterns and scales of phytoplankton variability in estuarine-coastal ecosystems. *Estuaries and Coasts*, 33(2):230–241.
- Cloern JE, Jassby AD. 2012. Drivers of change in estuarine-coastal ecosystems: Discoveries from four decades of study in San Francisco Bay. *Reviews of Geophysics*, 50(4):1–33.
- Cloern JE, Jassby AD, Thompson JK, Hieb KA. 2007. A cold phase of the East Pacific triggers new phytoplankton blooms in San Francisco Bay. *Proceedings of the National Academy of Sciences of the United States of America*, 104(47):18561–18565.
- Crauder JS, Thompson JK, Parchaso F, Anduaga RI, Pearson SA, Gehrts K, Fuller H, Wells E. 2016. Bivalve effects on the food web supporting delta smelt - a long-term study of bivalve recruitment, biomass, and grazing rate patterns with varying freshwater outflow. Technical Report Open-File Report 2016-1005, US Geological Survey, Reston, Virginia.
- Hirsch RM, Archfield SA, De Cicco LA. 2015. A bootstrap method for estimating uncertainty of water quality trends. *Environmental Modelling and Software*, 73:148–166.
- Hirsch RM, Moyer DL, Archfield SA. 2010. Weighted regressions on time, discharge, and season (WRTDS), with an application to Chesapeake Bay river inputs. *Journal of the American Water Resources Association*, 46(5):857–880.
- Jassby AD. 2008. Phytoplankton in the Upper San Francisco Estuary: Recent biomass trends, their causes, and their trophic significance. *San Francisco Estuary and Watershed Science*, 6(1):1–24.
- Jassby AD, Cloern JE. 2000. Organic matter sources and rehabilitations of the Sacramento-San Joaquin Delta (California, USA).
- Jassby AD, Cloern JE, Cole BE. 2002. Annual primary production: Patterns and mechanisms of change in a nutrient-rich tidal ecosystem. *Limnology and Oceanography*, 47(3):698–712.




Fig. 1: Conceptual diagram illustrating use of WRTDS to decompose trends in observed nitrogen time series and potential forcing factors that can explain model output. Results from the model are described as annual and seasonal trends, changes in flow-nutrient dynamics for different time periods, and residual variation independent of time, flow, and season. Relationships between environmental factors (climate, local, regional/historical) and nitrogen trends are more easily related to the separate components of the observed time series using results from the model.




Fig. 2: Percent changes in nitrogen analytes for (a) annual and (b) seasonal (monthly) periods in the record. Changes are based on the difference between the ending and starting estimates for the flow-normalized estimates within each period. Points are colored for direction (red increasing, green decreasing) and sized for relative magnitude. Station names are shown in the top left panel.




Fig. 3: Time series of DIN and flow at station C10. Subfigure (a) shows the observed DIN time series and subfigure (b) shows the annual (water year starting in October) predictions from WRTDS at different conditional quantiles ($\tau = 0.1, 0.5, 0.9$). The points in subfigure (b) are predictions of observed DIN and the lines are flow-normalized predictions. Subfigure (c) shows the difference between the model predictions and flow-normalized predictions at the fiftieth conditional quantile. Subfigure (d) shows the flow time series of the San Joaquin River with a locally-estimated (loess) smooth to emphasize the long-term trend.




Fig. 4: Modelled relationships between DIN, flow, and time at station C10. The top figure shows the annual and seasonal variation over the entire time series and the bottom figure shows annual variation for selected months to remove seasonal variation. Warmer colors indicate higher DIN concentrations. The y-axis on the bottom figure is truncated by the fifth and ninety-fifth percentiles of flow within each month. Model results are for the fiftieth conditional quantile of DIN.




Fig. 5: Nitrogen load measurements (kg d⁻¹) at the City of Tracy Wastewater Treatment Plant, San Joaquin County. Wastewater discharge requirements were implemented in May, 2007 to include nitrification/denitrification and tertiary filtration causing a reduction in total nitrogen effluent discharged to the Delta. Reductions were primarily observed for ammonium.




Fig. 6: Nitrogen trends at P8 as observed (a, top), predicted and flow-normalized estimates from WRTDS (a, bottom), and relationships with flow over time (b). Nitrite/nitrate trends are on the left and ammonium trends are on the right. Wastewater treatment plant upgrades at the City of Tracy (San Joaquin County), were completed in May 2007 (Fig. 5), coincident with a dramatic decrease in ammonium at P8.




Fig. 7: Trends in clam abundance and chl-*a* concentration from 1976 to 2014 at station D7 in Suisun Bay. Invasion by *Potamocorbula amurensis* clams in the late 1980s and displacement of *Corbicula fluminea* was shown by changes in clam density (a, annual means). A coincident decrease in chl-*a* concentration was also observed (c). A weak but significant ($p < 0.001$) relationship between clam biomass and chl-*a* concentration is shown in subfigure (b). Flow relationships with chl-*a* concentration have also changed over time (d, observations from June). Chlorophyll shows a slight positive then dominantly negative association with increasing flow (decreasing salinity) early in the time series, whereas the trend is reversed in recent years.




Fig. 8: Flow-normalized trends of annual (left) and seasonal (right) variation in DIN, chl-*a*, and SiO₂ at station D7 (top) and C10 (bottom). Covariation between nutrients, chl-*a*, and SiO₂ is observed at D7 but not C10, although an overall decrease in SiO₂ at C10 is shown. Seasonal changes at D7 are most pronounced during the summer.

Table 1: Summaries of flow-normalized trends in nitrogen analytes for all stations and different time periods. Summaries are averages (mg L⁻¹) and percent changes in parentheses (increasing in bold-italic). Changes are based on the difference between the ending and starting estimates for the estimates within each period. See Fig. 2 for a summary of spatial trends.

Analyte/Station	Annual			Seasonal		
	1976-1988	1989-2000	2001-2012	JFMA	MJJA	SOND
DIN						
C10	1.2 (38.4)	1.3 (16.2)	1.2 (-39.3)	1.1 (-25.7)	1.1 (-10.1)	1.5 (1.7)
C3	0.3 (33.6)	0.4 (27.4)	0.4 (-12.7)	0.4 (30.7)	0.3 (54.8)	0.5 (42.8)
D19	0.4 (-20.2)	0.4 (6.5)	0.4 (-22.6)	0.6 (-28.6)	0.3 (-35.1)	0.4 (-15.6)
D26	0.4 (-2.4)	0.4 (10.2)	0.4 (-15.4)	0.5 (-10.9)	0.3 (-3.4)	0.5 (-2.6)
D28	0.4 (-20.5)	0.4 (2.7)	0.4 (-32.6)	0.6 (-32.9)	0.2 (-53.3)	0.4 (-48.8)
D4	0.3 (-9.1)	0.4 (20.7)	0.4 (-4.3)	0.5 (11.4)	0.3 (4.8)	0.4 (10.7)
D6	0.4 (-9.6)	0.4 (21.9)	0.5 (-7.1)	0.5 (-4.7)	0.4 (45.6)	0.4 (34.7)
D7	0.4 (-26.4)	0.4 (20.4)	0.5 (-2.4)	0.5 (-6.9)	0.4 (19)	0.4 (21.4)
P8	1.1 (74.1)	1.5 (37.9)	1.6 (-31)	1.4 (31.6)	1 (52.8)	1.8 (45.4)
NH₄⁺						
C10	0.1 (-18.7)	0.1 (-58.6)	0 (-43.1)	0.1 (-78)	0 (-96.2)	0.1 (-91.7)
C3	0.2 (74.6)	0.2 (43)	0.3 (-18.6)	0.2 (81.8)	0.2 (95.8)	0.3 (74.6)
D19	0.1 (14.5)	0.1 (7.3)	0 (-11.6)	0.1 (-16)	0 (-21.8)	0.1 (-4.8)
D26	0.1 (32)	0.1 (9.3)	0.1 (-17.6)	0.1 (-0.9)	0.1 (7.9)	0.1 (18.7)
D28	0.1 (-9.3)	0 (-18.2)	0 (-12.5)	0.1 (-42.3)	0 (-24.7)	0.1 (-34.7)
D4	0.1 (10.5)	0.1 (47.8)	0.1 (0.2)	0.1 (66.9)	0 (33.6)	0.1 (68.3)
D6	0.1 (19.9)	0.1 (33.3)	0.1 (-7.3)	0.1 (36)	0.1 (56.4)	0.1 (45.3)
D7	0.1 (-18.1)	0.1 (34.4)	0.1 (-0.3)	0.1 (21.2)	0 (-16)	0.1 (34)
P8	0.2 (178.6)	0.3 (49.2)	0.2 (-86.1)	0.3 (-52.7)	0.1 (-23.9)	0.3 (-61.1)
NO₂⁻/NO₃²⁻						
C10	1 (48.1)	1.2 (23)	1.1 (-38.1)	1 (-12.5)	1 (-0.2)	1.4 (20.4)
C3	0.1 (-4.4)	0.1 (11.7)	0.2 (-0.8)	0.2 (-2.3)	0.1 (29.4)	0.2 (3.3)
D19	0.4 (-22)	0.4 (11.1)	0.3 (-23.7)	0.5 (-28.6)	0.2 (-34)	0.4 (-16.3)
D26	0.3 (-9)	0.3 (12.4)	0.3 (-14.8)	0.4 (-13.4)	0.2 (-11.8)	0.3 (-4.7)
D28	0.4 (-29.1)	0.4 (7)	0.3 (-34.7)	0.5 (-34.5)	0.2 (-59.1)	0.4 (-53.8)
D4	0.3 (-8.2)	0.3 (16.6)	0.3 (-3.7)	0.4 (5.4)	0.3 (6.6)	0.3 (7.3)
D6	0.3 (-9.5)	0.4 (16.5)	0.4 (-7.5)	0.4 (-17.9)	0.3 (42.6)	0.3 (27.6)
D7	0.3 (-7.5)	0.4 (20.5)	0.4 (-7.9)	0.4 (-12.7)	0.3 (41.9)	0.3 (24.4)
P8	0.9 (61.3)	1.2 (32.7)	1.3 (-12.5)	1.1 (52.2)	0.9 (60.6)	1.4 (67.3)

Table 2: Summaries of flow-normalized trends in nitrite/nitrate and ammonium (mg L^{-1}) concentrations before and after WWTP upgrades upstream of station P8. Upgrades were completed in May 2007 at the City of Tracy WWTP (San Joaquin County, Fig. 5). Summaries are means and percent changes based on annual means within the pre- and post-upgrade time periods (1976-2007, 2008-2013). Increasing values are in bold-italics.

Period	$\text{NO}_2^-/\text{NO}_3^{2-}$		NH_4^+	
	Mean	% change	Mean	% change
Annual				
1976-2007	1.27	75.7	0.33	71.3
2008-2013	1.47	-6	0.11	-62.6
Seasonal, pre				
JFM	1.46	77.7	0.49	57.1
AMJ	1	75.3	0.13	63.4
JAS	1.08	97.9	0.13	151.9
OND	1.53	81.9	0.54	102.4
Seasonal, post				
JFM	1.76	-3.4	0.14	-47
AMJ	0.94	-5.5	0.07	-45.3
JAS	1.17	-10	0.07	-50.3
OND	1.99	-4.6	0.15	-58.5

Table 3: Summaries of flow-normalized trends in dissolved inorganic nitrogen (mg L^{-1}), chlorophyll ($\mu\text{g L}^{-1}$), and silicon dioxide (mg L^{-1}) concentrations for different time periods at station D7. Summaries are means and percent changes based on annual means within the time periods. Increasing values are in bold-italics.

Period	DIN		Chl- <i>a</i>		SiO_2	
	Mean	% change	Mean	% change	Mean	% change
All						
1976-2013	0.4	27.8	4.4	-81.6	12.6	30.2
Annual						
1976-1985	0.4	10.2	9.8	-63.9	10.8	9
1986-1994	0.4	14.9	3.4	-48.8	12.2	14.2
1995-2003	0.5	4	2.1	-4.8	13.5	3.7
2004-2013	0.5	-8.6	2.5	28.5	13.7	-1.8
Seasonal						
JFM	0.5	-4.9	2.7	-56	15	4.4
AMJ	0.4	17.4	6.1	-74.3	12.9	20.9
JAS	0.3	40.6	6.2	-89	10.9	79.1
OND	0.5	13.1	2.7	-83.8	11.7	33.7

Table 4: Summaries of flow-normalized trends in dissolved inorganic nitrogen ($\text{mg } L^{-1}$), chlorophyll ($\mu\text{g } L^{-1}$), and silicon dioxide ($\text{mg } L^{-1}$) concentrations for different time periods at station C10. Summaries are means and percent changes based on annual means within the time periods. Increasing values are in bold-italics.

Period	DIN		Chl- <i>a</i>		SiO ₂	
	Mean	% change	Mean	% change	Mean	% change
All						
1976-2013	1.3	-16.5	27.5	9	14.4	-21.7
Annual						
1976-1985	1.3	12.1	26.1	-3.3	15.7	-4.3
1986-1994	1.4	6.3	27.5	23.5	14.8	-4
1995-2003	1.4	-0.8	29.3	-15.3	14.1	-5
2004-2013	1.2	-29.4	27.1	5.1	13.1	-8.7
Seasonal						
JFM	1.4	-20.5	12	14.8	14.3	-16.9
AMJ	1	-26.4	43	3.1	13.1	-27.5
JAS	1.3	-3.7	46.6	-13.6	14.8	-26.1
OND	1.6	-6.8	8.7	-18.5	15.6	-16