

Improving spatial resolution in estimates of seagrass depth of colonization

Marcus W. Beck

October 27, 2014

1 Outline

- Needs
 - Seagrass related to habitat quality and strongly affected by water clarity
 - Extensive datasets describing historical and current seagrass growth patterns and distribution in Florida estuaries
 - No consistent approach for estimating depth of colonization (DoC) to establish restoration targets
 - WBID has been considered appropriate management unit although considerable spatial heterogeneity in seagrass growth
 - Reproducible and empirical approaches can be developed that leverage multiple types of information to provide more consistent estimates for restoration targets or nutrient criteria
- Objectives
 - Use information-rich datasets to estimate seagrass DoC by incorporating spatially referenced information
 - Provide a basis for using these estimates to inform nutrient criteria development using empirical relationships with water clarity
- Approach
 - Describe Hagy method and/or WBID approach, emphasis on situations where seagrass growth is spatially variable or when restoration target is misinformed
 - Describe spatially-referenced method, case studies
 - Compare/contrast the two, with emphasis on relation to secchi data

- Implications for criteria development and/or restoration targets
- To do
 - Rectify seagrass depth bin procedures
 - Tidal datum correction
 - Segment specific relationship of seagrass depth of col w/ Secchi
 - Compare cumulative sum approach with binning
 - Quantitative evaluation of grid spacing, grid location, and radius

2 Methods

The following describes methods used to estimate seagrass DoC that incorporate spatial information to improve resolution. Methods build extensively on those in Hagy et al. (in prep).

2.1 Within-segment variation in seagrass depth of colonization estimates

First, an example is provided that illustrates spatial heterogeneity within an individual estuary segment to highlight a need for improved resolution. Segments are commonly used as a basis for quantifying nutrient criteria and it is shown that these spatial scales may not be sufficient for characterizing variation in seagrass growth. Hagy et al. (in prep) describe methods for estimating seagrass DoC by segments using historical and current records of seagrass coverage combined with bathymetric data. The approach begins by combining bathymetric sounding data with coverage data to create a Geographic Information System (GIS) layer containing both sets of information. The points are grouped into depth bins and the proportion of points within each depth bin that contain seagrass are quantified. The maximum DoC for each segment is estimated using a plot of proportion of points occupied against depth bin. In general, the plot is characterized by a decreasing trend such that the proportion of occupied points by depth bin decreases and flattens with increasing depth. A regression is fit on this descending portion of the curve such that the intercept point on the x-axis is considered the maximum depth of colonization. The median portion of this curve is considered the median depth of the deepwater edge of seagrass. Estimates are obtained for seagrass coverage layers that describe continuous and continuous with patchy (all) seagrass.

Figure 1 illustrates spatial variation in seagrass distribution on a latitudinal gradient in Old Tampa Bay, Florida. Using methods in Hagy et al. (in prep), the estimate for median

seagrass DoC for the segment is an over- and under-estimate for northern and southern portions, respectively. Figure 2 provides a similar example for the a segment in the Big Bend region of Florida. Again, seagrass depth of colonization is over- and under-estimated for different areas of the segment. In particular, DoC is greatly over-estimated at the outflow of the Steinhatchee where high concentrations of dissolved organic matter naturally limit seagrass growth. These examples suggest that estimates of DoC are needed at finer spatial scales to provide a more robust determination of restoration targets and nutrient criteria.

2.2 Estimating seagrass depth of colonization using spatial information

An alternative approach for estimating seagrass DoC was developed that incorporates spatial information for improved resolution. The new approach has a similar theoretical foundation as the original, although the methods for estimation are slightly different. The first difference is that the maximum DoC is estimated from a logistic growth curve fit through the data. The second and more important difference is that the estimates are specific to locations using a grid-based approach. These main differences are described below using an example from the Big Bend region of Florida.

Input data for estimating spatially-referenced depth of colonization require only two shapefiles: a segment polygon and a point layer of bathymetric soundings and seagrass presence/absence. The latter shapefile was created using methods in described in Hagy et al. (in prep). In general, the point layer contains coordinates of bathymetric sounding points that were within 1 km of a seagrass coverage layer. The sounding points were merged with the coverage layer to identify points occurring within seagrass beds.

The spatially-referenced approach for estimating depth of colonization begins by selecting a single sampling point within a grid of sampling points at a set distance within the segment polygon. Bathymetric soundings that occur within a set radius from each sampling point are selected (fig. 3). An estimate of seagrass DoC is obtained using the selected points and assigned as a spatially-referenced value to each sampling point. The seagrass DoC estimate for each sample point is quantified from the proportion of bathymetric soundings that contain seagrass at each depth bin in the data (fig. 4a). A buffer around a sample point that is sufficient to quantify depth of colonization typically has a plot similar to fig. 4a such that the proportion of points that are occupied by seagrass decreases continuously with increasing depth.

A decreasing logistic growth curve was then fit to the points to create a monotonic and asymptotic function to estimate depth of colonization. This curve is fit using non-linear regression to characterize the reduction in points occupied by seagrass as a function of depth. The logistic growth curve is fit using standard residual sums-of-squares and user-supplied starting parameters that are an approximate estimate of the curve characteristics.

The model has the following form:

$$Proportion = \frac{\alpha}{1 + e^{(\beta - Depth)/\gamma}} \quad (1)$$

where the proportion of points occupied by seagrass at each depth is defined by a logistic curve with an asymptote α , a midpoint inflection β , and a scale parameter γ .

Finally, a simple linear curve is fit through the inflection point of the logistic curve to estimate depth of colonization (fig. 4c). The inflection point is the depth at which seagrass are decreasing at a maximum rate and is assigned to the slope of the linear curve. Three measures are obtained from the linear curve. The maximum depth of seagrass colonization, DOC_{max} , is the x-axis intercept of the linear curve. The depth of maximum seagrass occupancy, SG_{max} is the location where the linear curve intercepts the asymptote of the logistic growth curve. Finally, the median depth of seagrass colonization, DOC_{med} , is the depth halfway between SG_{max} and DOC_{max} .

The method of estimating each of the three measures applies only under specific scenarios. First, estimates can only be provided if a sufficient number of points are present within the radius of the sample point to estimate a logistic growth curve. Second, estimates are only provided if an inflection point is present on the logistic curve within the range of the depth data. Finally, the estimate for SG_{max} is set to zero if the linear curve through the inflection point intercepts the asymptote at x-axis values less than zero. The estimate for DOC_{med} is also shifted if SG_{max} is zero. Adding exceptions for instances when depth estimates cannot be calculated provides a safety measure that ensures a sufficient amount of data are used in the calculations.

2.3 Sensitivity analysis and comparison with segment-based approach

2.4 Developing a spatially coherent relationship of water clarity with depth of colonization




Figure 1: Example of over- and under-estimates for seagrass depth of colonization in Old Tampa Bay, Florida. The top-left figure indicates over-estimation and the bottom-left indicates under-estimation. Bathymetric points are color-coded by the median depth of colonization estimate for all seagrass (patchy and continuous) in the segment.




Figure 2: Example of over- and under-estimates for seagrass depth of colonization for a segment in the Big Bend region, Florida. The top-left figure indicates over-estimation and the bottom-left indicates under-estimation. Bathymetric points are color-coded by the median depth of colonization estimate for continuous seagrass in the segment.







Figure 3: Sample grid used to estimate spatially-referenced depth of colonization. The selected bathymetric/seagrass points that were within a radius of 0.05 decimal degrees from the test point were selected for the estimate. Estimates can be obtained for each sample grid point and an arbitrary radius.



(a) Proportion of points with seagrass by depth



(b) Logistic growth curve fit through points



(c) Depth estimates

Figure 4: Methods for estimating seagrass depth of colonization. Figure 4a is the proportion of points with seagrass by depth using depth points within the buffer of the test point in fig. 3. Figure 4b adds a decreasing logistic growth curve fit through the points. Figure 4c shows three depth estimates based on a linear curve fit through the inflection point of logistic growth curve.