

1 **Seagrass light requirements using an algorithm to spatially**
2 **resolve depth of colonization**

3 **Marcus W. Beck¹, James D. Hagy III²**

¹ *ORISE Research Participation Program*

USEPA National Health and Environmental Effects Research Laboratory

Gulf Ecology Division, 1 Sabine Island Drive, Gulf Breeze, FL 32561

Phone: 850-934-2480, Fax: 850-934-2401, Email: beck.marcus@epa.gov

² *USEPA National Health and Environmental Effects Research Laboratory*

Gulf Ecology Division, 1 Sabine Island Drive, Gulf Breeze, FL 32561

Phone: 850-934-2455, Fax: 850-934-2401, Email: hagy.jim@epa.gov

1 Introduction

Issues related to excessive nutrient pollution have motivated a substantial amount of research to understand and address impacts on coastal waters. Eutrophication, defined as an increase in the rate of supply of organic matter to an ecosystem (Nixon 1995), is primarily caused by anthropogenic inputs of limiting nutrients that exceed background concentrations of receiving waters. Adverse impacts on aquatic resources are well-documented and have included increased occurrence in the frequency and severity of harmful algal blooms (Cloern 1996), reduction of dissolved oxygen necessary to support heterotrophic organisms (Justic et al. 1987, Diaz and Rosenberg 2008), and loss of ecosystem functioning through food web simplification (Tewfik et al. 2007). Although management activities have been successful in mitigating or reversing eutrophication impacts (e.g., Greening and Janicki 2006), the evaluation of response endpoints remains an important topic given that ecosystem changes in relation to different nutrient regimes are not fully understood nor anticipated (Duarte et al. 2009). The most appropriate indicators of ecosystem response may be those that exhibit clear biological linkages with water quality changes, such that the potential effects of management actions can be unambiguously characterized through known cause and effect pathways. Critical management decisions may be forced by tentative assessments, political or societal pressures, or qualitative criteria in the absence of empirical methods to identify adequate indicators of ecosystem response (Duarte et al. 2009).

The ecosystem services provided by seagrasses as well as their sensitivity to water quality changes has contributed to their proliferation as biological response endpoints for eutrophication. Seagrasses are ecosystem engineers (Jones et al. 1994, Koch 2001) that serve a structural and

26 functional role in altering aquatic habitat often through multiple feedback mechanisms with other
27 ecosystem components. For example, seagrass beds create habitat for juvenile fish and crabs by
28 reducing wave action and stabilizing sediment (Williams and Heck 2001, Hughes et al. 2009).
29 Seagrasses also respond to changes in water clarity through direct physiological linkages with
30 light availability. In short, increased nutrient loading contributes to reductions in water clarity
31 through increased algal concentrations, inhibiting the growth of seagrass through light limitation
32 (Duarte 1995). Empirical relationships between nutrient loading, water clarity, light requirements,
33 and the maximum depth of seagrass colonization have been identified (Duarte 1991, Kenworthy
34 and Fonseca 1996, Choice et al. 2014), such that quantitative standards can be developed to
35 maintain light regimes sufficient for seagrass growth targets (Steward et al. 2005). Conversely,
36 seagrass depth limits have formed the basis of quantitative criteria for nutrient load targets
37 (Janicki and Wade 1996). Contrasted with numeric standards for nutrients and phytoplankton,
38 seagrass-based criteria may be more practical for developing water quality standards given that
39 seagrasses are integrative of system-wide conditions over time and less variable with changes in
40 nutrient regimes (Duarte 1995).

41 The development of numeric criteria and standards for coastal waters has been a
42 management priority within the United States (USEPA, 1998) and internationally (WFD 2000).
43 Numerous agencies and management programs have developed a variety of techniques for
44 estimating seagrass depth limits as a basis for establishing numeric criteria, either as restoration
45 targets or for identifying critical load limits. Such efforts have been useful for site-specific
46 approaches where the analysis needs are driven by a particular management or research context
47 (e.g., Iverson and Bittaker 1986, Hale et al. 2004). However, a lack of standardization among
48 methods has prevented broad-scale comparisons between regions and has even contributed to

49 discrepancies between measures of depth limits based on the chosen technique. For example,
50 seagrass depth limits based on in situ techniques can vary with the sampling device (Spears et al.
51 2009). Despite the availability of data, techniques for estimating seagrass depth of colonization
52 using remotely sensed data have not been extensively developed. Such techniques have the
53 potential to facilitate broad-scale comparisons between regions given the spatial coverage and
54 annual availability of many products. For example, recent analyses by Hagy, In review have
55 shown that standardized techniques using seagrass coverage maps and bathymetric data can be
56 developed to compare growth patterns over time among different coastal regions of Florida. Such
57 methods show promise, although further development to improve the spatial resolution of the
58 analysis are needed. Specifically, methods for estimating seagrass depth limits should be
59 reproducible for broad-scale comparisons, while also maintaining flexibility for site-specific
60 estimates depending on management needs.

61 Reproducible and empirical approaches can be developed to provide more consistent
62 estimates of seagrass depth limits for restoration targets or criteria development. We describe a
63 method for estimating seagrass depth of colonization using information-rich datasets to create a
64 spatially explicit and repeatable estimate. In particular, methods described in Hagy, In review are
65 improved upon by creating a flexible and repeatable technique for estimating seagrass depth
66 limits from coverage maps and bathymetric data. The specific objectives are to 1) describe the
67 method for estimating seagrass depth limits within a relevant spatial context, 2) apply the
68 technique to four distinct regions of Florida to illustrate improved clarity of description for
69 seagrass growth patterns, and 3) develop a spatially coherent relationship between depth limits
70 and water clarity for the case studies. Overall, these methods are expected to inform the
71 development of water quality criteria based on empirical relationships of seagrass depth limits

72 with water clarity over time. The method is applied to data from Florida although the technique is
73 transferable to other regions with comparable data.

74 **2 Methods**

75 Development of a spatially-resolved approach to estimate seagrass depth of {acro:doc}
76 colonization (Z_c) relied extensively on data and partially on methods described in Hagy, In
77 review. The following is a summary of data sources, methods and rationale for improving spatial
78 resolution in seagrass Z_c estimates, and evaluation of the approach including relationships with
79 water clarity.

80 **2.1 Data sources**

81 **2.1.1 Study sites**

82 Four locations in Florida were chosen for the analysis: Choctawhatchee Bay (Panhandle),
83 Big Bend region (northeast Gulf of Mexico), Tampa Bay (central Gulf Coast of Florida), and
84 Indian River Lagoon (east coast) (Table 1 and Fig. 1). These locations represent different
85 geographic regions in the state, in addition to having available data and observed gradients in
86 water clarity that contribute to heterogeneity in seagrass growth patterns. For example, the Big
87 Bend region was chosen based on location near an outflow of the Steinhatchee River where higher
88 concentrations of dissolved organic matter are observed. Seagrasses near the outflow were
89 observed to grow at shallower depths as compared to locations far from the river source. Coastal
90 regions and estuaries in Florida are partitioned as distinct spatial units based on a segmentation
91 scheme developed by US Environmental Protection Agency (EPA) for the development of {acro:EPA}
92 numeric nutrient criteria. One segment from each geographic location was used to estimate
93 seagrass Z_c and to evaluate variation in growth patterns. The segments included portions of

94 Choctawhatchee Bay (segment 303), the big bend region (820), Old Tampa Bay (902), and Indian
95 River Lagoon (1502) (Fig. 1).

96 **2.1.2 Seagrass coverage and bathymetry**

{sec: data_}

97 Spatial data describing seagrass aerial coverage combined with co-located bathymetric
98 depth information were used to estimate Z_c . These geospatial data products are publicly
99 available in coastal regions of Florida through the US Geological Survey, Florida Department of
100 Environmental Protection, Florida Fish and Wildlife Conservation Commission, and watershed
101 management districts. Seagrass coverage maps were obtained for recent years in each of the study
102 sites described above (Table 1). Coverage maps were produced using photo-interpretations of
103 aerial images to categorize seagrass as absent, discontinuous (patchy), or continuous. For this
104 analysis, we considered seagrass as only present (continuous and patchy) or absent since
105 differences between continuous and patchy coverage were often inconsistent between data
106 sources.

107 Seagrass coverage maps were combined with bathymetric depth layers to characterize
108 location and depth of growth in each location. Bathymetric depth layers for each location were
109 obtained from the National Oceanic and Atmospheric Administration's (NOAA) National
110 Geophysical Data Center (<http://www.ngdc.noaa.gov/>) as either Digital Elevation
111 Models (DEMs) or raw sounding data from hydroacoustic surveys. Tampa Bay data provided by
112 the Tampa Bay National Estuary Program are described in Tyler et al. (2007). Bathymetric data
113 for the Indian River Lagoon were obtained from the St. John's Water Management District

{acro: DEM}

114 (Coastal Planning and Engineering 1997). NOAA products were referenced to mean lower low
115 water, whereas Tampa Bay data were referenced to the North American Vertical Datum of
116 1988 (NAVD88) and the Indian River Lagoon data were referenced to mean sea level. Depth

{acro: NAVD}

117 layers were combined with seagrass coverage layers using standard union techniques for raster
118 and vector layers in ArcMap 10.1 (Environmental Systems Research Institute 2012). To reduce
119 computation time, depth layers were first masked using a 1 km buffer of the seagrass coverage
120 layer. The final layer used for analysis was a point layer with attributes describing location
121 (latitude, longitude, segment), depth (m), and seagrass (present, absent). All spatial data were
122 referenced to the North American Datum of 1983 as geographic coordinates. Depth values in
123 each seagrass layer were further adjusted from the relevant vertical reference datum to local mean sea level (MSL) {acro:MSL}
124 using the NOAA VDatum tool (<http://vdatum.noaa.gov/>).

125 **2.1.3 Water clarity**

126 Seagrass light requirements can be estimated by evaluating spatial relationships between
127 depth of colonization and water clarity. Secchi measurements provide a precise estimate of water
128 clarity and have been obtained at numerous locations documented in the Florida Department of
129 Environmental Protection's Impaired Impaired Waters Rule (IWR) database. Secchi data (as
130 depth in meters, Z_{secchi}) for Florida coastal waters were obtained from update 40 of the IWR {acro:IWR}
131 database for all of Tampa Bay (2010 coverage) and the Indian River Lagoon (2009 coverage)
132 given the spatial extent of secchi observations for the two locations relative to the Big Bend and
133 Choctawhatchee segments. Secchi data within the previous ten years of the seagrass coverage
134 data were evaluated to capture water quality trends from the most recent decade (i.e., 1999–2009
135 for the Indian River Lagoon and 2000–2010 for Tampa Bay). Stations with less than five
136 observations and observations that were flagged indicating that the value was lower than the
137 maximum depth of the observation point were removed. Secchi data were also compared with
138 bathymetric data to verify unflagged values were not missed by initial screening. Secchi
139 observations that were measured at the same geographic location were averaged across all dates.

140 This approach was preferred given that seagrass depth patterns are more representative of
141 long-term trends in water clarity as opposed to individual secchi measures that may be highly
142 variable (Dennison 1987, Dennison et al. 1993).

143 **2.2 Segment-based estimates of seagrass depth of colonization**

144 [Hagy, In review](#) describe an approach to estimate seagrass Z_c as a segment-wide median.
145 Seagrass depth data described above can be used to estimate maximum ($Z_{c,max}$) and median
146 ($Z_{c,med}$) seagrass Z_c , where the maximum depth is defined as the deepest depth at which a
147 “significant” coverage of mappable seagrasses occurred in a segment and the median depth is
148 defined as the median depth occurring at the deep water edge. The seagrass depth points are
149 grouped into bins and the proportion of points within each depth bin that contain seagrass are
150 quantified. Both seagrass Z_c estimates are obtained from a plot of proportion of points occupied
151 at each depth bin. In general, the plot is characterized by a decreasing trend such that the
152 proportion of occupied points by depth bin decreases and eventually flattens with increasing
153 depth. A regression is fit on this descending portion of the curve such that the intercept point on
154 the x-axis is considered the maximum depth of colonization. The median portion of this curve is
155 considered the median depth of the deepwater edge of seagrass.

156 A segment-wide average of seagrass Z_c , although unbiased, may potentially reduce the
157 ability to relate patterns in Z_c to relevant water quality variables. Considerable spatial
158 heterogeneity in the observed seagrass growth patterns suggests that a segment-wide estimate of
159 seagrass Z_c may not fully describe variation at relevant spatial scales. Fig. 2a illustrates variation
160 in seagrass distribution for a location in the Big Bend region of Florida. Using methods in [Hagy,](#)
161 [In review](#), the segment-wide estimate for maximum depth of seagrass colonization (shown as a

162 red contour line) does not adequately describe within-segment variation in depth limits. Z_c is
163 greatly over-estimated at the outflow of the Steinhatchee River where high concentrations of
164 dissolved organic matter reduce water clarity and naturally limit seagrass growth. This example
165 suggests that it may be useful to have improved spatial resolution in estimates of Z_c , particularly
166 when site-specific characteristics may require a more detailed description of seagrass growth
167 patterns. Although the current example is immediately relevant for the Big Bend region of
168 Florida, the remaining examples discussed throughout also provide a justification for a more
169 comprehensive assessment of seagrass growth patterns.

170 **2.3 Estimating seagrass depth of colonization for finite and arbitrary areas**

171 The approach used to estimate seagrass Z_c with improved spatial resolution has several
172 key differences that make it distinct from the original method. As before, seagrass Z_c estimates
173 are based on empirical measures of the frequency occurrence of seagrass with increasing depth.
174 The first difference is that maximum Z_c is estimated using a logistic growth curve fit through the
175 data, as compared to a simple linear regression in the previous example. Second, a third measure
176 describing the minimum depth of colonization was defined, in addition to median and maximum
177 depth of growth. The third and most important difference is that the estimates are assigned to
178 discrete cartesian locations, using either a grid of points or as a single location of interest. The
179 area around each point within which seagrass depth data are used to estimate Z_c is based on a
180 fixed, arbitrary radius that can be increased or decreased depending on the question of interest.
181 Methods and implications of these differences are described below.

182 The spatially-resolved approach for estimating Z_c begins by creating a grid of points
183 within the segment where the same process for estimating Z_c is used for each point. Alternatively,

184 a single location of interest can be chosen rather than a grid-based design. Seagrass depth data
185 (i.e., merged bathymetric and seagrass coverage data) that are located within a set radius from the
186 chosen locations are selected for estimating seagrass Z_c values (Fig. 2). The estimate for each
187 location is quantified from a plot of the proportion of bathymetric soundings that contain seagrass
188 at each depth bin (Fig. 3a). Although the chosen radius for selecting depth points is
189 problem-specific, the minimum radius must sample a sufficient number of points for estimating
190 Z_c . In general, an appropriate radius will produce a plot that indicates a decrease in the proportion
191 of points that are occupied by seagrass with increasing depth. An appropriate radius is also one
192 that creates a sample area around each point that has minimal overlap with the seagrass depth data
193 sampled by adjacent points.

194 A curve is fit to the sampled depth points using non-linear regression to characterize the
195 reduction in seagrass as a function of depth (Fig. 3b). Specifically, a decreasing logistic growth
196 curve is used with the assumption that seagrass decline with increasing depth is monotonic and
197 asymptotic at the maximum depth of colonization. The curve is fit by minimizing the residual
198 sums-of-squares with the Gauss-Newton algorithm (Bates and Chambers 1992) and user-supplied
199 starting parameters that are an approximate estimate of the curve characteristics. The model has
200 the following form:

$$\text{Proportion} = \frac{\alpha}{1 + e^{(\beta - Z)/\gamma}} \quad (1) \quad \{\text{eqn:prop}\}$$

201 where the proportion of points occupied by seagrass at each depth, Z , is defined by a logistic
202 curve with an asymptote α , a midpoint inflection β , and a scale parameter γ . Starting values α , β ,
203 and γ were estimated empirically from the observed data.

204 Finally, a simple linear curve is fit through the inflection point (β) of the logistic curve to

205 estimate the three measures of depth of colonization (Fig. 3c). The inflection point is considered
206 the depth at which seagrass are decreasing at a maximum rate and is used as the slope of the
207 linear curve. Three measures describing seagrass growth characteristics are obtained. The
208 maximum depth of seagrass colonization, $Z_{c,max}$, is the x-axis intercept of the linear curve. The
209 minimum depth of seagrass growth, $Z_{c,min}$, is the location where the linear curve intercepts the
210 asymptote of the logistic growth curve. This depth can be considered the start of the decline in
211 seagrass coverage with increasing depth. The median depth of seagrass colonization, $Z_{c,med}$, is
212 the depth halfway between $Z_{c,min}$ and $Z_{c,max}$. $Z_{c,med}$ was typically the inflection point of the
213 logistic growth curve. Functionally, each measure has specific ecological significance. The
214 median and maximum depth estimates describe the growth limitations of seagrasses as a function
215 of water clarity, whereas minimum depth of growth was often where the highest percentage of
216 seagrass coverage was observed in the sample. Median and maximum depth estimates differ in
217 that the former describes the median depth of the deep water edge, whereas the latter describes a
218 nominal characterization of maximum depth independent of outliers.

219 Estimates for each of the three Z_c measures are obtained only if specific criteria are met.
220 These criteria were implemented as a safety measure that ensures a sufficient amount and
221 appropriate quality of data were used. First, estimates were provided only if a sufficient number
222 of seagrass depth points were present within the radius of the grid point to estimate a logistic
223 growth curve. This criteria applies to the sample size as well as the number of points with
224 seagrass in the sample. The curve could not be estimated for small samples or if an insufficient
225 number of points contained seagrass regardless of sample size. Second, estimates were provided
226 only if an inflection point was present on the logistic curve within the range of the sampled depth
227 data. This criteria applied under two scenarios where the curve was estimated but a trend was not

adequately described by the sampled data. That is, a curve could be estimated that described only the initial decrease in points occupied as a function of depth but the observed points do not occur at depths deeper than the predicted inflection point. The opposite scenario occurred when a curve was estimated but only the deeper locations beyond the inflection point were present in the sample. Third, the estimate for $Z_{c,min}$ was set to zero depth if the linear curve through the inflection point intercepted the asymptote at x-axis values less than zero. The estimate for $Z_{c,med}$ was also shifted to the depth value halfway between $Z_{c,min}$ and $Z_{c,max}$ if $Z_{c,min}$ was fixed at zero. Finally, estimates were considered invalid if the 95% confidence interval for $Z_{c,max}$ included zero. Methods used to determine confidence bounds on Z_c estimates are described below.

All estimates were obtained using custom-made functions in program R that were based on the `nls` and `SSlogis` functions to fit nonlinear least squares using a self-starting logistic growth model (Bates and Chambers 1992, R Development Core Team 2014). All seagrass depth shapefiles were imported and processed in R using functions in the `rgeos` and `sp` packages (Bivand et al. 2008, Bivand and Rundel 2014).

2.4 Comparison with segment-based approach and sensitivity analysis

Spatially-resolved estimates for seagrass Z_c were obtained for each of the four segments described above. Segment-wide estimates obtained using methods in Hagy, In review were used as a basis of comparison such that departures from these values were evidence of spatial heterogeneity in seagrass growth patterns and improved clarity of description in depth estimates using the new approach. A sampling grid of locations for estimating each of the three depth values in Fig. 3 was created for each segment. The grid was masked by the segment boundaries, whereas seagrass depth points used to estimate Z_c extended beyond the segment boundaries to

allow sampling by grid points that occurred near the edge of the segment. Initial spacing between sample points was chosen arbitrarily as 0.02 decimal degrees, which is approximately 2 km at 30 degrees N latitude. The sampling radius around each sampling location in the grid was also chosen as 0.02 decimal degrees to allow for complete coverage of seagrass within the segment while also minimizing redundancy of information described by each location. In other words, radii were chosen such that the seagrass depth points sampled by each grid location were only partially overlapped by those sampled by neighboring points.

The ability to characterize heterogeneity in seagrass growth patterns using the grid-based approach can be informed by evaluating the level of confidence associated with Z_c estimates. Confidence intervals for non-linear regression can be estimated using a Monte Carlo simulation approach that considers the variance and covariance between the model parameters and the depth measurements (Hilborn and Mangel 1997). For simplicity, we assume that the variability associated with parameter estimates is the dominant source of uncertainty. A 95% confidence interval for each Z_c estimate was constructed by repeated sampling of a multivariate normal distribution followed by prediction of the proportion of points occupied by seagrass as in eq. (1).

The sampling distribution assumes:

$$x \sim N(\mu, \Sigma) \quad (2)$$

where x is a predictor variable used in eq. (1) (depth) that follows a multivariate normal distribution with mean μ , and variance-covariance matrix Σ . The mean values are set at the depth value corresponding to the inflection point on the logistic curve and the predicted model parameters (i.e., α , β , and γ), whereas Σ is the variance-covariance matrix of the model parameters. A large number of samples ($n = 10000$) were drawn from the distribution to

271 characterize the uncertainty of the depth value at the inflection point. The 2.5th and 97.5th quantile
272 values of the sample were considered bounds on the 95% confidence interval.

273 The uncertainty associated with the Z_c estimates were based on the upper and lower limits
274 of the estimated inflection point on the logistic growth curve. This approach was used because
275 uncertainty in the inflection point is directly related to uncertainty in each of the Z_c estimates that
276 are based on the linear curve fit through the inflection point. Specifically, linear curves were fit
277 through the upper and lower estimates of the depth value at the inflection point to identify upper
278 and lower limits for the estimates of $Z_{c,min}$, $Z_{c,med}$, and $Z_{c,max}$. These values were compared
279 with the initial estimates from the linear curve that was fit through the inflection point on the
280 predicted logistic curve (i.e., Fig. 3c). This approach provided an indication of uncertainty for
281 individual estimates for the chosen radius. Uncertainty estimates were obtained for each Z_c
282 estimate for the grids in each segment.

283 **2.5 Developing a spatially coherent relationship of water clarity with depth
284 of colonization**

285 The relationship between seagrass depth limits and secchi measurements were explored
286 using established light requirements and attenuation equations. The traditional Lambert-Beer
287 equation describes the exponential decrease of light availability with depth:

$$I_z = I_O \cdot \exp(-K_Z \cdot Z) \quad (3) \quad \{\text{eqn:lambda}\}$$

288 such that the irradiance of incident light at depth Z (I_z) can be estimated from the irradiance at
289 the surface (I_O) and a light extinction coefficient (K_Z). Duarte (1991) indicate that minimum
290 light requirements for seagrass are on average 11% of surface irradiance. Light requirements may

291 also be species-specific and variable by latitude such that value may range from less than 5% to
292 greater than 30% (Dennison et al. 1993). Light requirements of seagrass at a specific location can
293 be estimated by rearranging eq. (3):

$$\%light = \exp(-K_Z \cdot Z_{c,max}) \quad (4) \quad \{eqn:perc\}$$

294 where the percent light requirements of seagrass at $Z_{c,max}$ are empirically related to light
295 extinction. A conversion factor is often used to estimate the light extinction coefficient from
296 secchi depth Z_{secchi} , such that such that $c = K_Z \cdot Z_{secchi}$, where c has been estimated as 1.7
297 (Poole and Atkins 1929, Idso and Gilbert 1974). Thus, K_Z can be replaced with the conversion
298 factor and the equation is rearranged to describe $Z_{c,max}$ as a function of secchi depth Z_{secchi} :

$$Z_{c,max} = \frac{-\log(0.20)}{1.7} \cdot Z_{secchi} \quad (5) \quad \{eqn:sgreg\}$$

299 A regression of seagrass depth estimates against secchi measurements is expected to have a slope
300 corresponding to the constant in eq. (5). For the current analysis, 20% light requirements were
301 assumed to be an approximate median requirement for seagrasses in Florida. Scatter in the
302 regression through these points can be considered biologically meaningful, such that points below
303 the curve are locations where seagrasses are observed at maximum depth with less irradiance than
304 expected given eq. (5), whereas points above the curve are those where seagrasses are growing
305 deeper than expected. The geographic coordinates for each secchi measurement in Tampa Bay
306 and the Indian River Lagoon were used as locations for estimating $Z_{c,max}$. These estimates were
307 compared with the averaged secchi estimates to identify light requirements at each location.

308 However, the relationship is expected to vary depending on the specific radius around each
309 sample point for estimating $Z_{c, max}$. An appropriate radius was chosen that minimized the
310 difference between the empirically estimated slope and that in eq. (5). The estimated light
311 requirements of each point were also plotted using the cartesian coordinates of each secchi
312 observation to evaluate spatial variation in seagrass growth as a function of light-limitation. Light
313 requirements were also summarized by individual segments in each bay to identify spatial trends
314 for relevant management units.

315 **3 Results**

316 **3.1 Segment characteristics and seagrass depth estimates**

317 Each of the four segments varied by several key characteristics that potentially explain
318 within-segment variation of seagrass growth patterns (Table 1). Mean surface area was 191.2
319 square kilometers, with area decreasing for the Big Bend (271.4 km), Indian River Lagoon (NA
320 km), Old Tampa Bay (205.5 km), and Choctawhatchee Bay (59.4 km) segments. Seagrass
321 coverage as a percentage of total surface area varied considerably by segment. Seagrasses covered
322 a majority of the surface area for the Big Bend segment (74.8 %), whereas coverage was much
323 less for Indian River Lagoon (NA %), Old Tampa Bay (11.9 %), and Choctawhatchee Bay (5.9
324 %). Visual examination of the seagrass coverage maps for the respective year of each segment
325 suggested that seagrasses were not uniformly distributed (Fig. 1). Seagrasses in the
326 Choctawhatchee Bay segments were generally sparse with the exception of a large patch located
327 to the west of the inlet connection with the Gulf of Mexico. Seagrasses in the Big Bend segment
328 were located throughout the segment with noticeable declines near the outflow of the
329 Steinhatchee River, whereas seagrasses in Old Tampa Bay and the Indian River Lagoon segment

330 were generally confined to shallow areas near the shore. Seagrass coverage showed a partial
331 decline toward the northern ends of both Old Tampa Bay and the Indian River Lagoon segments.
332 Mean depth was less than 5 meters for each segment, excluding Choctawhatchee Bay which was
333 slightly deeper than the other segments on average (5.3 m). Maximum depths were considerably
334 deeper for Choctawhatchee Bay (11.9 m) and Old Tampa Bay (10.4 m), as compared to the Big
335 Bend (3.6 m) and Indian River Lagoon (NA m) segments. Water clarity as indicated by average
336 secchi depths was similar between the segments (1.5 m), although Choctawhatchee Bay had a
337 slightly higher average (2.1 m).

338 Estimates of seagrass Z_c using a segment-wide approach that did not consider spatially
339 explicit locations indicated that seagrasses generally did not grow deeper than three meters in any
340 of the segments (Table 2). Maximum and median depth of colonization were deepest for the Big
341 Bend segment (3.7 and 2.5 m, respectively) and shallowest for Old Tampa Bay (1.1 and 0.9 m),
342 whereas the minimum depth of colonization was deepest for Choctawhatchee Bay (1.8 m) and
343 shallowest for Old Tampa Bay (0.6 m). Averages of all grid-based estimates for each segment
344 were different than the segment wide estimates, which suggests potential bias associated with
345 using a whole segment as a relevant spatial unit for estimating depth of colonization. In most
346 cases, the averages of all grid-based estimates were less than the whole segment estimates,
347 suggesting the latter provided an over-estimate of seagrass growth limits. For example, the
348 average of all grid estimates for $Z_{c, max}$ in the Big Bend region suggested seagrasses grew to
349 approximately 2 m, which was 1.6 m less than the whole segment estimate. This reduction is
350 likely related to improved resolution of seagrass depth limits near the outflow of the Steinhatchee
351 river. Although reductions were not as severe for the average grid estimates for the remaining
352 segments, considerable within-segment variation was observed depending on grid location. For

353 example, the deepest estimate for $Z_{c, min}$ (2 m) in the Indian River Lagoon exceeded the average
354 of all grid locations for $Z_{c, max}$ (1.7 m). $Z_{c, min}$ also had minimum values of zero meters for the
355 Big Bend and Old Tampa Bay segments, suggesting that seagrasses declined continuously from the
356 surface for several locations.

357 Visual interpretations of seagrass depth estimates using the grid-based approach provided
358 further information on the distribution of seagrasses in each segment (Fig. 4). Spatial
359 heterogeneity in depth limits was particularly apparent for the Big Bend and Indian River Lagoon
360 segments. As expected, depth estimates indicated that seagrasses grew deeper at locations far
361 from the outflow of the Steinhatchee River in the Big Bend segment. Similarly, seagrasses were
362 limited to shallower depths at the north end of the Indian River Lagoon segment near the Merrit
363 Island National Wildlife Refuge. Seagrasses were estimated to grow at maximum depths up to 2.1
364 m on the eastern portion of the Indian River Lagoon segment. Spatial heterogeneity was less
365 distinct for the remaining segments. Seagrasses in Old Tampa Bay grew deeper in the northeast
366 portion of the segment and declined to shallower depths near the inflow at the northern edge.
367 Spatial variation in the Choctawhatchee Bay segment was not apparent, although the maximum
368 Z_c estimate was observed in the northeast portion of the segment. Z_c values were not available for
369 all grid locations given the limitations imposed in the estimation method. Z_c could not be
370 estimated in locations where seagrasses were sparse or absent, nor where seagrasses were present
371 but the sampled points did not exhibit a sufficient decline with depth. The latter scenario was
372 most common in Old Tampa Bay and Choctawhatchee Bay where seagrasses were unevenly
373 distributed or confined to shallow areas near the shore. The former scenario was most common in
374 the Big Bend segment where seagrasses were abundant but locations near the shore were
375 inestimable given that seagrasses did not decline appreciably within the depths that were sampled.

376 Uncertainty for estimates of $Z_{c,max}$ indicated that confidence intervals were generally
377 acceptable (i.e., greater than zero), although the ability to discriminate between the three depth
378 estimates varied by segment (Fig. 5 and Table 3). Mean uncertainty for all estimates in each
379 segment measured as the width of a 95% confidence interval was 0.2 m. Greater uncertainty was
380 observed for Choctawhatchee Bay (mean width of all confidence intervals was 0.7 m) and Old
381 Tampa Bay (0.4 m), compared to the Big Bend (0.1 m) and Indian River Lagoon (0.1 m)
382 segments. The largest confidence interval for each segment was 1 m for Old Tampa Bay, 2.5 m for
383 Choctawhatchee Bay, 0.4 m for the Big Bend, and 0.3 m for the Indian River Lagoon segments.
384 However, most confidence intervals for the remaining grid locations were much smaller than the
385 maximum in each segment. A comparison of overlapping confidence intervals for $Z_{c,min}$, $Z_{c,med}$,
386 and $Z_{c,max}$ at each grid location indicated that not every measure was unique. Specifically, only
387 12.5% of grid points in Choctawhatchee Bay and 38.9% in Old Tampa Bay had significantly
388 different estimates, whereas 84% of grid points in the Indian River Lagoon and 94.1% of grid
389 points in the Big Bend segments had estimates that were significantly different. By contrast, all
390 grid estimates in Choctawhatchee Bay and Indian River Lagoon had $Z_{c,max}$ estimates that were
391 significantly greater than zero, whereas all but 10% of grid points in Old Tampa Bay and 5.6% of
392 grid points in the Big Bend segment had $Z_{c,max}$ estimates significantly greater than zero.

393 **3.2 Evaluation of seagrass light requirements**

394 Estimates of seagrass depth limits and corresponding light requirements for all segments
395 of Tampa Bay and the Indian River Lagoon indicated substantial variation, both between and
396 within the different bays (Table 4 and Figs. 6 and 7). Seagrass Z_c estimates were obtained for 61
397 locations in Tampa Bay and 50 locations in the Indian River Lagoon where secchi observations

were available in the Florida IWR database. Mean secchi depth for all recorded observations was 1.9 m ($n = 61$) for Tampa Bay and 1 m ($n = 50$) for Indian River Lagoon. Mean light requirements were significantly different between the bays (two-sided t-test, $t = 8.5$, $df = 109$, $p < 0.001$) with a mean requirement of 23% for Tampa Bay and 10.6% for Indian River Lagoon. Within each bay, light requirements were significantly different between segments (ANOVA, $F = 5.6$, $df = 3, 57$, $p = 0.00$ for Tampa Bay, $F = 5.2$, $df = 7, 42$, $p = 0.000$ for Indian River Lagoon). However, post-hoc evaluation of all pair-wise comparisons of mean light requirements indicated that significant differences were only observed between a few segments within each bay. Significant differences in Tampa Bay were observed between Old Tampa Bay and Hillsborough Bay (Tukey multiple comparisons, $p = 0.032$). Significant differences in the Indian River Lagoon were observed between the Upper Indian River Lagoon and Banana River ($p = 0.915$), the Upper Indian River Lagoon and Lower Indian River Lagoon ($p = 0.140$), and Upper Indian River Lagoon and Lower St. Lucie ($p = 0.103$) segments. In general, spatial variation of light requirements in Tampa Bay suggested that seagrasses were less light-limited (i.e., lower percent light requirements at $Z_{c, max}$) in Hillsborough Bay and western areas of Lower Tampa Bay near the Gulf of Mexico (Fig. 6). Seagrassess in the Indian River Lagoon were generally less light-limited towards the south and in the Banana River segment (Fig. 7).

4 Discussion

416 **References**

- 417 Bates DM, Chambers JM. 1992. Nonlinear models. In: Chambers JM, Hastie TJ, editors,
418 Statistical Models in S, pages 421–454. Wadsworth and Brooks/Cole, Pacific Grove, California.
- 419 Bivand R, Rundel C. 2014. rgeos: Interface to Geometry Engine - Open Source (GEOS). R
420 package version 0.3-8.
- 421 Bivand RS, Pebesma EJ, Gómez-Rubio V. 2008. Applied Spatial Data Analysis with R. Springer,
422 New York, New York.
- 423 Choice ZD, Frazer TK, Jacoby CA. 2014. Light requirements of seagrasses determined from
424 historical records of light attenuatoin along the Gulf coast of peninsular Florida. Marine
425 Pollution Bulletin, 81(1):94–102.
- 426 Cloern JE. 1996. Phytoplankton bloom dynamics in coastal ecosystems: A review with some
427 general lessons from sustained investigation of San Francisco Bay, California. Review of
428 Geophysics, 34(2):127–168.
- 429 Coastal Planning and Engineering. 1997. Indian River Lagoon bathymetric survey. A final report
430 to St. John's River Water Management District. Technical Report Contract 95W142, Coastal
431 Planning and Engineering, Palatka, Florida.
- 432 Dennison WC. 1987. Effects of light on seagrass photosynthesis, growth and depth distribution.
433 Aquatic Botany, 27(1):15–26.
- 434 Dennison WC, Orth RJ, Moore KA, Stevenson JC, Carter V, Kollar S, Bergstrom PW, Batiuk RA.
435 1993. Assessing water quality with submersed aquatic vegetation. BioScience, 43(2):86–94.
- 436 Diaz RJ, Rosenberg R. 2008. Spreading dead zones and consequences for marine ecosystems.
437 Science, 321:926–929.
- 438 Duarte CM. 1991. Seagrass depth limits. Aquatic Botany, 40(4):363–377.
- 439 Duarte CM. 1995. Submerged aquatic vegetation in relation to different nutrient regimes.
440 Ophelia, 41:87–112.
- 441 Duarte CM, Conley DJ, Carstensen J, Sánchez-Camacho M. 2009. Return to *Neverland*: Shifting
442 baseline affect eutrophication restoration targets. Estuaries and Coasts, 32(1):29–36.
- 443 Environmental Systems Research Institute. 2012. ArcGIS v10.1. ESRI, Redlands, California.
- 444 Greening H, Janicki A. 2006. Toward reversal of eutrophic conditions in a subtropical estuary:
445 Water quality and seagrass response to nitrogen loading reductions in Tampa Bay, Florida,
446 USA. Environmental Management, 38(2):163–178.
- 447 Hagy JD. In review. Seagrass depth of colonization in Florida estuaries.
- 448 Hale JA, Frazer TK, Tomasko DA, Hall MO. 2004. Changes in the distribution of seagrass species
449 along Florida's central gulf coast: Iverson and Bittaker revisited. Estuaries, 27(1):36–43.

- 450 Hilborn R, Mangel M. 1997. *The Ecological Detective: Confronting Models with Data*.
451 Princeton University Press, Princeton, New Jersey.
- 452 Hughes AR, Williams SL, Duarte CM, Heck KL, Waycott M. 2009. Associations of concern:
453 declining seagrasses and threatened dependent species. *Frontiers in Ecology and the
454 Environment*, 7(5):242–246.
- 455 Idso SB, Gilbert RG. 1974. On the universality of the Poole and Atkins secchi disk-light
456 extinction equation. *Journal of Applied Ecology*, 11(1):399–401.
- 457 Iverson RL, Bittaker HF. 1986. Seagrass distribution and abundance in eastern Gulf of Mexico
458 coastal waters. *Estuarine, Coastal and Shelf Science*, 22(5):577–602.
- 459 Janicki A, Wade D. 1996. Estimating critical external nitrogen loads for the Tampa Bay estuary:
460 An empirically based approach to setting management targets. Technical Report 06-96, Tampa
461 Bay National Estuary Program, St. Petersburg, Florida.
- 462 Jones CG, Lawton JH, Shachak M. 1994. Organisms as ecosystem engineers. *OIKOS*,
463 69(3):373–386.
- 464 Justić D, Legović T, Rottini-Sandrini L. 1987. Trends in oxygen content 1911–1984 and
465 occurrence of benthic mortality in the northern Adriatic Sea. *Estuarine, Coastal and Shelf
466 Science*, 25(4):435–445.
- 467 Kenworthy WJ, Fonseca MS. 1996. Light requirements of seagrasses *Halodule wrightii* and
468 *Syringodium filiforme* derived from the relationship between diffuse light attenuation and
469 maximum depth distribution. *Estuaries*, 19(3):740–750.
- 470 Koch EW. 2001. Beyond light: Physical, geological, and geochemical parameters as possible
471 submersed aquatic vegetation habitat requirements. *Estuaries*, 24(1):1–17.
- 472 Nixon SW. 1995. Coastal marine eutrophication: A definition, social causes, and future concerns.
473 *Ophelia*, 41:199–219.
- 474 Poole HH, Atkins WRG. 1929. Photo-electric measurements of submarine illumination
475 throughout the year. *Journal of the Marine Biological Association of the United Kingdom*,
476 16:297–324.
- 477 R Development Core Team. 2014. R: A language and environment for statistical computing,
478 v3.1.2. R Foundation for Statistical Computing, Vienna, Austria. <http://www.R-project.org>.
- 479 Spears BM, Gunn IDM, Carvalho L, Winfield IJ, Dudley B, Murphy K, May L. 2009. An
480 evaluation of methods for sampling macrophyte maximum colonisation depth in Loch Leven,
481 Scotland. *Aquatic Botany*, 91(2):75–81.
- 482 Steward JS, Virnstein RW, Morris LJ, Lowe EF. 2005. Setting seagrass depth, coverage, and light
483 targets for the Indian River Lagoon system, Florida. *Estuaries*, 28(6):923–935.

- 484 Tewfik A, Rasmussen JB, McCann KS. 2007. Simplification of seagrass food webs across a
485 gradient of nutrient enrichment. Canadian Journal of Fisheries and Aquatic Sciences,
486 64(7):956–967.
- 487 Tyler D, Zawada DG, Nayegandhi A, Brock JC, Crane MP, Yates KK, Smith KEL. 2007.
488 Topobathymetric data for Tampa Bay, Florida. Technical Report Open-File Report 2007-1051
489 (revised), US Geological Survey, US Department of the Interior, St. Petersburg, Florida.
- 490 USEPA (US Environmental Protection Agency). 1998. National strategy for the development of
491 regional nutrient criteria. Technical Report EPA-822-R-98-002, Office of Water, Office of
492 Research and Development, US Environmental Protection Agency, Washington, DC.
- 493 WFD. 2000. Water framework directive, 2000/60/ec. european communities official journal l327
494 22.12.2000, p. 73. <http://eur-lex.europa.eu/legal-content/EN/TXT/?uri=celex:32000L0060>.
- 495 Williams SL, Heck KL. 2001. Seagrass community ecology. In: Bertness MD, Gaines SD, Hay
496 ME, editors, Marine Community Ecology. Sinauer Associates, Sunderland, Massachusetts.

Table 1: Characteristics of coastal segments used to evaluate seagrass depth of colonization estimates (see Fig. 1 for spatial distribution). Year is the date of the seagrass coverage and bathymetric data. Latitude and longitude are the geographic centers of each segment. Area and depth values are meters and square kilometers, respectively. Secchi measurements (m) were obtained from the Florida Department of Environmental Protection’s Impaired Waters Rule (IWR), update number 40. Secchi mean and standard errors are based on all observations within the ten years preceding each seagrass survey.^{tab:seg_summ}

	Big Bend	Choctawhatchee Bay	Old Tampa Bay	Upper Indian R. Lagoon
Year ^a	2006	2007	2010	2009
Latitude	29.61	30.43	27.94	28.61
Longitude	-83.48	-86.54	-82.62	-80.77
Surface area	271.37	59.41	205.50	228.52
Seagrass area	203.02	3.51	24.48	74.89
Depth (mean)	1.41	5.31	2.56	1.40
Depth (max)	3.60	11.90	10.40	3.70
Secchi (mean)	1.34	2.14	1.41	1.30
Secchi (se)	0.19	0.08	0.02	0.02

^a Seagrass coverage data sources, see section 2.1.2 for bathymetry data sources:

Big Bend: http://atoll.floridamarine.org/Data/metadata/SDE_Current/seagrass_bigbend_2006_poly.htm

Choctawhatchee Bay: http://atoll.floridamarine.org/data/metadata/SDE_Current/seagrass_chotawhatchee_2007_poly.htm

Tampa Bay: http://www.swfwmd.state.fl.us/data/gis/layer_library/category/swim

Indian R. Lagoon: <http://www.sjrwmd.com/gisdevelopment/docs/themes.html>

Table 2: Summary of seagrass depth estimates (m) for each segment using all grid locations in Fig. 4. Whole segment estimates were obtained from all seagrass depth data for each segment.^{tab:est_summ}

Segment ^a	Whole segment	Mean	St. Dev.	Min	Max
BB					
$Z_{c,min}$	1.25	1.33	0.82	0.00	2.64
$Z_{c,med}$	2.46	1.68	0.77	0.66	2.85
$Z_{c,max}$	3.66	2.03	0.80	0.86	3.31
CB					
$Z_{c,min}$	1.82	1.57	0.72	0.00	2.27
$Z_{c,med}$	2.16	1.98	0.46	1.19	2.48
$Z_{c,max}$	2.50	2.40	0.32	1.86	2.74
OTB					
$Z_{c,min}$	0.61	0.52	0.29	0.05	0.98
$Z_{c,med}$	0.88	0.85	0.27	0.30	1.24
$Z_{c,max}$	1.15	1.18	0.39	0.37	1.81
UIRL					
$Z_{c,min}$	1.25	1.32	0.23	1.00	2.02
$Z_{c,med}$	1.51	1.49	0.21	1.12	2.08
$Z_{c,max}$	1.77	1.66	0.21	1.23	2.14

^aBB: Big Bend, CB: Choctawhatchee Bay, OTB: Old Tampa Bay, UIRL: Upper Indian River Lagoon.

Table 3: Summary of uncertainty for seagrass depth estimates (m) for each segment using all grid locations in Fig. 5. The uncertainty values are equally applicable to each seagrass depth measure ($Z_{c,min}$, $Z_{c,med}$, $Z_{c,max}$).^{tab:sens_summ}

Segment ^a	Mean	St. Dev	Min	Max
BB	0.11	0.10	0.01	0.35
CB	0.72	0.74	0.22	2.52
OTB	0.36	0.28	0.11	1.04
UIRL	0.09	0.06	0.01	0.30

^aBB: Big Bend, CB: Choctawhatchee Bay, OTB: Old Tampa Bay, UIRL: Upper Indian River Lagoon.

Table 4: Summary of water clarity data (Z_{secchi}), depth of colonization ($Z_{c,max}$), and estimated light requirements for bay segments with available data for the Indian River Lagoon and Tampa Bay. Water clarity data were obtained from secchi observations in the Florida Impaired Waters Rule database for all available locations and dates within ten years of the seagrass survey in each bay. Values are minimum and maximum years of secchi data, sample size of secchi data ($n_{Z_{secchi}}$), mean values (m) of secchi data, sample size of seagrass depth estimates ($n_{Z_{c,max}}$) at each unique secchi location, mean $Z_{c,max}$, and estimated % light requirements for each segment. See Figs. 6 and 7 for spatial distribution of the results.^a

Segment ^a	Min year	Max year	$n_{Z_{secchi}}$	Z_{secchi}	$n_{Z_{c,max}}$	$Z_{c,max}$	% light
Indian River Lagoon							
BR	2000	2009	899	1.06	2	1.38	11.96
LCIRL	2000	2009	644	1.02	12	1.41	9.23
LIRL	2000	2005	111	0.93	6	1.84	4.06
LML	2000	2009	217	1.14	4	1.14	17.84
LSL	2000	2005	52	0.94	3	2.37	2.02
UCIRL	2000	2009	1148	1.14	18	1.19	10.84
UIRL	2000	2009	593	1.30	1	1.15	20.32
UML	2000	2009	258	1.03	4	1.21	19.08
Tampa Bay							
HB	2001	2003	412	1.25	10	1.36	15.32
LTB	2001	2009	807	2.47	22	2.14	22.60
MTB	2001	2009	570	2.19	14	1.64	28.03
OTB	2001	2003	671	1.44	15	1.18	24.05

^aBR: Banana R., LCIRL: Lower Central Indian R. Lagoon, LIRL: Lower Indian R. Lagoon, LML: Lower Mosquito Lagoon, LSL: Lower St. Lucie, UCIRL: Upper Central Indian R. Lagoon, UIRL: Upper Indian R. Lagoon, UML: Upper Mosquito Lagoon, HB: Hillsborough Bay, LTB: Lower Tampa Bay, MTB: Middle Tampa Bay, OTB: Old Tampa Bay.






Fig. 1: Locations and seagrass coverage of estuary segments used to evaluate depth of colonization estimates. Seagrass coverage layers are from 2007 (CB: Choctawhatchee Bay), 2006 (BB: Big Bend), 2010 (OTB: Old Tampa Bay), and 2009 (UIRL: Upper Indian R. Lagoon).

{fig:seg_a}


(a) Seagrass coverage and bathymetry for the segment



(b) Grid of locations and sample areas for estimates



(c) Sampled seagrass data for a test point



- Seagrass coverage
- 2 m depth contours
- - Estimated depth limit for segment


- △ Seagrass absent
- Seagrass present

- Estimation grid
- Test point
- Sample area


Fig. 2: Examples of data and grid locations for estimating seagrass depth of colonization for a region of the Big Bend, Florida. Fig. 2a shows the seagrass coverage and depth contours at 2 meter intervals, including the whole segment estimate for depth of colonization. Fig. 2b shows a grid of sampling locations with sampling radii for estimating Z_c and seagrass depth points derived from bathymetry and seagrass coverage layers. Fig. 2c shows an example of sampled seagrass depth points for a test location. Estimates in Fig. 3 were obtained from the test location in Fig. 2c.

{fig:buff_}

(a) Proportion of points with seagrass by depth



(b) Logistic growth curve fit through points



(c) Depth estimates




Fig. 3: Methods for estimating seagrass depth of colonization using sampled seagrass depth points around a single location. Fig. 3a is the proportion of points with seagrass by depth using depth points within the buffer of the test point in Fig. 2. Fig. 3b adds a decreasing logistic growth curve fit through the points. Fig. 3c shows three depth estimates based on a linear curve fit through the inflection point of logistic growth curve.

{fig:est_e}




Fig. 4: Spatially-resolved estimates of seagrass depth limits (m) for four coastal segments of Florida. Estimates include minimum ($Z_{c, \text{min}}$), median ($Z_{c, \text{med}}$), and maximum depth of colonization ($Z_{c, \text{max}}$). Estimates are assigned to grid locations for each segment, where grid spacing was fixed at 0.02 decimal degrees. Radii for sampling seagrass bathymetric data around each grid location were fixed at 0.06 decimal degrees. BB: Big Bend, CB: Choctawhatchee Bay, OTB: Old Tampa Bay, UIRL: Upper Indian R. Lagoon.

{fig:all_e}




Fig. 5: Size of confidence intervals (m) for depth of colonization estimates in Fig. 4. Points are colored and sized based on the difference between the upper and lower bounds of a 95% confidence interval for all three Z_c estimates ($Z_{c,min}$, $Z_{c,med}$, $Z_{c,max}$). Bounds were obtained using Monte Carlo simulations to estimate uncertainty associated with the inflection point of the estimated logistic curve (Fig. 3) for each sample. BB: Big Bend, CB: Choctawhatchee Bay, OTB: Old Tampa Bay, UIRL: Upper Indian R. Lagoon.

{fig:all_s}




Fig. 6: Estimated maximum depths of seagrass colonization and light requirements for multiple locations in Tampa Bay, Florida. Map locations are georeferenced observations of water clarity from secchi measurements in the Florida Impaired Waters Rule database, update 40. Estimates are also summarized by bay segment as boxplots where the dimensions are the 25th percentile, median, and 75th percentile. Whiskers extend beyond the boxes as 1.5 multiplied by the interquartile range. Light requirements are based on averaged secchi values within ten years of the seagrass coverage data and estimated maximum depth of colonization using a radius of 0.02 decimal degrees for each secchi location to sample seagrass depth points. HB: Hillsborough Bay, LTB: Lower Tampa Bay, MTB: Middle Tampa Bay, OTB: Old Tampa Bay.

{fig:light}




Fig. 7: Estimated maximum depths of seagrass colonization and light requirements for multiple locations in Indian River Lagoon, Florida. Map locations are georeferenced observations of water clarity in the Florida Impaired Waters Rule database, update 40. Estimates are also summarized by bay segment as boxplots as in Fig. 6. Light requirements are based on averaged secchi values within ten years of the seagrass coverage data and estimated maximum depth of colonization using a radius of 0.02 decimal degrees for each secchi location to sample seagrass depth points. BR: Banana R., LCIRL: Lower Central Indian R. Lagoon, LIRL: Lower Indian R. Lagoon, LML: Lower Mosquito Lagoon, LSL: Lower St. Lucie, UCIRL: Upper Central Indian R. Lagoon, UIRL: Upper Indian R. Lagoon, UML: Upper Mosquito Lagoon.

{fig:light}