

NERRS / SWMP

Training Workshop: *R*, *SWMP*r, *SWMP*rats

Williamsburg, VA, Nov 13, 2016

Time series topic 3: Seasonal Kendall Tests

Marcus W. Beck¹

¹USEPA NHEERL Gulf Ecology Division
Email: beck.marcus@epa.gov

Objectives for the session (4:15 - 5:00)

- What is and why would we use a Kendall test
- What is a why would we use a *Seasonal* Kendall test
- Application with NERRS data
 - ▶ Data prep
 - ▶ Execution
 - ▶ Interpretation

Interactive portion

Follow along as we go:

- flash drive
- online: swmprats.net 2016 workshop tab

Interactive portion

Follow along as we go:

- flash drive
- online: swmprats.net 2016 workshop tab

You will run examples whenever you see this guy:



Is everything installed?

We will use functions in the EnvStats package

Option 1, from the R Console prompt:

```
install.packages('EnvStats')  
library(EnvStats)
```

Is everything installed?

We will use functions in the EnvStats package

Option 1, from the R Console prompt:

```
install.packages('EnvStats')  
library(EnvStats)
```

Option 2, install the source file from the flash drive:

```
# change as needed  
path_to_file <- 'C:/Users/mbeck/Desktop/EnvStats_2.1.1.tar.gz'  
  
# install, load  
install.packages(path_to_file, repos = NULL, type="source")  
library(SWMPPr)
```

Theory and background

Use these tests to answer the question:

Is there a *monotonic trend* and what is the *significance*?

Theory and background

Use these tests to answer the question:

Is there a *monotonic trend* and what is the *significance*?

Both tests:

- Are non-parametric (no distribution assumptions, based on signs)

Theory and background

Use these tests to answer the question:

Is there a *monotonic trend* and what is the *significance*?

Both tests:

- Are non-parametric (no distribution assumptions, based on signs)
- Work with a user-defined time interval

Theory and background

Use these tests to answer the question:

Is there a *monotonic trend* and what is the *significance*?

Both tests:

- Are non-parametric (no distribution assumptions, based on signs)
- Work with a user-defined time interval
- Provide a p-value indicating the probability due to chance alone

Theory and background

Use these tests to answer the question:

Is there a *monotonic trend* and what is the *significance*?

Both tests:

- Are non-parametric (no distribution assumptions, based on signs)
- Work with a user-defined time interval
- Provide a p-value indicating the probability due to chance alone
- Provide a direction of the trend as τ ('tau')

Theory and background

Use these tests to answer the question:

Is there a *monotonic trend* and what is the *significance*?

Both tests:

- Are non-parametric (no distribution assumptions, based on signs)
- Work with a user-defined time interval
- Provide a p-value indicating the probability due to chance alone
- Provide a direction of the trend as τ ('tau')
- Provide a slope as the rate of change

Theory and background

The ***Kendall test*** for time series:

$$\tau = \sum_{i=1}^{n-1} \sum_{j=i+1}^n \text{sign}[(X_j - X_i)(Y_j - Y_i)]$$

τ will vary from -1 to 1 similar to a correlation coefficient, follows an approximate normal-distribution for hypothesis-testing

Theory and background

The ***Kendall test*** for time series:

$$\tau = \sum_{i=1}^{n-1} \sum_{j=i+1}^n \text{sign}[(X_j - X_i)(Y_j - Y_i)]$$

τ will vary from -1 to 1 similar to a correlation coefficient, follows an approximate normal-distribution for hypothesis-testing

$$\hat{\beta}_1 = \text{Median} \left(\frac{Y_j - Y_i}{X_j - X_i} \right), i < j$$

$\hat{\beta}_1$ is the Theil (Sen) non-parametric estimate of slope or the rate of change in the interval

Theory and background

The *Seasonal Kendall test* is exactly the same...

...except separate tests by month across years (January 1981, 1982, ..., February 1981, 1982, ...), results are pooled.

Theory and background

The *Seasonal Kendall test* is exactly the same...

...except separate tests by month across years (January 1981, 1982, ..., February 1981, 1982, ...), results are pooled.

- Overall τ is the weighted average of the seasonal estimates

Theory and background

The *Seasonal Kendall test* is exactly the same...

...except separate tests by month across years (January 1981, 1982, ..., February 1981, 1982, ...), results are pooled.

- Overall τ is the weighted average of the seasonal estimates
- Overall $\hat{\beta}_1$ is the median of all two-point slope estimates within each season

Theory and background

The *Seasonal Kendall test* is exactly the same...

...except separate tests by month across years (January 1981, 1982, ..., February 1981, 1982, ...), results are pooled.

- Overall τ is the weighted average of the seasonal estimates
- Overall $\hat{\beta}_1$ is the median of all two-point slope estimates within each season

More info in help documentation for `kendallTrendTest`,
`kendallSeasonalTrendTest` in `EnvStats`,
[Hirsch et al., 1982, Millard, 2013]

Using decomp with NERRS data

Load some water quality data

Summary

Things to ask before Seasonal Kendall

- test

What does Seasonal Kendall not tell us

NERRS / SWMP

Training Workshop: *R*, *SWMP*_r, *SWMP*_r*ats*

Williamsburg, VA, Nov 13, 2016

Up next... nothing!

Questions??

References

Hirsch RM, Slack JR, Smith RA. 1982.
Techniques of trend analysis for monthly water quality data.
Water Resources Research, 18:107–121.

Millard SP. 2013.
EnvStats: An R Package for Environmental Statistics.
Springer, New York.