

*Tools for primal degenerate linear programs*

+

*Dual-guided pivot rules for LP*

+

*Vector Space Decomposition for LP*

J. Desrosiers<sup>1</sup>

J.B. Gauthier<sup>1</sup>

M.E. Lübbecke<sup>2</sup>

---

(<sup>1</sup>) HEC Montréal and GERAD

(<sup>2</sup>) RWTH Aachen University

**School on Column Generation, Paris**

March 10-14, 2014

- PS : **Primal Simplex** and degenerate solutions  
perturbation, pivot rules, etc. (Terlaky and Zhang 1993)

- PS : **Primal Simplex** and degenerate solutions  
perturbation, pivot rules, etc. (Terlaky and Zhang 1993)
- CG : **Column Generation** for *huge* problems  $\text{CG} \equiv \text{PS}$   
perturbation of the master problem (1985-2000)

- PS : **Primal Simplex** and degenerate solutions  
perturbation, pivot rules, etc. (Terlaky and Zhang 1993)
- CG : **Column Generation** for *huge* problems  $CG \equiv PS$   
perturbation of the master problem (1985-2000)
- DVS : **Dual Variable Stabilization** (du Merle et al. 1999)  
Oukil et al (2007) : cpu time reduction by factors of 10 to 100

- PS : **Primal Simplex** and degenerate solutions  
perturbation, pivot rules, etc. (Terlaky and Zhang 1993)
- CG : **Column Generation** for *huge* problems  $CG \equiv PS$   
perturbation of the master problem (1985-2000)
- DVS : **Dual Variable Stabilization** (du Merle et al. 1999)  
Oukil et al (2007) : cpu time reduction by factors of 10 to 100
- DCA : **Dynamic Constraints Aggregation** for Set Partitioning Problems  
(Elhallaoui, Metrane, Desaulniers, Soumis 2005-08)  
cpu time reduction by factors of 10 to 100

- PS : **Primal Simplex** and degenerate solutions  
perturbation, pivot rules, etc. (Terlaky and Zhang 1993)
- CG : **Column Generation** for *huge* problems  $CG \equiv PS$   
perturbation of the master problem (1985-2000)
- DVS : **Dual Variable Stabilization** (du Merle et al. 1999)  
Oukil et al (2007) : cpu time reduction by factors of 10 to 100
- DCA : **Dynamic Constraints Aggregation** for Set Partitioning Problems  
(Elhallaoui, Metrane, Desaulniers, Soumis 2005-08)  
cpu time reduction by factors of 10 to 100
- IPS : **Improved Primal Simplex** (Vincent, Soumis, Metrane, Orban 2008-11)  
generalizes DCA to LP ; non-degenerate pivots ; *positive edge rule*

- PS : **Primal Simplex** and degenerate solutions  
perturbation, pivot rules, etc. (Terlaky and Zhang 1993)
- CG : **Column Generation** for *huge* problems  $CG \equiv PS$   
perturbation of the master problem (1985-2000)
- DVS : **Dual Variable Stabilization** (du Merle et al. 1999)  
Oukil et al (2007) : cpu time reduction by factors of 10 to 100
- DCA : **Dynamic Constraints Aggregation** for Set Partitioning Problems  
(Elhallaoui, Metrane, Desaulniers, Soumis 2005-08)  
cpu time reduction by factors of 10 to 100
- IPS : **Improved Primal Simplex** (Vincent, Soumis, Metrane, Orban 2008-11)  
generalizes DCA to LP ; non-degenerate pivots ; *positive edge rule*
- MMCC : **minimum mean cost cycle canceling** algorithm for networks  
70%–90% of the PS pivots are degenerate ; MMCC strongly polynomial.

- PS : **Primal Simplex** and degenerate solutions  
perturbation, pivot rules, etc. (Terlaky and Zhang 1993)
- CG : **Column Generation** for *huge* problems  $CG \equiv PS$   
perturbation of the master problem (1985-2000)
- DVS : **Dual Variable Stabilization** (du Merle et al. 1999)  
Oukil et al (2007) : cpu time reduction by factors of 10 to 100
- DCA : **Dynamic Constraints Aggregation** for Set Partitioning Problems  
(Elhallaoui, Metrane, Desaulniers, Soumis 2005-08)  
cpu time reduction by factors of 10 to 100
- IPS : **Improved Primal Simplex** (Vincent, Soumis, Metrane, Orban 2008-11)  
generalizes DCA to LP ; non-degenerate pivots ; *positive edge rule*
- MMCC : **minimum mean cost cycle canceling** algorithm for networks  
70%–90% of the PS pivots are degenerate ; MMCC strongly polynomial.

What are the links between PS, DCA, IPS, and MMCC ?




## LP IN STANDARD FORM

$$\min \quad \mathbf{c}^T \mathbf{x}$$

$$\mathbf{Ax} = \mathbf{b} \quad [\pi]$$

$$\mathbf{x} \geq \mathbf{0}$$

FROM  $\mathbf{x}^0$  TO  $\mathbf{x}^1$ 

- ① Find a *potential* improving direction  $\mathbf{y}^0 \in \mathbb{R}^n$ .
- ② Determine step-size  $\rho^0 \in \mathbb{R}$ .
- ③ Compute  $\mathbf{x}^1 := \mathbf{x}^0 + \rho^0 \mathbf{y}^0$ .

$\bar{c}_j = 0, \forall j \in B$  : PRICING FOR  $j \in N$  (NON-BASIC VARIABLES)

Selection of an entering variable into basis  $\mathbf{A}_B$  relies on the minimum reduced cost of non-basic variables

$$\boldsymbol{\pi}^\top = \mathbf{c}_B^\top \mathbf{A}_B^{-1} \quad \bar{c}_j = c_j - \boldsymbol{\pi}^\top \mathbf{a}_j, \quad \forall j \in N.$$

$\bar{c}_j = 0, \forall j \in B$  : PRICING FOR  $j \in N$  (NON-BASIC VARIABLES)

Selection of an entering variable into basis  $\mathbf{A}_B$  relies on the minimum reduced cost of non-basic variables

$$\boldsymbol{\pi}^\top = \mathbf{c}_B^\top \mathbf{A}_B^{-1} \quad \bar{c}_j = c_j - \boldsymbol{\pi}^\top \mathbf{a}_j, \quad \forall j \in N.$$

FIND THE MINIMUM REDUCED COST VALUE  $\mu$  (OPTIMAL IF  $\mu \geq 0$ )

max  $\mu$

$$\mu \leq c_j - \boldsymbol{\pi}^\top \mathbf{a}_j, \quad \forall j \in N \quad [y_j]$$

\*\*\*  $\mu$  is the smallest reduced cost (given  $\boldsymbol{\pi}$ ). \*\*\*

$\bar{c}_j = 0, \forall j \in B$  : PRICING FOR  $j \in N$  (NON-BASIC VARIABLES)

Selection of an entering variable into basis  $\mathbf{A}_B$  relies on the minimum reduced cost of non-basic variables

$$\boldsymbol{\pi}^\top = \mathbf{c}_B^\top \mathbf{A}_B^{-1} \quad \bar{c}_j = c_j - \boldsymbol{\pi}^\top \mathbf{a}_j, \quad \forall j \in N.$$

FIND THE MINIMUM REDUCED COST VALUE  $\mu$  (OPTIMAL IF  $\mu \geq 0$ )

$$\max \quad \mu$$

$$\mu \leq c_j - \boldsymbol{\pi}^\top \mathbf{a}_j, \quad \forall j \in N \quad [y_j]$$

\*\*\*  $\mu$  is the smallest reduced cost (given  $\boldsymbol{\pi}$ ). \*\*\*

EQUIVALENT TO FINDING A CONVEX COMBINATION OF NON-BASIC VARIABLES

$$\begin{aligned} \mu = \min \quad & \sum_{j \in N} \bar{c}_j y_j \\ & \sum_{j \in N} y_j = 1 \quad [\mu] \\ & y_j \geq 0, \quad \forall j \in N \end{aligned}$$

$\mathbf{y}^0$

Entering variable only impacts the current basic variables, at most  $m$  variables.

All but one non-basic variables remain at zero.

$\mathbf{y}^0 \in \mathbb{R}^n$  makes changes on at most  $m + 1$  components.

$\mathbf{y}^0$

Entering variable only impacts the current basic variables, at most  $m$  variables.

All but one non-basic variables remain at zero.

$\mathbf{y}^0 \in \mathbb{R}^n$  makes changes on at most  $m + 1$  components.

Direction  $\mathbf{y}^0 \in \mathbb{R}^n$  :  
the selected entering variable,  
the non-selected non-basic variables,  
and the basic ones.

Step size computed such that  $\mathbf{x}^1 := \mathbf{x}^0 + \rho^0 \mathbf{y}^0 \geq \mathbf{0}$ .



LEFT MULTIPLICATION BY BASIS INVERSE  $A_B^{-1}$ 


	$x_1$	$x_2$	$x_3$	$\lambda_4$	$\lambda_5$	$\lambda_6$	$x_4$	$x_5$	$x_6$	$x_7$	$x_8$		$\theta_w$
$c$	2	3	1				10	17	-20	14	-4		-5
	1						2	2	1	-5	7	=	30
		1					4	3	-5	10	-10	=	25
			1				-3	1	2	3	11	=	50
				1					6	5	-13	=	0
					1				3	4	-8	=	0
						1			3	-4	0	=	0
$x^0$	30	25	50	0	0	0	0	0	0	0	0	$185 = c^T x^0$	
							0.4	0.3	0.3				

$\psi$	
2	1
3	-2
1	5
?	0
?	0
?	0




## PRIMAL SIMPLEX TABLEAU




\*\*\* Changes on at most  $m + 1$  components. \*\*\*

## POSITIVE EDGE STRATEGY



\*\*\* Changes on at most  $p + 1$  components. \*\*\*  
Non-degenerate pivot.

SPEEDUP OF 2.31 FOR LPs WITH A DEGENERACY LEVEL ABOVE 25%



The *Positive Edge* is a pricing rule for the Primal Simplex :  
it identifies variables  $x_j, j \notin B$  such that  $\bar{a}_{zj} = \mathbf{0} \dots$   
*without computing  $\bar{a}_{zj}$ .*

The *Positive Edge* is a pricing rule for the Primal Simplex :  
 it identifies variables  $x_j, j \notin B$  such that  $\bar{a}_{zj} = 0 \dots$   
*without computing  $\bar{a}_{zj}$ .*

If such a variable  $x_j$  has a negative reduced cost  $\bar{c}_j < 0$ ,  
 it strictly improves the objective function when entered into the basis.

Let  $\mathbf{v} \neq \mathbf{0}$  be a random vector of  $|Z|$  non-zero components.

Let  $\mathbf{v} \neq \mathbf{0}$  be a random vector of  $|Z|$  non-zero components.

$$\mathbf{v}^T \bar{\mathbf{a}}_{Zj} = 0 \quad \Longleftrightarrow \quad \bar{\mathbf{a}}_{Zj} = \mathbf{0} \quad \text{or} \quad \mathbf{v} \perp \bar{\mathbf{a}}_{Zj}$$

Let  $\mathbf{v} \neq \mathbf{0}$  be a random vector of  $|Z|$  non-zero components.

$$\mathbf{v}^\top \bar{\mathbf{a}}_{Zj} = 0 \iff \bar{\mathbf{a}}_{Zj} = \mathbf{0} \quad \text{or} \quad \mathbf{v} \perp \bar{\mathbf{a}}_{Zj}$$

$$\mathbf{v}^\top \bar{\mathbf{a}}_{Zj} = \mathbf{v}^\top [\mathbf{A}_B^{-1}]_Z \mathbf{a}_{Zj} = \mathbf{v}^\top [\mathbf{A}_B^{-1}]_Z \mathbf{a}_{Zj}$$

Pre-compute once  $\mathbf{w}^\top := \mathbf{v}^\top [\mathbf{A}_B^{-1}]_Z$ , then compute  $\mathbf{w}^\top \mathbf{a}_{Zj}$ , on the **original data**.



Let  $\mathbf{v} \neq \mathbf{0}$  be a random vector of  $|Z|$  non-zero components.

$$\mathbf{v}^\top \bar{\mathbf{a}}_{Zj} = 0 \iff \bar{\mathbf{a}}_{Zj} = \mathbf{0} \quad \text{or} \quad \mathbf{v} \perp \bar{\mathbf{a}}_{Zj}$$

$$\mathbf{v}^\top \bar{\mathbf{a}}_{Zj} = \mathbf{v}^\top [\mathbf{A}_B^{-1}]_Z \mathbf{a}_{Zj} = \mathbf{v}^\top [\mathbf{A}_B^{-1}]_Z \mathbf{a}_{Zj}$$

Pre-compute once  $\mathbf{w}^\top := \mathbf{v}^\top [\mathbf{A}_B^{-1}]_Z$ , then compute  $\mathbf{w}^\top \mathbf{a}_{Zj}$ , on the **original data**.

Determining if  $x_j$  satisfies the Positive Edge rule is done in  $O(m)$ ,  
as for  $\bar{c}_j = c_j - \pi^\top \mathbf{a}_j, j \notin B$ , where  $\pi = \mathbf{c}_B^\top \mathbf{A}_B^{-1}$ .

Let  $\mathbf{v} \neq \mathbf{0}$  be a random vector of  $|Z|$  non-zero components.

$$\mathbf{v}^\top \bar{\mathbf{a}}_{Zj} = 0 \iff \bar{\mathbf{a}}_{Zj} = \mathbf{0} \quad \text{or} \quad \mathbf{v} \perp \bar{\mathbf{a}}_{Zj}$$

$$\mathbf{v}^\top \bar{\mathbf{a}}_{Zj} = \mathbf{v}^\top [\mathbf{A}_B^{-1}]_Z \mathbf{a}_{Zj} = \mathbf{v}^\top [\mathbf{A}_B^{-1}]_Z \mathbf{a}_{Zj}$$

Pre-compute once  $\mathbf{w}^\top := \mathbf{v}^\top [\mathbf{A}_B^{-1}]_Z$ , then compute  $\mathbf{w}^\top \mathbf{a}_{Zj}$ , on the **original data**.

Determining if  $x_j$  satisfies the Positive Edge rule is done in  $O(m)$ ,  
as for  $\bar{c}_j = c_j - \pi^\top \mathbf{a}_j, j \notin B$ , where  $\pi = \mathbf{c}_B^\top \mathbf{A}_B^{-1}$ .

On large-scale instances ( $m \approx 100\,000$ ,  $n \approx 450\,000$ ),  
cpu time to identify all variables satisfying the PE rule :

usual definition

$$O(m^2 n)$$

Positive Edge

$$O(mn)$$

Let  $\mathbf{v} \neq \mathbf{0}$  be a random vector of  $|Z|$  non-zero components.

$$\mathbf{v}^\top \bar{\mathbf{a}}_{Zj} = 0 \iff \bar{\mathbf{a}}_{Zj} = \mathbf{0} \quad \text{or} \quad \mathbf{v} \perp \bar{\mathbf{a}}_{Zj}$$

$$\mathbf{v}^\top \bar{\mathbf{a}}_{Zj} = \mathbf{v}^\top [\mathbf{A}_B^{-1}]_Z \mathbf{a}_{Zj} = \mathbf{v}^\top [\mathbf{A}_B^{-1}]_Z \mathbf{a}_{Zj}$$

Pre-compute once  $\mathbf{w}^\top := \mathbf{v}^\top [\mathbf{A}_B^{-1}]_Z$ , then compute  $\mathbf{w}^\top \mathbf{a}_{Zj}$ , on the **original data**.

Determining if  $x_j$  satisfies the Positive Edge rule is done in  $O(m)$ ,  
as for  $\bar{c}_j = c_j - \pi^\top \mathbf{a}_j, j \notin B$ , where  $\pi = \mathbf{c}_B^\top \mathbf{A}_B^{-1}$ .

On large-scale instances ( $m \approx 100\,000$ ,  $n \approx 450\,000$ ),  
cpu time to identify all variables satisfying the PE rule :

usual definition

Positive Edge

$$O(m^2 n)$$

$$O(mn)$$

**2500 seconds**

Let  $\mathbf{v} \neq \mathbf{0}$  be a random vector of  $|Z|$  non-zero components.

$$\mathbf{v}^\top \bar{\mathbf{a}}_{Zj} = 0 \iff \bar{\mathbf{a}}_{Zj} = \mathbf{0} \quad \text{or} \quad \mathbf{v} \perp \bar{\mathbf{a}}_{Zj}$$

$$\mathbf{v}^\top \bar{\mathbf{a}}_{Zj} = \mathbf{v}^\top [\mathbf{A}_B^{-1}]_Z \mathbf{a}_{Zj} = \mathbf{v}^\top [\mathbf{A}_B^{-1}]_Z \mathbf{a}_{Zj}$$

Pre-compute once  $\mathbf{w}^\top := \mathbf{v}^\top [\mathbf{A}_B^{-1}]_Z$ , then compute  $\mathbf{w}^\top \mathbf{a}_{Zj}$ , on the **original data**.

Determining if  $x_j$  satisfies the Positive Edge rule is done in  $O(m)$ ,  
as for  $\bar{c}_j = c_j - \pi^\top \mathbf{a}_j, j \notin B$ , where  $\pi = \mathbf{c}_B^\top \mathbf{A}_B^{-1}$ .

On large-scale instances ( $m \approx 100\,000$ ,  $n \approx 450\,000$ ),  
cpu time to identify all variables satisfying the PE rule :

usual definition

Positive Edge


$$O(m^2 n)$$

$$O(mn)$$

**2500 seconds**

**0.5 seconds**

BLOCK DIAGONAL : KEEP DEGENERATE VARIABLES IN FIRST STEP (IPS)



\*\*\* Changes on at most  $m + 1$  components. \*\*\*

Vectors and matrices are written in **bold face**.

$\mathbf{I}_\ell$  : the  $\ell \times \ell$  identity matrix.

$\mathbf{0}$  ( $\mathbf{1}$ ) : a vector/matrix with all zeros (ones) entries of appropriate dimensions.

$\mathbf{A}_{RC}$  : sub-matrix of  $\mathbf{A}$  containing the rows and columns indexed by  $R$  and  $C$ .

Standard notation : basis  $\mathbf{A}_B$ , inverse  $\mathbf{A}_B^{-1}$ ,  $\mathbf{c}_B^T \mathbf{x}_B$ ,  $\mathbf{A}_B \mathbf{x}_B$ ,  $\boldsymbol{\pi} = \mathbf{c}_B \mathbf{A}_B^{-1} \dots$

$\mathbf{l}_F < \mathbf{x}_F < \mathbf{u}_F$ ,  $\mathbf{x}_L = \mathbf{l}_L$ ,  $\mathbf{x}_U = \mathbf{u}_U$

USEFUL DECOMPOSITION OF  $\mathbf{x} \in \mathbb{R}^n$  IN  $\mathbf{Ax} = \mathbf{b}$ ,  $\mathbf{l} \leq \mathbf{x} \leq \mathbf{u}$

$$\mathbf{x} = \begin{bmatrix} \mathbf{x}_F \\ \mathbf{x}_L \\ \mathbf{x}_U \end{bmatrix} \quad \mathbf{x} = \begin{bmatrix} \mathbf{x}_B \\ \mathbf{x}_N \end{bmatrix} = \begin{bmatrix} \mathbf{x}_F \\ \mathbf{x}_{B_L} \\ \mathbf{x}_{B_U} \\ \mathbf{x}_{N_L} \\ \mathbf{x}_{N_U} \end{bmatrix}$$

$$\text{For } \emptyset \subseteq S \subseteq B : \mathbf{x} = \begin{bmatrix} \mathbf{x}_S \\ \mathbf{x}_{\bar{S}} \end{bmatrix} = \begin{bmatrix} \mathbf{x}_{S_F} \\ \mathbf{x}_{S_L} \\ \mathbf{x}_{S_F} \\ \mathbf{x}_{\bar{S}_U} \\ \mathbf{x}_{\bar{S}_L} \\ \mathbf{x}_{\bar{S}_U} \end{bmatrix} \quad \mathbf{x} = \begin{bmatrix} \mathbf{x}_\emptyset \\ \mathbf{x} \end{bmatrix} = \begin{bmatrix} \mathbf{x}_F \\ \mathbf{x}_L \\ \mathbf{x}_U \end{bmatrix}$$

LINEAR PROGRAM  $LP$

$$\begin{array}{ll} z^* := & \min \quad \mathbf{c}^T \mathbf{x} \\ \text{st.} & \mathbf{Ax} = \mathbf{b} \\ & \mathbf{l} \leq \mathbf{x} \leq \mathbf{u} \end{array}$$

## LINEAR PROGRAM $LP$

$$\begin{aligned} z^* := & \min \quad \mathbf{c}^T \mathbf{x} \\ \text{st.} \quad & \mathbf{Ax} = \mathbf{b} \\ & \mathbf{l} \leq \mathbf{x} \leq \mathbf{u} \end{aligned}$$

## GENERIC ALGORITHM WITH SINGLE PARAMETER SET $S$ , $\emptyset \subseteq S \subseteq B$

- 1 Let  $k = 0$  and assume a feasible basic solution  $\mathbf{x}^k$  to LP.
- 2 For  $\emptyset \subseteq S \subseteq B$ , construct the residual problem  $LP_S(\mathbf{x}^k)$ .
- 3 Fix a subset of the dual variables in row-set  $R$ :  $\boldsymbol{\pi}_R^T = \mathbf{c}_S^T \mathbf{A}_{RS}^{-1}$ .
- 4 Determine the value of the smallest reduced cost  $\mu_S^k$ .  
If  $\mu_S^k \geq 0$ , STOP. Current solution  $\mathbf{x}^k$  is optimal for LP.
- 5 Retrieve direction  $\mathbf{y}_S^k \in \mathbb{R}^n$  and compute its maximum step-size  $\rho_S^k$ .
- 6 Update  $\begin{aligned} \mathbf{x}^{k+1} &:= \mathbf{x}^k + \rho_S^k \mathbf{y}_S^k; \\ z^{k+1} &:= z^k + \rho_S^k \mu_S^k; \\ k &:= k + 1. \end{aligned}$
- 7 Goto Step 2.



ASSUME A FEASIBLE SOLUTION  $\mathbf{x}^k = \begin{bmatrix} \mathbf{x}_F^k \\ \mathbf{x}_L^k \\ \mathbf{x}_U^k \end{bmatrix}$

$$\mathbf{x} := \mathbf{x}^k + \mathbf{y}, \quad \mathbf{y} \in \mathbb{R}^n$$

ASSUME A FEASIBLE SOLUTION  $\mathbf{x}^k = \begin{bmatrix} \mathbf{x}_F^k \\ \mathbf{x}_L^k \\ \mathbf{x}_U^k \end{bmatrix}$

$$\mathbf{x} := \mathbf{x}^k + \mathbf{y}, \quad \mathbf{y} \in \mathbb{R}^n$$

$$= \mathbf{x}^k + (\vec{\mathbf{y}} - \tilde{\mathbf{y}}),$$

$$\vec{\mathbf{y}}, \tilde{\mathbf{y}} \geq \mathbf{0}, \quad \vec{\mathbf{y}}^\top \tilde{\mathbf{y}} = 0, \quad \vec{\mathbf{y}} \leq \vec{\mathbf{r}}^k, \quad \tilde{\mathbf{y}} \leq \tilde{\mathbf{r}}^k$$


# STEP 2 CONSTRUCT RESIDUAL PROBLEM LP( $\mathbf{x}^k$ )

ASSUME A FEASIBLE SOLUTION  $\mathbf{x}^k = \begin{bmatrix} \mathbf{x}_F^k \\ \mathbf{x}_L^k \\ \mathbf{x}_U^k \end{bmatrix}$

$$\mathbf{x} := \mathbf{x}^k + \mathbf{y}, \quad \mathbf{y} \in \mathbb{R}^n$$

$$= \mathbf{x}^k + (\vec{\mathbf{y}} - \bar{\mathbf{y}}), \quad \vec{\mathbf{y}}, \bar{\mathbf{y}} \geq \mathbf{0}, \quad \vec{\mathbf{y}}^\top \bar{\mathbf{y}} = 0, \quad \vec{\mathbf{y}} \leq \vec{\mathbf{r}}^k, \quad \bar{\mathbf{y}} \leq \bar{\mathbf{r}}^k$$

$$= \mathbf{x}^k + \left( \begin{bmatrix} \vec{\mathbf{y}}_F \\ \vec{\mathbf{y}}_L \\ \vec{\mathbf{y}}_U \end{bmatrix} - \begin{bmatrix} \bar{\mathbf{y}}_F \\ \bar{\mathbf{y}}_L \\ \bar{\mathbf{y}}_U \end{bmatrix} \right); \quad \vec{\mathbf{y}}, \bar{\mathbf{y}} \geq \mathbf{0}, \quad \vec{\mathbf{y}}^\top \bar{\mathbf{y}} = 0, \quad \vec{\mathbf{y}} \leq \vec{\mathbf{r}}^k, \quad \bar{\mathbf{y}} \leq \bar{\mathbf{r}}^k$$



RESIDUAL PROBLEM  $LP(\mathbf{x}^k)$ ;  $\mathbf{x} := \mathbf{x}^k + (\vec{\mathbf{y}} - \tilde{\mathbf{y}})$  (CHANGE OF VARIABLES)

$$\begin{aligned}
 z^* := \min \quad & \mathbf{c}^\top \mathbf{x}^k + \mathbf{c}^\top (\vec{\mathbf{y}} - \tilde{\mathbf{y}}) \\
 \text{st.} \quad & \mathbf{A} \mathbf{x}^k + \mathbf{A}(\vec{\mathbf{y}} - \tilde{\mathbf{y}}) = \mathbf{b} \\
 & \mathbf{0} \leq \vec{\mathbf{y}} \leq \vec{\mathbf{r}}^k \\
 & \mathbf{0} \leq \tilde{\mathbf{y}} \leq \tilde{\mathbf{r}}^k
 \end{aligned}$$

## STEP 2 CONSTRUCT RESIDUAL PROBLEM $LP(\mathbf{x}^k)$

RESIDUAL PROBLEM  $LP(\mathbf{x}^k)$ ;  $\mathbf{x} := \mathbf{x}^k + (\vec{\mathbf{y}} - \tilde{\mathbf{y}})$  (CHANGE OF VARIABLES)

$$\begin{aligned} z^* := \min \quad & \mathbf{c}^\top \mathbf{x}^k + \mathbf{c}^\top (\vec{\mathbf{y}} - \tilde{\mathbf{y}}) \\ \text{st.} \quad & \mathbf{A} \mathbf{x}^k + \mathbf{A}(\vec{\mathbf{y}} - \tilde{\mathbf{y}}) = \mathbf{b} \\ & \mathbf{0} \leq \vec{\mathbf{y}} \leq \vec{\mathbf{r}}^k \\ & \mathbf{0} \leq \tilde{\mathbf{y}} \leq \tilde{\mathbf{r}}^k \end{aligned}$$

RESIDUAL PROBLEM  $LP(\mathbf{x}^k)$

$$\begin{aligned} z^* := \mathbf{c}^\top \mathbf{x}^k + \min \quad & \mathbf{c}^\top (\vec{\mathbf{y}} - \tilde{\mathbf{y}}) \\ \text{st.} \quad & \mathbf{A}(\vec{\mathbf{y}} - \tilde{\mathbf{y}}) = \mathbf{0} \\ & \mathbf{0} \leq \vec{\mathbf{y}} \leq \vec{\mathbf{r}}^k \\ & \mathbf{0} \leq \tilde{\mathbf{y}} \leq \tilde{\mathbf{r}}^k \end{aligned}$$

RESIDUAL PROBLEM WITH  $S = F$ ,  $\bar{S} = L \cup U$

$$z^* := \mathbf{c}^T \mathbf{x}^k +$$

$$\begin{array}{llllll} \min & \mathbf{c}_F^T(\vec{\mathbf{y}}_F - \tilde{\mathbf{y}}_F) & + & \mathbf{c}_L^T(\vec{\mathbf{y}}_L) & - & \mathbf{c}_U^T(\tilde{\mathbf{y}}_U) \\ \text{st.} & \mathbf{A}_F(\vec{\mathbf{y}}_F - \tilde{\mathbf{y}}_F) & + & \mathbf{A}_L(\vec{\mathbf{y}}_L) & - & \mathbf{A}_U(\tilde{\mathbf{y}}_U) = \mathbf{0} \\ & \vec{\mathbf{y}}_F \geq \mathbf{0}, \tilde{\mathbf{y}}_F \geq \mathbf{0}, & & \vec{\mathbf{y}}_L \geq \mathbf{0}, & & \tilde{\mathbf{y}}_U \geq \mathbf{0} \\ & \vec{\mathbf{y}}_F \leq \vec{\mathbf{r}}_F, \tilde{\mathbf{y}}_F \leq \tilde{\mathbf{r}}_F, & & \vec{\mathbf{y}}_L \leq \vec{\mathbf{r}}_L, & & \tilde{\mathbf{y}}_U \leq \tilde{\mathbf{r}}_U \end{array}$$

RESIDUAL PROBLEM – GENERAL CASE :  $\emptyset \subseteq S \subseteq B$

$$\begin{array}{ll}
 \mathbf{c}^\top \mathbf{x}^k + & \min \quad \mathbf{c}_S^\top (\vec{\mathbf{y}}_S - \tilde{\mathbf{y}}_S) + \mathbf{c}_{\bar{S}}^\top (\vec{\mathbf{y}}_{\bar{S}} - \tilde{\mathbf{y}}_{\bar{S}}) \\
 \text{st.} & \mathbf{A}_S (\vec{\mathbf{y}}_S - \tilde{\mathbf{y}}_S) + \mathbf{A}_{\bar{S}} (\vec{\mathbf{y}}_{\bar{S}} - \tilde{\mathbf{y}}_{\bar{S}}) = \mathbf{0} \\
 & \vec{\mathbf{y}}_S \geq \mathbf{0}, \tilde{\mathbf{y}}_S \geq \mathbf{0} \qquad \vec{\mathbf{y}}_{\bar{S}} \geq \mathbf{0}, \tilde{\mathbf{y}}_{\bar{S}} \geq \mathbf{0} \\
 & \qquad \qquad \qquad \vec{\mathbf{y}}_{\bar{S}_L} \leq \mathbf{0}, \vec{\mathbf{y}}_{\bar{S}_U} \leq \mathbf{0} \\
 & \vec{\mathbf{y}}_{S_F} \leq \vec{\mathbf{r}}_{S_F}, \tilde{\mathbf{y}}_{S_F} \leq \vec{\mathbf{r}}_{S_F} \qquad \vec{\mathbf{y}}_{\bar{S}_F} \leq \vec{\mathbf{r}}_{\bar{S}_F}, \tilde{\mathbf{y}}_{\bar{S}_F} \leq \vec{\mathbf{r}}_{\bar{S}_F} \\
 & \vec{\mathbf{y}}_{S_L} \leq \mathbf{0}, \vec{\mathbf{y}}_{S_U} \leq \mathbf{0} \qquad \vec{\mathbf{y}}_{\bar{S}_L} \leq \vec{\mathbf{r}}_{\bar{S}_L}, \tilde{\mathbf{y}}_{\bar{S}_U} \leq \vec{\mathbf{r}}_{\bar{S}_U}
 \end{array}$$

$$S = S_F \cup S_L \cup S_U; \quad \bar{S} = \bar{S}_F \cup \bar{S}_L \cup \bar{S}_U$$

RESIDUAL PROBLEM WITH  $S = B$  (BASIC),  $\bar{S} = N$  (NON-BASIC)

$$\begin{aligned}
 & \mathbf{c}^T \mathbf{x}^k + \min \quad \mathbf{c}_B^T (\vec{\mathbf{y}}_B - \tilde{\mathbf{y}}_B) + \mathbf{c}_N^T (\vec{\mathbf{y}}_N - \tilde{\mathbf{y}}_N) \\
 \text{st.} \quad & \mathbf{A}_B (\vec{\mathbf{y}}_B - \tilde{\mathbf{y}}_B) + \mathbf{A}_N (\vec{\mathbf{y}}_N - \tilde{\mathbf{y}}_N) = \mathbf{0} \\
 & \vec{\mathbf{y}}_B \geq \mathbf{0}, \tilde{\mathbf{y}}_B \geq \mathbf{0} \qquad \vec{\mathbf{y}}_N \geq \mathbf{0}, \tilde{\mathbf{y}}_N \geq \mathbf{0} \\
 & \qquad \qquad \qquad \vec{\mathbf{y}}_{N_L} \leq \mathbf{0}, \vec{\mathbf{y}}_{N_U} \leq \mathbf{0} \\
 & \vec{\mathbf{y}}_F \leq \vec{\mathbf{r}}_F, \tilde{\mathbf{y}}_F \leq \tilde{\mathbf{r}}_F \\
 & \vec{\mathbf{y}}_{B_L} \leq \mathbf{0}, \vec{\mathbf{y}}_{B_U} \leq \mathbf{0} \qquad \vec{\mathbf{y}}_{N_L} \leq \vec{\mathbf{r}}_{N_L}, \vec{\mathbf{y}}_{N_U} \leq \vec{\mathbf{r}}_{N_U}
 \end{aligned}$$

$$B = F \cup B_L \cup B_U; \quad N = N_L \cup N_U$$



## STEP 3 FIX SOME DUAL VARIABLES IN ROW-SET $R$

For  $\emptyset \subseteq S \subseteq B$ , find  $\mathbf{y}^k = \begin{bmatrix} \mathbf{y}_S^k \\ \mathbf{y}_{\bar{S}}^k \end{bmatrix} = \begin{bmatrix} (\bar{\mathbf{y}}_S^k - \underline{\mathbf{y}}_S^k) \\ (\bar{\mathbf{y}}_{\bar{S}}^k - \underline{\mathbf{y}}_{\bar{S}}^k) \end{bmatrix}$  of min reduced cost  $\mu^k$ .

For basic columns  $\mathbf{A}_S$ ,  
select a  $s \times s$  square matrix  $\mathbf{A}_{RS}$ ,  
a set of  $s$  independent rows.

$$\mathbf{T} = \begin{bmatrix} \mathbf{A}_{RS} & \mathbf{0} \\ \mathbf{A}_{ZS} & \mathbf{I}_{m-s} \end{bmatrix} \quad \mathbf{T}^{-1} = \begin{bmatrix} \mathbf{A}_{RS}^{-1} & \mathbf{0} \\ -\mathbf{A}_{ZS}\mathbf{A}_{RS}^{-1} & \mathbf{I}_{m-s} \end{bmatrix}$$

# STEP 3 FIX SOME DUAL VARIABLES IN ROW-SET $R$

$$\emptyset \subseteq S \subseteq B$$

PERFORM  $\mathbf{T}^{-1}\mathbf{A}(\vec{\mathbf{y}} - \tilde{\mathbf{y}}) = \mathbf{0}$ . GENERAL CASE  $\pi^\top = \psi^\top \mathbf{T}^{-1}$

$$\begin{aligned} \mathbf{c}^\top \mathbf{x}^k + \min \quad & \mathbf{c}_S^\top (\vec{\mathbf{y}}_S - \tilde{\mathbf{y}}_S) + \mathbf{c}_{\bar{S}}^\top (\vec{\mathbf{y}}_{\bar{S}} - \tilde{\mathbf{y}}_{\bar{S}}) \\ \text{st.} \quad & (\vec{\mathbf{y}}_S - \tilde{\mathbf{y}}_S) + \bar{\mathbf{A}}_{R\bar{S}} (\vec{\mathbf{y}}_{\bar{S}} - \tilde{\mathbf{y}}_{\bar{S}}) = \mathbf{0} \quad [\psi_R = \mathbf{c}_S] \\ & \vec{\mathbf{y}}_S \geq \mathbf{0}, \tilde{\mathbf{y}}_S \geq \mathbf{0} \\ & \bar{\mathbf{A}}_{Z\bar{S}} (\vec{\mathbf{y}}_{\bar{S}} - \tilde{\mathbf{y}}_{\bar{S}}) = \mathbf{0} \quad [\psi_Z \text{ unknown}] \\ & \vec{\mathbf{y}}_{\bar{S}} \geq \mathbf{0}, \tilde{\mathbf{y}}_{\bar{S}} \geq \mathbf{0} \\ & \vec{\mathbf{y}}_S \leq \vec{\mathbf{r}}_S, \tilde{\mathbf{y}}_S \leq \tilde{\mathbf{r}}_S \quad \vec{\mathbf{y}}_{\bar{S}} \leq \vec{\mathbf{r}}_{\bar{S}}, \tilde{\mathbf{y}}_{\bar{S}} \leq \tilde{\mathbf{r}}_{\bar{S}} \end{aligned}$$

Observe  $\bar{\mathbf{c}}_S = \mathbf{0}$  (basic variables).

PERFORM  $\mathbf{T}^{-1}\mathbf{A}(\vec{\mathbf{y}} - \tilde{\mathbf{y}}) = \mathbf{0}$ . GENERAL CASE : DETAILS

$$\boldsymbol{\pi}^T = \boldsymbol{\psi}^T \mathbf{T}^{-1}$$

$$\begin{aligned} & \mathbf{c}^T \mathbf{x}^k + \min \mathbf{c}_S^T (\vec{\mathbf{y}}_S - \tilde{\mathbf{y}}_S) + \mathbf{c}_{\bar{S}}^T (\vec{\mathbf{y}}_{\bar{S}} - \tilde{\mathbf{y}}_{\bar{S}}) \\ & \text{st. } (\vec{\mathbf{y}}_S - \tilde{\mathbf{y}}_S) + \bar{\mathbf{A}}_{R\bar{S}} (\vec{\mathbf{y}}_{\bar{S}} - \tilde{\mathbf{y}}_{\bar{S}}) = \mathbf{0} \quad [\boldsymbol{\psi}_R = \mathbf{c}_S] \\ & \quad \vec{\mathbf{y}}_S \geq \mathbf{0}, \tilde{\mathbf{y}}_S \geq \mathbf{0} \\ & \quad \quad \quad \bar{\mathbf{A}}_{Z\bar{S}} (\vec{\mathbf{y}}_{\bar{S}} - \tilde{\mathbf{y}}_{\bar{S}}) = \mathbf{0} \quad [\boldsymbol{\psi}_Z \text{ unknown}] \\ & \quad \quad \quad \vec{\mathbf{y}}_{\bar{S}} \geq \mathbf{0}, \tilde{\mathbf{y}}_{\bar{S}} \geq \mathbf{0} \\ & \quad \quad \quad \vec{\mathbf{y}}_{\bar{S}_L} \leq \mathbf{0}, \vec{\mathbf{y}}_{\bar{S}_U} \leq \mathbf{0} \\ & \vec{\mathbf{y}}_{S_F} \leq \vec{\mathbf{r}}_{S_F}, \tilde{\mathbf{y}}_{S_F} \leq \vec{\mathbf{r}}_{S_F} \quad \vec{\mathbf{y}}_{\bar{S}_F} \leq \vec{\mathbf{r}}_{\bar{S}_F}, \tilde{\mathbf{y}}_{\bar{S}_F} \leq \vec{\mathbf{r}}_{\bar{S}_F} \\ & \quad \vec{\mathbf{y}}_{S_L} \leq \mathbf{0}, \vec{\mathbf{y}}_{S_U} \leq \mathbf{0} \quad \vec{\mathbf{y}}_{\bar{S}_L} \leq \vec{\mathbf{r}}_{\bar{S}_L}, \vec{\mathbf{y}}_{\bar{S}_U} \leq \vec{\mathbf{r}}_{\bar{S}_U} \end{aligned}$$

$$S = S_F \cup S_L \cup S_U; \quad \bar{S} = \bar{S}_F \cup \bar{S}_L \cup \bar{S}_U$$

## STEP 4 FIND MINIMUM REDUCED COST $\mu_S^k$

### PRICING OF THE VARIABLES

$S \subseteq B$ , hence  $\bar{c}_S = 0$ .

Therefore pricing of  $\bar{y}_S$  and  $\bar{y}_S$  needed to get partial direction  $(\bar{y}_S^k - \bar{y}_S^k)$

followed by impact on  $(\bar{y}_S^k - \bar{y}_S^k)$  to complete direction  $y^k = \begin{bmatrix} y_S^k \\ y_S^k \end{bmatrix}$ .

### PRIMAL/DUAL FORMULATIONS OF THE PRICING

$$\psi_R^T = c_S^T A_{RS}^{-1}$$

$$\begin{aligned} \max \mu \quad \text{st.} \quad \mu \mathbf{1}^T &\leq c_S^T - \psi_R^T A_{RS} - \psi_Z^T \bar{A}_{ZS} & [\bar{y}_S] \\ \mu \mathbf{1}^T &\leq -(c_S^T - \psi_R^T A_{RS} - \psi_Z^T \bar{A}_{ZS}) & [\bar{y}_S] \end{aligned}$$

### A CONVEX COMBINATION OF THE VARIABLES $\bar{y}_S$ AND $\bar{y}_S$

$$\begin{aligned} \equiv \mu &= \min (c_S^T - \psi_R^T A_{RS})(\bar{y}_S - \bar{y}_S) \\ \text{st.} \quad \mathbf{1}^T \bar{y}_S + \mathbf{1}^T \bar{y}_S &= 1 & [\mu] \\ A_{ZS}(\bar{y}_S - \bar{y}_S) &= 0 & [\psi_Z] \\ \bar{y}_S, \bar{y}_S &\geq 0, \bar{y}_{S_L} \leq 0, \bar{y}_{S_U} \leq 0 \end{aligned}$$

\*\*\* Optimal solution :  $\mu_S^k$ ,  $\bar{y}_S^k$  and  $\bar{y}_S^k$ . \*\*\*

... COMPLETE DIRECTION  $\mathbf{y}_S^k$

Given  $\mu^k < 0$  and  $(\vec{\mathbf{y}}_S^k - \bar{\mathbf{y}}_S^k)$ , find impacts on other variables  $(\vec{\mathbf{y}}_S^k - \bar{\mathbf{y}}_S^k)$ .

$$\begin{aligned}
 (\vec{\mathbf{y}}_S - \bar{\mathbf{y}}_S) + \bar{\mathbf{A}}_{R\bar{S}}(\vec{\mathbf{y}}_S^k - \bar{\mathbf{y}}_S^k) &= \mathbf{0} \\
 \vec{\mathbf{y}}_S \geq \mathbf{0}, \bar{\mathbf{y}}_S \geq \mathbf{0} & \qquad \qquad \qquad \vec{y}_j \bar{y}_j = 0, \quad \forall j \in S_F
 \end{aligned}$$

DIRECTION  $\mathbf{y}_S^k$

$$\mathbf{y}_S^k = \begin{bmatrix} (\vec{\mathbf{y}}_S^k - \bar{\mathbf{y}}_S^k) \\ (\vec{\mathbf{y}}_S^k - \bar{\mathbf{y}}_S^k) \end{bmatrix} = \begin{bmatrix} -\bar{\mathbf{A}}_{R\bar{S}}(\vec{\mathbf{y}}_S^k - \bar{\mathbf{y}}_S^k) \\ (\vec{\mathbf{y}}_S^k - \bar{\mathbf{y}}_S^k) \end{bmatrix} = \begin{bmatrix} \text{Impact level} \\ \text{Pricing level} \end{bmatrix}$$

*Positive part; Negative part.*

### STEP 5 COMPUTE MAXIMUM STEP-SIZE $\rho_S^k$

$$\rho \begin{bmatrix} \vec{y}_{S_F}^k \\ \vec{y}_{S_L}^k \\ \vec{y}_{S_U}^k \end{bmatrix} \leq \begin{bmatrix} \vec{r}_{S_F}^k \\ \vec{r}_{S_L}^k \\ \mathbf{0} \end{bmatrix} ; \quad \rho \begin{bmatrix} \vec{y}_{S_F}^k \\ \vec{y}_{S_L}^k \\ \vec{y}_{S_U}^k \end{bmatrix} \leq \begin{bmatrix} \vec{r}_{S_F}^k \\ \mathbf{0} \\ \vec{r}_{S_U}^k \end{bmatrix}$$

$$\rho \begin{bmatrix} \vec{y}_{S_F}^k \\ \vec{y}_{S_L}^k \\ \mathbf{0} \end{bmatrix} \leq \begin{bmatrix} \vec{r}_{S_F}^k \\ \vec{r}_{S_L}^k \\ \mathbf{0} \end{bmatrix} ; \quad \rho \begin{bmatrix} \vec{y}_{S_F}^k \\ \mathbf{0} \\ \vec{y}_{S_U}^k \end{bmatrix} \leq \begin{bmatrix} \vec{r}_{S_F}^k \\ \mathbf{0} \\ \vec{r}_{S_U}^k \end{bmatrix}$$

10 out of 12 types of residual upper bonds to verify.  $\emptyset \subseteq S \subseteq B$

### STEP 5 COMPUTE MAXIMUM STEP-SIZE $\rho_S^k$

$$\rho \begin{bmatrix} \vec{y}_F^k \\ \vec{y}_{B_L}^k \\ \vec{y}_{B_U}^k \end{bmatrix} \leq \begin{bmatrix} \vec{r}_F^k \\ \vec{r}_{B_L}^k \\ \mathbf{0} \end{bmatrix} ; \quad \rho \begin{bmatrix} \vec{y}_F^k \\ \vec{y}_{B_L}^k \\ \vec{y}_{B_U}^k \end{bmatrix} \leq \begin{bmatrix} \vec{r}_F^k \\ \mathbf{0} \\ \vec{r}_{B_U}^k \end{bmatrix}$$

$$\rho \begin{bmatrix} \vec{y}_{N_L}^k \end{bmatrix} \leq \begin{bmatrix} \vec{r}_{N_L}^k \end{bmatrix} ; \quad \rho \begin{bmatrix} \vec{y}_{N_U}^k \end{bmatrix} \leq \begin{bmatrix} \vec{r}_{N_U}^k \end{bmatrix}$$

8 types of residual upper bonds to verify for Primal Simplex.  $S = B$

STEP 5 COMPUTE MAXIMUM STEP-SIZE  $\rho_S^k$

$$\rho \begin{bmatrix} \bar{\mathbf{y}}_P^k \\ \bar{\mathbf{y}}_Z^k \end{bmatrix} \leq \begin{bmatrix} \bar{\mathbf{r}}_P^k \\ \mathbf{0} \end{bmatrix}$$

Only 2 types of residual upper bounds to verify for PS in standard form.  
 $S = B = P \cup Z$  (positive and zero variables)



STEP 5 COMPUTE MAXIMUM STEP-SIZE  $\rho_S^k$

$$\rho \begin{bmatrix} \vec{y}_F^k \\ \vec{y}_L^k \end{bmatrix} \leq \begin{bmatrix} \vec{r}_F^k \\ \vec{r}_L^k \end{bmatrix}; \quad \rho \begin{bmatrix} \vec{y}_F^k \\ \vec{y}_U^k \end{bmatrix} \leq \begin{bmatrix} \vec{r}_F^k \\ \vec{r}_U^k \end{bmatrix}$$

4 types of strictly positive residual upper bonds to verify in MMCC.  $S = \emptyset$

STEP 5 COMPUTE MAXIMUM STEP-SIZE  $\rho_S^k$

$$\rho \left[ \vec{y}_F^k \right] \leq \left[ \vec{r}_F^k \right]; \quad \rho \left[ \vec{y}_F^k \right] \leq \left[ \vec{r}_F^k \right]$$

$$\rho \left[ \vec{y}_L^k \right] \leq \left[ \vec{r}_L^k \right]; \quad \rho \left[ \vec{y}_U^k \right] \leq \left[ \vec{r}_U^k \right]$$

4 types of strictly positive residual upper bonds to verify in IPS.  $S = F$

$$\mathbf{T} = [\mathbf{A}_B \quad \emptyset], \quad \mathbf{T}^{-1} = \begin{bmatrix} \mathbf{A}_B^{-1} \\ \emptyset \end{bmatrix}$$

$$B = F \cup B_L \cup B_U; \quad N = N_L \cup N_U$$

$$\mathbf{T} = [\mathbf{A}_B \quad \emptyset], \quad \mathbf{T}^{-1} = \begin{bmatrix} \mathbf{A}_B^{-1} \\ \emptyset \end{bmatrix}$$

$$B = F \cup B_L \cup B_U; \quad N = N_L \cup N_U$$

## PRIMAL SIMPLEX METHOD (DANTZIG 1945)

$$\begin{aligned} \mathbf{c}^\top \mathbf{x}^k + \min \quad & \mathbf{c}_B^\top (\vec{\mathbf{y}}_B - \tilde{\mathbf{y}}_B) + \mathbf{c}_N^\top (\vec{\mathbf{y}}_N - \tilde{\mathbf{y}}_N) \\ \text{st.} \quad & (\vec{\mathbf{y}}_B - \tilde{\mathbf{y}}_B) + \bar{\mathbf{A}}_N (\vec{\mathbf{y}}_N - \tilde{\mathbf{y}}_N) = \mathbf{0} \quad [\boldsymbol{\psi}^\top = \mathbf{c}_B] \\ & \vec{\mathbf{y}}_B \geq \mathbf{0}, \tilde{\mathbf{y}}_B \geq \mathbf{0} \end{aligned}$$

$$\vec{\mathbf{y}}_N \geq \mathbf{0}, \tilde{\mathbf{y}}_N \geq \mathbf{0}$$

$$\vec{\mathbf{y}}_{N_L} = \mathbf{0}, \vec{\mathbf{y}}_{N_U} = \mathbf{0}$$

$$\vec{\mathbf{y}}_F \leq \vec{\mathbf{r}}_F, \tilde{\mathbf{y}}_F \leq \vec{\mathbf{r}}_F$$

$$\vec{\mathbf{y}}_{B_L} \leq \mathbf{0}, \tilde{\mathbf{y}}_{B_U} \leq \mathbf{0}$$

$$\vec{\mathbf{y}}_{N_L} \leq \vec{\mathbf{r}}_{N_L}, \tilde{\mathbf{y}}_{N_U} \leq \vec{\mathbf{r}}_{N_U}$$

$$*\rho_B \geq 0*$$

## PROPERTIES OF PS

No equality constraints in the pricing problem.

Pricing contains • convex combination of the non-basic variables  
• non-negativity restrictions (a cone).

Due to the step-size  $*\rho_B \geq 0*$ , possible degenerate pivots.

Oscillation of  $\mu_B$ ; it may even not converge towards 0.

$$\mathbf{T} = \begin{bmatrix} \mathbf{A}_{RF} & \mathbf{0} \\ \mathbf{A}_{ZF} & \mathbf{I}_{m-r} \end{bmatrix}, \quad \mathbf{T}^{-1} = \begin{bmatrix} \mathbf{A}_{RF}^{-1} & \mathbf{0} \\ -\mathbf{A}_{ZF}\mathbf{A}_{RF}^{-1} & \mathbf{I}_{m-r} \end{bmatrix}.$$

## SPECIAL CASE #2 : $S = F \Rightarrow \bar{S} = L \cup U$

$$\mathbf{T} = \begin{bmatrix} \mathbf{A}_{RF} & \mathbf{0} \\ \mathbf{A}_{ZF} & \mathbf{I}_{m-r} \end{bmatrix}, \quad \mathbf{T}^{-1} = \begin{bmatrix} \mathbf{A}_{RF}^{-1} & \mathbf{0} \\ -\mathbf{A}_{ZF}\mathbf{A}_{RF}^{-1} & \mathbf{I}_{m-r} \end{bmatrix}.$$

### IMPROVED PRIMAL SIMPLEX METHOD (ELHALLAOUI ET AL. 2011)

$$\begin{aligned} & \mathbf{c}^T \mathbf{x}^k + \min \quad \mathbf{c}_F^T (\bar{\mathbf{y}}_F - \tilde{\mathbf{y}}_F) + \mathbf{c}_L^T \bar{\mathbf{y}}_L - \mathbf{c}_U^T \tilde{\mathbf{y}}_U \\ \text{st.} \quad & (\bar{\mathbf{y}}_F - \tilde{\mathbf{y}}_F) + \bar{\mathbf{A}}_{RL} \bar{\mathbf{y}}_L - \bar{\mathbf{A}}_{RU} \tilde{\mathbf{y}}_U = \mathbf{0} \quad [\psi_R = \mathbf{c}_F] \\ & \bar{\mathbf{y}}_F \geq \mathbf{0}, \tilde{\mathbf{y}}_F \geq \mathbf{0} \\ & \bar{\mathbf{A}}_{ZL} \bar{\mathbf{y}}_L - \bar{\mathbf{A}}_{ZU} \tilde{\mathbf{y}}_U = \mathbf{0} \quad [\psi_Z] \\ & \bar{\mathbf{y}}_L \geq \mathbf{0}, \tilde{\mathbf{y}}_U \geq \mathbf{0} \\ & \bar{\mathbf{y}}_F \leq \bar{\mathbf{r}}_F, \tilde{\mathbf{y}}_F \leq \bar{\mathbf{r}}_F, \quad \bar{\mathbf{y}}_L \leq \bar{\mathbf{r}}_L, \tilde{\mathbf{y}}_U \leq \bar{\mathbf{r}}_U \quad * \rho > 0 * \end{aligned}$$

### PROPERTIES OF IPS

$f$  equality constraints in the master problem,  $m - f$  in the pricing problem.

Non-degenerate pivots only ( $\rho_F > 0$ ).

$z^0 > z^1 > z^2 > \dots = z^*$  cost **strictly decreasing** at each iteration ( $\rho_F > 0$ ).

Oscillation of  $\mu_F$  but converging towards 0.

Select  $S = \emptyset$ .  $\mathbf{T} = \begin{bmatrix} \emptyset & \mathbf{I}_m \end{bmatrix}$ ,  $\mathbf{T}^{-1} = \begin{bmatrix} \emptyset \\ \mathbf{I}_m \end{bmatrix}$

Select  $S = \emptyset$ .  $\mathbf{T} = \begin{bmatrix} \emptyset & \mathbf{I}_m \end{bmatrix}$ ,  $\mathbf{T}^{-1} = \begin{bmatrix} \emptyset \\ \mathbf{I}_m \end{bmatrix}$

## MINIMUM MEAN CYCLE-CANCELING ALGORITHM ADAPTED FOR LP

$$\begin{aligned} & \mathbf{c}^T \mathbf{x}^k + \min \quad \mathbf{c}^T (\vec{\mathbf{y}} - \tilde{\mathbf{y}}) \\ & \text{st.} \quad \mathbf{A}(\vec{\mathbf{y}} - \tilde{\mathbf{y}}) = \mathbf{0} \quad \text{Directions} \\ & \quad \quad \vec{\mathbf{y}}, \tilde{\mathbf{y}} \geq \mathbf{0} \quad \text{in the cone} \\ & \quad \quad \vec{\mathbf{y}}_L, \vec{\mathbf{y}}_U \leq \mathbf{0} \quad \text{at vertex } \mathbf{x}^k \\ & \quad \quad \vec{\mathbf{y}}_F, \tilde{\mathbf{y}}_F, \vec{\mathbf{y}}_L, \tilde{\mathbf{y}}_U \leq \vec{\mathbf{r}}_F^k, \tilde{\mathbf{r}}_F^k, \vec{\mathbf{r}}_L^k, \tilde{\mathbf{r}}_U^k \quad * \text{Step size } \rho_\emptyset > 0 * \end{aligned}$$

## PROPERTIES OF MMCC

All equality constraints in the pricing problem.

Upper bounds in the master problem.

$z^0 > z^1 > z^2 > \dots = z^*$  cost **strictly decreasing** at each iteration ( $\rho_\emptyset > 0$ ).


$\mu^0 \leq \mu^1 \leq \mu^2 \leq \dots = 0$  smallest reduced cost **non decreasing**.

MMCC is strongly polynomial for network flow problems in  $O(mn)$  phases.


Goldberg and Tarjan (1989), Radzick and Goldberg (1994)




## COMPUTATIONAL RESULTS




# ILLUSTRATION OF $\mu$ (NETWORK WITH $n=1025$ , $m=91,220$ )




## COMPUTATIONAL RESULTS



## COMPUTATIONAL RESULTS



## COMPUTATIONAL RESULTS





## LINEAR PROGRAM $LP$

$$\begin{aligned} z^* := & \min \quad \mathbf{c}^T \mathbf{x} \\ \text{st.} \quad & \mathbf{Ax} = \mathbf{b} \\ & \mathbf{l} \leq \mathbf{x} \leq \mathbf{u} \end{aligned}$$

## GENERIC ALGORITHM WITH SINGLE PARAMETER SET $S$ , $\emptyset \subseteq S \subseteq B$

- 1 Let  $k = 0$  and assume a feasible basic solution  $\mathbf{x}^k$  to LP.
- 2 For  $\emptyset \subseteq S \subseteq B$ , construct the residual problem  $LP_S(\mathbf{x}^k)$ .
- 3 Fix a subset of the dual variables in row-set  $R$ :  $\boldsymbol{\pi}_R^T = \mathbf{c}_S^T \mathbf{A}_{RS}^{-1}$ .
- 4 Determine the value of the smallest reduced cost  $\mu_S^k$ .  
If  $\mu_S^k \geq 0$ , STOP. Current solution  $\mathbf{x}^k$  is optimal for LP.
- 5 Retrieve direction  $\mathbf{y}_S^k \in \mathbb{R}^n$  and compute its maximum step-size  $\rho_S^k$ .
- 6 Update  $\begin{aligned} \mathbf{x}^{k+1} &:= \mathbf{x}^k + \rho_S^k \mathbf{y}_S^k; \\ z^{k+1} &:= z^k + \rho_S^k \mu_S^k; \\ k &:= k + 1. \end{aligned}$
- 7 Goto Step 2.







- $S \cap \{L \cup U\} \neq \emptyset$  : It may come up with degenerate pivots and not converge.
- Primal Simplex method (PS).

$$S = B, * \rho_B \geq 0 *$$







- $S \cap \{L \cup U\} \neq \emptyset$  : It may come up with degenerate pivots and not converge.
  - Primal Simplex method (PS).  $S = B, * \rho_B \geq 0 *$
- $\emptyset \subseteq S \subseteq F$  : It ensures a non-degenerate pivot at every iteration.
  - Improved Primal Simplex algorithm (IPS).  $S = F, * \rho_F > 0 *$



- $S \cap \{L \cup U\} \neq \emptyset$  : It may come up with degenerate pivots and not converge.
  - Primal Simplex method (PS).  $S = B, * \rho_B \geq 0 *$
- $\emptyset \subseteq S \subseteq F$  : It ensures a non-degenerate pivot at every iteration.
  - Improved Primal Simplex algorithm (IPS).  $S = F, * \rho_F > 0 *$
  - Minimum mean cycle-canceling algorithm (MMCC)  $S = \emptyset, * \rho_\emptyset > 0 *$



- $S \cap \{L \cup U\} \neq \emptyset$  : It may come up with degenerate pivots and not converge.
  - Primal Simplex method (PS).  $S = B, * \rho_B \geq 0 *$
- $\emptyset \subseteq S \subseteq F$  : It ensures a non-degenerate pivot at every iteration.
  - Improved Primal Simplex algorithm (IPS).  $S = F, * \rho_F > 0 *$
  - Minimum mean cycle-canceling algorithm (MMCC)  $S = \emptyset, * \rho_\emptyset > 0 *$   
*Strongly polynomial for network flow problems.*



- $S \cap \{L \cup U\} \neq \emptyset$  : It may come up with degenerate pivots and not converge.
  - Primal Simplex method (PS).  $S = B, * \rho_B \geq 0 *$
- $\emptyset \subseteq S \subseteq F$  : It ensures a non-degenerate pivot at every iteration.
  - Improved Primal Simplex algorithm (IPS).  $S = F, * \rho_F > 0 *$
  - Minimum mean cycle-canceling algorithm (MMCC)  $S = \emptyset, * \rho_\emptyset > 0 *$   
*Strongly polynomial for network flow problems.*
  - \*  $S \subset F$  : Optimal direction  $y_s^k$  can be an interior ray.

\*\*\* Imagine the same transformation is kept for a while...

- $\mathbf{T} := [\mathbf{\Lambda}, \mathbf{\Lambda}^\perp]$

- $\mathbf{\Lambda} = \begin{bmatrix} \mathbf{\Lambda}_R \\ \mathbf{\Lambda}_Z \end{bmatrix}$ , where  $\mathbf{\Lambda}_R$  is a set of  $s$  independent rows.

- $\mathbf{T} = \begin{bmatrix} \mathbf{\Lambda}_R & \mathbf{0} \\ \mathbf{\Lambda}_Z & \mathbf{I}_{m-s} \end{bmatrix} \quad \mathbf{T}^{-1} = \begin{bmatrix} \mathbf{\Lambda}_R^{-1} & \mathbf{0} \\ \mathbf{\Lambda}_Z \mathbf{\Lambda}_R^{-1} & \mathbf{I}_{m-s} \end{bmatrix}$

- $\mathbf{T}^{-1}$  splits row-space  $\mathbb{R}^m$  of  $LP(\mathbf{x}^k)$  into two vector subspaces  $\mathbf{V}$  and  $\mathbf{V}^\perp$ .

- Vector subspace basis  $\mathbf{\Lambda}$  spans  $\mathbf{V}$  of dimension  $0 \leq s \leq m$ .

- Vector  $\mathbf{a} \in \mathbf{V}$  if and only if  $\bar{\mathbf{a}}_Z = \mathbf{0}$ , where  $\bar{\mathbf{a}} = \mathbf{T}^{-1}\mathbf{a} = \begin{bmatrix} \bar{\mathbf{a}}_R \\ \mathbf{0} \end{bmatrix}$ .

- $\emptyset \subseteq S \subseteq B$  : index subset of basic columns spanned by  $\mathbf{\Lambda}$ .

- Algorithmic properties derived according to subset  $S$ .



# DYNAMIC CONSTRAINT AGGREGATION FOR SET PARTITIONING (ELHALLAOUI ET AL. 2005)


The partition of the row-set is derived from the  $f$  groups of identical rows of  $A_F$ .

$$\begin{array}{c} \mathbf{A}_F \\ \left( \begin{array}{cccc} 1 & 1 & & \\ & 1 & 1 & \\ & & 1 & 1 \\ 1 & & 1 & \\ 1 & & & 1 \\ 1 & 1 & & \\ & 1 & 1 & \\ 1 & & & 1 \end{array} \right) \end{array} \mapsto \begin{array}{c} \mathbf{\Lambda} \\ \left( \begin{array}{cccc} 1 & & & \\ & 1 & & \\ & & 1 & \\ & & & 1 \\ \hline 1 & & & 1 \\ & 1 & & \\ & & & 1 \end{array} \right) \end{array} .$$


## PROPERTIES

$F \subseteq S \subseteq B$  for fractional solutions but  $S = F$  (or  $S = \emptyset$ ) for binary solutions.  
 Degenerate pivots may occur,  $\rho \geq 0$ , if  $r > f$ .

### Vector Space Decomposition of the Equality Constraints







# MMCC:

*non-extreme ray* a-c-d

