

# Asymptotic Analysis

Alfred Rossi, Ph.D.

# Why Asymptotic Analysis?

*Algorithm A is **2x faster** on My Computer than Algorithm B is  
**on yours.***

# Why Asymptotic Analysis?

What about this?

▲ *Algorithm A is 2x faster than Algorithm B on Hardware 1*

# Why Asymptotic Analysis?

What about this?

- ▲ *Algorithm A is 2x faster than Algorithm B on Hardware 1*
- ▲ *Algorithm B is 5x faster than Algorithm A on Hardware 2*

# Why Asymptotic Analysis?

Which algorithm is faster?

- ▲ The **method** behind *Algorithm C* is **universally slower** than the method of *Algorithm D*.

# Why Asymptotic Analysis?

Which algorithm is faster?

- ▲ The **method** behind *Algorithm C* is **universally slower** than the method of *Algorithm D*.
- ▲ So the authors of *Algorithm C* have precomputed a lookup table for all results up to size **10000**.

# Why Asymptotic Analysis?

Which algorithm is faster?

- ▲ The **method** behind *Algorithm C* is **universally slower** than the method of *Algorithm D*.
- ▲ So the authors of *Algorithm C* have precomputed a lookup table for all results up to size **10000**.
- ▲ This size is big enough (and therefore instant) for almost all applications.

# Why Asymptotic Analysis?

- ▲ We want to compare algorithms directly without reference to hardware, so we ***should not be sensitive to multiplicative constants.***



# Why Asymptotic Analysis?

- ▲ We want to compare algorithms directly without reference to hardware, so we ***should not be sensitive to multiplicative constants***.
- ▲ In other words, ***we want to count steps*** and not worry about how long each step takes.

# Why Asymptotic Analysis?

- ▲ We want to compare algorithms directly without reference to hardware, so we ***should not be sensitive to multiplicative constants.***
- ▲ In other words, ***we want to count steps*** and not worry about how long each step takes.
- ▲ Because your  $10n$  steps are comparable to my  $5n$  steps if your computer is twice as fast as mine we need to ignore the constants and, instead, focus on the ***rate of growth of the number of steps as the input size grows.***

# Why Asymptotic Analysis?

- ▲ We want to compare algorithms directly without reference to hardware, so we ***should not be sensitive to multiplicative constants***.
- ▲ In other words, ***we want to count steps*** and not worry about how long each step takes.
- ▲ Because your  $10n$  steps are comparable to my  $5n$  steps if your computer is twice as fast as mine we need to ignore the constants and, instead, focus on the ***rate of growth of the number of steps as the input size grows***.
- ▲ We need to ***rule out the lookup table trick***, so we choose to compare them on very large inputs ( $n \rightarrow \infty$ ).

In other words, in the asymptotic analysis of algorithms, we don't care about how long individual steps take, we ***care about how the number of steps blows up as the input size goes to infinity.***

Let's leave algorithms behind until next week and refresh on the math.

# Notation

# Notation

Some common sets include:

- ▲ the empty set, denoted  $\emptyset$ , is the unique set with 0 elements

# Notation

Some common sets include:

- ▲ the empty set, denoted  $\emptyset$ , is the unique set with 0 elements
- ▲ the set of natural numbers,  $\mathbb{N} = \{1, 2, \dots\}$



# Notation

Some common sets include:

- ▲ the empty set, denoted  $\emptyset$ , is the unique set with 0 elements
- ▲ the set of natural numbers,  $\mathbb{N} = \{1, 2, \dots\}$
- ▲ the set of natural numbers, with 0,  $\mathbb{N}_0 = \{0, 1, 2, \dots\}$

# Notation

Some common sets include:

- ▲ the empty set, denoted  $\emptyset$ , is the unique set with 0 elements
- ▲ the set of natural numbers,  $\mathbb{N} = \{1, 2, \dots\}$
- ▲ the set of natural numbers, with 0,  $\mathbb{N}_0 = \{0, 1, 2, \dots\}$
- ▲ the set  $[n]$  consisting of the natural numbers 1 through  $n$ .  
By convention we take  $[0] = \emptyset$ .

# Notation

Some common sets include:

- ▲ the empty set, denoted  $\emptyset$ , is the unique set with 0 elements
- ▲ the set of natural numbers,  $\mathbb{N} = \{1, 2, \dots\}$
- ▲ the set of natural numbers, with 0,  $\mathbb{N}_0 = \{0, 1, 2, \dots\}$
- ▲ the set  $[n]$  consisting of the natural numbers 1 through  $n$ .  
By convention we take  $[0] = \emptyset$ .
- ▲ the set of integers,  $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

# Notation

Some common sets include:

- ▲ the empty set, denoted  $\emptyset$ , is the unique set with 0 elements
- ▲ the set of natural numbers,  $\mathbb{N} = \{1, 2, \dots\}$
- ▲ the set of natural numbers, with 0,  $\mathbb{N}_0 = \{0, 1, 2, \dots\}$
- ▲ the set  $[n]$  consisting of the natural numbers 1 through  $n$ .  
By convention we take  $[0] = \emptyset$ .
- ▲ the set of integers,  $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$
- ▲ the set of rational numbers,  $\mathbb{Q} = \{p/q : p, q \in \mathbb{Z}, q \neq 0\}$

# Notation

Some common sets include:

- ▲ the empty set, denoted  $\emptyset$ , is the unique set with 0 elements
- ▲ the set of natural numbers,  $\mathbb{N} = \{1, 2, \dots\}$
- ▲ the set of natural numbers, with 0,  $\mathbb{N}_0 = \{0, 1, 2, \dots\}$
- ▲ the set  $[n]$  consisting of the natural numbers 1 through  $n$ .  
By convention we take  $[0] = \emptyset$ .
- ▲ the set of integers,  $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$
- ▲ the set of rational numbers,  $\mathbb{Q} = \{p/q : p, q \in \mathbb{Z}, q \neq 0\}$
- ▲ the set of real numbers,  $\mathbb{R}$

# Notation

Some common sets include:

- ▲ the empty set, denoted  $\emptyset$ , is the unique set with 0 elements
- ▲ the set of natural numbers,  $\mathbb{N} = \{1, 2, \dots\}$
- ▲ the set of natural numbers, with 0,  $\mathbb{N}_0 = \{0, 1, 2, \dots\}$
- ▲ the set  $[n]$  consisting of the natural numbers 1 through  $n$ .  
By convention we take  $[0] = \emptyset$ .
- ▲ the set of integers,  $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$
- ▲ the set of rational numbers,  $\mathbb{Q} = \{p/q : p, q \in \mathbb{Z}, q \neq 0\}$
- ▲ the set of real numbers,  $\mathbb{R}$
- ▲ the set of positive real numbers,  $\mathbb{R}^+ = \{x : x \in \mathbb{R}, x > 0\}$

# Notation

We will also use  $\log(n)$  in here to mean  $\log_2(n)$ .


**Big  $O$**





Let  $f$  and  $g$  be a pair of functions defined on an unbounded subset of positive real numbers.


## Definition

We say  $f \in O(g)$  if there exists  $c \in \mathbb{R}^+$ ,  $n_0 \in \mathbb{N}_0$ , such that for all  $n \geq n_0$ ,  $0 \leq f(n) \leq c \cdot g(n)$ .









# Set Notation

Note that the use of set notation in  $f \in O(g)$  is not accidental. We can think of  $O(g)$  as the **set of functions** whose tail behavior is asymptotically dominated by a **rescaling** of  $g$ .

# Transitivity of $O$

## Theorem (Transitivity of $O$ )

*Let  $f$ ,  $g$ , and  $h$  be functions defined on an unbounded subset of positive real numbers. If  $f \in O(g)$  and  $g \in O(h)$  then  $f \in O(h)$ .*



Recall, our goal is to show  $f \in O(h)$ .

Proof.

- ▲ Since  $f \in O(g)$  there exists constants  $c_g \in \mathbb{R}^+$ ,  $n_g \in \mathbb{N}_0$  such that, for all  $n \geq n_g$ ,  $0 \leq f(n) \leq c_g g(n)$ .



Recall, our goal is to show  $f \in O(h)$ .

Proof.

- ▲ Since  $f \in O(g)$  there exists constants  $c_g \in \mathbb{R}^+$ ,  $n_g \in \mathbb{N}_0$  such that, for all  $n \geq n_g$ ,  $0 \leq f(n) \leq c_g g(n)$ .
- ▲ Since  $g \in O(h)$  there exists constants  $c_h \in \mathbb{R}^+$ ,  $n_h \in \mathbb{N}_0$  such that, for all  $n \geq n_h$ ,  $0 \leq g(n) \leq c_h h(n)$ .



Recall, our goal is to show  $f \in O(h)$ .

**Proof.**

- ▲ Since  $f \in O(g)$  there exists constants  $c_g \in \mathbb{R}^+$ ,  $n_g \in \mathbb{N}_0$  such that, for all  $n \geq n_g$ ,  $0 \leq f(n) \leq c_g g(n)$ .
- ▲ Since  $g \in O(h)$  there exists constants  $c_h \in \mathbb{R}^+$ ,  $n_h \in \mathbb{N}_0$  such that, for all  $n \geq n_h$ ,  $0 \leq g(n) \leq c_h h(n)$ .
- ▲ So, provided  $n \geq n_h$ , it follows that  $0 \leq c_g g(n) \leq c_g c_h h(n)$ .



Recall, our goal is to show  $f \in O(h)$ .

**Proof.**

- ▲ Since  $f \in O(g)$  there exists constants  $c_g \in \mathbb{R}^+$ ,  $n_g \in \mathbb{N}_0$  such that, for all  $n \geq n_g$ ,  $0 \leq f(n) \leq c_g g(n)$ .
- ▲ Since  $g \in O(h)$  there exists constants  $c_h \in \mathbb{R}^+$ ,  $n_h \in \mathbb{N}_0$  such that, for all  $n \geq n_h$ ,  $0 \leq g(n) \leq c_h h(n)$ .
- ▲ So, provided  $n \geq n_h$ , it follows that  $0 \leq c_g g(n) \leq c_g c_h h(n)$ .
- ▲ It follows that  $0 \leq f(n) \leq c_g g(n) \leq c_g c_h h(n)$ , whenever  $n$  is *also* greater than  $n_g$ .



Recall, our goal is to show  $f \in O(h)$ .

**Proof.**

- ▲ Since  $f \in O(g)$  there exists constants  $c_g \in \mathbb{R}^+$ ,  $n_g \in \mathbb{N}_0$  such that, for all  $n \geq n_g$ ,  $0 \leq f(n) \leq c_g g(n)$ .
- ▲ Since  $g \in O(h)$  there exists constants  $c_h \in \mathbb{R}^+$ ,  $n_h \in \mathbb{N}_0$  such that, for all  $n \geq n_h$ ,  $0 \leq g(n) \leq c_h h(n)$ .
- ▲ So, provided  $n \geq n_h$ , it follows that  $0 \leq c_g g(n) \leq c_g c_h h(n)$ .
- ▲ It follows that  $0 \leq f(n) \leq c_g g(n) \leq c_g c_h h(n)$ , whenever  $n$  is *also* greater than  $n_g$ .
- ▲ Thus, picking  $c = c_g c_h$  and  $n_0 = \max(n_g, n_h)$  gives the desired result.



# Big $O$ : Problem 1

Show that  $5n^2 \in O(n^2)$ .

▲ Note that  $f(n) = 5n^2$  and  $g(n) = n^2$ .

Show that  $5n^2 \in O(n^2)$ .

- ▲ Note that  $f(n) = 5n^2$  and  $g(n) = n^2$ .
- ▲ We need to find  $c \in \mathbb{R}^+$  and  $n_0 \in \mathbb{N}_0$  such that  $5n^2 \leq cn^2$  for all  $n \geq n_0$ .



Show that  $5n^2 \in O(n^2)$ .

- ▲ Note that  $f(n) = 5n^2$  and  $g(n) = n^2$ .
- ▲ We need to find  $c \in \mathbb{R}^+$  and  $n_0 \in \mathbb{N}_0$  such that  $5n^2 \leq cn^2$  for all  $n \geq n_0$ .
- ▲ Observe we may simply pick  $c = 5$ .

Show that  $5n^2 \in O(n^2)$ .

- ▲ Note that  $f(n) = 5n^2$  and  $g(n) = n^2$ .
- ▲ We need to find  $c \in \mathbb{R}^+$  and  $n_0 \in \mathbb{N}_0$  such that  $5n^2 \leq cn^2$  for all  $n \geq n_0$ .
- ▲ Observe we may simply pick  $c = 5$ .
- ▲ Plugging in for  $c$ , it holds that  $5n^2 \leq 5n^2$ , for any  $n$ .

Show that  $5n^2 \in O(n^2)$ .

- ▲ Note that  $f(n) = 5n^2$  and  $g(n) = n^2$ .
- ▲ We need to find  $c \in \mathbb{R}^+$  and  $n_0 \in \mathbb{N}_0$  such that  $5n^2 \leq cn^2$  for all  $n \geq n_0$ .
- ▲ Observe we may simply pick  $c = 5$ .
- ▲ Plugging in for  $c$ , it holds that  $5n^2 \leq 5n^2$ , for any  $n$ .
- ▲ So we can just pick  $n_0 = 1$ .

## Big $O$ : Problem 2

Show that  $5n^2 \notin O(n)$ .

Show that  $5n^2 \notin O(n)$ .

**Proof.**

Suppose  $5n^2 \in O(n)$ .

Show that  $5n^2 \notin O(n)$ .

**Proof.**

Suppose  $5n^2 \in O(n)$ . Then there exists a  $c \in \mathbb{R}^+$  and  $n_0 \in \mathbb{N}_0$  such that  $5n^2 \leq cn$  for all  $n \geq n_0$ .

Show that  $5n^2 \notin O(n)$ .

**Proof.**

Suppose  $5n^2 \in O(n)$ . Then there exists a  $c \in \mathbb{R}^+$  and  $n_0 \in \mathbb{N}_0$  such that  $5n^2 \leq cn$  for all  $n \geq n_0$ . Dividing through by  $n$  we see that  $5n \leq c$ .



Show that  $5n^2 \notin O(n)$ .

**Proof.**

Suppose  $5n^2 \in O(n)$ . Then there exists a  $c \in \mathbb{R}^+$  and  $n_0 \in \mathbb{N}_0$  such that  $5n^2 \leq cn$  for all  $n \geq n_0$ . Dividing through by  $n$  we see that  $5n \leq c$ . In other words, the constant  $c$  must be bigger than  $n$  for all  $n \geq n_0$ .

Show that  $5n^2 \notin O(n)$ .

**Proof.**

Suppose  $5n^2 \in O(n)$ . Then there exists a  $c \in \mathbb{R}^+$  and  $n_0 \in \mathbb{N}_0$  such that  $5n^2 \leq cn$  for all  $n \geq n_0$ . Dividing through by  $n$  we see that  $5n \leq c$ . In other words, the constant  $c$  must be bigger than  $n$  for all  $n \geq n_0$ . This is **impossible**, so no such  $c$  exists and the definition cannot be satisfied!  $\square$

## Big $O$ : Problem 3

Show that  $\sqrt{8n^{10} + 3n^3 + 8} \in O(n^5)$ .

▲ Observe that for  $n \geq 1$ ,

$$\sqrt{8n^{10} + 3n^3 + 8} \leq \sqrt{8n^{10} + 3n^{10} + 8n^{10}} = \sqrt{19n^{10}} = \sqrt{19}n^5$$

Show that  $\sqrt{8n^{10} + 3n^3 + 8} \in O(n^5)$ .

▲ Observe that for  $n \geq 1$ ,

$$\sqrt{8n^{10} + 3n^3 + 8} \leq \sqrt{8n^{10} + 3n^{10} + 8n^{10}} = \sqrt{19}n^5$$

▲ Recall, we need to find  $c \in \mathbb{R}^+$  and  $n_0 \in \mathbb{N}_0$  such that  $\sqrt{8n^{10} + 3n^3 + 8} \leq cn^5$  for all  $n \geq n_0$ .

Show that  $\sqrt{8n^{10} + 3n^3 + 8} \in O(n^5)$ .

▲ Observe that for  $n \geq 1$ ,

$$\sqrt{8n^{10} + 3n^3 + 8} \leq \sqrt{8n^{10} + 3n^{10} + 8n^{10}} = \sqrt{19}n^5$$

▲ Recall, we need to find  $c \in \mathbb{R}^+$  and  $n_0 \in \mathbb{N}_0$  such that  $\sqrt{8n^{10} + 3n^3 + 8} \leq cn^5$  for all  $n \geq n_0$ .

▲ Thus if we pick  $c = \sqrt{19}$  and  $n_0 = 1$  we are done.

## Big $O$ : Problem 4

Show that  $\log(n^5) \in O(\log(n))$ .

- ▲ By the property of logs that  $\log_b(n^a) = a \log_b(n)$ , we have  $\log(n^5) = 5 \log(n)$ .



Show that  $\log(n^5) \in O(\log(n))$ .

- ▲ By the property of logs that  $\log_b(n^a) = a \log_b(n)$ , we have  $\log(n^5) = 5 \log(n)$ .
- ▲ Thus if we pick  $c = 5$  and  $n_0 = 1$  we are done.

## Big $O$ : Problem 5

Show that  $\log_b(n) \in O(\log_c(n))$ .

- ▲ By the change of base formula for logs,  
 $\log_b(n) = \log_c(n) / \log_c(b)$ .


Show that  $\log_b(n) \in O(\log_c(n))$ .


- ▲ By the change of base formula for logs,  
 $\log_b(n) = \log_c(n) / \log_c(b)$ .
- ▲ Thus if we pick  $c = 1 / \log_c(b)$  and  $n_0 = 1$  we are done.

**Big  $\Omega$**


## Definition


We say  $f \in \Omega(g)$  if there exists  $c \in \mathbb{R}^+$ ,  $n_0 \in \mathbb{N}_0$ , such that for all  $n \geq n_0$ ,  $0 \leq c \cdot g(n) \leq f(n)$ .











## Set Notation

Again, the use of set notation in  $f \in \Omega(g)$  is not accidental. We can think of  $\Omega(g)$  as the **set of functions** whose tail behavior is asymptotically dominates  $g$ .

# Transitivity of $\Omega$

## Theorem (Transitivity of $\Omega$ )

*Let  $f$ ,  $g$ , and  $h$  be functions defined on an unbounded subset of positive real numbers. If  $f \in \Omega(g)$  and  $g \in \Omega(h)$  then  $f \in \Omega(h)$ .*

## Relationship Between $O$ and $\Omega$

### Theorem

*Let  $f$  and  $g$  be a pair of functions defined on an unbounded subset of positive real numbers, then  $f \in O(g)$  if and only if  $g \in \Omega(f)$ .*

Proof.

- ▲ Since  $f \in O(g)$  there exists constants  $c_g \in \mathbb{R}^+$ ,  $n_g \in \mathbb{N}_0$  such that, for all  $n \geq n_g$ ,  $0 \leq f(n) \leq c_g g(n)$ .



Proof.

- ▲ Since  $f \in O(g)$  there exists constants  $c_g \in \mathbb{R}^+$ ,  $n_g \in \mathbb{N}_0$  such that, for all  $n \geq n_g$ ,  $0 \leq f(n) \leq c_g g(n)$ .
- ▲ That is, for all  $n \geq n_g$ ,  $0 \leq (1/c_g) f(n) \leq g(n)$ .



## Proof.

- ▲ Since  $f \in O(g)$  there exists constants  $c_g \in \mathbb{R}^+$ ,  $n_g \in \mathbb{N}_0$  such that, for all  $n \geq n_g$ ,  $0 \leq f(n) \leq c_g g(n)$ .
- ▲ That is, for all  $n \geq n_g$ ,  $0 \leq (1/c_g) f(n) \leq g(n)$ .
- ▲ But this is just the definition of  $g \in \Omega(f)$  with  $c = 1/c_g \in \mathbb{R}^+$ , and  $n_0 = n_g \in \mathbb{N}_0$ .








**Big  $\ominus$**


## Definition

We say  $f \in \Theta(g)$  if there exists  $c_1, c_2 \in \mathbb{R}^+$ ,  $n_0 \in \mathbb{N}_0$ , such that for all  $n \geq n_0$ ,  $0 \leq c_1 \cdot g(n) \leq f(n) \leq c_2 \cdot g(n)$ .









## Relationship with $O$ and $\Omega$

## Theorem

*Let  $f$  and  $g$  be a pair of functions defined on an unbounded subset of positive real numbers, then  $f \in \Theta(g)$  if and only if  $f \in O(g)$  and  $g \in O(f)$ .*



## **Other Asymptotic Relations**

## Little $o$

Let  $f$  and  $g$  be a pair of functions defined on an unbounded subset of positive real numbers.

### Definition (Little $o$ )

We say  $f \in o(g)$  whenever  $f \in O(g)$  but  $f \notin \Theta(g)$ .

# Little $o$

Let  $f$  and  $g$  be a pair of functions defined on an unbounded subset of positive real numbers.

## Definition (Alternative Definition of Little $o$ )

We say  $f \in o(g)$  if, for every  $c \in \mathbb{R}^+$ , there exists  $n_0 \in \mathbb{N}_0$  such that for all  $n \geq n_0$ ,  $0 \leq f(n) < c \cdot g(n)$ .

## Little $o$

In other words,  $f \in o(g)$  if and only if, for any constant  $c > 0$ , and all large enough  $n$ ,

$$0 \leq \frac{f(n)}{g(n)} < c.$$

## Little $o$

In *other* other words,  $f \in o(g)$  is equivalent to saying

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 0.$$

## Little $\omega$

Let  $f$  and  $g$  be a pair of functions defined on an unbounded subset of positive real numbers.

### Definition (Little $\omega$ )

We say  $f \in \omega(g)$  whenever  $f \in \Omega(g)$  but  $f \notin \Theta(g)$ .

## Little $\omega$

Analogously,  $f \in \omega(g)$  if and only if, for all  $c \in \mathbb{R}^+$ , and all large enough  $n$ , we have that

$$\frac{f(n)}{g(n)} > c.$$

## Little $\omega$

Analogously,  $f \in \omega(g)$  if and only if, for all  $c \in \mathbb{R}^+$ , and all large enough  $n$ , we have that

$$\frac{f(n)}{g(n)} > c.$$

In other words,  $f \in \omega(g)$  is equivalent to saying

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \infty.$$



# Comparator Analogy

In other words, we can think of the relationships as:

- ▲  $f \in o(g)$  is like " $f < g$ " ( $f$  blows up strictly slower than  $g$ )
- ▲  $f \in O(g)$  is like " $f \leq g$ " ( $f$  blows up at most as fast as  $g$ )
- ▲  $f \in \Theta(g)$  is like " $f = g$ " ( $f$  blows up at the same rate as  $g$ )
- ▲  $f \in \Omega(g)$  is like " $f \geq g$ " ( $f$  blows up at least as fast as  $g$ )
- ▲  $f \in \omega(g)$  is like " $f > g$ " ( $f$  blows up strictly faster than  $g$ )

where the middle three states hold up to a constant factor.

# Comparator Analogy

Example  $f(n) = n^2$  and  $g(n) = n^2 + n$ , then  $f \in \Theta(g)$  and  $f$  is blowing up at the same rate as  $g$  (and vice versa), while  $f(n) = n^2$  and  $g(n) = n^3$  then  $f \in o(g)$  and  $f$  is blowing up strictly slower than  $g$ .

# Limit Theorems

## Theorem (Limit Theorems)

1.  $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 0$  if and only if  $f \in o(g)$ .
2. If  $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \in \mathbb{R}^+$ , then  $f \in \Theta(g)$ .
3.  $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \infty$  if and only if  $f \in \omega(g)$ .

## Theorem (Limit Theorems)

1.  $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 0$  if and only if  $f \in o(g)$ .
2. If  $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \in \mathbb{R}^+$ , then  $f \in \Theta(g)$ .
3.  $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \infty$  if and only if  $f \in \omega(g)$ .

Please note that the middle theorem is a bit more nuanced than the other two. If the limit fails to exist (say because the ratio oscillates), then the theorem does not apply. This also means the converse does not hold.

# **Advanced Topics**

What about limit theorems that test for big  $O$  and  $\Omega$ ?


# Supremum

Informally, the supremum of a set is a kind of natural maximum.




# Supremum

For example, what is the maximum value of the set  $[0, 1)$ ?



# Supremum


For example, what is the maximum value of the set  $[0, 1)$ ?



Is it  $0.9$ ?

# Supremum


For example, what is the maximum value of the set  $[0, 1)$ ?



Is it 0.9? 0.99?

# Supremum


For example, what is the maximum value of the set  $[0, 1)$ ?



Is it 0.9? 0.99? 0.999?


# Supremum

For example, what is the maximum value of the set  $[0, 1)$ ?



# Supremum

For example, what is the maximum value of the set  $[0, 1)$ ?



We want to say it's 1, but  $1 \notin [0, 1)$ .

# Supremum

Let  $P$  be an ordered set, and  $S \subset P$ .

A value  $x \in P$  is an **upper bound** of  $S$  if  $s \leq x$  for all  $s \in S$ .



# Supremum

Let  $P$  be an ordered set, and  $S \subset P$ .

A value  $x \in P$  is an **upper bound** of  $S$  if  $s \leq x$  for all  $s \in S$ .





# Supremum

Let  $P$  be an ordered set, and  $S \subset P$ .

A value  $x \in P$  is an **upper bound** of  $S$  if  $s \leq x$  for all  $s \in S$ .



# Supremum

Let  $P$  be an ordered set, and  $S \subset P$ .

A value  $x \in P$  is an **upper bound** of  $S$  if  $s \leq x$  for all  $s \in S$ .



# Supremum

Let  $P$  be an ordered set, and  $S \subset P$ .

A value  $x \in P$  is an **upper bound** of  $S$  if  $s \leq x$  for all  $s \in S$ .



# Supremum

Let  $P$  be an ordered set, and  $S \subset P$ .

A value  $x \in P$  is an **upper bound** of  $S$  if  $s \leq x$  for all  $s \in S$ .



# Supremum

Let  $P$  be an ordered set, and  $S \subset P$ .

The **supremum** of  $S$  in  $P$  is the **least** upper bound of  $S$  in  $P$ .



# Supremum

Let  $P$  be an ordered set, and  $S \subset P$ .

The **supremum** of  $S$  in  $P$  is the **least** upper bound of  $S$  in  $P$ . That is, it is the *smallest* element of  $P$  that is greater than or equal to any element of  $S$ .



# Supremum

Let  $P$  be an ordered set, and  $S \subset P$ .

The **supremum** of  $S$  in  $P$  is the **least** *upper bound* of  $S$  in  $P$ . That is, it is the *smallest* element of  $P$  that is greater than or equal to any element of  $S$ .



We write  $\sup_P(S)$  or  $\sup(S)$  when  $P$  is clear from context.

# Supremum

Let  $P$  be an ordered set, and  $S \subset P$ .

The **supremum** of  $S$  in  $P$  is the **least** upper bound of  $S$  in  $P$ . That is, it is the *smallest* element of  $P$  that is greater than or equal to any element of  $S$ .



**Note!** When the  $S$  has a maximum element,  
 $\sup(S) = \max(S)$ .



# Infimum

Let  $P$  be an ordered set, and  $S \subset P$ .

Similarly, the **infimum** of  $S$  in  $P$  is the **greatest** lower bound of  $S$  in  $P$ .



# Infimum

Let  $P$  be an ordered set, and  $S \subset P$ .

Similarly, the **infimum** of  $S$  in  $P$  is the **greatest** lower bound of  $S$  in  $P$ .



# Infimum

Let  $P$  be an ordered set, and  $S \subset P$ .

Similarly, the **infimum** of  $S$  in  $P$  is the **greatest** lower bound of  $S$  in  $P$ .



We write  $\inf_P(S)$  or  $\inf(S)$  when  $P$  is clear from context.

# Infimum

Let  $P$  be an ordered set, and  $S \subset P$ .

Similarly, the **infimum** of  $S$  in  $P$  is the **greatest** lower bound of  $S$  in  $P$ .




**Note!** When the  $S$  has a **minimum element**,  
 $\inf(S) = \min(S)$ .

# Notation


We will sometimes write  $\{x_i\}_{i=1}^{\infty}$  to denote a sequence of numbers  $x_1, x_2, \dots$

## Question?



How can we capture the idea that the upper envelope remains bounded, even though the sequence  $\{x_i\}_{i=1}^{\infty}$  doesn't have a limit?


## Question?



How can we capture the idea that the upper envelope remains bounded, even though the sequence  $\{x_i\}_{i=1}^{\infty}$  doesn't have a limit?

# Answer!


Consider the "maximum value" (**supremum**) **of the tail** region as we slide the start of the tail further and further to the right.






# Answer!

Consider the "maximum value" (**supremum**) **of the tail** region as we slide the start of the tail further and further to the right.




# Answer!

Consider the "maximum value" (**supremum**) **of the tail** region as we slide the start of the tail further and further to the right.




# Answer!

Consider the "maximum value" (**supremum**) **of the tail** region as we slide the start of the tail further and further to the right.



# Answer!

Consider the "maximum value" (**supremum**) **of the tail** region as we slide the start of the tail further and further to the right.



# Limit Superior

Let  $\{x_i\}_{i=1}^{\infty}$  be a sequence of real numbers.

The **limit superior** of  $\{x_i\}_{i=1}^{\infty}$  (denoted  $\limsup_{n \rightarrow \infty} x_n$ ) is

$$\lim_{n \rightarrow \infty} \sup(\{x_i : i \geq n\}),$$

and the **limit inferior** of  $\{x_i\}_{i=1}^{\infty}$  (denoted  $\liminf_{n \rightarrow \infty} x_n$ ) is

$$\lim_{n \rightarrow \infty} \inf(\{x_i : i \geq n\}).$$

## Theorem (Limit Theorems)

- ▲ If  $\limsup_{n \rightarrow \infty} \frac{f(n)}{g(n)} \in \mathbb{R}^+$ , then  $f \in O(g)$ .
- ▲ If  $\liminf_{n \rightarrow \infty} \frac{f(n)}{g(n)} \in \mathbb{R}^+$ , then  $f \in \Omega(g)$ .