

Statistical Test	Complete Dataset (<i>n</i> = 304) A = Female, B = Male	Constitutive Costs (<i>n</i> = 171, uninfected) A = Female, B = Male	Infection Dominance (<i>n</i> = 304, by status) A = Uninfected, B = Infected
Model Comparisons			
Model comparison: H1 vs H0	0.07936.	0.13830	0.16200
Model comparison: H2 vs H0	0.94510	0.89830	0.00211**
Model comparison: H3 vs H1	0.33710	0.75340	0.00348**
Model comparison: H3 vs H2 ¹	0.02668*	0.17300	0.22850
Individual Hypothesis Tests			
H0: Overall hybrid effect ($\alpha \neq 0$) ²	0.01742*	0.54460	0.19800
H1: Group difference in hybrid effects	0.05504.	0.75400	0.29360
H2: Group A-specific hybrid effect ³	0.18930	0.90810	0.54460
H2: Group B-specific hybrid effect ⁴	0.03808*	0.43150	0.29290
H3: Advanced Group A model	0.20790	0.90850	0.75400
H3: Advanced Group B model	0.05075.	0.35280	0.26320
Complete & Constitutive: A = Female, B = Male Infection: A = Uninfected, B = Infected with Eimeria spp.			
Significance: *** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$, . $p < 0.1$ Bold = $p < 0.05$			
Color scheme matches nested hypothesis structure: H0 H1 H2 H3			
¹ Critical comparison for detecting group-specific effects (H3 vs H2)			
² α represents deviation from additivity (positive = increased costs with M. m. musculus ancestry)			
³ Group A = Female (Complete/Constitutive) OR Uninfected (Infection analysis)			
⁴ Group B = Male (Complete/Constitutive) OR Infected (Infection analysis) - Males show significant effect ($p = 0.038^*$)			