

① Is the vector $(3, -1, 0, -1)$ in the subspace spanned by $(2, -1, 3, 2)$, $(-1, 1, 1, -3)$ & $(1, 1, 9, -5)$?

Solution:

$$\alpha_1 = (2, -1, 3, 2)$$

$$\alpha_2 = (-1, 1, 1, -3) \quad \alpha = (3, -1, 0, -1)$$

$$\alpha_3 = (1, 1, 9, -5)$$

$$\text{Let } c_1 \alpha_1 + c_2 \alpha_2 + c_3 \alpha_3 = \alpha$$

If c_1 , c_2 and c_3 exists and are unique, we can say that the vector α is in the subspace spanned by $\alpha_1, \alpha_2, \alpha_3$.

$$\therefore c_1 (2, -1, 3, 2) + c_2 (-1, 1, 1, -3) + c_3 (1, 1, 9, -5) = (3, -1, 0, -1)$$

$$2c_1 - c_2 + c_3 = 3$$

$$-c_1 + c_2 + c_3 = -1$$

$$3c_1 + c_2 + 9c_3 = 0$$

$$2c_1 - 3c_2 - 5c_3 = -1$$

$$\Rightarrow \begin{bmatrix} 2 & -1 & 1 & 3 \\ -1 & 1 & 1 & -1 \\ 3 & 1 & 9 & 0 \\ 2 & -3 & -5 & -1 \end{bmatrix}$$

Finding Row reduced matrix

$$R_1 \rightarrow \frac{R_1}{2} = \begin{bmatrix} 1 & -1/2 & 1/2 & 3/2 \\ -1 & 1 & 1 & -1 \\ 3 & 1 & 9 & 0 \\ 2 & -3 & -5 & -1 \end{bmatrix} \quad R_2 \rightarrow R_1 + R_2 = \begin{bmatrix} 1 & -1/2 & 1/2 & 3/2 \\ 0 & 1/2 & 3/2 & 1/2 \\ 3 & 1 & 9 & 0 \\ 2 & -3 & -5 & -1 \end{bmatrix}$$

$$-3R_1 + R_3 = \begin{bmatrix} 1 & -1/2 & 1/2 & 3/2 \\ 0 & 1/2 & 3/2 & 1/2 \\ 0 & 5/2 & 15/2 & -9/2 \\ 2 & -3 & -5 & -1 \end{bmatrix} \quad -2R_1 + R_4 = \begin{bmatrix} 1 & -1/2 & 1/2 & 3/2 \\ 0 & 1/2 & 3/2 & 1/2 \\ 0 & 5/2 & 15/2 & -9/2 \\ 0 & -2 & -6 & -4 \end{bmatrix}$$

$$R_2 \rightarrow 2R_2 = \begin{bmatrix} 1 & -1/2 & 1/2 & 3/2 \\ 0 & 1 & 3 & 1 \\ 0 & 5/2 & 15/2 & -9/2 \\ 0 & -2 & -6 & -4 \end{bmatrix} \quad R_3 \rightarrow \frac{-5R_2}{R_2} + R_3 = \begin{bmatrix} 1 & -1/2 & 1/2 & 3/2 \\ 0 & 1 & 3 & 1 \\ 0 & 0 & 0 & -7 \\ 0 & -2 & -6 & -4 \end{bmatrix}$$

$$R_4 \rightarrow 2R_2 + R_4 = \begin{bmatrix} 1 & -1/2 & 1/2 & 3 \\ 0 & 1 & 3 & 1 \\ 0 & 0 & 0 & -7 \\ 0 & 0 & 0 & -2 \end{bmatrix} \quad R_3 \rightarrow \frac{-R_3}{7} = \begin{bmatrix} 1 & -1/2 & 1/2 & 3/2 \\ 0 & 1 & 3 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & -2 \end{bmatrix}$$

$$R_4 \rightarrow 2R_3 + R_4 = \begin{bmatrix} 1 & -1/2 & 1/2 & 3/2 \\ 0 & 1 & 3 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad R_2 \rightarrow -R_3 + R_2 = \begin{bmatrix} 1 & -1/2 & 1/2 & 0 \\ 0 & 1 & 3 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$R_1 \rightarrow \frac{-3R_3}{2} + R_1 = \begin{bmatrix} 1 & -1/2 & 1/2 & 0 \\ 0 & 1 & 3 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$R_1 \rightarrow \frac{R_2}{2} + R_1 = \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 1 & 3 & 0 \\ 0 & 0 & \boxed{0} & \boxed{1} \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

C_1 and C_2 are basic variable whereas C_3 is Free variable.

\therefore Infinitely many solutions.

$\therefore \mathcal{L}$ is the subspace spanned by α_1, α_2 and α_3 .

② Are the vectors linearly independent in R^4 ? Find the basis for the subspace of R^4 spanned by 4 vectors.

$$\alpha_1 = (1, 1, 2, 4) \quad \alpha_2 = (2, -1, -5, 2)$$

$$\alpha_3 = (1, -1, -4, 0) \quad \alpha_4 = (2, 1, 1, 6)$$

Q: $\alpha_1, \alpha_2, \alpha_3, \alpha_4$ are said to be linearly independent if the reduced row echelon form of the matrix $[\alpha_1 \alpha_2 \alpha_3 \alpha_4]$ has only basic variables and no free variables.

$$A = \begin{bmatrix} 1 & 1 & 2 & 4 \\ 2 & -1 & -5 & 2 \\ 1 & -1 & -4 & 0 \\ 2 & 1 & 1 & 6 \end{bmatrix} \quad R_2 \rightarrow -2R_1 + R_2 = \begin{bmatrix} 1 & 1 & 2 & 4 \\ 0 & -3 & -9 & -6 \\ 1 & -1 & -4 & 0 \\ 2 & 1 & 1 & 6 \end{bmatrix}$$

$$R_3 \rightarrow -R_1 + R_3 = \begin{bmatrix} 1 & 1 & 2 & 4 \\ 0 & -3 & -9 & -6 \\ 0 & -2 & -6 & -4 \\ 2 & 1 & 1 & 6 \end{bmatrix}$$

$$R_4 \rightarrow -2R_1 + R_4 = \begin{bmatrix} 1 & 1 & 2 & 4 \\ 0 & -3 & -9 & -6 \\ 0 & -2 & -6 & -4 \\ 0 & -1 & -3 & -2 \end{bmatrix}$$

$$R_2 \rightarrow -\frac{R_2}{3} = \begin{bmatrix} 1 & 1 & 2 & 4 \\ 0 & -3 & -9 & -6 \\ 0 & -2 & -6 & -4 \\ 0 & -1 & -3 & -2 \end{bmatrix}$$

$$R_3 \rightarrow 2R_2 + R_3 = \begin{bmatrix} 1 & 1 & 2 & 4 \\ 0 & 1 & 3 & 2 \\ 0 & 0 & 0 & 0 \\ 0 & -1 & -3 & -2 \end{bmatrix}$$

$$R_4 \rightarrow R_2 + R_4 = \begin{bmatrix} 1 & 1 & 2 & 4 \\ 0 & 1 & 3 & 2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad R_1 \rightarrow -R_2 + R_1 = \begin{bmatrix} \textcircled{1} & 0 & -1 & 2 \\ 0 & \textcircled{1} & 3 & 2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Two basic variables, two free.

$\therefore \alpha_1, \alpha_2, \alpha_3, \alpha_4$ are not linearly independent \Rightarrow

α_1 and α_2 are the vectors with basic variables.

$\therefore \alpha_1 \in \alpha_2$ form the basis for the vector space \mathbb{R}^4 .

$$\text{Basis} = \left(\begin{bmatrix} 1 \\ 1 \\ 2 \\ 4 \end{bmatrix}, \begin{bmatrix} 2 \\ -1 \\ -5 \\ 2 \end{bmatrix} \right)$$

3) Let V be the vector space of polynomials of degree 3, over \mathbb{R} .
 Determine whether u, v & w are linearly independent or not.

$$u = x^3 - 3x^2 + 5x + 1$$

$$v = x^3 - x^2 + 8x + 2$$

$$w = 2x^3 - 4x^2 + 9x + \underline{\underline{5}}$$

$$\Rightarrow \begin{bmatrix} 1 & -3 & 5 & 1 \\ 1 & -1 & 8 & 2 \\ 2 & -4 & 9 & 5 \end{bmatrix}$$

$$R_2 \rightarrow -R_1 + R_2 = \begin{bmatrix} 1 & -3 & 5 & 1 \\ 0 & 2 & 3 & 1 \\ 2 & -4 & 9 & 5 \end{bmatrix} \quad R_3 \rightarrow -2R_1 + R_3 = \begin{bmatrix} 1 & -3 & 5 & 1 \\ 0 & 2 & 3 & 1 \\ 0 & 2 & -1 & 3 \end{bmatrix}$$

$$R_2 \rightarrow R_2/2 = \begin{bmatrix} 1 & -3 & 5 & 1 \\ 0 & 1 & \frac{3}{2} & \frac{1}{2} \\ 0 & 0 & -4 & 2 \end{bmatrix} \quad R_3 \rightarrow -\frac{R_3}{4} = \begin{bmatrix} 1 & -3 & 5 & 1 \\ 0 & 1 & \frac{3}{2} & \frac{1}{2} \\ 0 & 0 & 1 & -\frac{1}{2} \end{bmatrix}$$

$$R_2 \rightarrow -3\frac{R_3}{2} + R_2 = \begin{bmatrix} 1 & -3 & 5 & 1 \\ 0 & 1 & \frac{3}{2} & \frac{1}{2} \\ 0 & 0 & 1 & -\frac{1}{2} \end{bmatrix}$$

$$-5R_3 + 3R_2 + 2R_1 \rightarrow R_1 = \begin{bmatrix} 1 & 0 & 0 & \frac{29}{4} \\ 0 & 1 & 0 & 5/4 \\ 0 & 0 & 1 & -1/2 \end{bmatrix} \quad \text{3 basic variables.}$$

$\therefore u, v$ & w are linearly Independent.

$$A = \begin{pmatrix} A1 & B1 & C1 & D1 \\ A2 & B2 & C2 & D2 \\ A3 & B3 & C3 & D3 \\ A4 & B4 & C4 & D4 \end{pmatrix}$$

$$\rightarrow \begin{pmatrix} 2 & -1 & 1 & 3 \\ -1 & 1 & 1 & -1 \\ 3 & 1 & 9 & 0 \\ 2 & -3 & -5 & -1 \end{pmatrix}$$

m1 = ReducedRowEchelonForm(A)

$$\rightarrow \begin{pmatrix} 1 & 0 & 2 & 0 \\ 0 & 1 & 3 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$



No Solutions. There fore the vector (3,-1,0,-1) doesnt exists in the subspace spanned by $\alpha^1, \alpha^2, \alpha^3$

	A	B	C	D
1	2	-1	1	3
2	-1	1	1	-1
3	3	1	9	0
4	2	-3	-5	-1
5				
6	Question 1			
7				
8				
9				
10				
11				
12				
13				
14				
15				
16				

$$m1 = \begin{pmatrix} A1 & B1 & C1 & D1 \\ A2 & B2 & C2 & D2 \\ A3 & B3 & C3 & D3 \\ A4 & B4 & C4 & D4 \end{pmatrix}$$

$$\rightarrow \begin{pmatrix} 1 & 1 & 2 & 4 \\ 2 & -1 & -5 & 2 \\ 1 & -1 & -4 & 0 \\ 2 & 1 & 1 & 6 \end{pmatrix}$$

$$m2 = \text{ReducedRowEchelonForm}(m1)$$

$$\rightarrow \begin{pmatrix} 1 & 0 & -1 & 2 \\ 0 & 1 & 3 & 2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

$$m3 = \begin{pmatrix} A7 & B7 & C7 & D7 \\ A8 & B8 & C8 & D8 \end{pmatrix}$$

$$\rightarrow \begin{pmatrix} 1 & 1 & 2 & 4 \\ 2 & -1 & -5 & 2 \end{pmatrix}$$

$$l1 = \text{Dimension}(m3)$$

$$\rightarrow \{2, 4\}$$

$$a = \text{MatrixRank}(m3)$$

$$\rightarrow 2$$

	A	B	C	D
1	1	1	2	4
2	2	-1	-5	2
3	1	-1	-4	0
4	2	1	1	6
5				
6				
7	1	1	2	4
8	2	-1	-5	2
9				
10	Question 2			
11				
12				
13				
14				
15				
16				
17				
18				
19				
20				
21				
22				
23				
24				
25				



$w(t) = 2t^3 - 4t^2 + 9t + 5$



$v(t) = t^3 - t^2 + 8t + 2$



$u(t) = t^3 - 3t^2 + 5t + 1$



$A = \begin{pmatrix} A1 & B1 & C1 & D1 \\ A2 & B2 & C2 & D2 \\ A3 & B3 & C3 & D3 \end{pmatrix}$

$\rightarrow \begin{pmatrix} 1 & -3 & 5 & 1 \\ 1 & -1 & 8 & 2 \\ 2 & -4 & 9 & 5 \end{pmatrix}$

$m1 = \text{ReducedRowEchelonForm}(A)$

$\rightarrow \begin{pmatrix} 1 & 0 & 0 & 7.25 \\ 0 & 1 & 0 & 1.25 \\ 0 & 0 & 1 & -0.5 \end{pmatrix}$



$U = 7.25$




$V = 1.25$



$W = -0.5$



Number of free variables = 0



	A	B	C
1	1	-3	5
2	1	-1	8
3	2	-4	9
4			
5	Question 3		
6			
7			
8			
9			
10			
11			
12			
13			
14			
15			
16			
17			
18			
19			
20			
21			
22			
23			
24			
25			
26			
27			
28			
29			