

[Home Page](#)

[Title Page](#)



Page 1 of 100

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)

EC540 Control Systems

Recalling Signals and Systems

(Dr. S. Patilkulkarni, 4/09/2021)



Home Page

Title Page



Page 2 of 100

Go Back

Full Screen

Close

Quit

Instructions:

1. Lecture session will be of one hour duration.
2. Fifteen minutes session will be made available for students after the lecture for any question and answers.
3. Regularly review Signals and Systems concepts
4. Regularly visit course webpage.
5. Everyday learn new functions from Octave/Python/MATLAB software
6. Email me on any queries at sudarshan_pk@sjce.ac.in

1. Basics of Signals

Signal is any *useful* information about a physical variable such as temperature, light intensity, sound level, stock market index etc with respect to independent variable time.

Signal is mathematically represented as function $f(t)$ where $-\infty < t < \infty$ indicates time.

Causal Signal: Signal having $f(t) = 0; t < 0$


Periodic Signal: Signal having $f(t) = f(t + nT)$ where n is any integer. $F_0 = \frac{1}{T}$ is called fundamental frequency of signal.

Standard functions:

Impulse Signal:

$$x_p(t) = \frac{1}{\tau} \quad \text{for } -\frac{\tau}{2} < t < \frac{\tau}{2}$$

$$\delta(t) = \lim_{\tau \rightarrow 0} x_p(t).$$



Standard functions:

Step Signal:

$$f(t) = u_s(t) = \begin{cases} 1 & t \geq 0, \\ 0 & \text{otherwise} \end{cases}$$

Ramp Signal:

$$f(t) = \begin{cases} t & t \geq 0, \\ 0 & \text{otherwise} \end{cases}$$

Exponential Signal:

$$f(t) = \begin{cases} e^{at} & t \geq 0, \\ 0 & \text{otherwise} \end{cases}$$

At $t = 0$ function value is 1. If $a > 0$ then function value increases exponentially; If $a < 0$ then function value decreases exponentially.

Standard functions:

Sin Signal:

$$f(t) = \begin{cases} \sin(\omega_0 t) & t \geq 0, \\ 0 & \text{otherwise} \end{cases}$$

cos Signal:

$$f(t) = \begin{cases} \cos(\omega_0 t) & t \geq 0, \\ 0 & \text{otherwise} \end{cases}$$

Complex Exponential Signal:

$$f(t) = \begin{cases} e^{j\omega_0 t} & t \geq 0, \\ 0 & \text{otherwise} \end{cases}$$

$$e^{j\omega_0 t} = \cos(\omega_0 t) + j \sin(\omega_0 t)$$

[Home Page](#)

[Title Page](#)

« »

◀ ▶

Page 7 of 100

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)

Any signal/function $f(t)$ can be represented using the fundamental function of time domain $\delta(t)$

$$f(t) = \int f(\tau)\delta(t - \tau)d\tau.$$

Any signal/function $f(t)$ can be represented using the fundamental function of frequency domain $e^{j\omega t}$

Fourier Series

For periodic signals

$$f(t) = \sum_{-\infty}^{\infty} c_k e^{j2\pi k F_0 t} = a_0 + \sum_{k=1}^{\infty} \{a_k \cos(2\pi k F_0 t) - b_k \sin(2\pi k F_0 t)\}$$

$$c_k = \frac{1}{T} \int_0^T f(t) e^{-j2\pi k F_0 t} dt.$$

Alternatively,

$$a_0 = \frac{1}{T} \int_0^T f(t) dt$$

$$a_k = \frac{2}{T} \int_0^T f(t) \cos(2\pi k F_0 t) dt \text{ for } k = 1, 2, \dots$$

$$b_k = \frac{2}{T} \int_0^T f(t) \sin(2\pi k F_0 t) dt \text{ for } k = 1, 2, \dots$$

Home Page

Title Page

◀◀

▶▶

◀

▶

Page 9 of 100

Go Back

Full Screen

Close

Quit

Fourier Transform: For aperiodic signals,

$$F(j\omega) = \mathcal{F}\{f(t)\} = \int f(t)e^{-j\omega t} dt$$

Inverse Fourier Transform:

$$f(t) = \frac{1}{2\pi} \int F(j\omega)e^{j\omega t} d\omega$$

here $-\infty < \omega < \infty$

Definition of Laplace Transform: For any signal,

$$F(s) = \int f(t)e^{-st}dt \quad \text{where } s = \sigma + j\omega$$

Region of Convergence Values of σ for which integral converges.

$$F(s) = \mathcal{F}\{f(t).e^{-\sigma t}\}$$

$$F(j\omega) = F(s) \text{ when } \sigma = 0$$

Laplace Transform of Standard Functions:

$\mathbf{f(t)}$	$\mathbf{F(s)}$
$\delta(t)$	1
$u_s(t)$	$\frac{1}{s}$
$tu_s(t)$	$\frac{1}{s^2}$
$\frac{t^2}{2}.u_s(t)$	$\frac{1}{s^3}$
$e^{at}.u_s(t)$	$\frac{1}{s-a}$
$t.e^{-at}.u_s(t)$	$\frac{1}{(s+a)^2}$
$\sin \omega_0 t.u_s(t)$	$\frac{\omega_0}{s^2+\omega_0^2}$
$\cos \omega_0 t.u_s(t)$	$\frac{s}{s^2+\omega_0^2}$

Home Page

Title Page



Page 11 of 100

Go Back

Full Screen

Close

Quit

Properties of Laplace Transform:

1. $\mathcal{L}\{\alpha f_1(t) + \beta f_2(t)\} = \alpha F_1(s) + \beta F_2(s)$. Principle of superposition
2. $\mathcal{L}\{f(t - T)\} = F(s)e^{-sT}$
3. $\mathcal{L}\{f(t)e^{at}\} = F(s - a)$;
4. $\mathcal{L}\{f(at)\} = \frac{1}{|a|}F\left(\frac{s}{a}\right)$

$$5. \mathcal{L}\{tf(t)\} = -\frac{dF(s)}{ds}$$

$$6. \mathcal{L}\left\{\frac{d^n f(t)}{dt^n}\right\} = s^n F(s) - s^{n-1}f(0) - s^{n-2}f'(0) - \dots - f^{n-1}(0)$$

$$7. \mathcal{L}\left\{\int f_1(\tau)f_2(t-\tau)d\tau\right\} = F_1(s)F_2(s)$$

$$8. \lim_{t \rightarrow 0} f(t) = \lim_{s \rightarrow \infty} sF(s)$$


$$9. \lim_{t \rightarrow \infty} f(t) = \lim_{s \rightarrow 0} sF(s)$$

2. Basics of Systems

System

$$y(t) = \mathcal{T}\{u(t)\}$$

transforms the input signal $u(t)$ to an output signal based on the operator \mathcal{T}



Home Page

Title Page



Page 15 of 100

Go Back

Full Screen

Close

Quit

Linear System

$$y(t) = \mathcal{T}\{u(t)\}$$

obeys principle of superposition:

$$\begin{aligned} y(t) &= \mathcal{T}\{u(t)\} \\ &= \mathcal{T}\{a_1 u_1(t) + a_2 u_2(t)\} \\ &= a_1 \mathcal{T}\{u_1(t)\} + a_2 \mathcal{T}\{u_2(t)\} \\ &= a_1 y_1(t) + a_2 y_2(t) \end{aligned}$$

Home Page

Title Page

◀◀

▶▶

◀

▶

Page 16 of 100

Go Back

Full Screen

Close

Quit

Example: $y(t) = Au(t)$ is linear system.

$$y_1(t) = \mathcal{T}\{u_1(t)\} = Au_1(t)$$

$$y_2(t) = \mathcal{T}\{u_2(t)\} = Au_2(t)$$

$$\begin{aligned} y(t) &= \mathcal{T}\{a_1u_1(t) + a_2u_2(t)\} = A\{a_1u_1(t) + a_2u_2(t)\} \\ &= a_1Au_1(t) + a_2Au_2(t) \\ &= a_1y_1(t) + a_2y_2(t) \end{aligned}$$

Example: $y(t) = A \frac{du(t)}{dt}$ is linear system.

$$y_1(t) = \mathcal{T}\{u_1(t)\} = A \frac{du_1(t)}{dt}$$

$$y_2(t) = \mathcal{T}\{u_2(t)\} = A \frac{du_2(t)}{dt}$$

$$\begin{aligned} y(t) &= \mathcal{T}\{a_1 u_1(t) + a_2 u_2(t)\} = A \frac{d\{a_1 u_1(t) + a_2 u_2(t)\}}{dt} \\ &= a_1 A \frac{du_1(t)}{dt} + a_2 A \frac{du_2(t)}{dt} \\ &= a_1 y_1(t) + a_2 y_2(t) \end{aligned}$$

Example:

$y(t) = \mathcal{T}\{u^2(t)\}$ is a Nonlinear system

$$y_1(t) = \mathcal{T}\{u_1(t)\} = u_1^2(t)$$

$$y_2(t) = \mathcal{T}\{u_2(t)\} = u_2^2(t)$$

$$\begin{aligned} y(t) &= \mathcal{T}\{a_1 u_1(t) + a_2 u_2(t)\} = (a_1 u_1(t) + a_2 u_2(t))^2 \\ &= a_1^2 u_1^2(t) + a_2^2 u_2^2(t) + 2a_1 a_2 u_1(t) u_2(t) \\ &\neq a_1 y_1(t) + a_2 y_2(t). \end{aligned}$$

Home Page

Title Page

◀◀

▶▶

◀

▶

Page 19 of 100

Go Back

Full Screen

Close

Quit

Example: Is $y(t) = mu(t) + c$ a linear system?


$$y_1(t) = \mathcal{T}\{u_1(t)\} = mu_1(t) + c$$

$$y_2(t) = \mathcal{T}\{u_2(t)\} = mu_2(t) + c$$

$$\begin{aligned} y(t) &= \mathcal{T}\{a_1u_1(t) + a_2u_2(t)\} \\ &= m(a_1u_1(t) + a_2u_2(t)) + c \\ &\neq a_1y_1(t) + a_2y_2(t). \end{aligned}$$

Time Varying/Invariant System

$$\begin{aligned}y(t) &= \mathcal{T}\{u(t)\} \\y_2(t) &= \mathcal{T}\{u(t - T)\} \\&\neq y(t - T) \text{ Time varying system} \\&= y(t - T) \text{ Time invariant system}\end{aligned}$$



Example: $y(t) = \mathcal{T}\{u(t)\} = tu(t)$ is time varying system.

$$y_2(t) = \mathcal{T}\{u(t - T)\} = tu(t - T)$$

$$y_2(t) \neq y(t - T) = (t - T)u(t - T).$$

$y(t) = \mathcal{T}\{u(t)\} = A \frac{du(t)}{dt}$ and $y(t) = \mathcal{T}\{u(t)\} = mu(t) + c$ are examples of Time invariant systems.

Home Page

Title Page



Page 22 of 100

Go Back

Full Screen

Close

Quit

Linear Time-Invariant System (LTI) satisfies both linearity and timeinvariance property.

$y(t) = Au(t)$, $y(t) = A\frac{d^nu(t)}{dt^n}$, $y(t) = \int Au(\tau)d\tau$ are examples.

Any linear combinations of above examples are also LTI systems.

Impulse Response and Convolution Let us denote $g(t) = \mathcal{T}\{\delta(t)\}$; is called impulse response of the system.

$$\begin{aligned} y(t) &= \mathcal{T}\{u(t)\} \\ &= \mathcal{T}\left\{\int u(\tau)\delta(t-\tau)d\tau\right\} \\ &= \int u(\tau)\mathcal{T}\{\delta(t-\tau)\}d\tau \quad \text{due to linearity} \\ &= \int u(\tau)g(t-\tau)d\tau \quad \text{due to time invariance.} \end{aligned}$$

is called convolution integral.

Contents

Home Page

Title Page

◀◀

▶▶

◀

▶

Page 24 of 100

Go Back

Full Screen

Close

Quit

$$a_0 \frac{d^n y(t)}{dt^n} + a_1 \frac{d^{n-1} y(t)}{dt^{n-1}} + \dots + a_n y(t) = b_0 \frac{d^m u(t)}{dt^m} + \dots + b_m$$

$$(a_0 s^n + a_1 s^{n-1} + \dots + a_n) Y(s) = (b_0 s^m + b_1 s^{m-1} + \dots + b_m)$$

$$G(s) = \frac{Y(s)}{U(s)} = \frac{b_0 s^m + b_1 s^{m-1} + \dots + b_m}{a_0 s^n + a_1 s^{n-1} + \dots + a_n}.$$

Home Page

Title Page

◀◀

▶▶

◀

▶

Page 25 of 100

Go Back

Full Screen

Close

Quit

$$G(s) = \mathcal{L}\{g(t)\}; \quad Y(s) = \mathcal{L}\{y(t)\} \quad U(s) = \mathcal{L}\{u(t)\}$$

$$\text{Since } y(t) = \int u(\tau)g(t - \tau)d\tau$$

By property of laplace transform of convolution integral:

$$Y(s) = G(s)U(s)$$

