



Spring 2021

SKKU Biostats and Big data

CHOONG-WAN WOO | COCOAN lab | <http://cocoanlab.github.io>



Lecture 04

Comparing distributions, Normal model

Key Points

Chapter 3: Displaying categorical data

- Bar chart for categorical data
- Pie chart for proportions of whole
- Faithful reporting and the area principle
- Contingency tables
- Simpson's paradox

Chapter 4: Displaying quantitative data

- Histograms, Stem-leaf, dot plots
- Shape (mode, symmetrical)
- Center (median, mean)
- Spread (range, IQR, variance, standard deviation)
- Box plots




Q&A

- What could be the difference between 학원 인강 vs. 대학교 online 강의?
- 1. 저는 여러 분야(과목)을 좋아하는 성격이어서 어떤 연구를 해야겠다라는 명확한 생각이 들지 않는데 혹시 교수님께서는 지금의 연구를 하게 되신 동력이 있으신가요? (석사를 정하는 것에 있어 고민이 많이 되는데 어떤 분야를 해야하는지 아직 잘 모르겠습니다. 연구 경험을 통해 파악해볼 수 있는 것인지 궁금합니다)
- 2. 교수님께서 해외에 나가셔서 박사 공부를 하신 것으로 알고 있는데 혹시 해외에 나가신 계기가 따로 있으신지 궁금합니다!

Comparing Groups with Histogram and Boxplots

Data:

- When: Days during 2011
- What: Average daily wind speed (mph)
- Where: Hopkins Forest in western Massachusetts
- Why: Long-term observations to study ecology and climate



Outliers

- Outliers may be the most important values.
- Or they maybe just errors.
- What to do with them?
 1. Correct them if possible
 2. Report summaries and analyses with and without the outliers (readers can decide)
 3. Some statistical methods: down-weight them (e.g., robust regression), smoothing, etc.
 4. *Never* do:
 - Leave them in place and proceed as if nothing were unusual
 - Omit an outlier from the analysis without comment

Re-expressing Data

- To improve symmetry:





FIGURE 5.9

Compensation paid to CEOs of the Fortune 500 companies in 2005. The distribution seems to have outliers. But maybe it just has a long tail on the right.


Logarithms of 2005 CEO compensations




- Very common in physio data preprocessing
 - E.g., logarithm, square root (normalization)

Re-expressing Data

- To equalize spread across groups





*ETS: exposed to smoke



- Normalization** is one of the key elements of recent successes of artificial intelligence (machine learning)



Timeplots

- Values against time: **timeplot**, **time-series data** (e.g., neural activity data, stock market, bio-sensor...)



Smoothing timeplots

- Lowess (locally weighted scatterplot smoothing)



Key Points

Chapter 5: Comparing distributions: considerations

- Outliers are context dependent
- Re-expressing data (log, sqrt)
 - to improve symmetry
 - to equalize spread
- Timeplots
- Moving-averages, smoothing

z-Scores






- To compare different values in different units,
- the values should be *standardized!*

$$z = \frac{y - \bar{y}}{s}$$

- z-scores: mean = 0, standard deviation = 1
 - standardized values
 - = using standard deviation as a **ruler!**
-
- Two elements: **shifting** and **scaling**

Shifting and rescaling

WHO	80 male participants of the NHANES survey between the ages of 19 and 24 who measured between 68 and 70 inches tall
WHAT	Their weights
UNIT	Kilograms
WHEN	2001-2002
WHERE	United States
WHY	To study nutrition, and health issues and trends
HOW	National survey



Quiz!

<https://forms.gle/BapezhpVbkebQ9mL7>

Normal models: When is a z-score big?

- Let's say you've got a z-score of 3. How surprising your observation is?
- To answer this question, you need a *model* of your data's distribution.
- “*All models are wrong, but some are useful.*” George Box
- Most popular model: **Normal models** (bell-shaped curves)
 - unimodal, symmetric
 - $N(\mu, \sigma)$, where μ is mean, σ is standard deviation
 - Why Greek? These are parameters of the *model*, not *numerical summaries* of data
 - Numerical summaries of the data: \bar{y} , and s
 - We still call the standardized value a *z-score*.
 - $z \sim N(0, 1)$: standard Normal model

$$z = \frac{y - \bar{y}}{s} \longrightarrow z = \frac{y - \mu}{\sigma}$$

Normality assumption


- All models make **assumptions**, which should be carefully examined.
- **Nearly normal condition** (it's sufficient):
 - shape: unimodal, symmetric
 - We can check it with *histogram* or a *normal probability plot*.

Quiz!

<https://forms.gle/ZH8WLse8K5D7gy1G9>


The 68-95-99.7 Rule

- 68% of the data fall within 1 standard deviation, 95% within 2 std, 99.7% within 3 std.
- z-score = 1 means, you are 84%! Why? $100 - (50 - 68/2) = 84\%$



Finding Normal percentiles

- A table of Normal percentiles (Table Z in Appendix D)



- Or google it!
- p-to-z is same: Table Z again.

Google Search

All Videos Images Maps News More Settings Tools

About 102,000,000 results (0.63 seconds)



z to P Calculator - VassarStats
www.vassarstats.net/textbook/ch6apx.html ▾
 z to P Calculator. For any given value of z, this page will calculate the respective one-tailed probabilities of $-z$ and $+z$; the two-tailed probability of $\pm z$; and the ...

z to P - VassarStats
vassarstats.net/tabs_z.html ▾
 the respective one-tailed probabilities of $-z$ and $+z$; the two-tailed probability of $\pm z$; and the proportion of the normal distribution falling between $-z$ and $+z$.

Statistical Tables Calculator - VassarStats

Normal probability plots

- Original value against normal scores (theoretically expected value):
- Similar to Q-Q plot (quantile-quantile plot)



Procedure

1. Sorting (draw a histogram)
2. Get percentile
3. P-to-Z
4. Make a scatter plot

y-axis = Original values

x-axis = Normal scores

Example

1. Sorting (draw a histogram)
2. Get percentile
3. P-to-Z
4. Make a scatter plot

y-axis = Original values

x-axis = Normal scores

Data

11.4 13.6 10.2 11.7 16.3 16.2 7.2 11.2 17.7 10.8 10.2 17.1 14.5 6.7 10.0 4.4 8.8 14.5 3.6 15.1



1. Sorting

1 ~ 12번 열

3.6000 4.4000 6.7000 7.2000 8.8000 10.0000 10.2000 10.2000 10.8000 11.2000 11.4000 11.7000

13 ~ 20번 열

13.6000 14.5000 14.5000 15.1000 16.2000 16.3000 17.1000 17.7000



2. Percentile,

If there are N numbers, percentile can be calculated as $100*((1-0.5)/N), 100*((2-0.5)/N), 100*((3-0.5)/N), \dots, 100*((N-0.5)/N)$

1 ~ 12번 열

Sorted value	3.6000	4.4000	6.7000	7.2000	8.8000	10.0000	10.2000	10.2000	10.8000	11.2000	11.4000	11.7000
Percentile	2.5000	7.5000	12.5000	17.5000	22.5000	27.5000	32.5000	37.5000	42.5000	47.5000	52.5000	57.5000

13 ~ 20번 열

13.6000	14.5000	14.5000	15.1000	16.2000	16.3000	17.1000	17.7000
62.5000	67.5000	72.5000	77.5000	82.5000	87.5000	92.5000	97.5000

Example

2. Percentile,

1. Sorting (draw a histogram)

If there are N numbers, percentile can be calculated as $100*((1-0.5)/N), 100*((2-0.5)/N), 100*((3-0.5)/N), \dots, 100*((N-0.5)/N)$

2. Get percentile

1 ~ 12번 열

3. P-to-Z

Sorted value	3.6000	4.4000	6.7000	7.2000	8.8000	10.0000	10.2000	10.2000	10.8000	11.2000	11.4000	11.7000
Percentile	2.5000	7.5000	12.5000	17.5000	22.5000	27.5000	32.5000	37.5000	42.5000	47.5000	52.5000	57.5000

4. Make a scatter plot

y-axis = Original values

13 ~ 20번 열

x-axis = Normal scores

13.6000	14.5000	14.5000	15.1000	16.2000	16.3000	17.1000	17.7000
62.5000	67.5000	72.5000	77.5000	82.5000	87.5000	92.5000	97.5000



3. P-to-Z

1 ~ 12번 열

Sorted value	3.6000	4.4000	6.7000	7.2000	8.8000	10.0000	10.2000	10.2000	10.8000	11.2000	11.4000	11.7000
Z-values	-1.9600	-1.4395	-1.1503	-0.9346	-0.7554	-0.5978	-0.4538	-0.3186	-0.1891	-0.0627	0.0627	0.1891

13 ~ 20번 열

13.6000	14.5000	14.5000	15.1000	16.2000	16.3000	17.1000	17.7000
0.3186	0.4538	0.5978	0.7554	0.9346	1.1503	1.4395	1.9600

Example

3. P-to-Z

1. Sorting (draw a histogram)

1 ~ 12번 열

2. Get percentile

	Sorted value	3.6000	4.4000	6.7000	7.2000	8.8000	10.0000	10.2000	10.2000	10.8000	11.2000	11.4000	11.7000
	Z-values	-1.9600	-1.4395	-1.1503	-0.9346	-0.7554	-0.5978	-0.4538	-0.3186	-0.1891	-0.0627	0.0627	0.1891

3. P-to-Z

13 ~ 20번 열

4. Make a scatter plot



	13.6000	14.5000	14.5000	15.1000	16.2000	16.3000	17.1000	17.7000
	0.3186	0.4538	0.5978	0.7554	0.9346	1.1503	1.4395	1.9600

y-axis = Original values



x-axis = Normal scores



4. Make a scatter plot (and also histogram)



Team 02 and 07



Quiz!

<https://forms.gle/wuVq3hz91B8vSM518>

Key Points

Chapter 5: Comparing distributions: considerations

- Outliers are context dependent
- Re-expressing data (log, sqrt)
 - to improve symmetry
 - to equalize spread
- Timeplots
- Moving-averages, smoothing

Chapter 6: Normal model

- z-score, shifting and rescaling
- Normal model
- Normality assumption; unimodal, symmetric
- 68-95-99.7 Rule
- z-to-p, p-to-z
- Normal probability plots