

# Looking for compensation at multiple scales in a wetland bird community

Frédéric Barraquand<sup>1,2,\*</sup>, Coralie Picoche<sup>1</sup>, Christelle Aluome<sup>1,3</sup>,  
Laure Carassou<sup>1,4</sup>& Claude Feigné<sup>5</sup>

October 17, 2018

<sup>1</sup> University of Bordeaux, Integrative and Theoretical Ecology, LabEx  
COTE Bât. B2 - Allée Geoffroy St-Hilaire, 33615 Pessac, France

<sup>2</sup> CNRS, Institute of Mathematics of Bordeaux 351 Cours de la Libération,  
33405 Talence, France

<sup>3</sup> ISPA, Bordeaux Sciences Agro, INRA, 33140, Villenave d'Ornon, France

<sup>4</sup> Irstea, UR EABX, 50 Avenue de Verdun, 33612 Cestas, France

<sup>5</sup> PNR Landes Gascognes, Teich Ornithological Reserve, Rue du Port BP  
11 33470 Le Teich, France

\* Corresponding author. Email: frederic.barraquand@u-bordeaux.fr

## Abstract

Compensatory dynamics, during which community composition shifts despite a near-constant total community size, are usually rare: synchronous dynamics prevail in natural communities. This is a puzzle for ecologists, because of the key role of compensation to explain the relation between biodiversity and ecosystem functioning. We take advantage of a long-term wetland bird community time series of 35 years, where we suspected that compensation might occur due to changes in water levels and known trends. We find that compensatory dynamics are still rare, likely due to the synchronizing influence of climate on birds, even after considering several temporal scales of covariation (during cold or warm seasons, above or below the seasonal scale). Negative covariation in abundance at the whole community level did only appear after a management change in the reserve, and at the scale of a few months or several years. Although most research has focused on the temporal scale of compensation vs. synchrony, we found that compensation varies with taxonomic and functional scale too: compensation appeared more frequently between rather than within guilds. This suggests that compensation has more potential to emerge between broad functional groups rather than between species.

## Introduction

Ecological theory suggests that within rich communities, where a number of species can have similar functions due to their proximity in morphological or phylogenetic space, they might exhibit compensatory dynamics (Gonzalez and Loreau, 2009). Compensation occurs when individuals of some species replace individuals of other species, either because of explicit competitive processes or shifts in some environmental driver that change selection pressures. This is particularly likely to occur when there is a space or resource constraint combined with temporal environmental variability. Which species “win” at any particular point in time may then depend on the fine-grained temporal environmental variation, or just on priority effects (i.e., who gets there first). Whatever the cause of compensatory dynamics, its main consequences for ecosystem functioning is that the community as a whole exhibits lower biomass variation than its constituent species (Gross et al., 2013). Compensation is therefore intertwined with community-level stability, at least when stability is understood as the reciprocal of variability. By contrast, another very frequent observed outcome on biodiversity time series is synchrony (Bush et al., 2017; Usinowicz et al., 2017). Synchrony occurs when all species fluctuate in phase, and therefore the biomass of the community may not fluctuate less than its constituent parts.

Early investigations of the frequency of synchronous vs. compensatory dynamics focused on the variance ratio, that is, the variance of the sum of the community biomass divided by the sum of the variance of the component species biomasses (Houlahan et al., 2007; Gonzalez and Loreau, 2009). Unfortunately, this metric is not appropriate for communities subjected to community-wide environmental forcing (Ranta et al., 2008), because a main environmental driver (e.g., temperature or light) may synchronize species abundances or growth rates at some scale, creating large variance in community-wide biomass, in spite of strongly competitive dynamics. Further research has therefore focused on specific timeframes where compensatory dynamics may be found (e.g., below the seasonal scale where temperature fluctuations

tend to synchronize species dynamics Vasseur et al., 2014).

Despite this effort to look for more meaningful temporal scales in community-level time series, temporal compensation has remained surprisingly elusive in the field (Houlihan et al., 2007; Vasseur et al., 2014); but see Ernest et al. (2008); Christensen et al. (2018). Most datasets used so far to evaluate temporal compensation vs synchrony involve planktonic organisms (Vasseur and Gaedke, 2007; Vasseur et al., 2014) or terrestrial plants (Houlihan et al., 2007; Gross et al., 2013; though see Bell et al. 2014. Here, we take advantage of a long-term bird time series record at the monthly scale (over 35 years <sup>1</sup>), in a natural reserve, that allows us to dig deeper into patterns of synchrony, at several temporal and taxonomic or functional scales.

Indeed, taxonomic scale should be a main modulator of synchrony/compensation, an explanatory factor that has been somewhat neglected for now. On the one hand, one could argue that compensation should be higher between closely related species, because functional and phylogenetic differences are generally correlated. For example, if species A and B are two duck species that share almost the same food niche as well as many traits, it makes little difference to the rest of the community whether one species gets replaced by the other (functional compensation, *sensu* Gonzalez and Loreau 2009), and priority effects could dominate. On the other hand, it could be argued that these two similar duck species will precisely respond in similar ways to environmental variables, which tends to obfuscate compensation. Under the latter scenario, more dissimilar species or groups of species - within the same trophic level nonetheless - could compensate each other within the whole community, precisely because they have different environmental preferences and the environment varies over time (e.g., groups of species preferring more open vs more closed habitats replacing each other as a function of changes in vegetation height). Surprisingly, compensation between guilds has been less well explored than within guilds, even though there is actually some empirical

---

<sup>1</sup>Considering only the years used in our analysis. Otherwise, the dataset begins in 1973.

evidence for compensation between dissimilar guilds (Sinclair et al., 2013). We therefore explore different ways to cluster the bird community, within or between guilds, along either taxonomic or functional classifications.

Our objective is therefore to examine how synchronous or compensatory bird communities are at different temporal and taxonomic (or functional) scales. Our dataset is ideally suited to the task given that (i) it is a highly temporally resolved time series with respect to the species typical generation times, but it also extends well beyond generation time (35 years) and (ii) the reserve where the data has been collected was subjected to a major management change c. 2006 (change in water levels), favouring different types of wetland birds (so that over long timescales, there is a real potential for changes in community composition).

## Material and Methods

### Data

The monthly time series used for the statistical analyses have been collected at the Teich Ornithological Reserve, Arcachon Bay, France (44.64°N / -1.02°E), by the staff of the Teich reserve. The reserve comprises 120 ha of wetlands, and the data have been aggregated at the reserve scale. We use for each species the maximum observed abundance over a month, which provides a “monthly snapshot” of the bird abundance. In the statistical analyses, we use both the original monthly data and seasonal averages. We defined two seasons based on observations of bird presence. We defined a ‘warm season’, from May to August, and a ‘cold season’ as the months between November and February of the following year. From an ecological viewpoint, this seasonal classification separates wintering birds from summer residents (some of whom are breeding). This makes sense biologically because the two communities have different requirements and respond differentially to abiotic drivers. It is also useful from a more statistical perspective, as there is a shift in composition between the seasons, though

winter and summer communities partially overlap due to a number of shared species.

Fig. 1 shows the patterns in abundance for key groups in the Teich reserve bird community, showing the marked signature of seasonality.




Figure 1: Time series of counts for ducks of the genus *Anas* (a), calidrids (b, *Calidris* genus), and all waders (c, including calidrids). The solid black lines represent trends in summed abundances for each guild, thin dotted lines represent putative trends (when some species are absent). The coloured symbols below the curves represent each species abundances, with species composition on the right side on the donut plots for the most abundant species (over 1% of relative abundance in the group considered).

## Statistical Analyses

We used for yearly analyses the synchrony index  $\eta$  defined by Gross et al. (2013), which is constructed as the mean cross-correlation between each species biomass and the summed

99 biomasses of the rest of the community (eq. 1).

$$\eta = \frac{1}{n} \sum_i \text{Corr}(P_i, \sum_{j \neq i} P_j) \quad (1)$$

100 where  $P_i$  is the abundance or biomass of species  $i$  in a community of  $n$  species. This  
101 synchrony index described in eq. 1 varies between -1 (perfect compensation, total biomass  
102 is constant) and 1 (complete synchrony), while 0 represents a case where all populations  
103 fluctuate independently. Contrary to other indices (e.g., Loreau and de Mazancourt 2008's  
104  $\phi$ ), this index is independent from the richness  $n$  of the community (or more generally the  
105 number of system components) and its overall stability (Blüthgen et al., 2016; Hallett et al.,  
106 2016). This is particularly important here as we perform analyses at different taxonomic  
107 scales, and therefore with a different  $n$  in eq. 1.

108 We computed synchrony indices at the year  $\times$  season scale (that is, for a given cold or  
109 warm season each year) using the *codyn* package in R (Hallett et al., 2016). We averaged  
110 monthly bird abundances, for each species, over the season duration, and computed the  
111 synchrony index using the year as our statistical unit. We also differentiated periods before  
112 and after 2006, given that a management change occurred within the reserve in 2006. We  
113 considered both the synchrony inside a given group (e.g., among species of the *Anas* genus)  
114 or between groups (e.g., between the summed abundances of the 9 species of genus *Anas*  
115 and the sum of the 12 *Calidris* species). In the latter case of between-groups comparisons,  
116 we summed species together before seasonal averaging, to consider seasonal averages of the  
117 monthly group density.

118 We used both taxonomic classifications of the species (i.e., between and within genera)  
119 and functional classifications of the species (e.g., 30 species of waders versus 34 species of  
120 ducks) as we suspected that a functional classification may allow to partition better the  
121 abiotic requirements of the species. We use “duck” as a shorthand for the larger functional

group of herbivorous divers, because the birds in that category are mostly ducks: this group includes nonetheless all anatids (geese and swans in particular) as well as the common coot (*Fulica atra*, an abundant species here).

We also “zoomed in” on a group of species that were known to exhibit potentially compensatory dynamics (through competition for roosting sites): the great cormorant (*Phalacrocorax carbo*), the little egret (*Egretta garzetta*) and the grey heron (*Ardea cinerea*). The little egret and the grey heron abundances were summed because of their similar requirements (i.e., they form a small functional group).

We computed statistical significance of synchrony index values using Monte Carlo randomizations (Gouhier and Guichard, 2014). For each set of time series (each combination year  $\times$  season), we kept the auto-correlation of the species time-series, but removed the cross-correlation between species by shifting each time series by a random lag (Purves and Law, 2002). We obtained 100 sets of randomized time series for each season and period of time considered and computed the corresponding synchrony index. We then compared the observed values of  $\eta$  to the values obtained with the randomized time-series. Independence of species was rejected at the Bonferroni-corrected 10% threshold.

In addition to the time-domain analyses above, we performed frequency-domain analyses, in particular for analyzing synchrony within the rich wader community, as well as the group formed by the great cormorant, grey heron and little egret. Based on the work by Keitt (2008) and follow-up by Vasseur et al. (2014), we used the wavelet transform of the time series to measure the coherency between time series

$$\rho(t, s) = \frac{\Lambda_{t,s}(|\sum_k w_k(\tau, s)|)}{\Lambda_{t,s}(\sum_k |w_k(\tau, s)|)} \quad (2)$$

where  $w_k(\tau, s)$  is the continuous Morlet wavelet transform of species  $k$  at time  $\tau$  for scale  $s$ ,  $\Lambda_{t,s}(\cdot) = \int_{-\infty}^{+\infty} e^{-\frac{1}{2}(\frac{t-\tau}{s})^2}(\cdot)d\tau$  and  $|\cdot|$  is the modulus of the complex number. The numerator corresponds to the total biomass variation while the denominator corresponds to



the variations of each species. This index is close to 0 when species compensate and reaches 1 when they are synchronous. As before, the significance of each value was tested at the 10%, Bonferroni-corrected, threshold by 100 phase-randomizations of each species time series, and computation of the corresponding  $\rho$  values.

All datasets and statistical analyses are available in a GitHub repository [https://github.com/fbarraquand/BirdTimeSeries\\_Teich](https://github.com/fbarraquand/BirdTimeSeries_Teich)<sup>2</sup>.

## Results

Using a taxonomic classification of the community (focusing on the genera *Calidris* and *Anas* as two key examples of contrasted birds), we can see that within-genus synchrony indices at the seasonal scale are always positive whenever significantly different from the null (no temporal correlation between species), i.e. there is no compensation within a genus (Fig. 2). This matches the patterns obtained within the entire wetland bird community (Fig A1 in Appendix 1).

For the cold season, *Calidris* and *Anas* exhibit opposite trends in synchrony in response to the management change in 2006. However, for the warm season, the management change, which consisted in lowering the water levels, created more synchronous communities of species within the *Anas* and *Calidris* genera. This increase in synchrony after 2006 is matched by the functional group classification.

Even though there is no widespread community-wide or genus-wide compensation at the yearly timescale (differentiating the seasons), there could be compensation at finer temporal scales, e.g. a month or two, or coarser scales, over several years. The wavelet plot (Fig. 3), that allows to consider a time-varying and scale-dependent strength of synchrony, suggests that there is synchrony even at a fine temporal scale throughout most of the time series.

---

<sup>2</sup>Made public upon acceptance and archived in Zenodo

<sup>169</sup> However, post-2006, there seems to be a possibility for overcompensation on a scale around  
<sup>170</sup> 5 years or around 3-4 months.




Figure 2: Gross' synchrony index ( $\eta$ ) as a function of the season (cold and warm seasons), calculated within (top, a-b) and between (bottom, c-d) groups. The groups considered were different functional groups (ducks vs. waders, right b-d) or taxonomic groups (*Anas* genus, *Calidris* genus, left a-b) groups. The index was computed in each panel on the whole dataset (black) or using two periods: before and after 2006 (light and dark blue), the year of the change in water level management. Red stars correspond to synchrony values significantly different from the null model (independent species), at the 10% threshold.

There are therefore relatively contrasted results regarding the effect of the management change on short-term synchrony within the wader community. At the yearly (season) timescale, it seems to increase the synchrony (though the Gross index and wavelets provide slightly different answers). At even shorter timescales though, it seems to decrease it.

More clear-cut results can be found when we examine the synchrony vs. compensation between functional groups (Fig. 2d). Since we consider only two functional groups, the Gross index reduces to a simple correlation<sup>3</sup>. Waders and ducks are negatively correlated during the cold season and positively correlated during the warm season. These patterns are in contrast unclear when using a taxonomic classification (no compensation, Fig. 2c).




Figure 3: Wavelet modulus ratio for the wader community, scaling from 0 (compensation, blue color) to 1 (synchrony, red color). Dashed black lines delineate regions significantly different from the null model with a false discovery rate controlled at the 10% threshold.

<sup>3</sup>It should be noted that with more species, the Gross et al. (2013) index has been specifically designed so as to be able to take the value -1 when there is a zero-sum dynamics, i.e., perfect compensation, unlike the classical correlation coefficient.




Figure 4: Time series of great cormorant abundance, as well as summed abundances of grey heron and little egret (logarithmic scale).




Figure 5: Time-domain (a) and frequency-domain (b) synchrony analyses of the group formed by cormorant, egret and heron

While compensation could be expected upon visual inspection of the time series of the two groups formed by cormorant on the one hand, and egret plus heron (summed as a small functional group) on the other hand (Fig. 4), we see on Fig. 5 that synchrony is in fact the rule around the annual scale and below, when considering the wavelet index. We wondered

if the patterns in Fig. 4 were caused by the use of a log scale, but we found that in fact the correlation was higher rather than lower on the log scale (Appendix 2). However, over long temporal scales ( $\sim 6$  years) there seems to be some compensation, traducing a progressive change in composition within this small community module, that was already visible on the time series plot (Fig. 4). There might be some compensation over very short timescale as well (within the season), but at very specific times and the biological mechanisms for this are unclear, since these species compete for roost sites, a process that it unlikely to manifest at such very short timescales.

## Discussion

Compensation was overall very rare at the yearly timescale (differentiating between the cold and warm season). At short timescales (below the season), and among taxonomically or functionally close species, some compensation could be found but only at certain periods. In other words, there was no widespread “functional compensation” (*sensu* Gonzalez and Loreau 2009) *within* genera or guilds at the annual scale or below.

Yet, summing species abundances within a guild and comparing the “biomass sums” of contrasted guilds, community composition did change in frequency in the long run; in other words, there was some compensation *between* guilds.

Given that we compare the level of synchrony/compensation within guilds (with many species) and between guilds (with only a handful of groups), we checked in Appendix 3 if changing the number of “compartments” ( $n$ ) in the Gross  $\eta$  index could affect its value: it did not. However, we found that if two groups respond in opposite ways to a shared driver, the stronger the response to the driver, the lesser the compensation indicated by  $\eta$  at the whole community level. This might explain the low levels of compensation that we found at the overall wetland bird community level (Appendix 1), in spite of the clear presence of two

groups reacting in opposite way to shared driver (here, water levels). Analyses at several taxonomic/functional scales are therefore warranted to be conclusive about compensation.

We used correlation between the summed abundances of closely related species (species within the *Anas* genus vs. species within the *Calidris* genus) or the summed abundances of functionally similar species (waders vs. ducks) to uncover compensation. The functional group classification produced much more clearly compensation between guilds than the taxonomic classification. We expected to see compensation at that “functional scale” irrespective of the season, because the requirements of these birds are different, but here waders and ducks were found to correlate negatively only during the cold (wintering) season. This may be because the summer is characterized by a broad inflow of birds, including non-resident individuals that somehow add random variation to the community dynamics (though other explanations are possible).

It may be better to say that we detected “compensation” rather than “compensatory dynamics” between bird species (Gonzalez and Loreau, 2009) as the observed long-term changes in species composition (more waders, proportionally less ducks; Appendix 4) might be due to an increased inflow of birds preferring low water levels, and outflow of birds preferring high water levels, under an overall space constraint. In other words, the shift in community dynamics is likely not directly due to birth and deaths. However, despite the importance of movements and habitat preference to the local community dynamics, there is certainly also an influence of the regional changes in births and deaths on these local dynamics.

Zooming in on the cormorant-heron-egret module, we find that compensation mostly occurs above the annual temporal scale, and predominantly in summer as well as before 2006. This occurs because of a slow replacement of species due to competition for resting/roosting sites in the summer season (C. Feigné, pers. obs.), which mostly occurred before 2006.

Overall, our results suggest to search for compensation more often *between* rather than *within* functional groups, and over relatively long timescales above that of the dominant



driver (e.g., above 5 years if the main driver is a seasonal climate). This goes against calls to search for compensation at very short timescales (Vasseur and Gaedke, 2007; Gonzalez and Loreau, 2009) in order to filter out the effect of the main seasonal driver. Although searching for compensation at temporal scales below the seasonal abiotic driver (e.g., temperature) was partly motivated by studies on plankton whose community dynamics are much faster, we could have expected compensation to manifest also that scale here as well (e.g., monthly). Movement of birds reacting to food availability can certainly occur that fast, and wetlands have a finite carrying capacity, which could promote short-term compensation. We suspect that instead, because many species share common abiotic drivers at short temporal scales (Loreau and de Mazancourt, 2008), compensation is bound to be quite rare below the dominant temporal scale of the environment.

In many ways, searching for compensation using biodiversity time series data is searching for needles in a haystack: only some specific temporal and functional/taxonomic scales allow to see compensation whilst numerous confounding factors make the community co-vary positively at all other scales (Vasseur et al., 2014). Although the knowledge of specific biological mechanisms increasing the densities of some species at the expense of others can help, synchrony will likely dominate community-level time series data for closely related species, even in species that compete strongly (Ranta et al., 2008). This is true even in cases of known mechanisms of competition or shifts in community composition due to abiotic changes as in this study. We suggest that “zooming out” taxonomically or functionally (considering summed abundances of dissimilar functional groups) and temporally (considering temporal scales well above the dominant driver) may often be the best strategy to see the compensation that will inevitably manifest if the community-level abundance is maintained within bounds.

## Acknowledgements

We thank the birdwatchers and staff of the Teich Reserve/Landes Gascogne regional park who contributed to data collection over the years, as well as LPO Aquitaine for helping us retrieve the raw data. The data collection was supported by the Landes Gascogne regional park as well as the Teich municipality, while data analysis was funded by LabEx COTE (ANR-10-LABX-45).

## References

- Bell, R., M. Fogarty, and J. Collie, 2014. Stability in marine fish communities. *Marine Ecology Progress Series* **504**:221–239.
- Blüthgen, N., N. K. Simons, K. Jung, D. Prati, S. C. Renner, S. Boch, M. Fischer, N. Hölzel, V. H. Klaus, T. Kleinebecker, M. Tschapka, W. W. Weisser, and M. M. Gossner, 2016. Land use imperils plant and animal community stability through changes in asynchrony rather than diversity. *Nature Communications* **7**:10697.
- Bush, E. R., K. A. Abernethy, K. Jeffery, C. Tutin, L. White, E. Dimoto, J.-T. Dikangadissi, A. S. Jump, and N. Bunnefeld, 2017. Fourier analysis to detect phenological cycles using long-term tropical field data and simulations. *Methods in Ecology and Evolution* **8**:530–540.
- Christensen, E., D. Harris, and S. Ernest, 2018. Long-term community change through multiple rapid transitions in a desert rodent community. *Ecology* **99**:1523–1529.
- Ernest, S. M., J. H. Brown, K. M. Thibault, E. P. White, and J. R. Goheen, 2008. Zero sum, the niche, and metacommunities: long-term dynamics of community assembly. *The American Naturalist* **172**:E257–E269.

- Gonzalez, A. and M. Loreau, 2009. The causes and consequences of compensatory dynamics in ecological communities. *Annu. Rev. Ecol. Evol. Syst* **40**:393–414.
- Gouhier, T. C. and F. Guichard, 2014. Synchrony: quantifying variability in space and time. *Methods in Ecology and Evolution* **5**:524–533.
- Gross, K., B. J. Cardinale, J. W. Fox, A. Gonzalez, M. Loreau, H. Wayne Polley, P. B. Reich, and J. van Ruijven, 2013. Species richness and the temporal stability of biomass production: a new analysis of recent biodiversity experiments. *The American Naturalist* **183**:1–12.
- Hallett, L. M., S. K. Jones, A. A. M. MacDonald, M. B. Jones, D. F. B. Flynn, J. Ripplinger, P. Slaughter, C. Gries, and S. L. Collins, 2016. codyn: An r package of community dynamics metrics. *Methods in Ecology and Evolution* **7**:1146–1151.
- Houlahan, J. E., D. J. Currie, K. Cottenie, G. S. Cumming, S. M. Ernest, C. S. Findlay, S. D. Fuhlendorf, U. Gaedke, P. Legendre, J. J. Magnuson, and others, 2007. Compensatory dynamics are rare in natural ecological communities. *Proceedings of the National Academy of Sciences* **104**:3273–3277.
- Keitt, T. H., 2008. Coherent ecological dynamics induced by large-scale disturbance. *Nature* **454**:331–334.
- Loreau, M. and C. de Mazancourt, 2008. Species Synchrony and Its Drivers: Neutral and Nonneutral Community Dynamics in Fluctuating Environments. *The American Naturalist* **172**:E48–E66.
- Purves, D. W. and R. Law, 2002. Fine-scale spatial structure in a grassland community: quantifying the plant’s-eye view. *Journal of Ecology* **90**:121–129.

- 301 Ranta, E., V. Kaitala, M. S. Fowler, J. Laakso, L. Ruokolainen, and R. O'Hara, 2008. De-  
 302 tecting compensatory dynamics in competitive communities under environmental forcing.  
 303 *Oikos* **117**:1907–1911.
- 304 Sinclair, A., K. L. Metzger, J. M. Fryxell, C. Packer, A. E. Byrom, M. E. Craft, K. Hampson,  
 305 T. Lembo, S. M. Durant, G. J. Forrester, et al., 2013. Asynchronous food-web pathways  
 306 could buffer the response of serengeti predators to el niño southern oscillation. *Ecology*  
 307 **94**:1123–1130.
- 308 Usinowicz, J., C.-H. Chang-Yang, Y.-Y. Chen, J. S. Clark, C. Fletcher, N. C. Garwood,  
 309 Z. Hao, J. Johnstone, Y. Lin, M. R. Metz, T. Masaki, T. Nakashizuka, I.-F. Sun, R. Valen-  
 310 cia, Y. Wang, J. K. Zimmerman, A. R. Ives, and S. J. Wright, 2017. Temporal coexistence  
 311 mechanisms contribute to the latitudinal gradient in forest diversity. *Nature* .
- 312 Vasseur, D. A., J. W. Fox, A. Gonzalez, R. Adrian, B. E. Beisner, M. R. Helmus, C. Johnson,  
 313 P. Kratina, C. Kremer, C. de Mazancourt, E. Miller, W. A. Nelson, M. Paterson, J. A.  
 314 Rusak, J. B. Shurin, and C. F. Steiner, 2014. Synchronous dynamics of zooplankton  
 315 competitors prevail in temperate lake ecosystems. *Proceedings of the Royal Society B:*  
 316 *Biological Sciences* **281**:20140633–20140633.
- 317 Vasseur, D. A. and U. Gaedke, 2007. Spectral analysis unmasks synchronous and compen-  
 318 satory dynamics in plankton communities. *Ecology* **88**:2058–2071.