

a
Mouse liver, rebuilt: 234,552 cells and nuclei
Cell type (n)
Hepatocyte (41,669)
Cholangiocyte (1,984)
Endothelial (51,229)
Fibroblast (3,026)
Mesothelial (359)
KC (48,350)
Monocyte (18,312)
Neutrophil (4,302)
Basophil (163)
cDC1 (7,558)
cDC2 (7,909)
Mig. cDCs (843)
Transitional DC (735)
pDCs (10,988)
B cell (8,905)
T cell (13,955)
NK cell (2,260)
ILC1 (5,146)
unresolved (6,859)
UMAP2
UMAP1
b
Suspension
cell (207,558)
nucleus (26,994)
UMAP2
UMAP1
c
Digestion
Ex Vivo (84,109)
In Vivo (123,449)
Nuclei (26,994)
UMAP2
UMAP1
d
paper 185,894
Fig 1B samples on GEO (46)
389,056
Authors' MAD QC + Scrublet
328,041
Doublet clusters removed
321,059
scDblFinder doublets
286,017
< 200 UMI after decontX
277,495
Between-type (bridge) cells
274,395
Sort-gate mismatches (final)
234,552
0
100
200
300
400
Cells and nuclei kept (thousands)
e
Hepatocyte
Cholangiocyte
Endothelial
Fibroblast
Mesothelial
KC
Monocyte
Neutrophil
Basophil
cDC1
cDC2
Mig. cDCs
Transitional DC
pDCs
B cell
T cell
NK cell
ILC1
unresolved
cell
nucleus
0
10
20
30
40
50
Cells and nuclei (thousands)
f
Table S1 top-3 marker genes per type (Guilliams et al. 2022)
Hepatocyte
Cholangiocyte
Endothelial
Fibroblast
Mesothelial
KC
Monocyte
Neutrophil
Basophil
cDC1
cDC2
Mig. cDCs
Transitional DC
pDCs
B cell
T cell
NK cell
ILC1
Fabp1
Apoa2
Mup20
Ankrd1
Ddit4l
1700011H14Rik
Mmrn2
Ptprb
Cldn5
Nrxn1
Carmn
Gli3
Msln
Upk3b
Lrrn4
Cd5l
Clec4f
Vsig4
Chil3
F13a1
Gm9733
Stfa2l1
S100a9
S100a8
Fcer1a
Cyp11a1
Cpa3
Xcr1
Itgae
Rab7b
Cd209a
Mgl2
Tnip3
Nudt17
Slco5a1
Cacnb3
Ly6c2
Siglech
Pgam2
Klk1
Cd79a
Ms4a1
Pax5
Cd3e
Cd3d
Themis
Klra4
Klra8
Eomes
Gzmc
Cd200r2
Klrb1b
1.0
0.5
0.0
Scaled mean
% expressing
25
50
100