

Testing production scenarios for (anti-)(hyper-)nuclei and exotica at ~~LHC~~ energies available at the CERN Large Hadron Collider

Francesca Bellini* and Alexander P. Kalweit†

European Organisation for Nuclear Research (CERN), [1211](#) Geneva, Switzerland

(Received 28 September 2018; Revised 21 March 2019)

Abstract

We present a detailed comparison of coalescence and thermal-statistical models for the production of (anti-)(hyper-)nuclei in high-energy collisions. For the first time, such a study is carried out as a function of the size of the object relative to the size of the particle emitting source. Our study reveals large differences between the two scenarios for the production of objects with extended wave functions. While both models give similar predictions and show similar agreement with experimental data for (anti-)deuterons and (anti-) ^3He nuclei, they largely differ in their description of (anti-)hyper-triton production. We propose to address experimentally the comparison of the production models by measuring the coalescence parameter systematically for different (anti-)(hyper-)nuclei in different collision systems and differentially in multiplicity. Such measurements are feasible with the current and upgraded ~~LHC~~Large Hadron Collider experiments. Our findings highlight the unique potential of ultra-relativistic heavy-ion collisions ~~as~~ain the laboratory to clarify the internal structure of exotic QCD objects and can serve as a basis for more refined calculations in the future.

* francesca.bellini@cern.ch

† alexander.kalweit@cern.ch

I. INTRODUCTION

Nuclei and hyper=~~nuclei~~ are special objects with respect to non=~~composite~~ hadrons (pions, protons, etc.), because their size is comparable to a fraction or the whole system created in high-energy proton-proton (pp), proton-nucleus (pA), and nucleus-nucleus (AA) collisions [1]. Their size is typically defined as the rms of their (charge) wave function, corresponding to about 2 fm for light (anti-)nuclei as obtained from electron scattering experiments. For the hyper=~~triton~~, theoretical calculations indicate a n rms of the wave function of about 5 fm [2], significantly larger than that of non=~~strange~~ nuclei with mass number $A = 3$ and driven by the average separation of the Λ relative to the ~~two-other~~other two nucleons. This difference in the wave functions results in dramatic consequences for the production scenarios, as discussed in the following. The properties of the objects under study here are summarized in Table I. TI

For about ~~sixty~~60 years, coalescence models have been used to describe the formation of composite objects [3–11]. Surprisingly, thermal-statistical models have been successful in describing the production of light (anti-)(hyper-)nuclei across a wide range of energies in AA collisions [12, 13]. In this approach, particles are produced from a fireball ~~in at~~ thermal equilibrium with temperatures of $T_{\text{chem}} \approx 156$ MeV.[COMP: Throughout file, please use standard PRC spacing.] Particle abundances are fixed at chemical freeze-out, when inelastic collisions cease. Further elastic and pseudo=~~elastic~~ collisions occur among the components of the expanding fireball, ~~that~~which affect the spectral shapes and the measurable yields of short-lived (strongly decaying) hadronic resonances. Once the mean free path for elastic collisions is larger than the system size, the fireball freezes out kinetically at $T_{\text{kin}} \approx 90$ MeV [14]. In such a dense and hot environment, composite objects with binding energies that are low with respect to the temperature of the system, appear as “fragile.” For instance, the binding energy of the deuteron is $B_{E,d} = 2.2 \text{ MeV} \ll T_{\text{chem}}, T_{\text{kin}}$. The cross section for pion-induced deuteron breakup is significantly larger than the typical (pseudo)-elastic cross sections for the rescattering of hadronic resonance decay products [15–18]. Similarly, the elastic cross=~~section~~ driving deuteron spectra to kinetic equilibration in central heavy-ion collisions [19] is smaller than the breakup cross section [15–18] (Anti-)nuclei produced at chemical freeze-out are not expected to survive the hadronic phase, yet their measured production is consistent with statistical-thermal model predictions and a

non-zero elliptic flow is observed [19, 20]. Several solutions have been proposed to solve this “(anti-)nuclei puzzle”: (a) a sudden freeze-out at the QGP-hadron phase boundary [21], (b) the thermal production of these objects as compact quark bags [13], (c) the continuous interplay of breakup and formation reactions resulting in the coincidence of thermal and kinetic equilibration [18, 22], [and](#) (d) the coincidence of coalescence and thermal production [7, 23]. Data from rescattering of short-lived hadronic resonances suggest a long-lasting hadronic phase [24], thus strongly disfavoring hypothesis a. Hypothesis b cannot presently be tested beyond the agreement of measured (anti-)nuclei yields with statistical-thermal model predictions. Calculations for [\(c.\)hypothesis c](#) are currently available only for deuterons. Hypothesis d is scrutinized in this [letterpaper](#). To this purpose, we propose a new method to compare models that also allows for a direct comparison with LHC data.

TABLE I. Properties of nuclei and hypernuclei with mass number $A \leq 3$. The nucleus size is given in terms of the (charge) rms radius of the wave function, λ_A . The size parameter of the wave function of the harmonic oscillator potential, r_A , is chosen to approximately reproduce the measured/expected rms, λ_A^{meas} (fm). The proton rms charge radius $\lambda_p = 0.879(8)$ fm [29] is subtracted from λ_A^{meas} according to $\lambda_A = \sqrt{(\lambda_A^{\text{meas}})^2 - \lambda_p^2}$ to account for the finite extension of the constituents. We assume $\lambda_\Lambda \approx \lambda_n \approx \lambda_p$.

Mass number	Nucleus	Composition	Binding energy (MeV)	Spin	λ_A^{meas} (fm)	r_A (fm)	Ref. No.(s.)
$A = 2$	d	pn	2.224575 (9)	1	2.1413 ± 0.0025	3.2	[25, 26]
$A = 3$	^3H	pnn	8.4817986 (20)	1/2	1.755 ± 0.086	2.15	[27]
	^3He	ppn	7.7180428 (23)	1/2	1.959 ± 0.030	2.48	[27]
	$^3_\Lambda\text{H}$	p Λ n	$0.13 \pm 0.05^{\text{a}}$	1/2	4.9–10.0	6.8–14.1	[2, 28]

^a For [the](#) hypertriton, we report here the separation energy of the Λ from the ~~two~~ other [two](#) nucleons.

For the first time, LHC data allow for the study of (anti-)(hyper-)nuclei production as a function of the system and object size. A quantitative comparison of the production scenarios has been proposed in [30], resulting in the idea [to study of studying](#) the production rates of nuclei with similar masses [es](#) but very different internal structures, as ^4He and ^4Li [10]. As ^4Li is not stable with respect to strong decay, its measurement is experimentally

difficult and probably less constraining than the systematic measurement of the (hyper-)nuclei coalescence parameters proposed here.

II. COALESCENCE APPROACH

In the coalescence picture, nucleons produced in the collision coalesce into nuclei if they are close in space and have similar velocities [3, 4]. For a nucleus with mass number $A = Z + N$, the coalescence probability is quantified by the coalescence parameter B_A . Considering that at LHC energies the numbers of produced protons and neutrons at midrapidity as well as their momentum distributions are expected to be equal, B_A is defined as **[COMP: Throughout file, please ignore extraneous coding for spacing in math, and align and space math per usual PRC style.]**

$$E_A \frac{d^3 N_A}{dp_A^3} = B_A \left(E_{p,n} \frac{d^3 N_{p,n}}{dp_{p,n}^3} \right)^A \Big|_{\vec{p}_p = \vec{p}_n = \frac{\vec{p}_A}{A}}, \quad (1)$$

where $p_{p,n}$ are the proton and neutron momenta and $E_{p,n}$ their energies. The LHC is particularly suited for the production of antinuclei, since the number of baryons and antibaryons is equal at midrapidity [31]. Consequently, the anti-particle-to-particle ratio for (hyper-)nuclei is consistent with unity [19, 32–35]. In a simple coalescence approach, B_A is expected to be independent of momentum and of the object size relative to the volume of particle emission (hereafter referred to as “source size”). While this picture is found to be approximately valid in pp and p-Pb collisions [32–34], it breaks down in Pb-Pb collisions, which exhibit a strong decrease of B_A with centrality [36]. The elliptic flow of deuterons cannot be explained by simple coalescence [19].

More advanced coalescence models [5–7] take into account the source size, as the coalescence probability naturally decreases for nucleons with similar momenta that are produced far apart in configuration space. We rely on the formalism proposed in [7]. As coalescence is a quantum-mechanical process, the classical definition of phase space is replaced by the Wigner formalism. The production probability of a nucleon cluster is given by the overlap of the Wigner function with the phase-space distributions of the constituents. The wave functions of the (hyper-)nuclei are approximated by the ground-state wave functions of an isotropic spherical harmonic oscillator as in [7] with one single characteristic-size parameter, r_A . For the deuteron wave function $\varphi_d(\vec{r})$, one obtains

$$\varphi_d(\vec{r}) = (\pi r_d^2)^{-3/4} \exp\left(-\frac{r^2}{2r_d^2}\right). \quad (2)$$

For nuclei with $A > 2$, analogous forms exist. The relation between r_A and the rms of the wave function was derived in [37] as

$$\lambda_A^2 = \frac{3}{2} \frac{A-1}{A} \frac{r_A^2}{2} \quad (3)$$

for point-like constituents. We follow the Gaussian ansatz to obtain fully analytic solutions. In Table I, we list the measured rms of the wave function, λ_A^{meas} , and the r_A parameter derived from these relations. We encourage future, more rigorous numerical studies that address the calculation of coalescence probabilities with more realistic wave functions, e.g., the Hulthen ~~parameterisation~~ [parametrization](#) for deuterons [6] or a Λ -deuteron ~~parameterisation~~ [parametrization](#) for hyper-tritons as done in [38] and [39].

The quantum-mechanical nature of the coalescence products is explicitly accounted for by means of an average correction factor, $\langle C_d \rangle$. For deuterons, $\langle C_d \rangle$ has been approximated as

$$\langle C_d \rangle \approx \frac{1}{\left[1 + \left(\frac{r_d}{2R_\perp(m_T)}\right)^2\right] \sqrt{1 + \left(\frac{r_d}{2R_\parallel(m_T)}\right)^2}}, \quad (4)$$

where r_d is the size parameter, R_\perp and R_\parallel are the lengths of homogeneity of the coalescence volume, and m_T is the transverse mass of the coalescing nucleons. The nucleus size enters ~~in~~ the calculation of B_2 via $\langle C_d \rangle$, as well as the homogeneity volume $R_\perp^2 R_\parallel$, according to the relation [7]

$$B_2 = \frac{3\pi^{3/2} \langle C_d \rangle}{2m_T R_\perp^2(m_T) R_\parallel(m_T)}. \quad (5)$$

The coalescence parameter decreases with increasing volume, as expected. $\langle C_d \rangle$ introduces a length scale defined by the deuteron size relative to the source size. If we assume that $R_\perp \approx R_\parallel \approx R$, Eqs. (4) and (5) simplify to

$$\langle C_d \rangle \approx \left[1 + \left(\frac{r_d}{2R(m_T)}\right)^2\right]^{-3/2} \quad (6)$$

and

$$B_2 = \frac{3\pi^{3/2} \langle C_d \rangle}{2m_T R^3(m_T)}. \quad (7)$$

Figure 1 shows the source radius (R) dependence of $\langle C_d \rangle$ and B_2 , calculated assuming (a) [F1](#)

$r_d = 0$, (b) $r_d = 0.3$ fm, (c) the actual value $r_d = 3.2$ fm [26], and (d) a larger, unrealistic $r_d = 10$ fm. As shown in Fig. 1, $\langle C_d \rangle$ suppresses significantly the production of objects with a radius larger than the source.

FIG. 1. The Quantum-mechanical correction factor $\langle C_d \rangle$ [top; see Eq. (6)] and the coalescence parameter B_2 for the deuteron [bottom; see Eq. (7)] as a function of the radius of the source R , calculated assuming a size parameter for the deuteron $r_d = 0, 0.3, 3.2$, and 10 fm. The inflection point of $\langle C_d \rangle$ corresponds to $R = r_d/\sqrt{6}$ and is indicated in the left panel by the vertical dotted line for $r_d = 3.2$ fm.

Following the discussion in [7] and [9], Eq. (4) may be generalized as

$$\langle C_A \rangle = \prod_{i=1,2,3} \left(1 + \frac{r^2}{4R_i^2} \right)^{-\frac{1}{2}(A-1)}. \quad (8)$$

Under the assumption $R_1 \approx R_2 \approx R_3 \approx R$ as in [9] and by combining Eq. 6.2 in [7] with our Eq. (8), we obtain:

$$B_A = \frac{2J_A + 1}{2^A} \frac{1}{\sqrt{A}} \frac{1}{m_T^{A-1}} \left(\frac{2\pi}{R^2 + (\frac{r_A}{2})^2} \right)^{\frac{3}{2}(A-1)}. \quad (9)$$

This general formula can be used to compare the predicted B_A with data directly. For small sources, as $R \rightarrow 0$, the coalescence probability is anti-proportional to the harmonic oscillator size parameter and, thus, proportional to the depth of the attractive potential in the harmonic oscillator picture (and, thus, to the nucleus binding energy). Quite naturally, the allowed momentum difference between the coalescing nucleons is larger for more attractive, i.e., deeper, potentials. For a large source where $R \gg r_A$, the coalescence probability is dominated by the classical phase-space separation and, thus, decreases for large distances in configuration space.

III. STATISTICAL-THERMAL APPROACH COMBINED WITH THE BLAST-WAVE MODEL

In the statistical-thermal approach [40–42], the yields (dN/dy) of light anti- and hypernuclei are very sensitive to T_{chem} due to their large mass and approximately scale approximately as $dN/dy \propto (2J_A + 1) \exp(-m/T_{\text{chem}})$. The thermal model implements

eigenvolume corrections by fixing the object radius as an external parameter. We refer to the literature for the extensive discussions on the validity of the eigenvolume correction for light (anti-)(hyper-)nuclei [43] and the relation with the possible production as compact quark bags [13]. In contrast to coalescence, the statistical-thermal models provide only p_T -integrated yields. Therefore, we use, in addition, a blast-wave [44] ~~parameterization~~parametrization to model the p_T -dependence, with parameters obtained from the simultaneous fit to pion, kaon, and proton spectra measured in Pb-Pb collisions by ALICE for several centralities [14]. As discussed in the following section, centrality-dependent parameters allow one to extract spectral distributions for a given source size. The object size does not enter in the formulation of the blast-wave model.

The normalization of the predicted blast-wave spectra for nuclei is fixed using the p_T -integrated deuteron-to-pion ratio and ^3He -to-pion ratio predicted by the GSI-Heidelberg implementation of the statistical-thermal model with $T_{\text{chem}} = 156$ MeV, multiplied by the measured pion yield [14]. This choice, as opposed to using the ratio to protons, is motivated by the fact that the measured proton yield is seen to be slightly overestimated by the thermal model [45]. For hypertritons, the normalization is extracted from the statistical-thermal model prediction of the strangeness population factor $S_3 = \frac{^3\text{H}/^3\text{He}}{\Lambda/p}$ multiplied by the measured Λ/p ratio [14, 46] and ^3He yield [36]. With the resulting spectra, we calculate B_A for a given p_T/A and compare it with coalescence expectations. In the following, we label this model ~~as~~ “thermal + blast wave.”

IV. MAPPING EVENT MULTIPLICITY INTO SOURCE SIZE

In order to compare the source radius-dependent predictions from coalescence with the centrality-dependent data and with predictions from the thermal + blast-wave model, we map the average charged particle multiplicity density ($\langle dN_{\text{ch}}/d\eta \rangle$) in each centrality (or multiplicity) event class into the system size. Experimentally, the source size can be controlled by selecting different collision geometries, i.e., different centralities [47]. This mapping is based on the ~~following parameterisation~~parametrization

$$R = a \langle dN_{\text{ch}}/d\eta \rangle^{1/3} + b, \quad (10)$$

where R is the source radius, $a = 0.473$ fm, and $b = 0$.

The value of the empirical parameter a is obtained by tuning the ~~parameterisation~~parametrization such that the measured (anti-)deuteron B_2 in the most central Pb-Pb class falls into the coalescence prediction. In this way, we constrain the coalescence volume with the more differential (anti-)deuteron data and assume that it is the same for all anti- and hyper-~~nuclei~~. We justify the choice of Eq. (10) by identifying the source volume as the effective subvolume of the whole system ~~that,~~ which is governed by the homogeneity length of the interacting nucleons (as in [7]) and experimentally accessible with Hanbury-Brown-Twiss (HBT) interferometry [48]. The HBT radii us scale with $\langle dN_{\text{ch}}/d\eta \rangle^{1/3}$ and we make the simplifying assumption that this scaling holds across collision systems, which is approximately fulfilled in the data [1, 49]. We also note that the HBT radii, and thus also the source size, depend on the pair average transverse momentum $\langle k_T \rangle$ [50]. In contrast to [9], we do not explicitly use the measured HBT radii in our study because using a linear fit to the ALICE HBT data [1, 51] would result in a smaller source size ($R \approx 4$) fm than required by the measured B_2 to agree with the coalescence prediction ($R \approx 5.5$ fm) in central Pb-Pb collisions. We do, however, take into account the experimentally observed $\langle k_T \rangle$ dependence of the source size, in contrast to a similar coalescence study reported in [39] (for a detailed discussion of f possible alternative source volume ~~parameterisations~~parametrizations, see the Appendix A). Production via coalescence could also be investigated by looking at the transverse momentum dependence of B_A . However, the advantage of studying these effects as a function of the multiplicity/centrality is that the system size offers a larger lever arm. For a fixed p_T/A , B_2 changes by ~~about~~ a factor of about 50 going from pp to central Pb-Pb collisions, whereas B_2 changes by a factor ~~two~~of 2 going from $p_T/A = 0.4$ GeV/ c to $p_T/A = 2.2$ GeV/ c in most central Pb-Pb collisions [36] and by ~~less than a factor two~~ a factor of less than 2 in the measured p_T/A range in pp collisions [32, 33].

V. COMPARISON WITH DATA

In Fig. 2, the available LHC data for (anti-)(hyper-)nuclei [32, 35, 36] are compared F2 to coalescence and to the thermal + blast-wave predictions. For the latter and for data, the radius ~~parameterisation~~parametrization given by Eq. (10) is used. For deuterons, both approaches lead to similar predictions and describe reasonably the data for $R \gtrsim 1.6$ fm. For ^3He , the models exhibit a qualitatively similar R dependence but differ by a factor of

FIG. 2. Comparison of the coalescence parameters measured by ALICE (~~solid markers~~filled symbols) for deuterons (upper panel), ^3He (middle panel), and $^3_\Lambda\text{H}$ (lower panel) in pp [32] and Pb-Pb [35, 36] collisions with the thermal + blast-wave model expectations (dotted line) and the coalescence predictions (solid lines). The dashed line in the lower-right panel corresponds to the coalescence prediction for the $^3_\Lambda\text{H}$ with a larger radius. We have rescaled the inelastic pp collision data in [32] to match the so-called INEL > 0 class by the ratio ~~of~~ $\langle dN_{\text{ch}}/d\eta \rangle$ in these two event classes; see [52] for details.

about 1.5 to 2. The currently available data are consistent with both models within 2σ to 3σ , where σ is the total uncertainty in the data. Both approaches show large differences (a factor of 5 to 6 for central Pb-Pb collisions and a factor ~~larger than~~of more than 50 for $R < 2$) fm for the $^3_\Lambda\text{H}$ caused by the significantly larger size of $^3_\Lambda\text{H}$ with respect to ^3He . The only data point available so far in Pb-Pb collisions is in agreement with the thermal + blast-wave model but differs by 6σ from our coalescence calculation. This discrepancy might differ for calculations that use a more realistic (non-Gaussian) wave function. In [38] it is argued that the difference between data and coalescence might be explained by a later formation through coalescence of Λ s and deuterons. A possible difference attributable to the presence of excited states with $J = 3/2$ of $^3_\Lambda\text{H}$ is not considered here, as there is no evidence for its existence [53].

Most importantly, Fig. 2 shows that the difference between the two approaches increases with decreasing R , ~~underlin~~highlighting the need for additional multiplicity-differential data to distinguish between the two production scenarios. For $^3_\Lambda\text{H}$ we considered also a prediction from coalescence for a wider wave function (see Fig. 2), which results in even lower production probabilities. This ~~behaviour~~behavior highlights the unique potential to constrain the wave function of (hyper-)nuclei via precise measurements of the R -dependent coalescence parameter. The curves presented here explicitly allow for an estimate of the hyper-~~triton~~ production in pp collisions, which is expected to be suppressed by about two orders of magnitude with respect to that of ^3He , making this measurement a prime candidate for future experimental studies.

VI. (ANTI-)TRITON PRODUCTION

For isobars with [the](#) same spin but different wave [functions](#), like ${}^3\text{H}({}^3\overline{\text{H}})$ and ${}^3\text{He}({}^3\overline{\text{He}})$, Eq. (9) provides a relation between the relative coalescence probabilities as [the](#) corollary

$$\rho(p_{\text{T}}) \equiv \frac{B_A({}^3\text{H})}{B_A({}^3\text{He})}(p_{\text{T}}) = \left(\frac{R(p_{\text{T}})^2 + \frac{r_3^2({}^3\text{He})}{4}}{R(p_{\text{T}})^2 + \frac{r_3^2({}^3\text{H})}{4}} \right)^3. \quad (11)$$

Under the assumption that the distributions of protons and neutrons are identical, this is equivalent to the ratio of ${}^3\text{H}$ yield relative to ${}^3\text{He}$ yield (or ${}^3\overline{\text{H}}/{}^3\overline{\text{He}}$) at a given transverse momentum. It is to be noted that the transverse momentum dependence of this ratio originates solely from the p_{T} dependence of the source volume. The ratio ρ is reported in Fig. 43 **[AU: Figure 3 (not Fig. 4) as meant here? If not, please correct. Thank you. F3]** for $p_{\text{T}}/A = 0.75 \text{ GeV}/c$ of the two isobars and as a function of the average charged particle multiplicity. [The](#) multiplicity is obtained from the system radius by inverting Eq. (10). The coalescence model predicts a dependence of ρ on the system size and thus on the charged particle multiplicity. In particular, the production of ${}^3\text{H}({}^3\overline{\text{H}})$ is predicted by our model to be ~~up to a factor two~~ more abundant [by up to a factor of 2](#) than the production of ${}^3\text{He}({}^3\overline{\text{He}})$ in small systems. Such an excess of (anti-)triton is not expected in thermal-statistical hadronization. Since the two isobars have the same mass, spin, and baryon number, the [grand canonical](#) thermal model [12, 13] predicts a ${}^3\text{H}/{}^3\text{He}$ ratio equal to unity. The same holds true when taking into account the potential suppression of (anti-)nucleus yields due to explicit conservation of [the](#) baryon number, as in [canonical statistical model](#) calculations [54]. Existing (anti-)triton measurements in pp collisions [32] are limited by statistical precision and thus do not ~~allow~~ yet [allow](#) for a distinction between the two scenarios.

FIG. 3. Yield of ${}^3\text{H}$ relative to ${}^3\text{He}$ for $p_{\text{T}}/A = 0.75 \text{ GeV}/c$ as defined by Eq. (11) in the coalescence picture (black line). The source radius has been mapped into [the](#) average charged particle multiplicity by using the ~~parameterisation~~[parametrization](#) constrained to the ALICE B_2 measurement described in Sec. IV. The ~~orange~~ dashed [orange](#) line represents the expectation from statistical hadronization models. Colored arrows highlight the multiplicity range spanned by ALICE measurements in different collision systems.

VII. CONCLUSIONS AND OUTLOOK

We ~~summarises~~summarize our main conclusions as follows:

1. For the production of $A = 2$ and $A = 3$ (anti-)nuclei in heavy-ion collisions, thermal + blast-wave and coalescence models give similar predictions for a source volume that is constrained by experimental data on d , \bar{d} production in central Pb-Pb collisions.
2. For hypertriton, the two models give very different predictions as a function of the source volume. In particular, the yield of hypertriton appears to be suppressed by about two orders of magnitude in pp collisions with respect to ${}^3\text{He}$ due to its wider wave function.
3. In Pb-Pb collisions, the very limited ~~amount~~number of data available favours the thermal + blast-wave model prediction within our assumptions.
4. Systematic measurements in pp, p-Pb, and Pb-Pb collisions at LHC energies have a unique potential to clarify the production mechanism and the nature of composite QCD objects. Ideally, such measurements are accompanied by systematic measurements of the HBT radii in the same multiplicity/centrality classes and collision systems.
5. Our findings suggest a clear experimental path to be pursued with high-precision measurements at the upcoming phase of the LHC in the next ~~ten~~10 years, which will finally provide sufficient integrated luminosity for the studies proposed here [55].

As our study is deliberately based on simplified assumptions that allow for a completely analytical treatment of the problem, future studies should be based on more realistic approximations (in particular, the wave function), which require numerical calculations. We plan to extend our study to explore further the p_T dependence as well as to investigate $A = 4$ systems and more exotic QCD objects like the $X(3872)$ [8, 56]. If the $X(3872)$ corresponds to a loosely bound $\bar{D}^{*0}-D^0$ molecule, the rms of its wave function can be as large as $4.9^{+13.4}_{-1.4}$ fm [57]. Thus, its possible production via coalescence in pp collisions would be subject to a similar suppression as the hypertriton. Setting a final word on the production mechanisms also has a broader application in astrophysics and dark-matter searches, by representing an essential input for the measurement of (anti-)nuclei in space ~~withi~~n ongoing [58, 59] and

future [60, 61] experiments. **[COMP: In heading below, please set "APPENDIX:" (not "Appendix A:").]**

Appendix A: COMPARISON OF ALTERNATIVE SOURCE VOLUME PARAMETRIZATIONS of alternative source volume parameterisations

For the considerations discussed in this work and in similar ~~onereports~~, one crucial aspect has to be taken into account, namely, how the source radius is parameterized as a function of the multiplicity and transverse momentum. As discussed in Sec. IV, for our main result we rely on the ~~parameterisation~~ parametrization given by Eq. (10), which assumes a linear dependence between the cubic root of the measured average charged particle multiplicity density and the radius of the source. In order to extract the a and b coefficients of Eq. (10), several solutions can be adopted. One possibility is to fix the coefficients by fitting the available HBT data from ALICE [1, 49, 51] with Eq. (10), as shown in Fig. 4 by ~~F4~~ the data points fitted with the ~~gray~~ dotted gray line. In order to preserve the momentum

FIG. 4. Comparison of different source volume ~~parameterisations~~ parametrizations. The dotted gray line is the result of a linear fit to the ALICE HBT data [1, 49, 51], according to Eq. (10) with $a = 0.339$ fm and $b = 0.128$ fm. The ~~blue~~ dashed blue line represents the ~~parameterisation~~ parametrization used in [39], corresponding to $a = 0.83$ fm and $b = 0$. The ~~continuous~~ solid black line represents the ~~parameterisation~~ parametrization used in our study ($a = 0.473$ fm and $b = 0$) and obtained by a constraint from the B_2 measured in central (0–10%) Pb-Pb collisions.

dependence of the source volume, the highest available k_T (≈ 0.9 GeV/c) from pion HBT is chosen, as it is closest in m_T to the lowest transverse momentum per nucleon ($p_T \approx 0.8$ GeV/c) accessible by ALICE for the measurement of nucleon production. Ideally, one would use the proton femtoscopic radii, but given the availability of these measurements in only some of the collision systems and centralities, we assume that m_T scaling holds. The fit to the ALICE data results in $a = 0.339$ fm and $b = 0.128$ fm. However, this approach does not describe the deuteron B_2 data: we observe that in order for the measured B_2 to agree with the coalescence prediction in most central Pb-Pb collisions, the larger source radius of $R \approx 5.5$ fm is required instead of the $R \approx 4$ fm as resulting from the HBT fit. This

might also indicate that the volume relevant for the coalescence process is slightly larger than the homogeneity volume that can be extracted from HBT studies. Because of this, we chose to constrain the ~~parameterisation~~parametrization to the measured B_2 in central Pb-Pb collisions for $p_T = 0.75$ GeV/ c as well as to the origin ($b = 0$). The resulting ~~parameterisation~~parametrization, with $a = 0.473$ fm, is ~~re~~presented in Fig. 4 ~~with a continuous~~by the solid black line, and it is used for the results of our study discussed in Sec. V.

The authors of [39] assume the same functional dependence as we do in Eq. (10). However, their approach deviates from ours, as they neglect the momentum dependence of the source volume and constrain it with HBT data at low momentum ($k_T = 0.25$ GeV/ c). The ~~parameterisation~~parametrization from [39] ($a = 0.83$ fm and $b = 0$) is shown for comparison in Fig. 4 as the ~~-blue-~~blue dashed line. While this choice allows for a good description of p_T -integrated yield ratios (e.g., d/p , ${}^3\text{He}/p$, ${}^3_\Lambda\text{H}/\Lambda$) [39], using this ~~parameterisation~~parametrization for the p_T -differential study would lead to a significantly lower coalescence probability than measured, caused by the much larger source volume (e.g., a B_2 lower by a factor of 5.6 for $p_T = 0.75$ GeV/ c , corresponding to a ~~a~~ factor of 2 larger radius). In other words, by integrating over the transverse momentum, one ~~loses~~ sensitivity to the p_T dependence of the coalescence probability as the object radius becomes dominated by the large source volumina for small k_T [see Eq. (9)].

ACKNOWLEDGMENTS

We would like to thank Kfir Blum for inspiring this work. We thank U. Heinz for the useful discussions and the clarification of the equivalence of the Bertsch-Pratt and Yano-Koonin-Podgoretskii ~~parameterisations~~parametrizations of the HBT radii. We further acknowledge discussions with Benjamin Doenigus, in particular, about the production of ${}^3_\Lambda\text{H}$ in pp collisions, and with Eulogio Serradilla Rodriguez. In addition, we would like to thank Juergen Schukraft, Peter Braun-Munzinger, Marco Van Leeuwen, Maximiliano Puccio, Roman Lietava, Natasha Sharma, Sebastian Hornung, and the colleagues from the ALICE

Collaboration for their valuable input.

-
- [1] **[COMP: Please space references per usual PRC style, including closing up final period to preceding character.]** J. Adam *et al.* (ALICE), Phys. Rev. C **93**, 024905 (2016),—
arXiv:1507.06842 [nucl-ex] [Issn: 2469-9985; [DOI: 10.1103/PhysRevC.93.024905] .
 - [2] H. Nemura, Y. Suzuki, Y. Fujiwara, and C. Nakamoto, Prog. Theor. Phys. **103**, 929 (2000)
[Issn: 0033-068X; Coden: PTPKAV] [DOI: 10.1143/PTP.103.929] .
 - [3] S. T. Butler and C. A. Pearson, Phys. Rev. **129**, 836 (1963) [Issn: 0031-899X; Coden: PHRVAO] [DOI: 10.1103/PhysRev.129.836] .
 - [4] J. I. Kapusta, Phys. Rev. C **21**, 1301 (1980) [Issn: 0556-2813; Coden: PRVCAN] [DOI: 10.1103/PhysRevC.21.1301] .
 - [5] H. Sato and K. Yazaki, Phys. Lett. B **98**, 153 (1981) [Issn: 0370-2693; Coden: PYLBAJ] [DOI: 10.1016/0370-2693(81)90976-X] .
 - [6] J. L. Nagle, B. S. Kumar, D. Kusnezov, H. Sorge, and R. Mattiello, Phys. Rev. C **53**, 367 (1996) [Issn: 0556-2813; Coden: PRVCAN] [DOI: 10.1103/PhysRevC.53.367] .
 - [7] R. Scheibl and U. W. Heinz, Phys. Rev. C **59**, 1585 (1999) [Issn: 0556-2813; Coden: PRVCAN] [DOI: 10.1103/PhysRevC.59.1585] .
 - [8] S. Cho *et al.* (ExHIC), Prog. Part. Nucl. Phys. **95**, 279 (2017) [Issn: 0146-6410; Coden: PPNPDB] [DOI: 10.1016/j.ppnp.2017.02.002] .
 - [9] K. Blum, K. C. Y. Ng, R. Sato, and M. Takimoto, Phys. Rev. D **96**, 103021 (2017) [Issn: 2470-0010; [DOI: 10.1103/PhysRevD.96.103021] .
 - [10] S. Bazak and S. Mrowczynski, arXiv:1802.08212 [nucl-th] .
 - [11] W. Zhao, L. Zhu, H. Zheng, C. M. Ko, and H. Song, arXiv:1807.02813 [nucl-th] .
 - [12] A. Andronic, P. Braun-Munzinger, J. Stachel, and H. Stocker, Phys. Lett. B **697**, 203 (2011)
[Issn: 0370-2693; Coden: PYLBAJ] [DOI: 10.1016/j.physletb.2011.01.053] .
 - [13] A. Andronic, P. Braun-Munzinger, K. Redlich, and J. Stachel, Nature **561**, 321 (2018) [Issn: 0028-0836; Coden: NATUAS] [DOI: 10.1038/s41586-018-0491-6] .
 - [14] B. Abelev *et al.* (ALICE), Phys. Rev. C **88**, 044910 (2013) [Issn: 0556-2813; Coden: PRVCAN] [DOI: 10.1103/PhysRevC.88.044910] .
 - [15] H. Garcilazo, Phys. Rev. Lett. **48**, 577 (1982) [Issn: 0031-9007; Coden: PRLTAO] [DOI:

- [10.1103/PhysRevLett.48.577](#)] .
- [16] S. A. Bass *et al.*, Prog. Part. Nucl. Phys. **41**, 255 (1998) [Issn: 0146-6410; Coden: PPNPDB] [DOI: [10.1016/S0146-6410\(98\)00058-1](#)] .
 - [17] J. Schukraft, *Proceedings, 26th International Conference on Ultra-relativistic Nucleus-Nucleus Collisions (Quark Matter 2017): Chicago, Illinois, USA, February 5–11, 2017*, Nucl. Phys. A **967**, 1 (2017) [Issn: 0375-9474; Coden: NUPABL] [DOI: [10.1016/j.nuclphysa.2017.05.036](#)] .
 - [18] D. Oliinychenko, L.-G. Pang, H. Elfner, and V. Koch, arXiv:1809.03071 [hep-ph] .
 - [19] S. Acharya *et al.* (ALICE), Eur. Phys. J. C **77**, 658 (2017) [Issn: 1434-6044; Coden: EPCFFB] [DOI: [10.1140/epjc/s10052-017-5222-x](#)] .
 - [20] M. Puccio (ALICE), *Proceedings, 27th International Conference on Ultrarelativistic Nucleus-Nucleus Collisions (Quark Matter 2018): Venice, Italy, May 14–19, 2018*, Nucl. Phys. A **982**, 447 (2019) [Issn: 0375-9474; Coden: NUPABL] [DOI: [10.1016/j.nuclphysa.2018.10.043](#)] .
 - [21] P. Castorina and H. Satz, arXiv:1901.10407 [hep-ph] .
 - [22] X. Xu and R. Rapp, arXiv:1809.04024 [nucl-th] .
 - [23] U. Heinz, Presentation at EMMI Workshop, Torino, November (2017) .
 - [24] B. B. Abelev *et al.* (ALICE), Phys. Rev. C **91**, 024609 (2015) [Issn: 0556-2813; Coden: PRVCAN] [DOI: [10.1103/PhysRevC.91.024609](#)] .
 - [25] C. Van Der Leun and C. Alderliesten, Nucl. Phys. A **380**, 261 (1982) [Issn: 0375-9474; Coden: NUPABL] [DOI: [10.1016/0375-9474\(82\)90105-1](#)] .
 - [26] P. J. Mohr, D. B. Newell, and B. N. Taylor, Rev. Mod. Phys. **88**, 035009 (2016) [Issn: 0034-6861; Coden: RMPHAT] [DOI: [10.1103/RevModPhys.88.035009](#)] .
 - [27] J. E. Purcell and C. G. Sheu, Nucl. Data Sheets **130**, 1 (2015) [Issn: 0090-3752; Coden: NDTSCA] [DOI: [10.1016/j.nds.2015.11.001](#)] .
 - [28] D. H. Davis, *Proceedings of 8th International Conference on Hypernuclear and Strange Particle Physics (HYP 2003): Newport News, USA, October 14–18, 2003*, Nucl. Phys. A **754**, 3 (2005) [Issn: 0375-9474; Coden: NUPABL] [DOI: [10.1016/j.nuclphysa.2005.01.002](#)] .
 - [29] J. C. Bernauer *et al.* (A1), Phys. Rev. Lett. **105**, 242001 (2010) [Issn: 0031-9007; Coden: PRLTAO] [DOI: [10.1103/PhysRevLett.105.242001](#)] .
 - [30] S. Mrowczynski, Acta Phys. Pol. B **48**, 707 (2017) [Issn: 0587-4254; Coden: APOBBB] [DOI: [10.5506/APhysPolB.48.707](#)] .
 - [31] E. Abbas *et al.* (ALICE), Eur. Phys. J. C **73**, 2496 (2013) [Issn: 1434-6044; Coden: EPCFFB]

- [DOI: 10.1140/epjc/s10052-013-2496-5] .
- [32] S. Acharya *et al.* (ALICE), Phys. Rev. C **97**, 024615 (2018) [Issn: 2469-9985; DOI: 10.1103/PhysRevC.97.024615] .
- [33] S. Acharya *et al.* (ALICE), [in press](#) (2019) **[AU: Please update [33] if now possible. Thank you.]** .
- [34] J. Anielski, [in](#) *Proceedings, Hot Quarks 2014: Workshop for Young Scientists on the Physics of Ultrarelativistic Nucleus–Nucleus Collisions, Andalucia, Spain*, J. Phys.: Conf. Ser. **612**, 012014 (2015) [Issn: 1742-6588; DOI: 10.1088/1742-6596/612/1/012014] .
- [35] J. Adam *et al.* (ALICE), Phys. Lett. B **754**, 360 (2016) [Issn: 0370-2693; Coden: PYLBAJ] [DOI: 10.1016/j.physletb.2016.01.040] .
- [36] J. Adam *et al.* (ALICE), Phys. Rev. C **93**, 024917 (2016) [Issn: 2469-9985; DOI: 10.1103/PhysRevC.93.024917] .
- [37] A. Shebeko, P. Papakonstantinou, and E. Mavrommatis, Eur. Phys. J. A **27**, 143 (2006) [Issn: 1434-6001; Coden: EPJAFV] [DOI: 10.1140/epja/i2005-10247-3] .
- [38] Z. Zhang and C. M. Ko, Phys. Lett. B **780**, 191 (2018) [Issn: 0370-2693; Coden: PYLBAJ] [DOI: 10.1016/j.physletb.2018.03.003] .
- [39] K.-J. Sun, C. M. Ko, and B. Doenigus, arXiv:1812.05175 [nucl-th] .
- [40] M. Petran, J. Letessier, J. Rafelski, and G. Torrieri, Comput. Phys. Commun. **185**, 2056 (2014) [Issn: 0010-4655; Coden: CPHCBZ] [DOI: 10.1016/j.cpc.2014.02.026] .
- [41] S. Wheaton and J. Cleymans, Comput. Phys. Commun. **180**, 84 (2009) [Issn: 0010-4655; Coden: CPHCBZ] [DOI: 10.1016/j.cpc.2008.08.001] .
- [42] A. Andronic, P. Braun-Munzinger, and J. Stachel, Nucl. Phys. A **772**, 167 (2006) [Issn: 0375-9474; Coden: NUPABL] [DOI: 10.1016/j.nuclphysa.2006.03.012] .
- [43] V. Vovchenko and H. Stoecker, in *Proceedings, 16th International Conference on Strangeness in Quark Matter (SQM 2016): Berkeley, California, United States*, J. Phys.: Conf. Ser. **779**, 012078 (2017) [Issn: 1742-6588; DOI: 10.1088/1742-6596/779/1/012078] .
- [44] E. Schnedermann, J. Sollfrank, and U. W. Heinz, Phys. Rev. C **48**, 2462 (1993) [Issn: 0556-2813; Coden: PRVCAN] [DOI: 10.1103/PhysRevC.48.2462] .
- [45] B. Abelev *et al.* (ALICE), Phys. Rev. Lett. **109**, 252301 (2012) [Issn: 0031-9007; Coden: PRLTAO] [DOI: 10.1103/PhysRevLett.109.252301] .
- [46] B. B. Abelev *et al.* (ALICE), Phys. Rev. Lett. **111**, 222301 (2013) [Issn: 0031-9007; Coden:

- PRLTAO] [DOI: 10.1103/PhysRevLett.111.222301] .
- [47] B. Abelev *et al.* (ALICE), Phys. Rev. C **88**, 044909 (2013) [Issn: 0556-2813; Coden: PRVCAN] [DOI: 10.1103/PhysRevC.88.044909] .
- [48] To match the Bertsch-Pratt ~~parameterisation~~[parametrization](#) ($R_{\text{out}}, R_{\text{side}}, R_{\text{long}}$) employed by experiments with the Yano-Koonin-Podgoretskii ($R_{\perp}, R_{\parallel}, R_0$) ~~parameterisation~~[parametrization](#) used in the coalescence model, we identify $R_{\perp} = R_{\text{side}}$, $R_{\parallel} = R_{\text{long}}$ and then take $R = (R_{\perp}^2 R_{\parallel})^{1/3} \approx (R_{\text{side}}^2 R_{\text{long}})^{1/3}$.
- [49] J. Adam *et al.* (ALICE), Phys. Rev. C **91**, 034906 (2015) [Issn: 0556-2813; Coden: PRVCAN] [DOI: 10.1103/PhysRevC.91.034906] .
- [50] K. Aamodt *et al.* (ALICE), Phys. Lett. B **696**, 328 (2011) [Issn: 0370-2693; Coden: PYLBAJ] [DOI: 10.1016/j.physletb.2010.12.053] .
- [51] B. Abelev *et al.* (ALICE), Phys. Rev. D **87**, 052016 (2013) [Issn: 1550-7998; Coden: PRVDAQ] [DOI: 10.1103/PhysRevD.87.052016] .
- [52] J. Adam *et al.* (ALICE), Eur. Phys. J. C **77**, 33 (2017) [Issn: 1434-6044; Coden: EPCFFB] [DOI: 10.1140/epjc/s10052-016-4571-1] .
- [53] T. Mart, L. Tiator, D. Drechsel, and C. Bennhold, Nucl. Phys. A **640**, 235 (1998) [Issn: 0375-9474; Coden: NUPABL] [DOI: 10.1016/S0375-9474(98)00441-2] .
- [54] V. Vovchenko, B. Doenigus, and H. Stoecker, Phys. Lett. B **785**, 171 (2018) [Issn: 0370-2693; Coden: PYLBAJ] [DOI: 10.1016/j.physletb.2018.08.041] .
- [55] Z. Citron *et al.*, in *HL/HE-LHC Workshop: Workshop on the Physics of HL-LHC, and Perspectives at HE-LHC Geneva, Switzerland, June 18–20, 2018* (2018) .
- [56] A. Esposito, A. L. Guerrieri, L. Maiani, F. Piccinini, A. Pilloni, A. D. Polosa, and V. Riquer, Phys. Rev. D **92**, 034028 (2015) [Issn: 1550-7998; Coden: PRVDAQ] [DOI: 10.1103/PhysRevD.92.034028] .
- [57] P. Artoisenet and E. Braaten, Phys. Rev. D **83**, 014019 (2011) [Issn: 1550-7998; Coden: PRVDAQ] [DOI: 10.1103/PhysRevD.83.014019] .
- [58] J. Alcaraz *et al.* (AMS), Phys. Lett. B **461**, 387 (1999) [Issn: 0370-2693; Coden: PYLBAJ] [DOI: 10.1016/S0370-2693(99)00874-6] .
- [59] V. Poulin, P. Salati, I. Cholis, M. Kamionkowski, and J. Silk, arXiv:1808.08961 [astro-ph.HE] .
- [60] S. Schael, Presentation at AMS Days, La Palma, April 2018 .

- [61] T. Aramaki, C. J. Hailey, S. E. Boggs, P. von Doetinchem, H. Fuke, S. I. Mognet, R. A. Ong, K. Perez, and J. Zweerink (GAPS), *Astropart. Phys.* **74**, 6 (2016) [[Issn: 0927-6505](#); [Coden: APHYEE](#)] [[DOI: 10.1016/j.astropartphys.2015.09.001](#)] .