

1    **Testing coalescence and statistical-thermal production scenarios**  
2    **for (anti-)(hyper-)nuclei and exotic QCD objects at LHC energies**  
3    **with recent and future experimental data**

4                    F. Bellini and A. Kalweit

5                    16th of May 2018

6            (Anti-)(hyper-)nuclei are unique probes of the medium created in proton-proton,  
7            proton-Pb, and Pb–Pb collisions at LHC energies. At LHC energies, their production  
8            is typically discussed within the framework of coalescence and thermal-statistical pro-  
9            duction models. While it is often argued that both approaches are not distinguishable,  
10           we present a detailed study of both theories which reveals largely different predictions  
11           between the two approach for the production of  $^3\text{He}$  and hyper-tritons. Confronting  
12           our results with recent ALICE measurements, the coalescence approach is found to  
13           provide a correct description of the data only in small systems such as pp collisions,  
14           while it fails for central Pb–Pb collisions. The thermal-statistical model on the other  
15           hand is in agreement with results in central Pb–Pb collisions even though such fragile  
16           objects should be destroyed in hadronic interactions after the chemical freeze-out of  
17           the system. Our findings thus indicate the existence of a novel production mechanism  
18           for these objects.

## 19 Contents

20	<b>1. Introduction</b>	<b>3</b>
21	<b>2. Coalescence approach</b>	<b>5</b>
22	2.1. Simple coalescence . . . . .	5
23	2.2. Full coalescence . . . . .	6
24	2.3. Source volume . . . . .	7
25	<b>3. Statistical-thermal approach and blast-wave</b>	<b>8</b>
26	<b>4. Comparison with experimental data</b>	<b>9</b>
27	4.1. (Anti-)nuclei with $A = 2, 3, 4$ . . . . .	11
28	4.2. (Anti-)hyper-nuclei . . . . .	11
29	4.3. Other exotic QCD objects . . . . .	11
30	<b>5. Projections for the LHC Run 3 and 4</b>	<b>11</b>
31	<b>6. Summary and conclusions</b>	<b>11</b>
32	<b>A. HBT radii in Bertsch-Pratt and Yano-Koonin-Podgoretskii parameterisation for com-</b>	
33	<b>parison with coalescence models</b>	<b>11</b>

# 1. Introduction

The formation of light anti- and hyper- nuclei in highly energetic proton-proton, proton-nucleus and nucleus-nucleus collisions provides unique observables for the study of the system created in these collisions. In this context, nuclei and hyper-nuclei are special objects with respect to non-composite hadrons, because their size (i.e. the extension of their wave-function) is comparable to a fraction or the whole system created in the collision [1]. Halo nuclei as  ${}^6\text{He}$  would be ideal for such studies, but they remain out of the experimental reach in high-energy experiments in the near future. It must be noted, that these studies thus do not only shed light on the characteristics of the system created in the collision (if (anti-)nuclei with well known properties are studied), but also on the internal structure of the produced objects if their properties are not well known. The relevant known properties of the objects under study are summarised in Table 1. As quantum-mechanical objects, their size is typically defined as the rms of their wave function, which ranges from  $xx \text{ fm}$  for the  ${}^3\text{He}$  up to  $yy \text{ fm}$  for the hyper-triton. For simplicity, we approximate the wave-function of the objects under study by the wave-function of a spherical harmonic oscillator as in [10]. This allows to derive in the following a set of analytic formulas which can be used for the calculation of coalescence predictions. We encourage future more rigorous studies which address the coalescence probabilities with more precise wave-functions.

Surprisingly, thermal-statistical models have been successful at describing not only light-flavour particle production, but also that of light (anti-)(hyper-)nuclei across a wide range of energies in nucleus-nucleus collisions [2, 3]. In this approach, particles are produced from a fireball in thermal and kinetic equilibrium with temperatures of the order of  $T_{chem} = 156 \text{ MeV}$  (near the temperature at the QCD phase transition boundary, as predicted by lattice QCD calculations [4]. Particle abundances are fixed at chemical freeze-out, when inelastic collisions cease. Further elastic and pseudo-elastic collisions occur among the components of the expanding fireball, that can affect the spectral shapes and the measurable yields of short-lived (strongly decaying) hadronic resonances. Once the particle density of the system is so low that the mean free path for elastic collisions is larger than the size of the system, the fireball freezes-out kinetically. This is seen to occur when the system has reached temperatures of the order of  $T_{kin} \approx 90 \text{ MeV}$ . In such a dense and hot environment, composite objects with binding energies that are small with respect to the temperature of the system, appear as “fragile” objects. For instance, the binding energy of the deuteron is  $E_{B,d} = 2.2 \text{ MeV} \ll T_{chem}, T_{kin}$ . As a matter of fact, the cross-section for pion-induced deuteron breakup is significantly larger than the typical (pseudo)-elastic cross-sections for the re-scattering of hadronic resonance decay products (check statement) [5, 6, 7]. Similarly, the elastic cross-section which drives the deuteron spectra to kinetic equilibration in central heavy-ion collisions [8] is smaller than the breakup cross-section [7]. Based on this, the deuterons produced at chemical freeze-out would be expected not to survive the hadronic phase of the medium expansion, yet their production is measured to be consistent to the predictions from statistical-thermal models and they develop also a non-zero elliptic flow which is consistent with a common radial expansion together with the non-composite hadrons [8]. Do similar estimates as Karel in Frascati. In addition, it was recently shown that the assumption of realistic eigenvolumina for light nuclei would lead to instabilities of the statistical/thermal model predictions [9]. Several solutions have been proposed to solve this “light (anti-)nuclei puzzle”: (a.) a sudden freeze-out at the QGP-hadron phase boundary, (b.) the thermal production of these objects as compact quark bags [2], and (c.) the coincidence of coalescence mechanism with that of thermal production [10]. Data from rescattering of short-lived hadronic resonances indicate that the system undergoes a long-lasting hadronic phase before decoupling [11], thus strongly disfavours hypothesis (a.). While hypothesis (b.) cannot presently be tested beyond the agreement of measured (anti-)nuclei production yields with statistical-thermal model predictions, hypothesis (c.) is scrutinised in the present work.

For the table: there seems to be a nice summary here (including that the hyper-triton is spin 1/2): Properties And Interactions Of Hyperons - Proceedings Of U.S.-japan Seminar heraus-

Mass number	Nucleus	Composition	$B_E$ (MeV)	Spin $J$	rms radius of wavefunction (fm)	harmonic oscillator width $r$ (fm)	Refs.
$A = 2$	d	pn	2.2	1	$2.1413 \pm 0.0025$	3.2	[25]
$A = 3$	$^3\text{H}$	pnn	8.4817986 (20)	$1/2$	$1.755 \pm 0.086$		[?]
	$^3\text{He}$	ppn		$1/2$			[?]
	$^3_\Lambda\text{H}$	$p\Lambda n$					[?]
	$^4_\Lambda\text{He}$	ppnn					
$A = 4$	$^4_\Lambda\text{H}$	$p\Lambda nn$					
	$^4_{\Lambda\Lambda}\text{H}$	$p\Lambda\Lambda n$					
	$^4_\Lambda\text{He}$	$pp\Lambda n$					

Table 1: Properties of nuclei and hyper-nuclei.  $B_E$  is the binding energy per nucleon in MeV and the radius is given in terms of the rms radius of the wavefunction. References are given in the last column.

gegeben von Barnes Peter D, Nakai K, Gibson Benjamin F

To this purpose, we compare to models the existing data from the Large Hadron Collider. For the first time, these data allow for the systematic study of the light (anti-)(hyper-)nuclei production as a function of the system and object size. In the nucleon-coalescence approach, nuclei are formed at kinetic freeze-out by coalescence of nucleons that are nearby in space and have similar velocities. The coalescence model is reviewed in Section 2, starting from its simplest form (uncorrelated nucleon emission from a point-like source) to the full space-time evolution picture as discussed in [10]. In section 3, a blast-wave parameterisation for particle transverse momentum spectra in combination with predictions from the statistical-thermal model for the yields is used as an alternative approach. The direct comparison of the two approaches and the comparison with data are discussed in section 4. We find that a systematic study of the coalescence parameter  $B_A$  provides an important discrimination power between the two approaches. A comparison of the production rates of nuclei with similar mass but very different internal structure has already been suggested for the case of  $^4\text{He}$  and  $^4\text{Li}$  [12]. However, as the  $^4\text{Li}$  is not stable with respect to strong decay, its measurement is experimentally very challenging and probably less constraining than the comparison with hyper-nuclei proposed here. In section 5 we propose that  $B_A$  is systematically measured in all collision systems by exploiting the large statistics sample that will be available with the LHC Run 3 and 4, in order to rule out or support the aforementioned scenarios. As a matter of fact, the upcoming years of LHC data taking provide a unique opportunity to for the final understanding of (anti-)(hyper-)nuclei production. Setting a final word on the production mechanisms is not only in the interest of the heavy-ion community, but has a broader application in astrophysics and dark-matter searches, by representing an essential input for the measurement of (anti-)nuclei in space with ongoing [13] and future [?, 14] experiments. In addition to this, the study of light(anti-)nuclei might serve as a baseline for understanding the debated nature of exotic states such as the  $X(3872)$ , that has been interpreted as tetraquark state or hadronic molecule [15, 16].

## 2. Coalescence approach

### 2.1. Simple coalescence

In simple coalescence, nucleons produced in the collision coalesce into nuclei if they are close in space and have similar velocities [17, 18]. For a nucleus with mass number  $A = Z + N$ , the coalescence probability is typically quantified in terms of the coalescence parameter  $B_A$ , which is defined as

$$E_A \frac{d^3 N_A}{dp_A^3} = B_A \left( E_p \frac{d^3 N_p}{dp_p^3} \right)^Z \left( E_n \frac{d^3 N_n}{dp_n^3} \right)^N \Big|_{\vec{p}_p = \vec{p}_n = \frac{\vec{p}_A}{A}} \quad , \quad (1)$$

where  $p_{p,n}$  are the momenta of the proton and neutron and  $E_{p,n}$  their energy. Since at LHC energies the number of produced protons and neutrons at midrapidity is expected to be equal, the equation simplifies to

$$E_A \frac{d^3 N_A}{dp_A^3} = B_A \left( E_p \frac{d^3 N_p}{dp_p^3} \right)^A \Big|_{\vec{p}_p = \frac{\vec{p}_A}{A}} \quad . \quad (2)$$

Moreover, the LHC is particularly suited for the production of anti-nuclei, since the number of baryons and anti-baryons is essentially equal at midrapidity [19]. In a simple coalescence approach, the coalescence parameter is expected to be independent of  $p_T$  and of the object size with respect to the volume of particle emission (hereafter referred to as “source volume” or “source size”). In this naive expectation, the number of nuclei produced by coalescence increases with increasing number of nucleons produced in the collision. If the nucleon number increases with the event multiplicity, so does the number of (anti-)nuclei. While this picture is found to be

approximately valid in pp and p-Pb collisions [20, 21], it breaks down in Pb-Pb collisions, that exhibit a strong decrease of  $B_A$  with the centrality of the collision [22]. In addition, the elliptic flow of deuteron cannot be explained by simple coalescence [23].

## 2.2. Full coalescence

In contrast to the simple approach described in the previous section, a more advanced coalescence model takes into account the size of the particle emission source, as the coalescence probability naturally decreases for two nucleons with similar momenta that are produced far apart in configuration space. While there are several approaches to address this effect [?], we rely in our study on the formalism proposed in [10]. **Consider mentioning the approximations done by Uli in his paper, if there is time and how Uli's paper relates to the Wigner formalism.** In this approach, the quantum mechanical nature of the coalescence products is explicitly accounted for by means of an average quantum mechanical correction factor,  $\langle C_A \rangle$ . In the case of the deuteron, the quantum mechanical correction factor  $\langle C_d \rangle$  has been approximated as

$$\langle C_d \rangle \approx \frac{1}{\left[1 + \left(\frac{r_d}{2R_\perp(m_T)}\right)^2\right] \sqrt{1 + \left(\frac{r_d}{2R_\parallel(m_T)}\right)^2}} \quad (3)$$

where  $r_d$  is the radius of the deuteron,  $R_\perp$  and  $R_\parallel$  are the lengths of homogeneity of the coalescence volume and  $m_T$  is the transverse mass of the coalescing nucleons. The size of the nucleus enters in the determination of the coalescence parameter  $B_2$  via the quantum-mechanical correction factor  $\langle C_d \rangle$ , as well as the homogeneity volume  $R_\perp^2 R_\parallel$ , according to the relation

$$B_2 = \frac{3\pi^{3/2}\langle C_d \rangle}{2m_T R_\perp^2(m_T) R_\parallel(m_T)} \quad (4)$$

which is the main result of [10]. It is interesting to note that the coalescence parameter decreases with increasing volume, as expected. In addition to this, the quantum mechanical correction factor introduces a length scale defined by the deuteron size relative to the source size in the calculation of  $B_2$ , which reflects the coalescence probability. If we assume that  $R_\perp \approx R_\parallel \approx R$ , Eqs. 3 and 4 simplify to

$$\langle C_d \rangle \approx \left[1 + \left(\frac{r_d}{2R(m_T)}\right)^2\right]^{-3/2} \quad (5)$$

and

$$B_2 = \frac{3\pi^{3/2}\langle C_d \rangle}{2m_T R^3(m_T)}. \quad (6)$$

Figure 1 shows the source radius ( $R$ ) dependence of the quantum-mechanical correction factor (on the left) and the coalescence parameter  $B_2$  (on the right), calculated assuming (a.) a point-like nucleus, (b.)  $r_d = 0.3$  fm as currently assumed in thermal model calculations [24], (c.) the current estimate of the rms radius of the deuteron  $r_d = 3.2$  fm [25], (d.) a larger, unrealistic value of  $r_d = 10$  fm. As can be seen in Fig. 1, the quantum-mechanical correction factor leads to a significant suppression in the production of those objects whose radius is large compared to that of the source.

Following the approach and discussion presented in [26], Eq. 3 may be generalised as

$$\langle C_A \rangle = \prod_{i=1,2,3} \left(1 + \frac{r^2}{4R_i^2}\right)^{-\frac{1}{2}(A-1)} \quad (7)$$

for mass number  $A$  and the radii  $R_i$  which describe the volume of the emitting source. Similarly, the coalescence parameter  $B_A$  for a nucleus with mass number  $A$  and spin  $J_A$  is generalised in




Figure 1: The quantum mechanical correction factor  $\langle C_d \rangle$  (left panel, see Eq. 5) and the coalescence parameter  $B_2$  for deuteron (right panel, see Eq. 6) as a function of the radius of the source,  $R$ , calculated assuming a radius of the deuteron  $r_d = 0, 0.3, 3.2$  and  $10$  fm.

160 [10] as

$$B_A = \frac{2J_A + 1}{2^A} \frac{1}{\sqrt{A}} \langle C_A \rangle \left( \frac{(2\pi)^{3/2}}{m_T \prod_{i=1,2,3} R_i} \right)^{A-1}. \quad (8)$$

161 In particular, for the case of  ${}^3\text{He}$  with  $A = 3$  and  $J = 1/2$ , Eq. 8 becomes Eq. 9 presented in  
162 [26]:

$$B_3 = \frac{(2\pi)^3}{4\sqrt{3}} \langle C_3 \rangle (m_T \prod_{i=1,2,3} R_i)^{-2} \quad (9)$$

163 where

$$\langle C_3 \rangle \approx \prod_{i=1,2,3} \left( 1 + \frac{r_3^2}{4R_i^2} \right)^{-1}. \quad (10)$$

164  $\rightarrow$  Mention (if true, please check) that this assumes a gaussian wave function of the produced  
165 object. This might not be true in reality for the hyper-triton, but it allows to show the sensitivity  
166 of the wave-function and gives easy analytic expressions.

### 167 2.3. Source volume

168 As in [10], we identify the source volume as the effective sub-volume of the whole system which  
169 is governed by the homogeneity length of the interacting nucleons. In addition, the same authors  
170 claim that this volume is experimentally accessible with Hanbury-Brown-Twiss (HBT) interfer-  
171 ometry. The experimental results are typically obtained following the Bertsch-Pratt (BP) pa-  
172 rameterisation  $(R_{out}, R_{side}, R_{long})$ , while the coalescence model described in Section 2 expresses

the volume in terms of the Yano-Koonin-Podgoretskii (YKP) parameterisation. As discussed in Appendix ??, we identify  $R_{\perp} = R_{side}$  and  $R_{\parallel} = R_{long}$ , thus  $R = (R_{\perp}^2 R_{\parallel})^{1/3} \approx (R_{side}^2 R_{long})^{1/3}$ .

Experimentally, the size of the effective volume can be controlled by selecting different collision geometries, i.e. different centrality classes [27]. In heavy-ion collisions the HBT radii are known to scale with the cubic root of the average charged particle multiplicity density  $\langle dN_{ch}/d\eta \rangle^{1/3}$  [1], and to depend on the pair average transverse momentum  $\langle k_T \rangle$  [28]. In the following, we make the simplifying assumption that the scaling with  $\langle dN_{ch}/d\eta \rangle^{1/3}$  holds across collision systems, which is approximately fulfilled in data [29]. In contrast to [26], we therefore do not explicitly use the measured HBT radii in our study, but we derive the radii from the measured  $\langle dN_{ch}/d\eta \rangle$  according to the following relation:

$$R = a \langle dN_{ch}/d\eta \rangle^{1/3} + b \quad (11)$$

The coefficients,  $a = 0.339$  and  $b = 0.128$ , have been determined by fitting linearly the ALICE data, and the parameterisation is reported in Fig. 2. These values are consistent with the radius from kaon femtoscopy for  $m_T \approx 1$  GeV/c in low-multiplicity pp collisions [30] and the radius from pion femtoscopy in high-multiplicity Pb-Pb collisions at the highest available  $k_T \approx 0.8$  GeV/c [1]. The highest  $k_T$  bin was chosen as it corresponds in  $m_T$  to the lowest transverse momentum per nucleon ( $p_T/A \approx 0.8$  GeV/c) accessible by ALICE for the measurement of nuclei production. Ideally, one would use the proton femtosopic radii for such study, but given that these measurements are not available in all collision systems, we assume that  $m_T$ -scaling holds for HBT radii [31].




Figure 2: Parameterisation of the dependence of the source radius on multiplicity assumed in this paper, compared to HBT data from [1, 30].

- excess binding energy needs to be released to the system to make coalescence work

### 3. Statistical-thermal approach and blast-wave

In the statistical-thermal approach, the yields ( $dN/dy$ ) of light anti- and hyper-nuclei are very sensitive to the chemical freeze-out temperature  $T_{chem}$  due to their large mass  $m$  and approximately scale as  $dN/dy \propto \exp(-m/T_{chem})$ . As a matter of fact, the chemical freeze-out temperature defines the only scale in this model, since at the LHC the chemical potentials which ensure the conservation of baryon number ( $\mu_B$ ), strangeness ( $\mu_S$ ), and electric charge ( $\mu_Q$ ) are negligible. In contrast to the coalescence approach, the current implementations [] of the statistical-thermal



model provide only  $p_T$ -integrated yields and not the full hadron spectra. In order to fill this gap, and for the purposes of this exercise,  $p_T$  spectra have been modeled using a blast-wave parameterization. When extracting the predicted spectrum for a given particle species (proton,  $\Lambda$ , deuteron,  $^3\text{He}$ ), the parameters of the blast-wave (average radial flow velocity  $\langle\beta_T\rangle$ , kinetic freeze-out temperature  $T_{kin}$ , and velocity profile  $n$ ) are fixed to the values obtained from the simultaneous fit of the pion, kaon and proton spectra measured in each collision system and multiplicity event class by ALICE. The normalisation is fixed using the  $p_T$ -integrated deuteron-to-pion ratio and  $^3\text{He}$ -to-pion ratio predicted by the GSI-Heidelberg model with  $T_{chem} = 156$  MeV, multiplied by the pion yield measured by ALICE. This choice is motivated by the fact that the measured proton yield is seen to be slightly underestimated by the thermal model predictions at the LHC. In the case of hyper-triton, a slightly different procedure is chosen, namely the normalisation for  $^3\text{H}$  is extracted from the statistical-thermal model prediction of the strangeness population factor  $S_3$  multiplied by the measured  $\Lambda/p$  ratio and the measured  $^3\text{He}$  yield. Based on the thus obtained spectra, we calculate the corresponding coalescence parameters for a given  $p_T/A$  and compare it with coalescence expectations. Because we use experimental data to constrain the blast-wave prediction as well as the normalisation, and because such data are provided for given centrality/multiplicity classes, we use the corresponding  $\langle dN_{ch}/d\eta \rangle$  in the same class to estimate the system radius based on the parameterisation discussed in Sec. 2.3. In contrast to the coalescence approach which explicitly depends on the size of the produced object with respect to the system size, the object size does not enter in the formulation of the blast-wave model, which as a simplified hydrodynamic model treats the system based as a continuum and is not particle based. The thermal model on the other hand, implements eigenvolume corrections where the object radius is fixed as an external parameter ( $r = 0.3$  fm in the case of baryons for the GSI-Heidelberg prediction used here). We refer to the literature for the extensive discussions on the validity of the eigenvolume correction for light anti- and hyper-nuclei [?] and the relation with the possible production of these objects as compact quark bags [2].

## 4. Comparison with experimental data

Data on anti- and hyper-nuclei production at LHC energies and different collision systems have been released by the ALICE Collaboration in recent years. In Fig. ?? we compare the coalescence parameter  $B_2$  from the coalescence model (see Eqs. 4 and 6, with  $r_d = 3.2$  fm) with the ALICE data. First, the parameterisation of the system radius fitted to the HBT data as described in Sec. 2.3 is used to map the  $\langle dN_{ch}/d\eta \rangle$  to the source size (datapoints shown with empty markers). We notice already at this point that the coalescence volume from our extracted parameterisation of the HBT radii leads to discrepancies with respect to the curve from the coalescence calculation, and in particular, we notice that the model would require a larger radius for a given value of  $B_2$ . Therefore, in a second step the radius is tuned such that the data points for (anti-) deuterons fall onto the coalescence prediction (solid markers). We find that the parameters of Eq. 11 turn out to be  $a = xxx$  and  $b = yyy$ . With this second parameterisation, we investigate also the agreement of the model with the measured coalescence parameter  $B_3$ . As shown in Fig. 3, also in the case of  $B_3$ , a tension between the model and the data is found which ameliorates with the parameterisation which is tuned to the (anti-)deuteron data.. It is also noteworthy that any change of the radii parameterisation would result in a shift along the x axis of both  $B_2$  and  $B_3$  data points in the same way, while the theoretical coalescence curve would not be affected.

- discuss sensitivity to the choice of the kT for the radii parameterisation.
- add a plot with different radii for hyper-triton from 10fm to 3fm
- add a plot for A=4
- add a plot for the X(3872)???? -> Look at the ExHic and Maiani rebuttal




Figure 3: Comparison of the ALICE data from [1] with the coalescence prediction for two different mappings of the  $\langle dN_{\text{ch}}/d\eta \rangle$  to the system size  $R$ : (left panels) the parameterisation fitted to the HBT radii, (right panels) the parameterisation tuned to the  $B_2$  values in central Pb-Pb collisions.

#### 4.1. (Anti-)nuclei with $A = 2, 3, 4$

#### 4.2. (Anti-)hyper-nuclei

- cite Che Ming Ko [32]

#### 4.3. Other exotic QCD objects

We can consider essentially all Objects proposed by the ExHIC collaboration if they are of a composite nature or might be of a composite nature. One of them is the X(3872) which might have a wide wave-function as described in <http://inspirehep.net/record/861896?ln=de>

### 5. Projections for the LHC Run 3 and 4

### 6. Summary and conclusions

We conclude that (c.) appears unlikely, thus leaving (b.) as a viable option, at least with our present knowledge of the hypertriton size.

### A. HBT radii in Bertsch-Pratt and Yano-Koonin-Podgoretskii parameterisation for comparison with coalescence models

The results of HBT analyses are typically presented in either the Bertsch-Pratt ( $R_{out}$ ,  $R_{side}$ ,  $R_{long}$ ) or the Yano-Koonin-Podgoretskii ( $R_{\perp}$ ,  $R_0$ ,  $R_{\parallel}$ ) parameterization. The ALICE HBT results [28, 1] are given in the Bertsch-Pratt convention, whereas the coalescence parameter is derived in [10] by expressing the dependence on the volume in terms of the Yano-Koonin-Podgoretskii (YKP) parameterisation. The transformation between the two parameterisations is best presented in [33] in the equations (W 3.48) to (W 3.52)<sup>1</sup>:

$$R_{side}^2 = R_{\perp}^2, \quad (12)$$

$$R_{diff}^2 = R_{out}^2 - R_{side}^2 = \beta_{\perp}^2 \gamma^2 (R_0^2 + v^2 R_{\parallel}^2), \quad (13)$$

$$R_{long}^2 = (1 - \beta_l^2) R_{\parallel}^2 + \gamma^2 (\beta_l - v)^2 (R_0^2 + v^2 R_{\parallel}^2), \quad (14)$$

$$R_{ol}^2 = \beta_{\perp} (-\beta_l R_{\parallel}^2 + \gamma^2 (\beta_l - v) (R_0^2 + v^2 R_{\parallel}^2)). \quad (15)$$

We immediately identify that  $R_{\perp}^2$  can be identified with  $R_{side}^2$ . Following the reasoning and the nomenclature in [33] (W 3.52-3.53), the above equations can be inverted and  $R_{\parallel}^2$  can be expressed as

$$R_{\parallel}^2 = B - v \cdot C, \quad (16)$$

$$= R_{long}^2 - 2 \frac{\beta_l}{\beta_{\perp}} R_{ol}^2 + \frac{\beta_l^2}{\beta_{\perp}^2} R_{diff}^2 - v \cdot \left( -\frac{1}{\beta_{\perp}} R_{ol}^2 + \frac{\beta_l}{\beta_{\perp}^2} R_{diff}^2 \right). \quad (17)$$

As it turns out all corrections which are subtracted from  $R_{long}^2$  can be neglected. First, we notice that the cross term  $R_{ol}$  vanishes if the measured fireball is longitudinally boost-invariant, which is a valid approximation for the rapidity ranges studied here. The remaining terms are all proportional to  $\beta_l$ , which is also  $\beta_l = 0$  by definition of the rest frame for a longitudinally-boosted invariant system. In summary, we can consider  $R_{\perp} = R_{side}$  and  $R_{\parallel} = R_{long}$  for the present study.

<sup>1</sup>The equations in [33] are denoted as (W...) in order to distinguish them from the equations presented in this paper.

## Acknowledgements

We thank U. Heinz for the useful discussions and the clarification on the equivalence of the Bertsch-Pratt and Yano-Koonin-Podgoretskii parameterisations of the HBT radii. The authors would like to thank themselves for the auto-critics.

## References

- [1] **ALICE** Collaboration, J. Adam *et al.*, “Centrality dependence of pion freeze-out radii in Pb-Pb collisions at  $\sqrt{s_{NN}} = 2.76$  TeV,” *Phys. Rev.* **C93** no. 2, (2016) 024905, [arXiv:1507.06842 \[nucl-ex\]](#).
- [2] A. Andronic, P. Braun-Munzinger, K. Redlich, and J. Stachel, “Decoding the phase structure of QCD via particle production at high energy,” [arXiv:1710.09425 \[nucl-th\]](#).
- [3] A. Andronic, P. Braun-Munzinger, J. Stachel, and H. Stocker, “Production of light nuclei, hypernuclei and their antiparticles in relativistic nuclear collisions,” *Physics Letters B* **697** no. 3, (2011) 203 – 207.  
<http://www.sciencedirect.com/science/article/pii/S0370269311001006>.
- [4] **HotQCD** Collaboration, A. Bazavov *et al.*, “Equation of state in ( 2+1 )-flavor QCD,” *Phys. Rev.* **D90** (2014) 094503, [arXiv:1407.6387 \[hep-lat\]](#).
- [5] H. Garcilazo, “PION DEUTERON BREAKUP IN THE REGION OF THE (3,3) RESONANCE,” *Phys. Rev. Lett.* **48** (1982) 577–580.
- [6] S. A. Bass *et al.*, “Microscopic models for ultrarelativistic heavy ion collisions,” *Prog. Part. Nucl. Phys.* **41** (1998) 255–369, [arXiv:nucl-th/9803035 \[nucl-th\]](#). [*Prog. Part. Nucl. Phys.*41,225(1998)].
- [7] J. Schukraft, “QM2017: Status and Key open Questions in Ultra-Relativistic Heavy-Ion Physics,” *Nucl. Phys.* **A967** (2017) 1–10, [arXiv:1705.02646 \[hep-ex\]](#).
- [8] **ALICE** Collaboration, S. Acharya *et al.*, “Measurement of deuteron spectra and elliptic flow in Pb-Pb collisions at  $\sqrt{s_{NN}} = 2.76$  TeV at the LHC,” *Eur. Phys. J.* **C77** no. 10, (2017) 658, [arXiv:1707.07304 \[nucl-ex\]](#).
- [9] V. Vovchenko and H. Stoecker, “Analysis of hadron yield data within hadron resonance gas model with multi-component eigenvolume corrections,” *J. Phys. Conf. Ser.* **779** no. 1, (2017) 012078, [arXiv:1610.02346 \[nucl-th\]](#).
- [10] R. Scheibl and U. W. Heinz, “Coalescence and flow in ultrarelativistic heavy ion collisions,” *Phys. Rev.* **C59** (1999) 1585–1602, [arXiv:nucl-th/9809092 \[nucl-th\]](#).
- [11] **ALICE** Collaboration, B. B. Abelev *et al.*, “ $K^*(892)^0$  and  $\Lambda(1020)$  production in Pb-Pb collisions at  $\sqrt{s_{NN}} = 2.76$  TeV,” *Phys. Rev.* **C91** (2015) 024609, [arXiv:1404.0495 \[nucl-ex\]](#).
- [12] S. Bazak and S. Mrowczynski, “ $^4\text{He}$  vs.  $^4\text{Li}$  and production of light nuclei in relativistic heavy-ion collisions,” [arXiv:1802.08212 \[nucl-th\]](#).
- [13] **AMS** Collaboration, J. Alcaraz *et al.*, “Search for anti-helium in cosmic rays,” *Phys. Lett.* **B461** (1999) 387–396, [arXiv:hep-ex/0002048 \[hep-ex\]](#).
- [14] **GAPS** Collaboration, T. Aramaki, C. J. Hailey, S. E. Boggs, P. von Doetinchem, H. Fuke, S. I. Mognet, R. A. Ong, K. Perez, and J. Zweerink, “Antideuteron Sensitivity for the GAPS Experiment,” *Astropart. Phys.* **74** (2016) 6–13, [arXiv:1506.02513 \[astro-ph.HE\]](#).

- [15] A. Esposito, A. L. Guerrieri, L. Maiani, F. Piccinini, A. Pilloni, A. D. Polosa, and V. Riquer, “Observation of light nuclei at ALICE and the X(3872) conundrum,” *Phys. Rev. D* **D92** no. 3, (2015) 034028, [arXiv:1508.00295 \[hep-ph\]](#).
- [16] **ExHIC** Collaboration, S. Cho *et al.*, “Exotic Hadrons from Heavy Ion Collisions,” *Prog. Part. Nucl. Phys.* **95** (2017) 279–322, [arXiv:1702.00486 \[nucl-th\]](#).
- [17] S. T. Butler and C. A. Pearson, “Deuterons from High-Energy Proton Bombardment of Matter,” *Phys. Rev.* **129** (1963) 836–842.
- [18] J. I. Kapusta, “Mechanisms for deuteron production in relativistic nuclear collisions,” *Phys. Rev.* **C21** (1980) 1301–1310.
- [19] **ALICE** Collaboration, E. Abbas *et al.*, “Mid-rapidity anti-baryon to baryon ratios in pp collisions at  $\sqrt{s} = 0.9, 2.76$  and 7 TeV measured by ALICE,” *Eur. Phys. J.* **C73** (2013) 2496, [arXiv:1305.1562 \[nucl-ex\]](#).
- [20] **ALICE** Collaboration, S. Acharya *et al.*, “Production of deuterons, tritons,  $^3\text{He}$  nuclei and their antinuclei in pp collisions at  $\sqrt{s} = 0.9, 2.76$  and 7 TeV,” *Phys. Rev.* **C97** no. 2, (2018) 024615, [arXiv:1709.08522 \[nucl-ex\]](#).
- [21] **ALICE** Collaboration, S. Acharya *et al.*, “**Multiplicity dependence of (anti-)nuclei production in p-Pb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV**,” *To be published* **xx** (2018) yy, [arXiv:xxxx.xxxx \[nucl-ex\]](#).
- [22] **ALICE** Collaboration, J. Adam *et al.*, “Production of light nuclei and anti-nuclei in pp and Pb-Pb collisions at energies available at the CERN Large Hadron Collider,” *Phys. Rev.* **C93** no. 2, (2016) 024917, [arXiv:1506.08951 \[nucl-ex\]](#).
- [23] **ALICE** Collaboration, S. Acharya *et al.*, “Measurement of deuteron spectra and elliptic flow in Pb-Pb collisions at  $\sqrt{s_{NN}} = 2.76$  TeV at the LHC,” *Eur. Phys. J.* **C77** no. 10, (2017) 658, [arXiv:1707.07304 \[nucl-ex\]](#).
- [24] A. Andronic, P. Braun-Munzinger, K. Redlich, and J. Stachel, “Hadron yields, the chemical freeze-out and the QCD phase diagram,” *J. Phys. Conf. Ser.* **779** no. 1, (2017) 012012, [arXiv:1611.01347 \[nucl-th\]](#).
- [25] P. J. Mohr, D. B. Newell, and B. N. Taylor, “CODATA Recommended Values of the Fundamental Physical Constants: 2014,” *Rev. Mod. Phys.* **88** no. 3, (2016) 035009, [arXiv:1507.07956 \[physics.atom-ph\]](#).
- [26] K. Blum, K. C. Y. Ng, R. Sato, and M. Takimoto, “Cosmic rays, antihelium, and an old navy spotlight,” *Phys. Rev.* **D96** no. 10, (2017) 103021, [arXiv:1704.05431 \[astro-ph.HE\]](#).
- [27] **ALICE** Collaboration, B. Abelev *et al.*, “Centrality determination of Pb-Pb collisions at  $\sqrt{s_{NN}} = 2.76$  TeV with ALICE,” *Phys. Rev.* **C88** no. 4, (2013) 044909, [arXiv:1301.4361 \[nucl-ex\]](#).
- [28] **ALICE** Collaboration, K. Aamodt *et al.*, “Two-pion Bose-Einstein correlations in central Pb-Pb collisions at  $\sqrt{s_{NN}} = 2.76$  TeV,” *Phys. Lett.* **B696** (2011) 328–337, [arXiv:1012.4035 \[nucl-ex\]](#).
- [29] **ALICE** Collaboration, J. Adam *et al.*, “Two-pion femtoscopy in p-Pb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV,” *Phys. Rev.* **C91** (2015) 034906, [arXiv:1502.00559 \[nucl-ex\]](#).

- 358 [30] **ALICE** Collaboration, B. Abelev *et al.*, “Charged kaon femtoscopic correlations in  $pp$   
359 collisions at  $\sqrt{s} = 7$  TeV,” *Phys. Rev.* **D87** no. 5, (2013) 052016, [arXiv:1212.5958](#)  
360 [\[hep-ex\]](#).
- 361 [31] **ALICE** Collaboration, J. Adam *et al.*, “One-dimensional pion, kaon, and proton  
362 femtoscopy in Pb-Pb collisions at  $\sqrt{s_{NN}} = 2.76$  TeV,” *Phys. Rev.* **C92** no. 5, (2015) 054908,  
363 [arXiv:1506.07884](#) [\[nucl-ex\]](#).
- 364 [32] Z. Zhang and C. M. Ko, “Hypertriton production in relativistic heavy ion collisions,” *Phys.*  
365 *Lett.* **B780** (2018) 191–195.
- 366 [33] U. A. Wiedemann and U. W. Heinz *Phys. Rept.* **319** (1999) 145.