

Chapter 3. Potentials

3.1	Laplace's Equation	
3.1.1	Introduction	
3.1.2	Laplace's Equation in One Dimension	
3.1.3	Laplace's Equation in Two Dimensions	
3.1.4	Laplace's Equation in Three Dimensions	
3.1.5	Boundary Conditions and Uniqueness Theorems	
3.1.6	Conductors and the Second Uniqueness Theorem	
3.2	The Method of Images	
3.2.1	The Classic Image Problem	
3.2.2	Induced Surface Charge	
3.2.3	Force and Energy	
3.2.4	Other Image Problems	
3.3	Separation of Variables	
3.3.1	Cartesian Coordinates	
3.3.2	Spherical Coordinates	
3.4	Multipole Expansion	
3.4.1	Approximate Potentials at Large Distances	
3.4.2	The Monopole and Dipole Terms	
3.4.3	Origin of Coordinates in Multipole Expansions	
3.4.4	The Electric Field of a Dipole	

3.3 Separation of Variables


Let's attack Laplace's equation directly using the method of **separation of variables**.

→ by *look for solutions that are products of functions, each of which depends on only one of the coordinates.*

→ Consider two cases: **Cartesian coordinates** and **Spherical coordinates**

3.3.1 Cartesian Coordinates

Example 3.3 Two infinite grounded metal plates lie parallel. Find the potential inside this "slot."




3.3.1 Cartesian Coordinates

Example 3.3 Find the potential inside.

$$\frac{\partial^2 V}{\partial x^2} + \frac{\partial^2 V}{\partial y^2} = 0$$

(Boundary conditions)

- (i) $V = 0$ when $y = 0$,
- (ii) $V = 0$ when $y = a$,
- (iii) $V = V_0(y)$ when $x = 0$,
- (iv) $V \rightarrow 0$ as $x \rightarrow \infty$.



Since the potential is specified on all boundaries, the answer is uniquely determined.

The first step is to look for solutions in the form of products:

$$V(x, y) = X(x)Y(y) \longrightarrow Y \frac{d^2 X}{dx^2} + X \frac{d^2 Y}{dy^2} = 0$$

The next step is to "separate the variables"

$$Y \frac{d^2 X}{dx^2} + X \frac{d^2 Y}{dy^2} = 0 \longrightarrow \frac{1}{X} \frac{d^2 X}{dx^2} + \frac{1}{Y} \frac{d^2 Y}{dy^2} = 0$$

→ The first term depends only on x and the second only on y .

→ ***They must both be constant.***


Cartesian Coordinates $\frac{\partial^2 V}{\partial x^2} + \frac{\partial^2 V}{\partial y^2} = 0$

Example 3.3

$$V(x, y) = X(x)Y(y)$$

$$\frac{1}{X} \frac{d^2 X}{dx^2} + \frac{1}{Y} \frac{d^2 Y}{dy^2} = 0 \rightarrow \text{Must both be constant.}$$

$$\begin{array}{l} \frac{1}{X} \frac{d^2 X}{dx^2} = C_1 \\ \frac{1}{Y} \frac{d^2 Y}{dy^2} = C_2 \end{array} \xrightarrow{C_1 + C_2 = 0} \begin{array}{l} \frac{d^2 X}{dx^2} = k^2 X \longrightarrow X(x) = Ae^{kx} + Be^{-kx} \\ \frac{d^2 Y}{dy^2} = -k^2 Y \longrightarrow Y(y) = C \sin ky + D \cos ky \end{array} \quad (k > 0)$$



$$\Rightarrow V(x, y) = (Ae^{kx} + Be^{-kx})(C \sin ky + D \cos ky)$$

$$\begin{array}{ll} \text{(iv) } V \rightarrow 0 \text{ as } x \rightarrow \infty & \longrightarrow A = 0 \\ \text{(i) } V = 0 \text{ when } y = 0 & \longrightarrow D = 0 \end{array} \Rightarrow V(x, y) = Ce^{-kx} \sin ky$$

$$\text{(ii) } V = 0 \text{ when } y = a \longrightarrow \sin ka = 0 \longrightarrow k = \frac{n\pi}{a} \quad (n = 1, 2, 3, \dots)$$


$$\text{(iii) } V = V_0(y) \text{ when } x = 0$$

Cartesian Coordinates $\frac{\partial^2 V}{\partial x^2} + \frac{\partial^2 V}{\partial y^2} = 0$

Example 3.3

$$V(x, y) = X(x)Y(y) \longrightarrow V(x, y) = C e^{-n\pi x/a} \sin(n\pi y/a)$$

$$(n = 1, 2, 3, \dots)$$



- An **infinite set of solutions** (one for each n)
- none of them *by itself* satisfies the final boundary condition

$$(iii) V = V_0(y) \text{ when } x = 0$$

Since Laplace's equation is **linear**,

$$V(x, y) = \sum_{n=1}^{\infty} C_n e^{-n\pi x/a} \sin(n\pi y/a) \rightarrow \text{This linear combination is also a solution.}$$

(“Completeness” of V function)

$$(iii) V = V_0(y) \text{ when } x = 0 \longrightarrow V(0, y) = \sum_{n=1}^{\infty} C_n \sin(n\pi y/a) = V_0(y)$$

→ It's a Fourier sine series.


Multiply **$\sin(n'\pi y/a)$** (where n' is a positive integer), and integrate from 0 to a :

$$\sum_{n=1}^{\infty} C_n \int_0^a \sin(n\pi y/a) \sin(n'\pi y/a) dy = \int_0^a V_0(y) \sin(n'\pi y/a) dy$$

Cartesian Coordinates $\frac{\partial^2 V}{\partial x^2} + \frac{\partial^2 V}{\partial y^2} = 0$

Example 3.3

$$V(x, y) = X(x)Y(y) \Rightarrow V(x, y) = \sum_{n=1}^{\infty} C_n e^{-n\pi x/a} \sin(n\pi y/a)$$



$$\sum_{n=1}^{\infty} C_n \int_0^a \sin(n\pi y/a) \sin(n'\pi y/a) dy = \int_0^a V_0(y) \sin(n'\pi y/a) dy$$

$$\Rightarrow \int_0^a \sin(n\pi y/a) \sin(n'\pi y/a) dy = \begin{cases} 0, & \text{if } n' \neq n, \\ \frac{a}{2}, & \text{if } n' = n. \end{cases} \quad (\text{Orthogonality})$$

$$\Rightarrow C_n = \frac{2}{a} \int_0^a V_0(y) \sin(n\pi y/a) dy$$



Suppose the strip at $x = 0$ is a metal plate with constant potential V_0 .
(insulated from the grounded plates at $y = 0$ and $y = a$)

$$C_n = \frac{2V_0}{a} \int_0^a \sin(n\pi y/a) dy = \frac{2V_0}{n\pi} (1 - \cos n\pi) = \begin{cases} 0, & \text{if } n \text{ is even,} \\ \frac{4V_0}{n\pi}, & \text{if } n \text{ is odd.} \end{cases}$$




$$\Rightarrow V(x, y) = \frac{4V_0}{\pi} \sum_{n=1,3,5\dots} \frac{1}{n} e^{-n\pi x/a} \sin(n\pi y/a).$$

Cartesian Coordinates $\frac{\partial^2 V}{\partial x^2} + \frac{\partial^2 V}{\partial y^2} = 0$

Example 3.3 $V(x, y) = X(x)Y(y)$




$$V(x, y) = \frac{4V_0}{\pi} \sum_{n=1,3,5,\dots} \frac{1}{n} e^{-n\pi x/a} \sin(n\pi y/a) = \frac{2V_0}{\pi} \tan^{-1} \left(\frac{\sin(\pi y/a)}{\sinh(\pi x/a)} \right)$$



→ It shows how the first few terms in the Fourier series combine to make a better and better approximation to the constant v_o .

Cartesian Coordinates

Example 3.4 Two infinitely long grounded metal plates, again at $y = 0$ and $y = a$, are connected at $x = \pm b$ by metal strips maintained at a constant potential V_0 . Find the potential inside the resulting rectangular pipe.



(Boundary conditions)

$$\frac{\partial^2 V}{\partial x^2} + \frac{\partial^2 V}{\partial y^2} = 0$$

- (i) $V = 0$ when $y = 0$,
- (ii) $V = 0$ when $y = a$,
- (iii) $V = V_0$ when $x = b$,
- (iv) $V = V_0$ when $x = -b$.

$$V(x, y) = X(x)Y(y) \quad \longrightarrow \quad V(x, y) = (Ae^{kx} + Be^{-kx})(C \sin ky + D \cos ky)$$

The situation is *symmetric* with respect to x , so $V(-x, y) = V(x, y) \Rightarrow \mathbf{A = B}$

$$e^{kx} + e^{-kx} = 2 \cosh kx \quad \longrightarrow \quad V(x, y) = \cosh kx (C \sin ky + D \cos ky)$$

$$\begin{array}{ll} \text{(i)} & V = 0 \text{ when } y = 0 \quad \longrightarrow \quad \mathbf{D = 0} \\ \text{(ii)} & V = 0 \text{ when } y = a \quad \longrightarrow \quad \mathbf{k = n\pi/a} \end{array} \quad \longrightarrow \quad V(x, y) = C \cosh(n\pi x/a) \sin(n\pi y/a)$$


$$\longrightarrow \quad V(x, y) = \sum_{n=1}^{\infty} C_n \cosh(n\pi x/a) \sin(n\pi y/a)$$

Cartesian Coordinates

$$\frac{\partial^2 V}{\partial x^2} + \frac{\partial^2 V}{\partial y^2} = 0$$

Example 3.4



$$V(x, y) = X(x)Y(y) \quad \Rightarrow \quad V(x, y) = \sum_{n=1}^{\infty} C_n \cosh(n\pi x/a) \sin(n\pi y/a)$$



$$(iii) \quad V = V_0 \text{ when } x = b \quad \longrightarrow \quad V(b, y) = \sum_{n=1}^{\infty} C_n \cosh(n\pi b/a) \sin(n\pi y/a) = V_0$$

→ This is the same problem in Fourier analysis.

$$C_n \cosh(n\pi b/a) = \begin{cases} 0, & \text{if } n \text{ is even} \\ \frac{4V_0}{n\pi}, & \text{if } n \text{ is odd} \end{cases}$$




$$V(x, y) = \frac{4V_0}{\pi} \sum_{n=1,3,5,\dots} \frac{1}{n} \frac{\cosh(n\pi x/a)}{\cosh(n\pi b/a)} \sin(n\pi y/a).$$

Cartesian Coordinates

Example 3.5 An infinitely long rectangular metal pipe is grounded, but one end, at $x = 0$, is maintained at a specified potential $V_0(y, z)$. Find the potential inside the pipe.

→ *This is a three-dimensional problem:*



(Boundary conditions)

- (i) $V = 0$ when $y = 0$,
- (ii) $V = 0$ when $y = a$,
- (iii) $V = 0$ when $z = 0$,
- (iv) $V = 0$ when $z = b$,
- (v) $V \rightarrow 0$ as $x \rightarrow \infty$,
- (vi) $V = V_0(y, z)$ when $x = 0$.

$$\frac{\partial^2 V}{\partial x^2} + \frac{\partial^2 V}{\partial y^2} + \frac{\partial^2 V}{\partial z^2} = 0$$

$$V(x, y, z) = X(x)Y(y)Z(z) \longrightarrow \frac{1}{X} \frac{d^2 X}{dx^2} + \frac{1}{Y} \frac{d^2 Y}{dy^2} + \frac{1}{Z} \frac{d^2 Z}{dz^2} = 0$$

$$\frac{d^2 X}{dx^2} = (k^2 + l^2)X \longrightarrow X(x) = Ae^{\sqrt{k^2 + l^2} x} + Be^{-\sqrt{k^2 + l^2} x}$$

$$\frac{d^2 Y}{dy^2} = -k^2 Y \longrightarrow Y(y) = C \sin ky + D \cos ky$$

$$\frac{d^2 Z}{dz^2} = -l^2 Z \longrightarrow Z(z) = E \sin lz + F \cos lz$$

Cartesian Coordinates

$$\frac{\partial^2 V}{\partial x^2} + \frac{\partial^2 V}{\partial y^2} + \frac{\partial^2 V}{\partial z^2} = 0 \quad V(x, y, z) = X(x)Y(y)Z(z)$$

Example 3.5

$$\begin{aligned} X(x) &= Ae^{\sqrt{k^2+l^2}x} + Be^{-\sqrt{k^2+l^2}x} \\ Y(y) &= C \sin ky + D \cos ky \\ Z(z) &= E \sin lz + F \cos lz \end{aligned}$$

From the boundary conditions from (i) to (v),

$$V(x, y, z) = Ce^{-\pi\sqrt{(n/a)^2+(m/b)^2}x} \sin(n\pi y/a) \sin(m\pi z/b)$$


The most general linear combination is a *double* sum:

$$V(x, y, z) = \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} C_{n,m} e^{-\pi\sqrt{(n/a)^2+(m/b)^2}x} \sin(n\pi y/a) \sin(m\pi z/b)$$


(vi) $V = V_0(y, z)$ when $x = 0 \rightarrow V(0, y, z) = \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} C_{n,m} \sin(n\pi y/a) \sin(m\pi z/b) = V_0(y, z)$

$$\Rightarrow C_{n,m} = \frac{4}{ab} \int_0^a \int_0^b V_0(y, z) \sin(n\pi y/a) \sin(m\pi z/b) dy dz$$

If the end of the tube is a conductor at *constant* potential V_0 ,



$$V(x, y, z) = \frac{16V_0}{\pi^2} \sum_{n,m=1,3,5\ldots}^{\infty} \frac{1}{nm} e^{-\pi\sqrt{(n/a)^2+(m/b)^2}x} \sin(n\pi y/a) \sin(m\pi z/b).$$



3.3.2 Spherical Coordinates $\nabla^2 V = 0$

$$\frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial V}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial V}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 V}{\partial \phi^2} = 0$$

Assume the problem has **azimuthal symmetry**, so that V is independent of ϕ ;

$$\frac{\partial}{\partial r} \left(r^2 \frac{\partial V}{\partial r} \right) + \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial V}{\partial \theta} \right) = 0$$

$$\boxed{V(r, \theta) = R(r)\Theta(\theta)} \Rightarrow \frac{1}{R} \frac{d}{dr} \left(r^2 \frac{dR}{dr} \right) + \frac{1}{\Theta \sin \theta} \frac{d}{d\theta} \left(\sin \theta \frac{d\Theta}{d\theta} \right) = 0$$

$$\frac{1}{R} \frac{d}{dr} \left(r^2 \frac{dR}{dr} \right) = l(l+1) \longrightarrow R(r) = Ar^l + \frac{B}{r^{l+1}}$$

$$\frac{1}{\Theta \sin \theta} \frac{d}{d\theta} \left(\sin \theta \frac{d\Theta}{d\theta} \right) = -l(l+1) \longrightarrow \Theta(\theta) = P_l(\cos \theta) \text{ (Legendre polynomials)}$$

The general solution is

$$\Rightarrow \boxed{V(r, \theta) = \sum_{l=0}^{\infty} \left(A_l r^l + \frac{B_l}{r^{l+1}} \right) P_l(\cos \theta).}$$

Spherical Coordinates $V(r, \theta) = R(r)\Theta(\theta)$

Example 3.6 $V_0(\theta)$ is specified on the surface of a hollow sphere, of radius R . Find the **potential inside** the sphere.

$$V(r, \theta) = \sum_{l=0}^{\infty} \left(A_l r^l + \frac{B_l}{r^{l+1}} \right) P_l(\cos \theta).$$

→ $B_l = 0$; otherwise the potential would blow up at the origin.

$$V(r, \theta) = \sum_{l=0}^{\infty} A_l r^l P_l(\cos \theta) \quad \rightarrow \text{At } r = R; \quad V(R, \theta) = \sum_{l=0}^{\infty} A_l R^l P_l(\cos \theta) = V_0(\theta)$$

Legendre polynomials (like the sines) are *orthogonal* functions:

$$\int_{-1}^1 P_l(x) P_{l'}(x) dx = \int_0^\pi P_l(\cos \theta) P_{l'}(\cos \theta) \sin \theta d\theta = \begin{cases} 0, & \text{if } l' \neq l, \\ \frac{2}{2l+1}, & \text{if } l' = l. \end{cases}$$

Thus, multiplying by $P_{l'}(\cos \theta) \sin \theta$ and integrating,

$$A_l = \frac{2l+1}{2R^l} \int_0^\pi V_0(\theta) P_l(\cos \theta) \sin \theta d\theta \quad \longrightarrow \quad V(r, \theta) = \sum_{l=0}^{\infty} A_l r^l P_l(\cos \theta)$$

Spherical Coordinates $V(r, \theta) = R(r)\Theta(\theta)$

Example 3.7 $V_0(\theta)$ is specified on the surface of a hollow sphere, of radius R . Find the **potential outside** the sphere.

$$V(r, \theta) = \sum_{l=0}^{\infty} \left(A_l r^l + \frac{B_l}{r^{l+1}} \right) P_l(\cos \theta).$$

→ Now, $A_l = 0$; otherwise the potential would not go to zero at infinity.

$$V(r, \theta) = \sum_{l=0}^{\infty} \frac{B_l}{r^{l+1}} P_l(\cos \theta)$$

→ At $r = R$; $V(R, \theta) = \sum_{l=0}^{\infty} \frac{B_l}{R^{l+1}} P_l(\cos \theta) = V_0(\theta)$

Thus, multiplying by $P_{l'}(\cos \theta) \sin \theta$ and integrating,

$$B_l = \frac{2l+1}{2} R^{l+1} \int_0^\pi V_0(\theta) P_l(\cos \theta) \sin \theta d\theta \longrightarrow V(r, \theta) = \sum_{l=0}^{\infty} \frac{B_l}{r^{l+1}} P_l(\cos \theta)$$

Spherical Coordinates $V(r, \theta) = R(r)\Theta(\theta)$

Example 3.8 An **uncharged metal sphere** of radius R is placed in an otherwise uniform electric field $E = E_0 \mathbf{z}$. The **induced charge**, in turn, distorts the field in the neighborhood of the sphere.

Find the potential in the region outside the sphere.

$$V(r, \theta) = \sum_{l=0}^{\infty} \left(A_l r^l + \frac{B_l}{r^{l+1}} \right) P_l(\cos \theta).$$

The sphere is an equipotential \rightarrow we may as well set it to zero.

Far from the sphere the field is $E_0 \mathbf{z}$, hence the **boundary conditions** are


$$(i) \quad V = 0 \quad \text{when } r = R \quad \longrightarrow \quad A_l R^l + \frac{B_l}{R^{l+1}} = 0 \quad \longrightarrow \quad B_l = -A_l R^{2l+1}$$

$$V(r, \theta) = \sum_{l=0}^{\infty} A_l \left(r^l - \frac{R^{2l+1}}{r^{l+1}} \right) P_l(\cos \theta)$$

$$(ii) \quad V \rightarrow -E_0 r \cos \theta \quad \text{for } r \gg R \quad \longrightarrow \quad \sum_{l=0}^{\infty} A_l r^l P_l(\cos \theta) = -E_0 r \cos \theta$$

since $P_1(\cos \theta) = \cos \theta$, $A_1 = -E_0$, all other A_l 's zero.

$$\Rightarrow V(r, \theta) = -E_0 \left(r - \frac{R^3}{r^2} \right) \cos \theta. \quad \Rightarrow \sigma(\theta) = -\epsilon_0 \frac{\partial V}{\partial r} \Big|_{r=R} = 3\epsilon_0 E_0 \cos \theta$$



Spherical Coordinates $V(r, \theta) = R(r)\Theta(\theta)$

Example 3.9 A specified charge density $\sigma_0(\theta)$ is glued over the surface of a spherical shell of radius R .

Find the potential inside and outside the sphere.


$$V(r, \theta) = \sum_{l=0}^{\infty} \left(A_l r^l + \frac{B_l}{r^{l+1}} \right) P_l(\cos \theta) \quad \Rightarrow \quad \begin{aligned} V(r, \theta) &= \sum_{l=0}^{\infty} A_l r^l P_l(\cos \theta) & (r \leq R) \\ V(r, \theta) &= \sum_{l=0}^{\infty} \frac{B_l}{r^{l+1}} P_l(\cos \theta) & (r \geq R) \end{aligned}$$

→ At $r = R$; the boundary conditions are:

$$(1) \quad \sum_{l=0}^{\infty} A_l R^l P_l(\cos \theta) = \sum_{l=0}^{\infty} \frac{B_l}{R^{l+1}} P_l(\cos \theta) \quad \longrightarrow \quad B_l = A_l R^{2l+1}$$

$$(2) \quad \mathbf{E}_{\text{above}} - \mathbf{E}_{\text{below}} = \frac{\sigma}{\epsilon_0} \hat{\mathbf{n}} \quad \longrightarrow \quad \left(\frac{\partial V_{\text{out}}}{\partial r} - \frac{\partial V_{\text{in}}}{\partial r} \right) \Big|_{r=R} = -\frac{1}{\epsilon_0} \sigma_0(\theta)$$

$$\sum_{l=0}^{\infty} (2l+1) A_l R^{l-1} P_l(\cos \theta) = \frac{1}{\epsilon_0} \sigma_0(\theta)$$



$$A_l = \frac{1}{2\epsilon_0 R^{l-1}} \int_0^\pi \sigma_0(\theta) P_l(\cos \theta) \sin \theta d\theta.$$

$$B_l = A_l R^{2l+1}.$$


3.4 Multipole Expansion

3.4.1 Approximate Potentials at Large Distances

Example 3.10 Find the **approximate potential** at points far from the dipole.

$$V(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \left(\frac{q}{r_+} - \frac{q}{r_-} \right)$$

$$r_{\pm}^2 = r^2 + (d/2)^2 \mp rd \cos \theta = r^2 \left(1 \mp \frac{d}{r} \cos \theta + \frac{d^2}{4r^2} \right)$$



In the regime $r \gg d$,

$$\frac{1}{r_{\pm}} \cong \frac{1}{r} \left(1 \mp \frac{d}{r} \cos \theta \right)^{-1/2} \cong \frac{1}{r} \left(1 \pm \frac{d}{2r} \cos \theta \right) \longrightarrow V(\mathbf{r}) \cong \frac{1}{4\pi\epsilon_0} \frac{qd \cos \theta}{r^2}.$$

Monopole
($V \sim 1/r$)


Dipole
($V \sim 1/r^2$)

Quadrupole
($V \sim 1/r^3$)

Octopole
($V \sim 1/r^4$)

Approximate Potentials by multiple expansion

Now, develop a **systematic expansion for the potential of an arbitrary localized charge distribution, in powers of $1/r$.**



$$V(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \int \frac{1}{z} \rho(\mathbf{r}') d\tau'$$

$$z^2 = r^2 + (r')^2 - 2rr' \cos \theta' = r^2 \left[1 + \left(\frac{r'}{r}\right)^2 - 2\left(\frac{r'}{r}\right) \cos \theta' \right]$$

$$z = r\sqrt{1 + \epsilon} \quad \text{where} \quad \epsilon \equiv \left(\frac{r'}{r}\right) \left(\frac{r'}{r} - 2 \cos \theta'\right) \rightarrow \text{much less than 1}$$

$$\frac{1}{z} = \frac{1}{r} (1 + \epsilon)^{-1/2} = \frac{1}{r} \left(1 - \frac{1}{2}\epsilon + \frac{3}{8}\epsilon^2 - \frac{5}{16}\epsilon^3 + \dots \right)$$

$$= \frac{1}{r} \left[1 - \frac{1}{2} \left(\frac{r'}{r}\right) \left(\frac{r'}{r} - 2 \cos \theta'\right) + \frac{3}{8} \left(\frac{r'}{r}\right)^2 \left(\frac{r'}{r} - 2 \cos \theta'\right)^2 - \frac{5}{16} \left(\frac{r'}{r}\right)^3 \left(\frac{r'}{r} - 2 \cos \theta'\right)^3 + \dots \right]$$

$$= \frac{1}{r} \left[1 + \left(\frac{r'}{r}\right) (\cos \theta') + \left(\frac{r'}{r}\right)^2 (3 \cos^2 \theta' - 1)/2 + \left(\frac{r'}{r}\right)^3 (5 \cos^3 \theta' - 3 \cos \theta')/2 + \dots \right]$$

$$\frac{1}{z} = \frac{1}{r} \sum_{n=0}^{\infty} \left(\frac{r'}{r}\right)^n P_n(\cos \theta') \longrightarrow V(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \sum_{n=0}^{\infty} \frac{1}{r^{(n+1)}} \int (r')^n P_n(\cos \theta') \rho(\mathbf{r}') d\tau',$$

$$V(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \left[\frac{1}{r} \int \rho(\mathbf{r}') d\tau' + \frac{1}{r^2} \int r' \cos \theta' \rho(\mathbf{r}') d\tau' + \frac{1}{r^3} \int (r')^2 \left(\frac{3}{2} \cos^2 \theta' - \frac{1}{2} \right) \rho(\mathbf{r}') d\tau' + \dots \right]$$

monopole
dipole
quadrupole

3.4.2 The Monopole and Dipole Terms

$$V(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \left[\overset{\text{monopole}}{\frac{1}{r} \int \rho(\mathbf{r}') d\tau'} + \overset{\text{dipole}}{\frac{1}{r^2} \int r' \cos \theta' \rho(\mathbf{r}') d\tau'} + \overset{\text{quadrupole}}{\frac{1}{r^3} \int (r')^2 \left(\frac{3}{2} \cos^2 \theta' - \frac{1}{2} \right) \rho(\mathbf{r}') d\tau'} + \dots \right]$$


→ Ordinarily, the **multipole expansion** is dominated (at large r) by the **monopole** term.

→ If the **total charge is zero**, the dominant term in the potential will be the **dipole**.

$$V_{\text{dip}}(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \frac{1}{r^2} \int r' \cos \theta' \rho(\mathbf{r}') d\tau'$$

$$\longrightarrow r' \cos \theta' = \hat{\mathbf{r}} \cdot \mathbf{r}'$$

$$\longrightarrow V_{\text{dip}}(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \frac{1}{r^2} \hat{\mathbf{r}} \cdot \int \mathbf{r}' \rho(\mathbf{r}') d\tau'$$



$$\boxed{\mathbf{p} \equiv \int \mathbf{r}' \rho(\mathbf{r}') d\tau'} \quad \Rightarrow \text{Dipole moment of the charge distribution}$$

$$\Rightarrow \boxed{V_{\text{dip}}(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \frac{\mathbf{p} \cdot \hat{\mathbf{r}}}{r^2}}$$

3.4.3 Origin of Coordinates in Multipole Expansions

$$V(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \left[\overset{\text{monopole}}{\frac{1}{r} \int \rho(\mathbf{r}') d\tau'} + \overset{\text{dipole}}{\frac{1}{r^2} \int r' \cos \theta' \rho(\mathbf{r}') d\tau'} + \overset{\text{quadrupole}}{\frac{1}{r^3} \int (r')^2 \left(\frac{3}{2} \cos^2 \theta' - \frac{1}{2} \right) \rho(\mathbf{r}') d\tau'} + \dots \right]$$


A point charge at the origin constitutes a "pure" monopole.

If it is not at the origin, it's no longer a pure monopole.

→ The monopole term, $(1/4\pi\epsilon_0)q/r$ is not quite correct.

→ The exact monopole potential is $(1/4\pi\epsilon_0)q/r$

→ So, moving the origin can radically alter the expansion.




But there is an important exception:

→ If the total charge is zero, the dipole moment is independent of the choice of origin.

Let's displace the origin by an amount **a**

$$\begin{aligned} \bar{\mathbf{p}} &= \int \bar{\mathbf{r}}' \rho(\mathbf{r}') d\tau' = \int (\mathbf{r}' - \mathbf{a}) \rho(\mathbf{r}') d\tau' \\ &= \int \mathbf{r}' \rho(\mathbf{r}') d\tau' - \mathbf{a} \int \rho(\mathbf{r}') d\tau' = \mathbf{p} - Q\mathbf{a} \end{aligned}$$

if $Q = 0$, then $\bar{\mathbf{p}} = \mathbf{p}$.



3.4.4 The Electric Field of a Dipole

$$V(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \left[\overset{\text{monopole}}{\frac{1}{r} \int \rho(\mathbf{r}') d\tau'} + \overset{\text{dipole}}{\frac{1}{r^2} \int r' \cos \theta' \rho(\mathbf{r}') d\tau'} + \overset{\text{quadrupole}}{\frac{1}{r^3} \int (r')^2 \left(\frac{3}{2} \cos^2 \theta' - \frac{1}{2} \right) \rho(\mathbf{r}') d\tau'} + \dots \right]$$

Now let's calculate the electric field of a (pure, nearly point) dipole.


If we choose coordinates so that \mathbf{p} lies at the origin and points in the z direction

$$V_{\text{dip}}(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \frac{\mathbf{p} \cdot \hat{\mathbf{r}}}{r^2} = \frac{p \cos \theta}{4\pi\epsilon_0 r^2}$$


$$E_r = -\frac{\partial V}{\partial r} = \frac{2p \cos \theta}{4\pi\epsilon_0 r^3},$$

$$E_\theta = -\frac{1}{r} \frac{\partial V}{\partial \theta} = \frac{p \sin \theta}{4\pi\epsilon_0 r^3},$$

$$E_\phi = -\frac{1}{r \sin \theta} \frac{\partial V}{\partial \phi} = 0.$$



$$\mathbf{E}_{\text{dip}}(r, \theta) = \frac{p}{4\pi\epsilon_0 r^3} (2 \cos \theta \hat{\mathbf{r}} + \sin \theta \hat{\boldsymbol{\theta}}).$$



$$\mathbf{E}_{\text{dip}}(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \frac{1}{r^3} [3(\mathbf{p} \cdot \hat{\mathbf{r}})\hat{\mathbf{r}} - \mathbf{p}]. \quad (\text{Prob. 3.33})$$

