

Leveraging allelic series to enhance QTL detection

Frederick J. Boehm^{1, }
✉ frederick.boehm@umassmed.edu
Clare M. Smith² Megan Proulx¹ Daniel M. Gatti³ Michael Kiritsy¹ Sherry Kurtz⁴ Karen Elkins⁴ Gillian Beamer⁵ Christopher M. Sassetti¹

¹ Microbiology and Physiological Systems, University of Massachusetts Medical School
² Duke University
³ College of the Atlantic
⁴ Food and Drug Administration
⁵ Tufts University

Introduction

- We examined Mtb-related traits in both a Collaborative Cross cohort (52 lines) and a Diversity Outbred cohort (763 mice)
- In CC and DO QTL mapping, we typically work with 8 founder allele probabilities

$$Trait = p_A b_A + p_B b_B + p_C b_C + p_D b_D + p_E b_E + p_F b_F + p_G b_G + p_H b_H + \epsilon$$

- Crouse and Valdar (personal communication) developed a method for inferring the allelic series at a QTL
- In DO and CC, some QTL have fewer than 8 alleles
- Using this fact in our QTL scans may enhance QTL detection ([Jansen, Jannink, and Beavis 2003](#))

Objectives

1. Infer QTL allelic series in CC mice
2. Use CC QTL allelic series to inform QTL mapping in DO mice

Methods





Figure 1: Experimental design in DO mice



Figure 2: Experimental design in CC mice

CC allelic series informs QTL mapping in DO



- We mapped QTL in CC cohort with standard methods ([Broman et al. 2019](#))

- We used Tree-based Inference of Multi-allelism with Bayesian Regression (TIMBR) ([Crouse 2018](#)) to infer allelic series at CC QTL

- For each CC QTL, we scanned the DO cohort with reduced models that reflected the CC QTL allelic series

- Example of reducing a model with inferred allelic series
For the allelic series: 0,1,1,0,2,1,0,0:
 1. Set $p_0 = p_A + p_D + p_G + p_H$,
 $p_1 = p_B + p_C + p_F$,
 $p_2 = p_E$
 2. Use model

$$Y = \beta_0 p_0 + \beta_1 p_1 + \beta_2 p_2 + \epsilon$$

in QTL scan

Results

1. Traditional QTL mapping in CC identified 14 suggestive peaks
2. For each CC QTL, we:
 1. inferred allelic series
 2. fitted reduced linear models (informed by allelic series) for QTL detection in DO
 3. identified any proximal DO QTL with LOD greater than 3

QTL peaks on Chr 4				
trait	cohort	chr	pos	lod
Spleen CFU in DO	DO	4	152.95	3.09
VEGF in CC	CC	4	154.07	6.83

Future research

1. Determine DO QTL p-values with approximate permutation tests ([Churchill and Doerge 1994](#))
2. Consider multiple allelic series per CC QTL

Image credits

<https://webcomicms.net/clipart-10020867-pictures-cartoon-mouse>

<https://www.latimes.com/science/story/2020-04-13/old-vaccines-for-other-germs-being-tested-against-the-new-coronavirus>

<https://www.lshtm.ac.uk/research/research-action/features/ending-tb-race-control-disease-hiding>

Vitruvian Mouse by Eleni Jaecklein

References

Broman, Karl W, Daniel M Gatti, Petr Simecek, Nicholas A Furlotte, Piotr Prins, Saunak Sen, Brian S Yandell, and Gary A Churchill. 2019. "R/Qtl2: Software for Mapping Quantitative Trait Loci with High-Dimensional Data and Multiparent Populations." *Genetics* 211 (2): 495–502.

Churchill, Gary A, and Rebecca W Doerge. 1994. "Empirical Threshold Values for Quantitative Trait Mapping." *Genetics* 138 (3): 963–71.

Crouse, Wesley. 2018. *TIMBR: Tree-Based Inference of Multiallelism via Bayesian Regression*.

Jansen, Ritser C, Jean-Luc Jannink, and William D Beavis. 2003. "Mapping Quantitative Trait Loci in Plant Breeding Populations." *Crop Science* 43 (3): 829–34.