

Leveraging allelic series to enhance QTL detection

Frederick J. Boehm¹, 
✉ frederick.boehm@umassmed.edu

Clare M. Smith² Megan Proulx¹ Daniel M. Gatti³ Michael Kiritsy¹ Sherry Kurtz⁴ Karen Elkins⁴ Gillian Beamer⁵ Christopher M. Sassetti¹

¹ Microbiology and Physiological Systems, University of Massachusetts Medical School
² Duke University
³ College of the Atlantic
⁴ Food and Drug Administration
⁵ Tufts University

Introduction

- In CC and DO, we typically work with 8 founder allele probabilities

$$Trait = p_A b_A + p_B b_B + p_C b_C + p_D b_D + p_E b_E + p_F b_F + p_G b_G + p_H b_H + \epsilon$$

- Standard QTL methods allow every founder to have its own allele
- However, founders often share alleles
- Thus, in DO and CC, some QTL have fewer than 8 alleles
- Using this fact in our QTL scans may enhance QTL detection (Jansen, Jannink, and Beavis 2003)

Objectives

1. Infer QTL allelic series in CC mice
2. Use CC QTL allelic series to map QTL in DO mice

Methods

- We measured *M. tuberculosis* burden in lungs and spleens in a DO cohort (763 mice) and a CC cohort (52 lines)
 - We also measured cytokine levels in the CC cohort

Allelic series informs QTL mapping in a second cohort

- We mapped QTL in CC cohort with standard methods (Broman et al. 2019)
- We used Tree-based Inference of Multi-allelism with Bayesian Regression (TIMBR) (Crouse 2018) to infer allelic series at CC QTL

- For each CC QTL, we scanned the DO cohort with reduced models that reflected the CC QTL allelic series
 - Example of reducing a model with inferred allelic series For the allelic series: 0,1,1,0,2,1,0,0:
 1. Set $p_0 = p_A + p_D + p_G + p_H$, $p_1 = p_B + p_C + p_F$
 $p_2 = p_E$
 2. Use model

$$Y = \beta_0 p_0 + \beta_1 p_1 + \beta_2 p_2 + \epsilon$$

in QTL scan

Results

1. Traditional QTL mapping in CC identified 14 suggestive peaks
2. For each CC QTL, we:
 1. inferred allelic series
 2. fitted reduced linear models (informed by allelic series) for QTL detection in DO
 3. identified any proximal DO QTL with LOD greater than 2

QTL peaks on Chr 4				
trait	cohort	chr	pos	lod
Spleen CFU in DO	DO	4	152.95	3.09
VEGF in CC	CC	4	154.07	6.83

Future research

1. Determine DO QTL p-values with approximate permutation tests (Churchill and Doerge 1994)
2. Consider multiple allelic series per CC QTL

References

Broman, Karl W, Daniel M Gatti, Petr Simecek, Nicholas A Furlotte, Piotr Prins, Šaunak Sen, Brian S Yandell, and Gary A Churchill. 2019. "R/Qtl2: Software for Mapping Quantitative Trait Loci with High-Dimensional Data and Multiparent Populations." *Genetics* 211 (2): 495–502.

Churchill, Gary A, and Rebecca W Doerge. 1994. "Empirical Threshold Values for Quantitative Trait Mapping." *Genetics* 138 (3): 963–71.

Crouse, Wesley. 2018. *TIMBR: Tree-Based Inference of Multiallelism via Bayesian Regression*.

Jansen, Ritsert C, Jean-Luc Jannink, and William D Beavis. 2003. "Mapping Quantitative Trait Loci in Plant Breeding Populations." *Crop Science* 43 (3): 829–34.