

Opinion Dynamics

Anahita Mirtabatabaei



Center for Control, Dynamical Systems & Computation
University of California at Santa Barbara

September 16, 2010



1 / 26

Opinion Dynamics



2 / 26

Motivations

- Economics (Sznajd-Weron, 2002)
- Politics (Ben-Naim, 2005)
- Physics (Ben-Naim et al., 2003)
- Sociology (Tavares, 2007)

Can we convince others to follow us?

1 / 26

Motivations

- Economics (Sznajd-Weron, 2002)
- Politics (Ben-Naim, 2005)
- Physics (Ben-Naim et al., 2003)
- Sociology (Tavares, 2007)

Can we convince others to follow us?




3 / 26

4 / 26

Synchronized Bounded Confidence (SBC) Model

- n agents, with n bounds of confidence $\{r_1, \dots, r_n\}$
- agent i has opinion $x_i(t) \in \mathbb{R}$
- j is i 's out-neighbor if $|x_i(t) - x_j(t)| \leq r_i$




$$x(t+1) = A(x(t))x(t)$$

$$a_{ij}(x(t)) = \begin{cases} \frac{1}{\# \text{ of } i\text{'s neighbors}} & \text{if } j \text{ is } i\text{'s out-neighbor} \\ 0 & \text{otherwise} \end{cases}$$


5 / 26

For equal initial opinion vector and bounds vector, an SBC system exhibits more complex behavior than an SBI system

SBC




SBI



Synchronized Bounded Influence (SBI) Model

- n agents, with n bounds of confidence $\{r_1, \dots, r_n\}$
- agent i has opinion $x_i(t) \in \mathbb{R}$
- j is i 's out-neighbor if $|x_i(t) - x_j(t)| \leq r_j$



$$x(t+1) = A(x(t))x(t)$$

$$a_{ij}(x(t)) = \begin{cases} \frac{1}{\# \text{ of } i\text{'s neighbors}} & \text{if } j \text{ is } i\text{'s out-neighbor} \\ 0 & \text{otherwise} \end{cases}$$


6 / 26

■ System with homogeneous bounds

converges in finite time

the order of opinions is preserved

in steady state agents are in agreement or disconnected




■ System with heterogeneous bounds

may converge in infinite time

pseudo-stable configurations

disconnected clusters may reconnect



7 / 26


8 / 26

■ Homogeneous SBC (Hegselman-Krause) Model

time complexity of convergence (Bullo et al. 2007)

consensus (Olshevsky et al. 2009)

final opinion vector (Blondel et al. 2009)




■ Heterogeneous SBC (Hegselman-Krause) Model

enhances the chances for consensus (Lorenz 2008)

classification of agents (Lorenz 2006)

product of infinite adjacency matrices (Lorenz 2006)




Conjectures

when does convergence start?



what is the final value?

who moves and who stops?

rate and direction of moving ones?





Conjectures



when does convergence start?

what is the final value?


who moves and who stops?




rate and direction of moving ones?

do all SBC and SBI systems converge?

Classification of Agents




Classification of Agents



13 / 26


Classification of Agents



14 / 26


Classification of Agents

Open-minded
Moderate-minded
Closed-minded



15 / 26

Conjectures



16 / 26

$$A(x) = \begin{bmatrix} C & 0 & 0 \\ 0 & M & 0 \\ \Theta_C & \Theta_M & \Theta \end{bmatrix}$$



Final value at constant topology

$$\begin{aligned} x^*(x) &= \lim_{t \rightarrow \infty} A(x)^t x \\ &= \begin{bmatrix} C & 0 & 0 \\ 0 & M^* & 0 \\ (I - \Theta)^{-1} \Theta_C C & (I - \Theta)^{-1} \Theta_M M^* & 0 \end{bmatrix} x \end{aligned}$$

If $A(x^*(x)) = A(x)$, then

- $x^*(x)$ is an equilibrium vector
- their proximity graph contains no moderate-minded

Conjectures



when does convergence start?

what is the final value?

who moves and who stops?

rate and direction of moving ones?

do all SBC and SBI systems converge?

Equi-topology neighborhood of $z \in \mathbb{R}^n$

The set of vectors $y \in \mathbb{R}^n$ such that for all $i \in \{1, \dots, n\}$

$$\begin{aligned} |y_j - z_j| &< \epsilon_i(z), & \text{if } \epsilon_i(z) > 0, \text{ and} \\ |y_j - z_j| &= \epsilon_i(z), & \text{if } \epsilon_i(z) = 0, \end{aligned}$$

where

$$\epsilon_i(z) = 0.5 \min\{||z_j - z_i| - R| \mid j \in \{1, \dots, n\} \setminus \{i\}, R \in \{r_i, r_j\}\}.$$

Invariant equi-topology neighborhood of $z \in \mathbb{R}^n$

The set of vectors $y \in \mathbb{R}^n$ such that for all $i \in \{1, \dots, n\}$

$$\begin{aligned} |y_j - z_j| &< \delta_i(z), & \text{if } \delta_i(z) > 0, \text{ and} \\ |y_j - z_j| &= \delta_i(z), & \text{if } \delta_i(z) = 0. \end{aligned}$$

where



$$\delta_i(z) = \min\{\epsilon_j(z) \mid j \text{ is a predecessor of } i\}.$$

Theorem (Sufficient condition for constant topology and convergence)

Consider a trajectory $x(t)$ of an SBC or SBI system. Assume that there exists an equilibrium opinion vector $z \in \mathbb{R}^n$ for the system such that $x(0) \in \mathbb{R}^n$ belongs to the invariant equi-topology neighborhood of z . Then, for all $t \geq 0$:

- 1 $x(t)$ takes value in the equi-topology neighborhood of z ;
- 2 the proximity digraphs $G_r(x(t))$ and $G_r(z)$ are equal;
- 3 $G_r(x(t))$ contains no moderate-minded component; and
- 4 $x(t)$ converges to the final value at constant topology of $x(0)$ as $t \rightarrow \infty$.

Conjectures





when does convergence start?

what is the final value?

who moves and who stops?


rate and direction of moving ones?

do all SBC and SBI systems converge?



Theorem (Evolution under Constant Topology after τ)

- $x(t)$ converges to $x^*(x(\tau))$
- no moderate minded at τ
- if $\rho(\Theta_4) \geq \rho(\Theta_{1,2,3})$, then for all $i \in G_{\Theta_1}$ and $j \in G_{\Theta_4}$
 - $\lim_{t \rightarrow \infty} k_i(x(t)) = \rho(\Theta_4)$
 - there exists $T \geq \tau$, after which $(x_i(t) - x_i^*)(x_j(t) - x_j^*) \geq 0$



Rate of Convergence

Agent's per-step convergence factor


$$k_i(x(t)) = \frac{x_i(t+1) - x_i^*(x(t))}{x_i(t) - x_i^*(x(t))}$$

$k_i = 1 - \text{rate of convergence of agent } i$



monotonic convergence toward final value $\equiv 0 \leq k_i \leq 1$

In a Real Society

- the initial opinion of an open-minded has no effect, $x^* = \begin{bmatrix} C & 0 \\ \Theta_C^* & 0 \end{bmatrix} x$.
- an individual converges to his final decision as slow as the slowest group.
- the leader govern followers direction and rate.
- one can become a leader by joining a large strongly connected group:



Conjectures





when does convergence start?

what is the final value?

who moves and who stops?

rate and direction of moving ones?

do all SBC and SBI systems converge?



Future Work

■ Convergence of SBC and SBI systems

1. what is the basin of attraction of x^* ?
2. what are possible x^* 's for a system?
3. does the system fall into one of those basins of attraction?

Theorem (Convergence of products of stochastic matrices, Lorenz '06)

$$\lim_{t \rightarrow \infty} A(t, 0) = \begin{bmatrix} C_1 & & & \\ & \ddots & & \\ & & C_m & \\ ? & & & 0 \end{bmatrix} A(t_0, 0)$$

■ How can one become a leader by changing his r ?