

# Contraction Theory for Networked Optimization and Control

Francesco Bullo



Center for Control,  
Dynamical Systems & Computation  
University of California at Santa Barbara  
<https://fbullo.github.io/ctds>

Workshop on Emerging Challenges of Network-Enabled Control and Optimization  
CDC Workshop, Singapore, Dec 12, 2023

Many thanks to: General Chair Jie Chen and General Co-Chair Huansui Zhang  
Shandong University of Science and Technology & Nanyang Technological University

# Acknowledgments



Veronica Centorrino  
Scuola Sup Meridionale



Lily Cothren  
UC Boulder



Emiliano Dall'Anese  
UC Boulder



Alex Davydov  
UC Santa Barbara



Anand Gokhale  
UC Santa Barbara



Giovanni Russo  
Univ Salerno


A. Davydov, V. Centorrino, A. Gokhale, G. Russo, and F. Bullo. Contracting dynamics for time-varying convex optimization. *IEEE Transactions on Automatic Control*, June 2023. Submitted

A. Gokhale, A. Davydov, and F. Bullo. Contractivity of distributed optimization and Nash seeking dynamics. *IEEE Control Systems Letters*, Sept. 2023. To appear

L. Cothren, F. Bullo, and E. Dall'Anese. Singular perturbation via contraction theory. *IEEE Transactions on Automatic Control*, Oct. 2023. Submitted


# Solving optimization problems via dynamical systems

- studies in linear and nonlinear programming (Arrow, Hurwicz, and Uzawa 1958)
- neural networks (Hopfield and Tank 1985)
- analog circuits (Kennedy and Chua 1988)
- optimization on manifolds (Brockett 1991)
- ...



## Continuous-time optimizations solvers

- ① online feedback optimization
- ② distributed optimization
- ③ parametric convex optimization
- ④ model predictive control
- ⑤ control barrier functions




Online feedback optimization


Slide courtesy of Emiliano Dall'Anese, University of Colorado Boulder

**contractivity = robust computationally-friendly stability**

fixed point theory + Lyapunov stability theory + geometry of metric spaces



**search for** contraction properties  
**design** engineering systems to be contracting  
**verify** correct/safe behavior via known Lipschitz constants



"Continuous improvement is better than delayed perfection" **Mark Twain**

- 2023 ACC Workshop "Contraction Theory for Systems, Control, and Learning" <http://motion.me.ucsb.edu/contraction-workshop-2023>
- 2021 IEEE CDC Tutorial session "Contraction Theory for Machine Learning" <https://sites.google.com/view/contractiontheory> (PDFs and youtube videos)
- 2022 IEEE CDC plenary presentation "Contraction Theory in Systems and Control" <https://fbullo.github.io/talks/2022-12-FBullo-ContractionSystemsControl-CDC.pdf>
- Textbook: Contraction Theory for Dynamical Systems, Francesco Bullo, rev 1.1, Mar 2023. (Book and slides freely available)  
<https://fbullo.github.io/ctds>
- 2023 Comprehensive tutorial slides: <https://fbullo.github.io/ctds>
- 2023 Sep: Youtube lectures: "Minicourse on Contraction Theory"  
<https://youtu.be/FQV5PrRHks8> 12h in 6 lectures

- Three CDC2023 invited sessions on **Contraction Theory for Analysis, Synchronization and Regulation**, tomorrow Wednesday!

# Outline

§1. Introduction

§2. Basic contractivity concepts

§3. Examples: Gradient systems defined by strongly convex functions are contracting



§4. Theory: Equilibrium tracking in parametric optimization

§5. Application: Online feedback optimization

§6. Conclusions

# Induced matrix norms

Vector norm	Induced matrix norm	Induced matrix log norm
$\ x\ _1 = \sum_{i=1}^n  x_i $	$\ A\ _1 = \max_{j \in \{1, \dots, n\}} \sum_{i=1}^n  a_{ij} $ = max column "absolute sum" of $A$	$\mu_1(A) = \max_{j \in \{1, \dots, n\}} \left( a_{jj} + \sum_{i=1, i \neq j}^n  a_{ij}  \right)$ absolute value only off-diagonal
$\ x\ _2 = \sqrt{\sum_{i=1}^n x_i^2}$	$\ A\ _2 = \sqrt{\lambda_{\max}(A^\top A)}$	$\mu_2(A) = \lambda_{\max}\left(\frac{A + A^\top}{2}\right)$
$\ x\ _\infty = \max_{i \in \{1, \dots, n\}}  x_i $	$\ A\ _\infty = \max_{i \in \{1, \dots, n\}} \sum_{j=1}^n  a_{ij} $ = max row "absolute sum" of $A$	$\mu_\infty(A) = \max_{i \in \{1, \dots, n\}} \left( a_{ii} + \sum_{j=1, j \neq i}^n  a_{ij}  \right)$ absolute value only off-diagonal



$\dot{x} = F(x)$       on  $\mathbb{R}^n$  with norm  $\|\cdot\|$  and induced log norm  $\mu(\cdot)$

## One-sided Lipschitz constant

$$\begin{aligned}\text{osLip}(F) &= \inf\{b \in \mathbb{R} \text{ such that } \|F(x) - F(y), x - y\| \leq b\|x - y\|^2 \text{ for all } x, y\} \\ &= \sup_x \mu(DF(x))\end{aligned}$$

For **scalar map**  $f$ ,  $\text{osLip}(f) = \sup_x f'(x)$

For **affine map**  $F_A(x) = Ax + a$


$$\text{osLip}_{2,P}(F_A) = \mu_{2,P}(A) \leq \ell \iff A^\top P + AP \preceq 2\ell P$$

$$\text{osLip}_{\infty,\eta}(F_A) = \mu_{\infty,\eta}(A) \leq \ell \iff a_{ii} + \sum_{j \neq i} |a_{ij}| \eta_i / \eta_j \leq \ell$$

## Banach contraction theorem for continuous-time dynamics:

If  $-c := \text{osLip}(F) < 0$ , then

- ①  $F$  is **infinitesimally contracting** = distance between trajectories decreases exp fast ( $e^{-ct}$ )
- ②  $F$  has a unique, glob exp stable equilibrium  $x^*$
- ③ global Lyapunov functions  $V_1(x) = \|x - x^*\|^2$  and  $V_2(x) = \|F(x)\|^2$



ball centered at  $x(t)$  with radius  $e^{-ct}$

## Euler discretization theorem for contracting dynamics

Given arbitrary norm  $\|\cdot\|$  and differentiable  $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$ , equivalent statements

- ①  $\dot{x} = F(x)$  is infinitesimally contracting
- ② there exists  $\alpha > 0$  such that  $x_{k+1} = x_k + \alpha F(x_k)$  is contracting

Interconnected subsystems:  $x_i \in \mathbb{R}^{N_i}$  and  $x_{-i} \in \mathbb{R}^{N-N_i}$ :

$$\dot{x}_i = F_i(x_i, x_{-i}), \quad \text{for } i \in \{1, \dots, n\}$$

**Network contraction theorem.** Assume

- **contractivity wrt  $x_i$ :**  $\text{osLip}_{x_i}(F_i) \leq -c_i < 0$ , uniformly in  $x_{-i}$
- **Lipschitz wrt  $x_j$ ,  $j \neq i$ :**  $\text{Lip}_{x_j}(F_i) \leq \ell_{ij}$ , uniformly in  $x_{-j}$

- the Lipschitz constants matrix  $\Gamma = \begin{bmatrix} -c_1 & \dots & \ell_{1n} \\ \vdots & & \vdots \\ \ell_{n1} & \dots & -c_n \end{bmatrix}$  is **Hurwitz**


$\implies$  **interconnected system** is contracting wrt rate  $|\alpha(\Gamma)|$

**contractivity = robust computationally-friendly stability**

fixed point theory + Lyapunov stability theory + geometry of metric spaces

**highly-ordered transient and asymptotic behavior, no anonymous constants/functions:**

- ① unique globally exponential stable equilibrium  
& two natural Lyapunov functions
- ② robustness properties
  - bounded input, bounded output (iss)
  - finite input-state gain
  - robustness margin wrt unmodeled dynamics
  - robustness margin wrt delayed dynamics
- ③ periodic input, periodic output
- ④ modularity and interconnection properties
- ⑤ accurate numerical integration and equilibrium point computation



**search for** contraction properties  
**design** engineering systems to be contracting  
**verify** correct/safe behavior via known Lipschitz constants

# Outline

§1. Introduction

§2. Basic contractivity concepts

§3. Examples: Gradient systems defined by strongly convex functions are contracting

§4. Theory: Equilibrium tracking in parametric optimization

§5. Application: Online feedback optimization

§6. Conclusions

**Kachurovskii's Theorem:** For differentiable  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ , equivalent statements:

- ①  $f$  is **strongly convex** with parameter  $\nu$  (and minimum  $x^*$ )
- ②  $-\nabla f$  is  **$\nu$ -strongly infinitesimally contracting** (with equilibrium  $x^*$ ), that is

$$(-\nabla f(x) + \nabla f(y))^\top (x - y) \leq -\nu \|x - y\|_2^2$$

R. I. Kachurovskii. Monotone operators and convex functionals. *Uspekhi Matematicheskikh Nauk*, 15(4):213–215, 1960

## Example #1: Gradient dynamics for strongly convex function

Given differentiable, strongly convex  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  with parameter  $\nu > 0$ , **gradient dynamics**

$$\dot{x} = F_G(x) := -\nabla f(x)$$

$F_G$  is infinitesimally contracting wrt  $\|\cdot\|_2$  with rate  $\nu$

unique globally exp stable point is global minimum

## Example #2: Primal-dual gradient dynamics

strongly convex function  $f$

s.t.  $0 \prec \nu_{\min} I_n \preceq \text{Hess } f \preceq \nu_{\max} I_n$

constraint matrix  $A$

s.t.  $0 \prec a_{\min} I_m \preceq AA^\top \preceq a_{\max} I_m$

(independent rows)

**linearly constrained optimization:**

$$\begin{aligned} & \min_{x \in \mathbb{R}^n} f(x) \\ & \text{subj. to } Ax = b \end{aligned}$$

**primal-dual gradient dynamics:**

$$\begin{bmatrix} \dot{x} \\ \dot{\lambda} \end{bmatrix} = F_{\text{PDG}}(x, \lambda) := \begin{bmatrix} -\nabla f(x) - A^\top \lambda \\ Ax - b \end{bmatrix}$$

$F_{\text{PDG}}$  is infinitesimally contracting wrt  $\|\cdot\|_{2,P^{1/2}}$  with rate  $c$

$$P = \begin{bmatrix} I_n & \alpha A^\top \\ \alpha A & I_m \end{bmatrix} \text{ with } \alpha = \frac{1}{2} \min \left\{ \frac{1}{\nu_{\max}}, \frac{\nu_{\min}}{a_{\max}} \right\} \quad \text{and} \quad c = \frac{1}{4} \min \left\{ \frac{a_{\min}}{\nu_{\max}}, \frac{a_{\min}}{a_{\max}} \nu_{\min} \right\}$$

## Example #3: Laplacian-based distributed gradient

Given  $\Pi_n = I_n - \mathbb{1}_n \mathbb{1}_n^\top / n$  = orthogonal projection onto  $\text{span}\{\mathbb{1}_n\}^\perp$ ,

$$0 \prec \lambda_2 \Pi_n \preceq L \preceq \lambda_n I_n$$

**decomposable cost:**  $\min_{x \in \mathbb{R}^n} \sum_{i=1}^n f_i(x)$  where each  $f_i$  is  $\nu_i$ -strongly convex

$$\begin{cases} \min_{x_{[i]} \in \mathbb{R}} & \sum_{i=1}^n f_i(x_{[i]}) \\ \text{subj. to} & \sum_{j=1}^n a_{ij}(x_i - x_j) = 0 \end{cases}$$

**Laplacian-based distributed gradient** (primal-dual gradient,  $2n$  vars):

$$\begin{cases} \dot{x}_{[i]} = -\nabla f_i(x_{[i]}) - \sum_{j=1}^n a_{ij}(\lambda_i - \lambda_j) & \text{for each node } i \\ \dot{\lambda}_i = \sum_{j=1}^n a_{ij}(x_i - x_j) & \text{for each node } i \end{cases}$$

$F_{\text{Laplacian-DistributedG}}$  is infinitesimally contracting<sup>†</sup> with  $c = \frac{1}{4} \left( \frac{\lambda_2}{\lambda_n} \right)^2 \min_i \nu_i$

## Detour: Composite optimization and the proximal operator

**composite minimization** (cost = sum of terms with structurally different properties):

$$x^* = \operatorname{argmin}_{x \in \mathbb{R}^n} f(x, u) + g(x)$$

$f(x, u)$  is convex and differentiable in  $x$        $g : \mathbb{R}^n \rightarrow \overline{\mathbb{R}}$  is convex, closed, and proper (ccp)

**proximal operator:**

$$\operatorname{prox}_{\gamma g}(z) := \operatorname{argmin}_{x \in \mathbb{R}^n} g(x) + \frac{1}{2\gamma} \|x - z\|_2^2$$

generalized form of projection for nonsmooth/constrained/large-scale/distributed optimization

## Detour: Composite optimization and the proximal operator

**composite minimization** (cost = sum of terms with structurally different properties):

$$x^* = \operatorname{argmin}_{x \in \mathbb{R}^n} f(x, u) + g(x)$$

$f(x, u)$  is convex and differentiable in  $x$        $g : \mathbb{R}^n \rightarrow \overline{\mathbb{R}}$  is convex, closed, and proper (ccp)

**proximal operator:**

$$\operatorname{prox}_{\gamma g}(z) := \operatorname{argmin}_{x \in \mathbb{R}^n} g(x) + \frac{1}{2\gamma} \|x - z\|_2^2$$

generalized form of projection for nonsmooth/constrained/large-scale/distributed optimization

### Equivalence property:

①  $x^*$  is minimizer for:

$$\min_{x \in \mathbb{R}^n} f(x, u) + g(x)$$

②  $x^*$  is fixed point for:

$$x = \operatorname{prox}_{\gamma g}(x - \gamma \nabla f(x, u)) \quad \text{for all } \gamma$$

## Example #4: Proximal gradient dynamics

Equivalence property motivates:

**proximal gradient dynamics:**

$$\dot{x} = F_{\text{ProxG}}(x) := -x + \text{prox}_{\gamma g}(x - \gamma \nabla f(x))$$

projected gradient descent is special case

$F_{\text{ProxG}}$  is infinitesimally contracting wrt  $\|\cdot\|_2$


$$\text{for } 0 < \gamma < \frac{2}{\ell}, \quad \text{with rate} \quad c = 1 - \max\{|1 - \gamma\nu|, |1 - \gamma\ell|\},$$

$$\text{for } \gamma^* = \frac{2}{\nu + \ell}, \quad \text{with maximal rate} \quad c^* = \frac{2\nu}{\nu + \ell}$$

## Example #5: Firing-rate recurrent neural network

$$\dot{x} = \mathsf{F}_{\text{FR}}(x) := -x + \Phi(Wx + Bu)$$

sigmoid, hyperbolic tangent  
 $\text{ReLU} = \max\{x, 0\} = (x)_+$   
 $0 \leq \Phi'_i(y) \leq 1$




$\mathsf{F}_{\text{FR}}$  is infinitesimally contracting wrt  $\|\cdot\|_\infty$  with rate  $1 - \mu_\infty(W)_+$  if

$$\mu_\infty(W) < 1 \quad (\text{i.e., } w_{ii} + \sum_j |w_{ij}| < 1 \text{ for all } i)$$

## Example #5: Firing-rate recurrent neural network

$$\dot{x} = \mathsf{F}_{\text{FR}}(x) := -x + \Phi(Wx + Bu)$$

sigmoid, hyperbolic tangent  
 $\text{ReLU} = \max\{x, 0\} = (x)_+$   
 $0 \leq \Phi'_i(y) \leq 1$



$\mathsf{F}_{\text{FR}}$  is infinitesimally contracting wrt  $\|\cdot\|_\infty$  with rate  $1 - \mu_\infty(W)_+$  if

$$\mu_\infty(W) < 1 \quad (\text{i.e., } w_{ii} + \sum_j |w_{ij}| < 1 \text{ for all } i)$$

Note: clear graphical interpretation + generalization to interconnection theorem

## Example #6: Firing-rate network with symmetric synapses

$$\dot{x} = \mathsf{F}_{\text{FR}}(x) := -x + \Phi(Wx + Bu)$$

$$0 \leq \Phi'_i(y) \leq 1 \quad \text{and} \quad W = W^\top \text{ with } \lambda_W = \lambda_{\max}(W)$$

For  $\lambda_W < 1$  and  $\lambda_W \neq 0$ ,  $\mathsf{F}_{\text{FR}}$  is infinitesimally contracting with rate  $-1 + (\lambda_W)_+$

For  $\lambda_W = 1$ ,  $\mathsf{F}_{\text{FR}}$  is weakly infinitesimally contracting

Note: when  $W = W^\top$ , sharper result, but no graph interpretation and hard to generalize

## Example #7: Saddle dynamics

Assume  $f : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}$

- $x \mapsto f(x, y)$  is  $\nu_x$ -strongly convex, uniformly in  $y$
- $y \mapsto f(x, y)$  is  $\nu_y$ -strongly concave, uniformly in  $x$

**saddle dynamics (primal-descent / dual-ascent):**

$$\begin{bmatrix} \dot{x} \\ \dot{y} \end{bmatrix} = \mathsf{F}_S(x, y) := \begin{bmatrix} -\nabla_x f(x, y) \\ \nabla_y f(x, y) \end{bmatrix}$$

$\mathsf{F}_S$  is infinitesimally contracting wrt  $\|\cdot\|_2$  with rate  $\min\{\nu_x, \nu_y\}$

unique globally exp stable point is saddle point (min in  $x$ , max in  $y$ )

## Example #8: Pseudogradient play

Each player  $i$  aims to minimize its own cost function  $J_i(x_i, x_{-i})$  (not a potential game)

**pseudogradient dynamics (aka gradient play in game theory):**

$$\begin{aligned}\dot{x} &= \mathsf{F}_{\text{PseudoG}}(x) = -(\nabla_1 J_1(x_1, x_{-1}), \dots, \nabla_n J_n(x_n, x_{-n})) \quad (\text{stacked vector}) \\ \iff \dot{x}_i &= -\nabla_i J_i(x_i, x_{-i})\end{aligned}$$

• **strong convexity wrt  $x_i$ :**  $J_i$  is  $\mu_i$  strongly convex wrt  $x_i$ , uniformly in  $x_{-i}$

• **Lipschitz wrt  $x_{-i}$ :**  $\text{Lip}_{x_j}(\nabla_i J_i) \leq \ell_{ij}$ , uniformly in  $x_{-j}$

•  $\mathsf{F}_{\text{PseudoG}}$  gain matrix is Hurwitz

⇒  $\mathsf{F}_{\text{PseudoG}}$  is infinitesimally contracting wrt appropriate diag-weighted  $\|\cdot\|_2$

## Example #9: Best response play

Each player  $i$  aims to minimize its own cost function  $J_i(x_i, x_{-i})$

$\text{BR}_i : x_{-i} \rightarrow \operatorname{argmin}_{x_i} J_i(x_i, x_{-i})$  best response of player  $i$  wrt other decisions  $x_{-i}$

**best response dynamics:**

$$\begin{aligned}\dot{x} &= F_{\text{BR}}(x) := \text{BR}(x) - x \\ \iff \dot{x}_i &= \text{BR}_i(x_{-i}) - x_i\end{aligned}$$

- **strong convexity wrt  $x_i$ :**  $J_i$  is  $\mu_i$  strongly convex wrt  $x_i$ , uniformly in  $x_{-i}$
- **Lipschitz wrt  $x_{-i}$ :**  $\text{Lip}_{x_j}(\nabla_i J_i) \leq \ell_{ij}$ , uniformly in  $x_{-j}$   
 $\implies \text{BR}_i$  is Lipschitz wrt  $x_j$  with constant  $\ell_{ij}/\mu_i$
- $F_{\text{BR}}$  gain matrix is Hurwitz  $\iff$  BR is a discrete-time contraction  
 $\implies \text{BR} - \text{Id}$  is infinitesimally contracting wrt appropriate diag-weighted  $\|\cdot\|_2$

## Equivalent statements:

①  $F_{\text{PseudoG}}$  gain matrix:

$$\begin{bmatrix} -\mu_1 & \dots & \ell_{1n} \\ \vdots & & \vdots \\ \ell_{n1} & \dots & -\mu_n \end{bmatrix} \text{ is Hurwitz}$$

②  $F_{\text{BR}}$  gain matrix:

$$\begin{bmatrix} -1 & \dots & \ell_{1n}/\mu_1 \\ \vdots & & \vdots \\ \ell_{n1}/\mu_n & \dots & -1 \end{bmatrix} \text{ is Hurwitz}$$

③ discrete-time  $F_{\text{BR}}$  gain matrix:

$$\begin{bmatrix} 0 & \dots & \ell_{1n}/\mu_1 \\ \vdots & & \vdots \\ \ell_{n1}/\mu_n & \dots & 0 \end{bmatrix} \text{ is Schur}$$

**Aggregative games:**  $J_i(x_i, x_{-i}) = f_i(x_i, \frac{1}{n} \sum_{j=1}^n x_j)$

assume  $f_i$  is  $\mu_i$ -strongly convex wrt  $x_i$  and  $\ell_i = \text{Lip}_y(\nabla_{x_i} f_i(x_i, y))$

$\mu_i > \ell_i$  for each agent  $i$   $\implies$  gain matrix is Hurwitz

# Outline

§1. Introduction

§2. Basic contractivity concepts

§3. Examples: Gradient systems defined by strongly convex functions are contracting

§4. Theory: Equilibrium tracking in parametric optimization

§5. Application: Online feedback optimization

§6. Conclusions

Many convex optimization problems can be solved with contracting dynamics

$$\dot{x} = F(x)$$

Many convex optimization problems can be solved with contracting dynamics

$$\dot{x} = F(x)$$

**contracting dynamics for parametric strongly-convex optimization**

$$\dot{x} = F(x, \theta)$$

**contracting dynamics for time-varying strongly-convex optimization**

$$\dot{x} = F(x, \theta(t))$$

# Equilibrium tracking

For parameter-dependent vector field  $\mathbf{F} : \mathbb{R}^n \times \mathbb{R}^d \rightarrow \mathbb{R}^n$  and differentiable  $\theta : \mathbb{R}_{\geq 0} \rightarrow \Theta \subset \mathbb{R}^d$

$$\dot{x}(t) = \mathbf{F}(x(t), \theta(t))$$

Assume there exist norms  $\|\cdot\|_{\mathcal{X}}$  and  $\|\cdot\|_{\Theta}$  s.t.

- **contractivity wrt  $x$ :**  $\text{osLip}_x(\mathbf{F}) \leq -c < 0$ , uniformly in  $\theta$
- **Lipschitz wrt  $\theta$ :**  $\text{Lip}_{\theta}(\mathbf{F}) \leq \ell$ , uniformly in  $x$

## Theorem: Equilibrium tracking for contracting dynamics

- ① for each fixed  $\theta$ , there exists a unique equilibrium  $x^*(\theta)$
- ② the equilibrium map  $x^*(\cdot)$  is Lipschitz with constant  $\frac{\ell}{c}$

# Equilibrium tracking

For parameter-dependent vector field  $\mathbf{F} : \mathbb{R}^n \times \mathbb{R}^d \rightarrow \mathbb{R}^n$  and differentiable  $\theta : \mathbb{R}_{\geq 0} \rightarrow \Theta \subset \mathbb{R}^d$

$$\dot{x}(t) = \mathbf{F}(x(t), \theta(t))$$

Assume there exist norms  $\|\cdot\|_{\mathcal{X}}$  and  $\|\cdot\|_{\Theta}$  s.t.

- **contractivity wrt  $x$ :**  $\text{osLip}_x(\mathbf{F}) \leq -c < 0$ , uniformly in  $\theta$
- **Lipschitz wrt  $\theta$ :**  $\text{Lip}_{\theta}(\mathbf{F}) \leq \ell$ , uniformly in  $x$

## Theorem: Equilibrium tracking for contracting dynamics

- ① for each fixed  $\theta$ , there exists a unique equilibrium  $x^*(\theta)$
- ② the equilibrium map  $x^*(\cdot)$  is Lipschitz with constant  $\frac{\ell}{c}$
- ③  $D^+ \|x(t) - x^*(\theta(t))\|_{\mathcal{X}} \leq -c \|x(t) - x^*(\theta(t))\|_{\mathcal{X}} + \frac{\ell}{c} \|\dot{\theta}(t)\|_{\Theta}$

## Consequences for tracking error

$$D^+ \|x(t) - x^*(\theta(t))\|_{\mathcal{X}} \leq -c \|x(t) - x^*(\theta(t))\|_{\mathcal{X}} + \frac{\ell}{c} \|\dot{\theta}(t)\|_{\Theta}$$

## Consequences for tracking error

$$D^+ \|x(t) - x^*(\theta(t))\|_{\mathcal{X}} \leq -c \|x(t) - x^*(\theta(t))\|_{\mathcal{X}} + \frac{\ell}{c} \|\dot{\theta}(t)\|_{\Theta}$$

- bounded input, bounded error  
with asymptotic bound:

$$\limsup_{t \rightarrow \infty} \|x(t) - x^*(\theta(t))\|_{\mathcal{X}} \leq \frac{\ell}{c^2} \limsup_{t \rightarrow \infty} \|\dot{\theta}(t)\|_{\Theta}$$

## Consequences for tracking error

$$D^+ \|x(t) - x^*(\theta(t))\|_{\mathcal{X}} \leq -c \|x(t) - x^*(\theta(t))\|_{\mathcal{X}} + \frac{\ell}{c} \|\dot{\theta}(t)\|_{\Theta}$$

- bounded input, bounded error  
with asymptotic bound:

$$\limsup_{t \rightarrow \infty} \|x(t) - x^*(\theta(t))\|_{\mathcal{X}} \leq \frac{\ell}{c^2} \limsup_{t \rightarrow \infty} \|\dot{\theta}(t)\|_{\Theta}$$

- bounded energy input, bounded energy error
- vanishing input, vanishing error
- exponentially vanishing input  $\sim e^{-ht}$ , exponentially vanishing error  $\sim e^{-\min\{c,h\}t}$
- periodic input, periodic error

# Outline

§1. Introduction

§2. Basic contractivity concepts


§3. Examples: Gradient systems defined by strongly convex functions are contracting

§4. Theory: Equilibrium tracking in parametric optimization

§5. Application: Online feedback optimization

§6. Conclusions

# Application: Online feedback optimization



## online feedback optimization

online optimization, optimization-based feedback, input/output regulation ...

$$\begin{cases} \min & \text{cost}_1(u) + \text{cost}_2(y) \\ \text{subj. to} & y = \text{Plant}(u, w(t)) \end{cases} \implies \begin{cases} \dot{u} = \text{Optimizer}(t, u, y) \\ y = \text{Plant}(u, w(t)) \end{cases}$$

A. Hauswirth, S. Bolognani, G. Hug, and F. Dorfler. Timescale separation in autonomous optimization. *IEEE Transactions on Automatic Control*, 66(2):611–624, 2021.

G. Bianchin, J. Cortés, J. I. Poveda, and E. Dall'Anese. Time-varying optimization of LTI systems via projected primal-dual gradient flows. *IEEE Transactions on Control of Network Systems*, 9(1):474–486, 2022.

## Example #10: Gradient controller

- fast/stable LTI plant with control input  $u$  and state/measurement disturbance  $w(t)$ :

$$\begin{aligned}\epsilon \dot{x} &= Ax + Bu + Ew(t) && A \text{ Hurwitz} \\ y &= Cx + Dw(t)\end{aligned}$$

- in singular perturbation limit as  $\epsilon \rightarrow 0^+$ , **steady state map** ( $Y_u$  and  $Y_w$ )

$$y = \underbrace{-CA^{-1}B u}_{=: Y_u} + \underbrace{(D - CA^{-1}E) w}_{=: Y_w}$$

## Example #10: Gradient controller

- fast/stable LTI plant with control input  $u$  and state/measurement disturbance  $w(t)$ :

$$\begin{aligned}\epsilon \dot{x} &= Ax + Bu + Ew(t) && A \text{ Hurwitz} \\ y &= Cx + Dw(t)\end{aligned}$$

- in singular perturbation limit as  $\epsilon \rightarrow 0^+$ , **steady state map (Y<sub>u</sub> and Y<sub>w</sub>)**

$$y = \underbrace{-CA^{-1}B u}_{=: Y_u} + \underbrace{(D - CA^{-1}E) w}_{=: Y_w}$$

- define **cost function  $\mathcal{E}$  on  $u$  and  $y$** :

$$\mathcal{E}(u, w) = \phi(u) + \psi(Y_u u + Y_w w), \quad (\phi \text{ is } \nu\text{-strongly convex and } \psi \text{ is convex})$$

and note

$$\begin{aligned}\nabla_u \mathcal{E}(u, w) &= \nabla \phi(u) + Y_u^\top \nabla \psi(Y_u u + Y_w w) \\ &= \nabla \phi(u) + Y_u^\top \nabla \psi(y) \quad (\text{no need to measure } w(t))\end{aligned}$$

## Example #10: Gradient controller

**equilibrium trajectory** let  $u^*(t)$  be solution to

$$\begin{aligned} \min_u \quad & \phi(u) + \psi(y(t)) && (\nu\text{-strongly convex } \phi, \text{ convex } \psi) \\ \text{subj to} \quad & y(t) = Y_u u + Y_w w(t) \end{aligned}$$

**gradient controller**

$$\dot{u} = F_{\text{GradCtrl}}(u, w) := -\nabla \mathcal{E}_u(u, w) = -\nabla \phi(u) - Y_u^\top \nabla \psi(Y_u u + Y_w w)$$

## Example #10: Gradient controller

**equilibrium trajectory** let  $u^*(t)$  be solution to

$$\begin{aligned} \min_u \quad & \phi(u) + \psi(y(t)) && (\nu\text{-strongly convex } \phi, \text{ convex } \psi) \\ \text{subj to} \quad & y(t) = Y_u u + Y_w w(t) \end{aligned}$$

**gradient controller**

$$\dot{u} = F_{\text{GradCtrl}}(u, w) := -\nabla \mathcal{E}_u(u, w) = -\nabla \phi(u) - Y_u^\top \nabla \psi(Y_u u + Y_w w)$$

### Equilibrium tracking for the gradient controller

- ①  $\text{osLip}_u(F_{\text{GradCtrl}}) \leq -\nu$  (gradient of  $\nu$ -strongly convex function)
- ②  $\text{Lip}_w(F_{\text{GradCtrl}}) = \ell_w := \|Y_u^\top\| \text{Lip}(\nabla \psi) \|Y_w\|$

$$\limsup_{t \rightarrow \infty} \|u(t) - u^*(t)\| \leq \frac{\ell_w}{\nu^2} \limsup_{t \rightarrow \infty} \|\dot{w}(t)\|$$

## Example #11: Projected gradient controller

### Constrained feedback optimization:

$$\begin{aligned} \min_u \quad & \mathcal{E}(u, w) = \phi(u) + \psi(Y_u u + Y_w w) \quad (\nu \text{ strongly convex}, \ell_u \text{ strongly smooth}, \ell_w) \\ \text{subj. to} \quad & u \in \mathcal{U} \quad (\text{nonempty, closed, convex. } P_{\mathcal{U}} = \text{orthogonal projection}) \end{aligned}$$

**Projected gradient controller** (example of proximal gradient dynamics):

$$\dot{u} = F_{\text{PGC}}(u, w) := -u + P_{\mathcal{U}}(u - \gamma \nabla_u \mathcal{E}(u, w))$$

**Equilibrium tracking for projected gradient controller** At  $\gamma = \frac{2}{\nu + \ell_u}$ ,

①  $\text{osLip}_u(F_{\text{PGC}}) \leq -c_{\text{PGC}} := -\frac{2\nu}{\nu + \ell_u}$  (contractivity prox gradient)

②  $\text{Lip}_w(F_{\text{PGC}}) = \ell_{\text{PGC}} := \frac{2}{\nu + \ell_u} \ell_w$

$$\limsup_{t \rightarrow \infty} \|u(t) - u^*(t)\| \leq \frac{\ell_{\text{PGC}}}{c_{\text{PGC}}^2} \limsup_{t \rightarrow \infty} \|\dot{w}(t)\| \quad (\text{eq tracking})$$

# Outline

§1. Introduction

§2. Basic contractivity concepts

§3. Examples: Gradient systems defined by strongly convex functions are contracting

§4. Theory: Equilibrium tracking in parametric optimization

§5. Application: Online feedback optimization

§6. Conclusions

**contractivity = robust computationally-friendly stability**

fixed point theory + Lyapunov stability theory + geometry of metric spaces

- theory
- examples
- control application

## Ongoing work

- ① applications to ML and biologically-inspired neural networks
- ② applications to optimization-based control designs:  
model predictive control, control barrier functions, low-gain integral control
- ③ equilibrium tracking with noise  
applications to optimization-based control