



# Contraction Theory in Systems and Control

Francesco Bullo



Center for Control,  
Dynamical Systems & Computation  
University of California at Santa Barbara  
<https://fbullo.github.io>

XIX Red Raider MiniSymposium, "Differential Geometry and Integrable Systems"  
Texas Tech University, Lubbock Texas, April 20-23, 2023



# Acknowledgments




Saber Jafarpour  
Georgia Tech



Alexander Davydov  
UC Santa Barbara



Kevin D. Smith  
UC Santa Barbara



Xiaoming Duan  
Shanghai Jiao Tong



Pedro Cisneros-Velarde  
University of Illinois



Veronica Centorrino  
Scuola Sup Meridionale



Robin Delabays  
HES-SO Sion



Anton Proskurnikov  
Politecnico Torino



Giovanni Russo  
Univ Salerno



John W. Simpson-Porco  
University of Toronto



AFOSR



ARO



DTRA/ERDC

Frederick Leve @AFOSR FA9550-22-1-0059  
Edward Palazzolo @ARO W911NF-22-1-0233  
Paul Tandy @DTRA W912HZ-22-2-0010  
Donald Wagner @AFOSR FA9550-21-1-0203

**contractivity = robust computationally-friendly stability**


fixed point theory + Lyapunov stability theory + geometry of metric spaces

**contractivity = robust computationally-friendly stability**

fixed point theory + Lyapunov stability theory + geometry of metric spaces

### highly-ordered transient and asymptotic behavior:

- ① unique globally exponential stable equilibrium  
& two natural Lyapunov functions
- ② robustness properties
  - bounded input, bounded output (iss)
  - finite input-state gain
  - robustness margin wrt unmodeled dynamics
  - robustness margin wrt delayed dynamics
- ③ periodic input, periodic output
- ④ modularity and interconnection properties
- ⑤ accurate numerical integration and equilibrium point computation




**contractivity = robust computationally-friendly stability**

fixed point theory + Lyapunov stability theory + geometry of metric spaces

### highly-ordered transient and asymptotic behavior:

- ① unique globally exponential stable equilibrium  
& two natural Lyapunov functions
- ② robustness properties
  - bounded input, bounded output (iss)
  - finite input-state gain
  - robustness margin wrt unmodeled dynamics
  - robustness margin wrt delayed dynamics
- ③ periodic input, periodic output
- ④ modularity and interconnection properties
- ⑤ accurate numerical integration and equilibrium point computation



search for contraction properties  
design engineering systems to be contracting

# Contraction theory: historical notes

## • Origins

S. Banach. Sur les opérations dans les ensembles abstraits et leur application aux équations intégrales. *Fundamenta Mathematicae*, 3(1):133–181, 1922. 



# Contraction theory: historical notes

- **Origins**

S. Banach. Sur les opérations dans les ensembles abstraits et leur application aux équations intégrales. *Fundamenta Mathematicae*, 3(1):133–181, 1922. 

- **Dynamics:**

G. Dahlquist. *Stability and error bounds in the numerical integration of ordinary differential equations*. PhD thesis, (Reprinted in Trans. Royal Inst. of Technology, No. 130, Stockholm, Sweden, 1959), 1958

S. M. Lozinskii. Error estimate for numerical integration of ordinary differential equations. I. *Izvestiya Vysshikh Uchebnykh Zavedenii. Matematika*, 5:52–90, 1958. URL <http://mi.mathnet.ru/eng/ivm2980>. (in Russian)



- **Computation:**

C. A. Desoer and H. Haneda. The measure of a matrix as a tool to analyze computer algorithms for circuit analysis. *IEEE Transactions on Circuit Theory*, 19(5):480–486, 1972. 

# Contraction theory: historical notes

- **Origins**

S. Banach. Sur les opérations dans les ensembles abstraits et leur application aux équations intégrales. *Fundamenta Mathematicae*, 3(1):133–181, 1922. doi:

- **Dynamics:**

G. Dahlquist. *Stability and error bounds in the numerical integration of ordinary differential equations*. PhD thesis, (Reprinted in Trans. Royal Inst. of Technology, No. 130, Stockholm, Sweden, 1959), 1958

S. M. Lozinskii. Error estimate for numerical integration of ordinary differential equations. I. *Izvestiya Vysshikh Uchebnykh Zavedenii. Matematika*, 5:52–90, 1958. URL <http://mi.mathnet.ru/eng/ivm2980>. (in Russian)



- **Computation:**

C. A. Desoer and H. Haneda. The measure of a matrix as a tool to analyze computer algorithms for circuit analysis. *IEEE Transactions on Circuit Theory*, 19(5):480–486, 1972. doi:

- **Systems and control:**

W. Lohmiller and J.-J. E. Slotine. On contraction analysis for non-linear systems. *Automatica*, 34(6):683–696, 1998. doi:

- **Incomplete list of contributors who influenced me**

Aminzare, Arcak, Chung, Coogan, Di Bernardo, Manchester, Margaliot, Pavlov, Pham, Proskurnikov, Russo, Sepulchre, Slotine, Sontag, ...

- **Incomplete list of contributors who influenced me**

Aminzare, Arcak, Chung, Coogan, Di Bernardo, Manchester, Margaliot, Pavlov, Pham, Proskurnikov, Russo, Sepulchre, Slotine, Sontag, ...


- **Surveys:**

Z. Aminzare and E. D. Sontag. Contraction methods for nonlinear systems: A brief introduction and some open problems. In *IEEE Conf. on Decision and Control*, pages 3835–3847, Dec. 2014. 

M. Di Bernardo, D. Fiore, G. Russo, and F. Scafuti. Convergence, consensus and synchronization of complex networks via contraction theory. In *Complex Systems and Networks*. Springer, 2016. 

H. Tsukamoto, S.-J. Chung, and J.-J. E. Slotine. Contraction theory for nonlinear stability analysis and learning-based control: A tutorial overview. *Annual Reviews in Control*, 52:135–169, 2021. 

P. Giesl, S. Hafstein, and C. Kawan. Review on contraction analysis and computation of contraction metrics. *Journal of Computational Dynamics*, 10(1):1–47, 2023. 



# Contraction Theory for Dynamical Systems

Francesco Bullo

**Contraction Theory for Dynamical Systems**, Francesco Bullo,  
KDP, 1.1 edition, 2023, ISBN 979-8836646806

- ➊ Textbook with exercises and answers. Format: textbook, slides, and paperbook
- ➋ Content:
  - Fixed point theory
  - Theory of contracting dynamics on vector spaces
  - Applications to nonlinear and interconnected systems
- ➌ Self-Published and Print-on-Demand at:  
<https://www.amazon.com/dp/B0B4K1BTF4>
- ➍ PDF Freely available at  
<https://fbullo.github.io/ctds>
- ➎ 10h minicourse on youtube:  
<https://youtu.be/RvR47ZbqJjc>
- ➏ Future version to include: systems on Riemannian manifolds, homogeneous spaces, and solid cones
  - "Continuous improvement is better than delayed perfection"
  - Mark Twain**

# Outline

- 1 Contractivity of dynamical systems
  - From discrete-time to continuous-time dynamics
  - Table of infinitesimal contractivity conditions
  - Application to recurrent neural networks
  - Connection with convex optimization
- 2 From closed to open, interconnected and optimal systems
- 3 From nominal to uncertain, local and weakly contracting systems
- 4 Contractivity on Riemannian manifolds
  - A contractivity conjecture
- 5 Conclusions and Future Research

# Linear algebra: induced norms

Vector norm

$$\|x\|_1 = \sum_{i=1}^n |x_i|$$

$$\|x\|_2 = \sqrt{\sum_{i=1}^n x_i^2}$$

$$\|x\|_\infty = \max_{i \in \{1, \dots, n\}} |x_i|$$

Induced matrix norm

$$\|A\|_1 = \max_{j \in \{1, \dots, n\}} \sum_{i=1}^n |a_{ij}|$$

$$\|A\|_2 = \sqrt{\lambda_{\max}(A^\top A)}$$

$$\|A\|_\infty = \max_{i \in \{1, \dots, n\}} \sum_{j=1}^n |a_{ij}|$$

Induced matrix log norm

$$\begin{aligned}\mu_1(A) &= \max_{j \in \{1, \dots, n\}} \left( a_{jj} + \sum_{i=1, i \neq j}^n |a_{ij}| \right) \\ &= \text{max column "absolute sum" of } A\end{aligned}$$

$$\mu_2(A) = \lambda_{\max}\left(\frac{A + A^\top}{2}\right)$$

$$\begin{aligned}\mu_\infty(A) &= \max_{i \in \{1, \dots, n\}} \left( a_{ii} + \sum_{j=1, j \neq i}^n |a_{ij}| \right) \\ &= \text{max row "absolute sum" of } A\end{aligned}$$

## Discrete-time dynamics and Lipschitz constants

$x_{k+1} = F(x_k)$       on  $\mathbb{R}^n$  with norm  $\|\cdot\|$  and induced norm  $\|\cdot\|$

$x_{k+1} = F(x_k)$       on  $\mathbb{R}^n$  with norm  $\|\cdot\|$  and induced norm  $\|\cdot\|$

## Lipschitz constant

$$\begin{aligned}\text{Lip}(F) &= \inf\{\ell > 0 \text{ such that } \|F(x) - F(y)\| \leq \ell\|x - y\| \text{ for all } x, y\} \\ &= \sup_x \|J_F(x)\|\end{aligned}$$

For **scalar map**  $f$ ,     $\text{Lip}(f) = \sup_x |f'(x)|$

$$x_{k+1} = \mathsf{F}(x_k) \quad \text{on } \mathbb{R}^n \text{ with norm } \|\cdot\| \text{ and induced norm } \|\cdot\|$$

## Lipschitz constant

$$\begin{aligned}\text{Lip}(\mathsf{F}) &= \inf\{\ell > 0 \text{ such that } \|\mathsf{F}(x) - \mathsf{F}(y)\| \leq \ell \|x - y\| \text{ for all } x, y\} \\ &= \sup_x \|\mathsf{J}_{\mathsf{F}}(x)\|\end{aligned}$$

For **scalar map**  $f$ ,  $\text{Lip}(f) = \sup_x |f'(x)|$

For **affine map**  $\mathsf{F}_A(x) = Ax + a$


$$\|x\|_{2,P} = (x^\top Px)^{1/2} \quad \text{Lip}_{2,P}(\mathsf{F}_A) = \|A\|_{2,P} \leq \ell \iff A^\top PA \preceq \ell^2 P$$

$$\|x\|_{\infty,\eta} = \max_i |x_i|/\eta_i \quad \text{Lip}_{\infty,\eta}(\mathsf{F}_A) = \|A\|_{\infty,\eta} \leq \ell \iff \eta^\top |A| \leq \ell \eta^\top$$

## Banach contraction theorem for discrete-time dynamics:

If  $\rho := \text{Lip}(F) < 1$ , then

- ①  $F$  is **contracting** = distance between trajectories decreases exp fast ( $\rho^k$ )
- ②  $F$  has a unique, glob exp stable equilibrium  $x^*$



## From discrete to continuous time

The **induced log norm** of  $A \in \mathbb{R}^{n \times n}$  wrt to  $\|\cdot\|$ :

$$\mu(A) := \lim_{h \rightarrow 0^+} \frac{\|I_n + hA\| - 1}{h}$$

subadditivity:

$$\mu(A + B) \leq \mu(A) + \mu(B)$$

scaling:

$$\mu(bA) = b\mu(A), \quad \forall b \geq 0$$

# From discrete to continuous time

The **induced log norm** of  $A \in \mathbb{R}^{n \times n}$  wrt to  $\|\cdot\|$ :



$$\mu(A) := \lim_{h \rightarrow 0^+} \frac{\|I_n + hA\| - 1}{h}$$

subadditivity:

$$\mu(A + B) \leq \mu(A) + \mu(B)$$

scaling:

$$\mu(bA) = b\mu(A), \quad \forall b \geq 0$$



## Example induced log norms

Vector norm	Induced matrix norm	Induced matrix log norm
$\ x\ _1 = \sum_{i=1}^n  x_i $	$\ A\ _1 = \max_{j \in \{1, \dots, n\}} \sum_{i=1}^n  a_{ij} $	$\mu_1(A) = \max_{j \in \{1, \dots, n\}} \left( a_{jj} + \sum_{i=1, i \neq j}^n  a_{ij}  \right)$ = max column "absolute sum" of $A$
$\ x\ _2 = \sqrt{\sum_{i=1}^n x_i^2}$	$\ A\ _2 = \sqrt{\lambda_{\max}(A^\top A)}$	$\mu_2(A) = \lambda_{\max}\left(\frac{A + A^\top}{2}\right)$
$\ x\ _\infty = \max_{i \in \{1, \dots, n\}}  x_i $	$\ A\ _\infty = \max_{i \in \{1, \dots, n\}} \sum_{j=1}^n  a_{ij} $	$\mu_\infty(A) = \max_{i \in \{1, \dots, n\}} \left( a_{ii} + \sum_{j=1, j \neq i}^n  a_{ij}  \right)$ = max row "absolute sum" of $A$

## Continuous-time dynamics and one-sided Lipschitz constants

$\dot{x} = F(x)$       on  $\mathbb{R}^n$  with norm  $\|\cdot\|$  and induced log norm  $\mu(\cdot)$

# Continuous-time dynamics and one-sided Lipschitz constants

$\dot{x} = F(x)$       on  $\mathbb{R}^n$  with norm  $\|\cdot\|$  and induced log norm  $\mu(\cdot)$

## One-sided Lipschitz constant

$$\begin{aligned}\text{osLip}(F) &= \inf\{b \in \mathbb{R} \text{ such that } \langle\langle F(x) - F(y), x - y \rangle\rangle \leq b\|x - y\|^2 \quad \text{for all } x, y\} \\ &= \sup_x \mu(J_F(x))\end{aligned}$$

For **scalar map**  $f$ ,     $\text{osLip}(f) = \sup_x f'(x)$

# Continuous-time dynamics and one-sided Lipschitz constants

$\dot{x} = F(x)$  on  $\mathbb{R}^n$  with norm  $\|\cdot\|$  and induced log norm  $\mu(\cdot)$

## One-sided Lipschitz constant

$$\begin{aligned}\text{osLip}(F) &= \inf\{b \in \mathbb{R} \text{ such that } \langle\langle F(x) - F(y), x - y \rangle\rangle \leq b\|x - y\|^2 \text{ for all } x, y\} \\ &= \sup_x \mu(J_F(x))\end{aligned}$$

For **scalar map**  $f$ ,  $\text{osLip}(f) = \sup_x f'(x)$

For **affine map**  $F_A(x) = Ax + a$


$$\text{osLip}_{2,P}(F_A) = \mu_{2,P}(A) \leq \ell \iff A^\top P + AP \preceq 2\ell P$$

$$\text{osLip}_{\infty,\eta}(F_A) = \mu_{\infty,\eta}(A) \leq \ell \iff a_{ii} + \sum_{j \neq i} |a_{ij}| \eta_i / \eta_j \leq \ell$$

## Banach contraction theorem for continuous-time dynamics:

If  $-c := \text{osLip}(F) < 0$ , then

- ①  $F$  is **infinitesimally contracting** = distance between trajectories decreases exp fast ( $e^{-ct}$ )
- ②  $F$  has a unique, glob exp stable equilibrium  $x^*$



## From inner products to weak pairings

$$\frac{1}{2} \frac{d}{dt} \|x(t)\|_2^2 = \dot{x}^\top x = \langle\langle \dot{x}, x \rangle\rangle$$

$$\implies \frac{1}{2} D^+ \|x(t)\|^2 =: [\![\dot{x}, x]\!]$$

- $D^+$  is upper-right Dini derivative

# From inner products to weak pairings

$$\frac{1}{2} \frac{d}{dt} \|x(t)\|_2^2 = \dot{x}^\top x = \langle\langle \dot{x}, x \rangle\rangle$$

$$\implies \frac{1}{2} D^+ \|x(t)\|^2 =: [\![\dot{x}, x]\!]$$

- $D^+$  is upper-right Dini derivative
- **weak pairing**  $[\![\cdot, \cdot]\!] : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$  exists for each norm, i.e.,

$$[\![y, x]\!]_1 := \|x\|_1 \text{sign}(x)^\top y \quad (\text{sign pairing})$$

$$[\![y, x]\!]_\infty := \max_{i \in \mathcal{A}_\infty(x)} x_i y_i \quad \text{for } \mathcal{A}_\infty(x) = \{i \text{ s.t. } |x_i| = \|x\|_\infty\} \quad (\text{max pairing})$$

# From inner products to weak pairings

$$\frac{1}{2} \frac{d}{dt} \|x(t)\|_2^2 = \dot{x}^\top x = \langle\langle \dot{x}, x \rangle\rangle$$

$$\implies \frac{1}{2} D^+ \|x(t)\|^2 =: [\![\dot{x}, x]\!]$$

- $D^+$  is upper-right Dini derivative
- **weak pairing**  $[\![\cdot, \cdot]\!] : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$  exists for each norm, i.e.,

$$[\![y, x]\!]_1 := \|x\|_1 \operatorname{sign}(x)^\top y \quad (\text{sign pairing})$$

$$[\![y, x]\!]_\infty := \max_{i \in \mathcal{A}_\infty(x)} x_i y_i \quad \text{for } \mathcal{A}_\infty(x) = \{i \text{ s.t. } |x_i| = \|x\|_\infty\} \quad (\text{max pairing})$$

theory of weak pairings: computational properties  
and applications to monotone operators

**Log norm  
bounds****Demidovich  
conditions****One-sided Lipschitz  
conditions**

$$\mu_{2,P}(\mathbf{J}_F(x)) \leq -c \quad P\mathbf{J}_F(x) + \mathbf{J}_F(x)^\top P \preceq -2cP \quad (x-y)^\top P(F(x) - F(y)) \leq -c\|x-y\|_{P^{1/2}}^2$$

$$\mu_1(\mathbf{J}_F(x)) \leq -c \quad \text{sign}(v)^\top \mathbf{J}_F(x)v \leq -c\|v\|_1 \quad \text{sign}(x-y)^\top (F(x) - F(y)) \leq -c\|x-y\|_1$$

$$\mu_\infty(\mathbf{J}_F(x)) \leq -c \quad \max_{i \in \mathcal{A}_\infty(v)} v_i (\mathbf{J}_F(x)v)_i \leq -c\|v\|_\infty^2 \quad \max_{i \in \mathcal{A}_\infty(x-y)} (x_i - y_i)(F_i(x) - F_i(y)) \leq -c\|x-y\|_\infty^2$$

Each row = three equivalent statements.

To be understood for all  $x, y \in \mathbb{R}^n$  and all  $v \in \mathbb{R}^n$ .

# One sided Lipschitz conditions

- ① **simple sufficient condition** for uniqueness of continuous ODEs in: A. F. Filippov. *Differential Equations with Discontinuous Righthand Sides*. Kluwer, 1988. ISBN 902772699X (Chapter 1, page 5, citing Krasnosel'skii and Krein 1955)
- ② **one-sided Lipschitz maps** in: E. Hairer, S. P. Nørsett, and G. Wanner. *Solving Ordinary Differential Equations I. Nonstiff Problems*. Springer, 1993. doi (Section I.10)
- ③ **uniformly decreasing maps** in: L. Chua and D. Green. A qualitative analysis of the behavior of dynamic nonlinear networks: Stability of autonomous networks. *IEEE Transactions on Circuits and Systems*, 23(6):355–379, 1976. doi
- ④ **maps with negative nonlinear measure** in: H. Qiao, J. Peng, and Z.-B. Xu. Nonlinear measures: A new approach to exponential stability analysis for Hopfield-type neural networks. *IEEE Transactions on Neural Networks*, 12(2):360–370, 2001. doi
- ⑤ **dissipative Lipschitz maps** in: T. Caraballo and P. E. Kloeden. The persistence of synchronization under environmental noise. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 461 (2059):2257–2267, 2005. doi
- ⑥ **maps with negative lub log Lipschitz constant** in: G. Söderlind. The logarithmic norm. History and modern theory. *BIT Numerical Mathematics*, 46(3):631–652, 2006. doi
- ⑦ **QUAD maps** in: W. Lu and T. Chen. New approach to synchronization analysis of linearly coupled ordinary differential systems. *Physica D: Nonlinear Phenomena*, 213(2):214–230, 2006. doi
- ⑧ **incremental quadratically stable maps** in: L. D'Alto and M. Corless. Incremental quadratic stability. *Numerical Algebra, Control and Optimization*, 3:175–201, 2013. doi

## Advantages of non-Euclidean approaches

- ① *well suited for certain class of systems*

$\ell_1$  for monotone flow systems

- ② *computational advantages*

$\ell_1/\ell_\infty$  constraints lead to LPs, whereas  $\ell_2$  constraints leads to LMIs

- ③ *robustness to structural perturbations*

$\ell_1/\ell_\infty$  contractions are connectively robust (i.e., edge removal)


- ④ *adversarial input-output analysis*

$\ell_\infty$  better suited for the analysis of adversarial examples than  $\ell_2$

- ⑤ *asynchronous distributed computation*

$\ell_\infty$  contractions converge under fully asynchronous distributed execution

## Application: $\ell_\infty$ -contracting neural networks



If


$$\mu_\infty(A) < 1 \quad \left( \text{i.e., } a_{ii} + \sum_j |a_{ij}| < 1 \text{ for all } i \right)$$

$$\dot{x} = -x + \Phi(Ax + Bu + b) \quad (\text{recurrent NN})$$

$$x = \Phi(Ax + Bu + b) \quad (\text{implicit NN})$$

$$x_{k+1} = (1 - \alpha)x_k + \alpha\Phi(Ax_k + Bu + b) \quad (\text{forward Euler})$$

# Application: $\ell_\infty$ -contracting neural networks



$$\begin{aligned}\dot{x} &= -x + \Phi(Ax + Bu + b) && (\text{recurrent NN}) \\ x &= \Phi(Ax + Bu + b) && (\text{implicit NN}) \\ x_{k+1} &= (1 - \alpha)x_k + \alpha\Phi(Ax_k + Bu + b) && (\text{forward Euler})\end{aligned}$$

If

$$\mu_\infty(A) < 1 \quad \left( \text{i.e., } a_{ii} + \sum_j |a_{ij}| < 1 \text{ for all } i \right)$$

- recurrent NN is contracting with rate  $1 - \mu_\infty(A)_+$
- implicit NN is well posed
- forward Euler is contracting with factor  $1 - \frac{1 - \mu_\infty(A)_+}{1 - \min_i(a_{ii})_-}$

$$\text{at } \alpha = \frac{1}{1 - \min_i(a_{ii})_-}$$

## Detour: convexity and fixed point theory

For differentiable  $V : \mathbb{R}^n \rightarrow \mathbb{R}$ , equivalent statements:

- ①  $V$  is **strongly convex** with parameter  $m$
- ②  $-\text{grad}V$  is  **$m$ -strongly infinitesimally contracting** with respect to  $\|\cdot\|_2$

## Forward Euler theorem for contracting dynamics

Given arbitrary norm  $\|\cdot\|$ , equivalent statements

- ①  $\dot{x} = F(x)$  is infinitesimally contracting
- ② there exists  $\alpha > 0$  such that  $x_{k+1} = x_k + \alpha F(x_k)$  is contracting

## Forward Euler theorem for contracting dynamics

Given arbitrary norm  $\|\cdot\|$ , equivalent statements

- ①  $\dot{x} = F(x)$  is infinitesimally contracting
- ② there exists  $\alpha > 0$  such that  $x_{k+1} = x_k + \alpha F(x_k)$  is contracting

Given *contraction rate*  $c$  and *Lipschitz constant*  $\ell$ , define *condition number*  $\kappa = \frac{\ell}{c} \geq 1$


- ①  $\text{Id} + \alpha F$  is contracting for

$$0 < \alpha < \frac{1}{c\kappa(1 + \kappa)}$$

- ② the optimal step size minimizing and minimum contraction factor:

$$\begin{aligned}\alpha^* &= \frac{1}{c} \left( \frac{1}{2\kappa^2} - \frac{3}{8\kappa^3} + \mathcal{O}\left(\frac{1}{\kappa^4}\right) \right) \\ \ell^* &= 1 - \frac{1}{4\kappa^2} + \frac{1}{8\kappa^3} + \mathcal{O}\left(\frac{1}{\kappa^4}\right)\end{aligned}$$

## Application: $\ell_\infty$ -contracting neural networks



$$\dot{x} = -x + \Phi(Ax + Bu + b)$$

(*recurrent NN*)

$$x = \Phi(Ax + Bu + b)$$

(*implicit NN*)

$$x_{k+1} = (1 - \alpha)x_k + \alpha\Phi(Ax_k + Bu + b)$$

(*forward Euler*)

If

$$\mu_\infty(A) < 1 \quad \left( \text{i.e., } a_{ii} + \sum_j |a_{ij}| < 1 \text{ for all } i \right)$$

- recurrent NN is contracting with rate  $1 - \mu_\infty(A)_+$


- implicit NN is well posed

- forward Euler is contracting with factor  $1 - \frac{1 - \mu_\infty(A)_+}{1 - \min_i(a_{ii})_-}$

$$\text{at } \alpha^* = \frac{1}{1 - \min_i(a_{ii})_-}$$

# Motivation: $\ell_\infty$ -contracting neural networks

While most ML architectures are feedforward,  
biological neural networks are recurrent and recent interest for implicit ML architectures




artificial neural network AlexNet '12



C. elegans connectome '17

# Motivation: $\ell_\infty$ -contracting neural networks

While most ML architectures are feedforward,  
biological neural networks are recurrent and recent interest for implicit ML architectures



artificial neural network AlexNet '12




C. elegans connectome '17

For recurrent NN,  $\ell_\infty$ -contractivity characterizes the synaptic weights to ensure:

- reproducible & robust behavior
- highly-ordered transient+asymptotic dynamic behavior
- efficient computational methods

# Motivation: $\ell_\infty$ -contracting neural networks

While most ML architectures are feedforward,  
biological neural networks are recurrent and recent interest for implicit ML architectures



artificial neural network AlexNet '12



C. elegans connectome '17

For recurrent NN,  $\ell_\infty$ -contractivity characterizes the synaptic weights to ensure:

- reproducible & robust behavior
- highly-ordered transient+asymptotic dynamic behavior
- efficient computational methods

A. Krizhevsky, I. Sutskever, and G. E. Hinton. Imagenet classification with deep convolutional neural networks. *Advances in Neural Information Processing Systems*, 25, 2012

G. Yan, P. E. Vértes, E. K. Towlson, Y. L. Chew, D. S. Walker, W. R. Schafer, and A.-L. Barabási. Network control principles predict neuron function in the Caenorhabditis elegans connectome. *Nature*, 550(7677):519–523, 2017. 😊

# Outline

- 1 Contractivity of dynamical systems
  - From discrete-time to continuous-time dynamics
  - Table of infinitesimal contractivity conditions
  - Application to recurrent neural networks
  - Connection with convex optimization
- 2 From closed to open, interconnected and optimal systems
- 3 From nominal to uncertain, local and weakly contracting systems
- 4 Contractivity on Riemannian manifolds
  - A contractivity conjecture
- 5 Conclusions and Future Research

# #1: From closed to open systems

Incremental ISS and input-state gain

Given normed spaces  $(\mathcal{X}, \|\cdot\|_{\mathcal{X}})$  and  $(\mathcal{U}, \|\cdot\|_{\mathcal{U}})$ , consider

$$\dot{x} = F(x, u(t)), \quad x_0 \in \mathcal{X}, \quad u(t) \in \mathcal{U}$$


Assume:

- **contractivity wrt  $x$ :**  $\text{osLip}_x(F) \leq -c < 0$ , uniformly in  $u$
- **Lipschitz wrt  $u$ :**  $\text{Lip}_u(F) \leq \ell$ , uniformly in  $x$

Then

- ① each soltns:  $x(t)$  with input  $u_x$  and  $y(t)$  with input  $u_y$


$$D^+ \|x(t) - y(t)\|_{\mathcal{X}} \leq -c \|x(t) - y(t)\|_{\mathcal{X}} + \ell \|u_x(t) - u_y(t)\|_{\mathcal{U}}$$



Then

- ① each soltns:  $x(t)$  with input  $u_x$  and  $y(t)$  with input  $u_y$

$$D^+ \|x(t) - y(t)\|_{\mathcal{X}} \leq -c \|x(t) - y(t)\|_{\mathcal{X}} + \ell \|u_x(t) - u_y(t)\|_{\mathcal{U}}$$



- ② F is **incrementally ISS**, that is, for all  $x_0, y_0$

$$\|x(t) - y(t)\|_{\mathcal{X}} \leq e^{-ct} \|x_0 - y_0\|_{\mathcal{X}} + \frac{\ell(1 - e^{-ct})}{c} \sup_{\tau \in [0, t]} \|u_x(\tau) - u_y(\tau)\|_{\mathcal{U}}$$

## #2: From closed to interconnected contracting systems

Networks of contracting systems



Consider  $n$  interconnected subsystems

$$\dot{x}_i = F_i(x_i, x_{-i}), \quad \text{for } i \in \{1, \dots, n\}$$

with state  $x_i \in \mathbb{R}^{N_i}$

with states of connected subsystems  $x_{-i} \in \mathbb{R}^{N-N_i}$ , and

consider  $n$  *local norms*  $\|\cdot\|_i$  on  $\mathbb{R}^{N_i}$



Assume for each node  $i$ :

- **contractivity wrt  $x_i$ :**  $\text{osLip}_{x_i}(\mathcal{F}_i) \leq -c_i < 0$ , uniformly in  $x_{-i}$
- **Lipschitz wrt  $x_j$ :**  $\text{Lip}_{x_j}(\mathcal{F}_i) \leq \ell_{ij}$ , uniformly in  $x_{-j}$

Assume for each node  $i$ :

- **contractivity wrt  $x_i$ :**  $\text{osLip}_{x_i}(\mathcal{F}_i) \leq -c_i < 0$ , uniformly in  $x_{-i}$
- **Lipschitz wrt  $x_j$ :**  $\text{Lip}_{x_j}(\mathcal{F}_i) \leq \ell_{ij}$ , uniformly in  $x_{-j}$

### Network contraction theorem

If the Lipschitz constants matrix  $\begin{bmatrix} -c_1 & \dots & \ell_{1n} \\ \vdots & & \vdots \\ \ell_{n1} & \dots & -c_n \end{bmatrix}$  is **Hurwitz**

⇒ the **interconnected system** is infinitesimally contracting

History: interconnection of stable systems, method of vector Lyapunov functions, connective stability via M-matrix theory  
– Matrosov and Bellman 1962, Ström, Siljak, Russo/DiBernardo/Sontag, ...

$$\begin{bmatrix} -c_1 & \dots & \ell_{1n} \\ \vdots & & \vdots \\ \ell_{n1} & \dots & -c_n \end{bmatrix} \text{ is Metzler}$$

### Hurwitzness depends upon both topology and edge weights

- Hurwitz iff there exists a positive  $\xi$  such that  $M\xi < \mathbb{0}_n$  (power method)
- Hurwitz iff Lyapunov diagonally stable

$$\begin{bmatrix} -c_1 & \dots & \ell_{1n} \\ \vdots & & \vdots \\ \ell_{n1} & \dots & -c_n \end{bmatrix} \text{ is Metzler}$$

## Hurwitzness depends upon both topology and edge weights

- Hurwitz iff there exists a positive  $\xi$  such that  $M\xi < \mathbb{0}_n$  (power method)
- Hurwitz iff Lyapunov diagonally stable
- for  $n = 2$ , Hurwitz if and only if **small gain condition**

$$\text{cycle gain} := \frac{\ell_{12}}{c_1} \frac{\ell_{21}}{c_2} < 1$$

and, for  $n \geq 3$ , **network small-gain theorem for Metzler matrices**


### #3: From closed to systems with optimal controls

For  $\dot{x} = F(x, u)$ , compute  $u : [0, T] \rightarrow \mathbb{R}^k$  to minimize  $\psi(x(T)) + \int_0^T \phi(x, u) dt$

### #3: From closed to systems with optimal controls

For  $\dot{x} = F(x, u)$ , compute  $u : [0, T] \rightarrow \mathbb{R}^k$  to minimize  $\psi(x(T)) + \int_0^T \phi(x, u) dt$

**Pontryagin Minimum Principle:**  $u = \mathcal{FBS}[u]$



$$\mathcal{F}: \quad \dot{x} = F(x, u)$$

$$\mathcal{B}: \quad \dot{\lambda} = -J_F^\top(x, u)\lambda - \phi_x(x, u)$$


$$\mathcal{S}: \quad u = \operatorname{argmin}_{\tilde{u}} \underbrace{\lambda^\top F(x, \tilde{u}) + \phi(x, \tilde{u})}_{\mathcal{H}(x, \tilde{u}, \lambda)}$$

To compute a solution to:

$$u = \mathcal{FBS}[u]$$

adopt

$$u^+ = \mathcal{FBS}[u]$$




To compute a solution to:

$$u = \mathcal{FBS}[u]$$

adopt

$$u^+ = \mathcal{FBS}[u]$$



If  $\text{osLip}_x(F) = -c$  and all other maps are Lipschitz,

①  $\text{osLip}_\lambda(\text{Adjoint}(F)) = \text{osLip}_x(F)$

②  $\text{Lip}(\mathcal{FBS}) = \text{constant} \times \frac{1 - e^{-cT}}{c}$

**$\mathcal{FBS}$  contracting for short  $T$  or large  $c$**

**contractivity = robust computationally-friendly stability**

fixed point theory + Lyapunov stability theory + geometry of metric spaces

From closed to open, interconnected and optimal systems:

- ① iISS
- ② network small gain theorems
- ③ numerical optimal control


**contractivity = robust computationally-friendly stability**

fixed point theory + Lyapunov stability theory + geometry of metric spaces

From closed to open, interconnected and optimal systems:

- ① iISS
- ② network small gain theorems
- ③ numerical optimal control

**Applications** coupled neural-synaptic dynamics and ML via optimal control



# Outline

- 1 Contractivity of dynamical systems
  - From discrete-time to continuous-time dynamics
  - Table of infinitesimal contractivity conditions
  - Application to recurrent neural networks
  - Connection with convex optimization
- 2 From closed to open, interconnected and optimal systems
- 3 From nominal to uncertain, local and weakly contracting systems
- 4 Contractivity on Riemannian manifolds
  - A contractivity conjecture
- 5 Conclusions and Future Research

# From nominal to uncertain systems

Given a norm  $\|\cdot\|$ , consider

$$\dot{x} = F(x) + \Delta(x)$$

Assume:

- **contractivity**:  $\text{osLip}(F) \leq -c < 0$
- **bounded disturbance**:  $\text{osLip}(\Delta) \leq d < c$

Then


- ①  $F + \Delta$  is strongly contracting with rate  $c - d$
- ② the unique equilibria  $x_F^*$  of  $F$  and  $x_{F+\Delta}^*$  of  $F + \Delta$  satisfy

$$\|x_F^* - x_{F+\Delta}^*\| \leq \frac{\|\Delta(x_F^*)\|}{c - d}$$

# From global to local contractivity

Given a norm  $\|\cdot\|$ , consider

$$\dot{x} = F(x)$$



Assume:

- **contractivity over closed set  $D$ :**  $\text{osLip}(F|_D) \leq -c < 0$
- **existence of almost equilibrium:**  $D$  contains the closed  $B$  at  $\bar{x}$  of radius  $r \geq \|F(\bar{x})\|/c$

Then

- ①  $B$  is forward invariant
- ②  $F|_B$  is strongly infinitesimally contracting





# From strongly to weakly contracting systems

Given a norm  $\|\cdot\|$ , consider

$$\dot{x} = F(x) \quad \text{satisfying} \quad \text{osLip}(F) = 0$$

## Dichotomy for weakly-contracting systems

- ① no equilibrium and every trajectory is unbounded, or
- ② at least one equilibrium, every trajectory is bounded, and local asy stability  $\implies$  global




# Outline

- 1 Contractivity of dynamical systems
  - From discrete-time to continuous-time dynamics
  - Table of infinitesimal contractivity conditions
  - Application to recurrent neural networks
  - Connection with convex optimization
- 2 From closed to open, interconnected and optimal systems
- 3 From nominal to uncertain, local and weakly contracting systems
- 4 Contractivity on Riemannian manifolds
  - A contractivity conjecture
- 5 Conclusions and Future Research

# From vector spaces to Riemannian manifold $(M, \mathbb{G})$

Assume existence and uniqueness of geodesic curve between each  $(x, y)$

$F$  **contracting** if geodesic distances from  $x$  to  $y$  diminishes along the flow of  $F$



**integral test:** the inner product between  $F$  and the geodesic velocity vector  $\gamma'$  at  $x$  and  $y$

**differential test:** condition on covariant differential of  $F$

Given vector field  $F$  on a Riemannian manifold  $(M, \mathbb{G})$  and  $c > 0$ , equivalent statements:

- ① for each  $x, y \in M$  and corresponding geodesic  $\gamma : [0, 1] \rightarrow M$  with  $\gamma(0) = x, \gamma(1) = y$ ,

$$\langle\langle F(y), \gamma'(1) \rangle\rangle_{\mathbb{G}} - \langle\langle F(x), \gamma'(0) \rangle\rangle_{\mathbb{G}} \leq -c d_{\mathbb{G}}(x, y)^2$$

or, equivalently, using the parallel transport map  $P_{y \rightarrow x} : T_y M \rightarrow T_x M$ ,

$$\langle\langle P_{y \rightarrow x} F(y) - F(x), \gamma'(0) \rangle\rangle_{\mathbb{G}} \leq -c d_{\mathbb{G}}(x, y)^2$$

Given vector field  $\mathbf{F}$  on a Riemannian manifold  $(M, \mathbb{G})$  and  $c > 0$ , equivalent statements:

- ① for each  $x, y \in M$  and corresponding geodesic  $\gamma : [0, 1] \rightarrow M$  with  $\gamma(0) = x, \gamma(1) = y$ ,

$$\langle\langle \mathbf{F}(y), \gamma'(1) \rangle\rangle_{\mathbb{G}} - \langle\langle \mathbf{F}(x), \gamma'(0) \rangle\rangle_{\mathbb{G}} \leq -c d_{\mathbb{G}}(x, y)^2$$

or, equivalently, using the parallel transport map  $P_{y \rightarrow x} : T_y M \rightarrow T_x M$ ,

$$\langle\langle P_{y \rightarrow x} \mathbf{F}(y) - \mathbf{F}(x), \gamma'(0) \rangle\rangle_{\mathbb{G}} \leq -c d_{\mathbb{G}}(x, y)^2$$

- ② for all  $v_x \in T_x M$

$$\langle\langle \nabla_{v_x} \mathbf{F}(x) v_x, v_x \rangle\rangle_{\mathbb{G}} \leq -c \|v_x\|_{\mathbb{G}}^2,$$

where  $\nabla$  is the Levi-Civita connection. In components:

$$\mathbb{G}(x) \mathbf{J}_{\mathbf{F}}(x) + \mathbf{J}_{\mathbf{F}}(x)^\top \mathbb{G}(x) + \mathcal{L}_{\mathbf{F}} \mathbb{G}(x) \preceq -2c \mathbb{G}(x)$$

Given vector field  $\mathbf{F}$  on a Riemannian manifold  $(M, \mathbb{G})$  and  $c > 0$ , equivalent statements:

- ① for each  $x, y \in M$  and corresponding geodesic  $\gamma : [0, 1] \rightarrow M$  with  $\gamma(0) = x, \gamma(1) = y$ ,

$$\langle\langle \mathbf{F}(y), \gamma'(1) \rangle\rangle_{\mathbb{G}} - \langle\langle \mathbf{F}(x), \gamma'(0) \rangle\rangle_{\mathbb{G}} \leq -c d_{\mathbb{G}}(x, y)^2$$

or, equivalently, using the parallel transport map  $P_{y \rightarrow x} : T_y M \rightarrow T_x M$ ,

$$\langle\langle P_{y \rightarrow x} \mathbf{F}(y) - \mathbf{F}(x), \gamma'(0) \rangle\rangle_{\mathbb{G}} \leq -c d_{\mathbb{G}}(x, y)^2$$

- ② for all  $v_x \in T_x M$

$$\langle\langle \nabla_{v_x} \mathbf{F}(x) v_x, v_x \rangle\rangle_{\mathbb{G}} \leq -c \|v_x\|_{\mathbb{G}}^2,$$

where  $\nabla$  is the Levi-Civita connection. In components:

$$\mathbb{G}(x) \mathbf{J}_{\mathbf{F}}(x) + \mathbf{J}_{\mathbf{F}}(x)^{\top} \mathbb{G}(x) + \mathcal{L}_{\mathbf{F}} \mathbb{G}(x) \preceq -2c \mathbb{G}(x)$$

- ③ for all solutions  $x(\cdot), y(\cdot)$

$$D^+ d_{\mathbb{G}}(x(t), y(t)) \leq -c d_{\mathbb{G}}(x(t), y(t))$$

# A contractivity conjecture

Given step size  $\alpha > 0$  and start point  $x_0 \in M$ , *forward Euler algorithm* for the vector field  $F$

$$x_{k+1} = \exp_{x_k}(\alpha F(x_k))$$

where  $\exp_x : T_x M \rightarrow M$  is the exponential map

**Riemannian forward Euler algorithm:** Consider a vector field  $F$  on a Riemannian manifold  $(M, G)$  with strong contraction rate  $c > 0$  and Lipschitz constant  $\ell > 0$

- ① there exists a unique equilibrium  $x^*$  of  $F$
- ② sequence  $\{x_k\}$  converges to  $x^*$  for  $0 < \alpha < 2 \frac{c}{\ell^2} = \frac{1}{c\kappa^2}$

# A contractivity conjecture

Given step size  $\alpha > 0$  and start point  $x_0 \in M$ , *forward Euler algorithm* for the vector field  $F$

$$x_{k+1} = \exp_{x_k}(\alpha F(x_k))$$

where  $\exp_x : T_x M \rightarrow M$  is the exponential map

**Riemannian forward Euler algorithm:** Consider a vector field  $F$  on a Riemannian manifold  $(M, G)$  with strong contraction rate  $c > 0$  and Lipschitz constant  $\ell > 0$

- ① there exists a unique equilibrium  $x^*$  of  $F$
- ② sequence  $\{x_k\}$  converges to  $x^*$  for  $0 < \alpha < 2 \frac{c}{\ell^2} = \frac{1}{c\kappa^2}$

**Conjecture:** The algorithm is a Banach contraction mapping

Known to be true for non-negative constant curvature, by Dongjun Wu.

# Outline

- 1 Contractivity of dynamical systems
  - From discrete-time to continuous-time dynamics
  - Table of infinitesimal contractivity conditions
  - Application to recurrent neural networks
  - Connection with convex optimization
- 2 From closed to open, interconnected and optimal systems
- 3 From nominal to uncertain, local and weakly contracting systems
- 4 Contractivity on Riemannian manifolds
  - A contractivity conjecture
- 5 Conclusions and Future Research

# Robust and computationally-friendly stability theory

- ① contractivity conditions on normed vector spaces
- ② convexity and fixed point methods
- ③ disturbances, interconnections and optimal control



	Lyapunov Theory	Contraction Theory for Dynamical Systems
existence of equilibrium	F admits global Lyapunov function	F is strongly contracting
Lyapunov function	assumed	implied + computational methods
inputs	arbitrary	distance to trajectory (+ norm of vector field)
	ISS via $\mathcal{KL}$ and $\mathcal{L}$ functions	iISS via explicit formulas

search for contraction properties  
design engineering systems to be contracting

# References (1/2)

## Contraction theory on normed spaces:

- A. Davydov, S. Jafarpour, and F. Bullo. Non-Euclidean contraction theory for robust nonlinear stability. *IEEE Transactions on Automatic Control*, 67(12):6667–6681, 2022b. doi: 
- S. Jafarpour, A. Davydov, and F. Bullo. Non-Euclidean contraction theory for monotone and positive systems. *IEEE Transactions on Automatic Control*, 2023. doi:  To appear

## Contractivity in optimal control:

- K. D. Smith and F. Bullo. Contractivity of the method of successive approximations for optimal control. *IEEE Control Systems Letters*, (7):919–924, 2023. doi: 

## Contracting neural networks and fixed point theory:

- S. Jafarpour, A. Davydov, A. V. Proskurnikov, and F. Bullo. Robust implicit networks via non-Euclidean contractions. In *Advances in Neural Information Processing Systems*, Dec. 2021. doi: 
- A. Davydov, A. V. Proskurnikov, and F. Bullo. Non-Euclidean contractivity of recurrent neural networks. In *American Control Conference*, pages 1527–1534, Atlanta, USA, May 2022d. doi: 
- F. Bullo, P. Cisneros-Velarde, A. Davydov, and S. Jafarpour. From contraction theory to fixed point algorithms on Riemannian and non-Euclidean spaces. In *IEEE Conf. on Decision and Control*, Dec. 2021. doi: 

# References (2/2)

## Here at CDC 2022


- A. Davydov, S. Jafarpour, A. V. Proskurnikov, and F. Bullo. Non-Euclidean monotone operator theory with applications to recurrent neural networks. In *IEEE Conf. on Decision and Control*, Cancún, México, Dec. 2022c. 
- A. Davydov, S. Jafarpour, M. Abate, F. Bullo, and S. Coogan. Comparative analysis of interval reachability for robust implicit and feedforward neural networks. In *IEEE Conf. on Decision and Control*, Cancún, México, 2022a. 
- V. Centorrino, F. Bullo, and G. Russo. Contraction analysis of Hopfield neural networks with Hebbian learning. In *IEEE Conf. on Decision and Control*, Cancún, México, Dec. 2022. 
- R. Ofir, F. Bullo, and M. Margaliot. Minimum effort decentralized control design for contracting network systems. *IEEE Control Systems Letters*, 6:2731–2736, 2022. 

## Resources on contraction theory for dynamics, control and learning

- ① tutorial session "Contraction Theory for Machine Learning" at the 2021 IEEE CDC conference:  
<https://sites.google.com/view/contractiontheory>
- ② free online book and 10h minicourse  
<https://fbullo.github.io/ctds>  
<https://youtu.be/RvR47ZbqJjc>
- ③ upcoming Workshop on "Contraction Theory for Systems, Control, and Learning" at the 2023 American Control Conference in San Diego, California (under review):  
<https://fbullo.github.io/contraction-workshop-2023>


## Theoretical frontiers

- higher order contraction
- relationship with monotone operator theory
- metric spaces: seminorms, Hilbert metrics ...




## Theoretical frontiers

- higher order contraction
- relationship with monotone operator theory
- metric spaces: seminorms, Hilbert metrics ...




**Limitations:** not all stable systems are contractive:

- Lyapunov-diagonally-stable networks
- multistable systems
- biochemical networks




## Theoretical frontiers

- higher order contraction
- relationship with monotone operator theory
- metric spaces: seminorms, Hilbert metrics ...



**Limitations:** not all stable systems are contractive:

- Lyapunov-diagonally-stable networks
- multistable systems
- biochemical networks



## Application to control and learning

- ① control: optimization-based control design
- ② ML: implicit models and energy-based learning
- ③ neuroscience: robust dynamical modeling

