

# Scheme Theory

*this is my note for scheme theory. this is loosely based on (1) Görtz - Wedhorn AG 1 (2) Borchers online lecture (3) Vakil FOAG*

Nguyen Ngoc Khanh

December 30, 2025

# Contents

<b>1</b>	<b>AFFINE SCHEME</b>	<b>2</b>
1.1	DEFINITION OF AFFINE SCHEME . . . . .	2
1.1.1	CONSTRUCT AFFINE SCHEME FROM RING . . . . .	2
1.1.2	SOME EXAMPLES OF AFFINE SCHEME . . . . .	3
1.2	MORPHISM OF AFFINE SCHEMES . . . . .	4
1.2.1	CONSTRUCT MORPHISM OF AFFINE SCHEMES FROM MAP OF RINGS . . . . .	6
1.2.2	SOME EXAMPLES OF MORPHISM OF AFFINE SCHEMES . . . . .	6
<b>2</b>	<b>SCHEME</b>	<b>8</b>
2.1	DEFINITION OF SCHEME AND MORPHISM OF SCHEMES . . . . .	8
2.2	GLUING OF MORPHISMS . . . . .	8
2.3	SOME BASIC MORPHISMS . . . . .	9
2.3.1	MORPHISM FROM AFFINE POINT . . . . .	9
2.3.2	MORPHISM INTO AFFINE POINT . . . . .	10
2.4	GLUING SCHEMES, DISJOINT UNION OF SCHEMES . . . . .	11
2.4.1	SOME EXAMPLES OF GLUING SCHEMES . . . . .	12
2.5	BASIC PROPERTIES OF SCHEME AND MORPHISM OF SCHEMES . . . . .	12
2.5.1	ON LOCAL PROPERTIES . . . . .	12
2.5.2	TOPOLOGICAL PROPERTIES . . . . .	13
2.5.3	NOETHERIAN SCHEMES . . . . .	14
2.5.4	GENERIC POINTS . . . . .	15
2.5.5	REDUCED SCHEME, INTEGRAL SCHEME, FUNCTION FIELD . . . . .	16

# Chapter 1

## AFFINE SCHEME

### 1.1 DEFINITION OF AFFINE SCHEME

**Definition 1.1.1** (ringed space, locally ringed space)

A ringed space  $(X, \mathcal{O})$  is a topological space  $X$  together with a sheaf of rings  $\mathcal{O}$ . A ringed space is a locally ringed space if for every point  $x \in X$ , the stalk  $\mathcal{O}_x$  is a local ring.

**Definition 1.1.2** (affine scheme)

An affine scheme is a locally ringed space that is isomorphic to the spectrum of some ring  $A$

#### 1.1.1 CONSTRUCT AFFINE SCHEME FROM RING

Let  $A$  be a ring, let  $X = \text{Spec } A$  be the set of prime ideals in  $A$

$$\text{Spec } A = \{\text{prime } \mathfrak{p} \subseteq A\}$$

We put a topology on  $\text{Spec } A$  generated by the basis of open sets

$$D(f) = \{\mathfrak{p} \in \text{Spec } A : f \notin \mathfrak{p}\}$$

for every  $f \in A$ . The constructed topology is called Zariski's topology and the open subsets of the form  $D(f)$  for some  $f \in A$  are called **principal open subsets**. We put a sheaf of rings  $\mathcal{O}$  on  $\text{Spec } A$  generated by

$$\mathcal{O}(D(f)) = A_f$$

If  $D(g) \subseteq D(f)$ , that is  $g \in \sqrt{(f)}$ ,  $g^n = fh$  for some  $h \in A$  and  $n \geq 1$ . Since  $f$  is a unit in  $A_g$ , the restriction map is well-defined and unique

$$\begin{aligned} A_f &\rightarrow A_g \\ \frac{x}{f^m} &\mapsto x \left( \frac{h}{g^n} \right)^m \end{aligned}$$

Under this construction,  $(X, \mathcal{O})$  is an affine scheme. An element  $f \in A$  is called **function**, a element  $x \in X$  is called

**point**, when referring  $x$  as a prime ideal in  $A$ , we write  $\mathfrak{p}_x \subseteq A$ , and function evaluation is equivalent to sending  $f$  to the residue field of stalk  $\mathcal{O}_X = A_{\mathfrak{p}_x}$ .

When  $A$  is a polynomial ring of  $n$  variables over ring  $R$ , we write  $\mathbb{A}_R^n = \text{Spec } A$

*(I see many proofs, people assumed that open set is just principle open set. maybe the argument generalizes easily to general open set. anyway, if needed, below is the definition of sheaf on open set)*

On  $\text{Spec } A$ , any open set is of the form

$$D(S) = \{\mathfrak{p} \in \text{Spec } A : \mathfrak{p} \cap S = \emptyset\}$$

for some multiplicatively closed subset  $S \subseteq A$ . The ring over  $D(S)$  is

$$\mathcal{O}(D(S)) = S^{-1}A$$

If  $D(T) \subseteq D(S)$ , then  $S \subseteq T$ , the restriction map is the canonical map

$$\mathcal{O}(D(S)) \rightarrow \mathcal{O}(D(T))$$

$$S^{-1}A \rightarrow T^{-1}A$$

$$\frac{a}{s} \mapsto \frac{a}{s}$$

Note that, this map corresponds to the inclusion of affine schemes

$$D(T) \hookrightarrow D(S)$$

## 1.1.2 SOME EXAMPLES OF AFFINE SCHEME

### **Remark 1.1.3** (irreducible)

A topological space  $X$  is reducible if it can be written as a union  $X = X_1 \cup X_2$  of two non-empty proper closed subsets.

1. Let  $X$  be a topological space, then the following are equivalent
  - (a)  $X$  is irreducible
  - (b)  $X$  is not reducible
  - (c) every non-empty open subset of  $X$  is dense
  - (d) any two non-empty open subsets of  $X$  have non-empty intersection
2. A subset  $F$  of topological space  $X$  is reducible/irreducible if  $F$  as a subspace is reducible/irreducible
3. An irreducible component is a maximal irreducible subset.

**Remark 1.1.4** (closed, generic, specialization, generization, maximal)

Let  $X$  be a topological space

1. A point  $x \in X$  is called closed if  $\{x\} \subseteq X$  is closed
2. A point  $\eta \in X$  is called generic if  $\overline{\{\eta\}} = X$
3. Let  $x, y \in X$ ,  $y$  is called a specialization of  $x$  if  $y \in \overline{\{x\}}$ . In that case,  $x$  is called generization of  $y$
4. A point  $x \in X$  is called maximal if  $\overline{\{x\}}$  is an irreducible component of  $X$

**example:** (integers) Let  $A = \mathbb{Z}$ , then

$$X = \text{Spec } \mathbb{Z} = \{(2), (3), (5), \dots, (0)\}$$

for each prime number  $p \in \mathbb{Z}$ ,  $(p)$  is a closed point.  $(0)$  is a generic point, and closure of  $(0)$  is the whole space  $X$ .

**example:** (field) Let  $k$  be a field. Let  $A = k$ , then

$$X = \text{Spec } k = \{(0)\}$$

is a singleton set.

**example:** (polynomial ring of two variables over an algebraically closed field) Let  $A$  be an arbitrary ring, maximal ideals are closed points. Moreover, the closure of a point  $\mathfrak{p}$  is the set of prime ideals containing  $\mathfrak{p}$

$$V(\mathfrak{p}) = \{\mathfrak{q} \in \text{Spec } A : \mathfrak{p} \subseteq \mathfrak{q}\}$$

In particular, let  $k$  be an algebraically closed field, let  $A = k[x, y]$ , by Nullstellensatz, the set of closed points are

$$\{(x - a, y - b) : a \in k, b \in k\}$$

$X$  admits other generic points  $(0)$  and  $(f)$  for every irreducible  $f \in k[x, y]$ . By dimensionality argument, the prime  $(f)$  is of height 1 and the closure of  $(f)$  consists of  $(f)$  and  $(x - a, y - b)$  for  $(a, b) \in k^2$  in the vanishing set of  $f$ . Similarly, closure of  $(0)$  is the whole space.

## 1.2 MORPHISM OF AFFINE SCHEMES

**Remark 1.2.1** (inverse image, direct image)

Let  $\psi : X \rightarrow Y$  be a continuous map. The inverse image functor  $\psi^{-1}$  and direct image functor  $\psi_*$  is an adjoint pair between the category of sheaves on  $X$  and the category of sheaves on  $Y$ . Let  $\mathcal{F}$  and  $\mathcal{G}$  be a sheaf on  $X$  and a sheaf on  $Y$  respectively, then

$$\mathrm{Hom}_{\mathrm{Sh}(X)}(\psi^{-1}\mathcal{G}, \mathcal{F}) \cong \mathrm{Hom}_{\mathrm{Sh}(Y)}(\mathcal{G}, \psi_*\mathcal{F})$$

The direct image functor  $\psi_*$  is defined as follows: for every open subset  $V \subseteq Y$ , then

$$(\psi_*\mathcal{F})(V) = \mathcal{F}(\psi^{-1}(V))$$

The inverse image functor  $\psi^{-1}$  is defined as follows: for every open subset  $U \subseteq X$ , then

$$(\psi^{-1}\mathcal{G})(U) = \mathrm{colim}_{V \subseteq Y: \psi(U) \subseteq V} \mathcal{G}(V)$$

**Definition 1.2.2** (morphism of ringed spaces)

A morphism of ringed spaces  $(X, \mathcal{O}_X) \rightarrow (Y, \mathcal{O}_Y)$  is defined by a continuous map  $\psi : X \rightarrow Y$  and a morphism of sheaves of rings  $\psi^\flat : \mathcal{O}_Y \rightarrow \psi_*\mathcal{O}_X$ . By adjunction between  $\psi^{-1}$  and  $\psi_*$ , this is equivalent to a morphism of sheaves of rings  $\psi^\# : \psi^{-1}\mathcal{O}_Y \rightarrow \mathcal{O}_X$ . Moreover,  $\psi^\#$  induces a map on stalks

$$\psi_x^\# : (\psi^{-1}\mathcal{O}_Y)_x = \mathcal{O}_{Y, \psi(x)} \rightarrow \mathcal{O}_{X, x}$$

**Remark 1.2.3** (compatible germs)

Given a sheaf of sets  $\mathcal{F}$  on  $X$ , for every open subset  $U \subseteq X$ , the natural map from sections into product of stalks

$$\mathcal{F}(U) \hookrightarrow \prod_{x \in U} \mathcal{F}_x$$

is injective. An element  $(s_x)_{x \in U} \subseteq \prod_{x \in U} \mathcal{F}_x$  in the image of this map is called compatible germs.

From the equivalence between set of sections and set of compatible germs, one can identify a morphism of ringed spaces by its topological space map  $\psi : X \rightarrow Y$  and its map on stalks  $\psi_x^\# : (\psi^{-1}\mathcal{O}_Y)_x = \mathcal{O}_{Y, \psi(x)} \rightarrow \mathcal{O}_{X, x}$ .

**Definition 1.2.4** (morphism of locally ringed spaces, morphism of affine schemes)

A morphism of ringed spaces  $(\psi, \psi^\#) : (X, \mathcal{O}_X) \rightarrow (Y, \mathcal{O}_Y)$  is a morphism of locally ringed spaces if for every  $x \in X$ , the induced map on stalks  $\psi_x^\# : (\psi^{-1}\mathcal{O}_Y)_x = \mathcal{O}_{Y, \psi(x)} \rightarrow \mathcal{O}_{X, x}$  is a local homomorphism. A morphism of affine schemes is a morphism of locally ringed spaces.

We obtain the category of affine schemes, denoted by  $\mathrm{AffSch}$ . The local homomorphism condition of the map between stalks ensures that any zero function  $g \in \mathcal{O}_{Y, \psi(x)}$  at  $\psi(x) \in Y$  will be sent to a zero function  $\psi_x^\#(g) \in \mathcal{O}_{X, x}$  at  $x \in X$ .

**Proposition 1.2.5** (equivalence between commutative rings and affine schemes)

The functor  $\mathrm{Spec}$  from the opposite category of commutative rings into the category of affine schemes is fully faithful and essentially surjective.

$$\mathrm{Spec} : \mathrm{CRing}^{\mathrm{op}} \xrightarrow{\sim} \mathrm{AffSch}$$

## 1.2.1 CONSTRUCT MORPHISM OF AFFINE SCHEMES FROM MAP OF RINGS

Let  $\phi : A \rightarrow B$  be a map of rings, let  $X = \text{Spec } B$  and  $Y = \text{Spec } A$  be the corresponding affine schemes. Then, the corresponding morphism of locally ringed space  $(\psi, \psi^\flat) : X \rightarrow Y$  is defined as follows: for every  $x \in X$

$$\begin{aligned}\psi &: X \rightarrow Y \\ x &\mapsto \phi^{-1}(x)\end{aligned}$$

The local homomorphism  $\psi_x^\# : \mathcal{O}_{Y, \psi(x)} = A_{\psi(x)} \rightarrow B_x$  is induced from  $\phi : A \rightarrow B$

$$\begin{array}{ccc} A & \xrightarrow{\phi} & B \\ \downarrow & & \downarrow \\ A_{\psi(x)} & \xrightarrow{\psi_x^\#} & B_x \end{array}$$

## 1.2.2 SOME EXAMPLES OF MORPHISM OF AFFINE SCHEMES

**example:** (quotient) Let  $A$  be a ring and  $\mathfrak{a} \subseteq A$  be an ideal. The map  $A \twoheadrightarrow A/\mathfrak{a}$  induces an injective map of sets

$$\text{Spec } A/\mathfrak{a} \hookrightarrow \text{Spec } A$$

As sets, we have  $\text{Spec } A/\sqrt{\mathfrak{a}} = \text{Spec } A/\mathfrak{a} = \text{Spec } A/\mathfrak{a}^n$  for every  $n \geq 1$ . However, as affine schemes, in general, they are different,  $A/\sqrt{\mathfrak{a}}$  is reduced, i.e. has no nilpotent while  $\text{Spec } A/\mathfrak{a}$  and  $\text{Spec } A/\mathfrak{a}^n$  might have nilpotents.

Furthermore, when  $\mathfrak{a} = \mathfrak{p}_x = x$  is prime, quotient by  $\mathfrak{p}_x$  is the action of taking closed subscheme <sup>1</sup> of  $X$

*keep all primes containing  $\mathfrak{p}_x$ . equivalently, keep all points contained in  $x$*

**example:** (localization) Let  $A$  be a ring and  $S \subseteq A$  be a multiplicatively closed subset. Localization at  $S$  induces an injective map of sets

$$\text{Spec } S^{-1}A \hookrightarrow \text{Spec } A$$

Furthermore, when  $A - S = \mathfrak{p}_x = x$  is prime, localization at  $\mathfrak{p}_x$  is the action of taking open subscheme <sup>2</sup> of  $X$

*keep all primes contained in  $\mathfrak{p}_x$ . equivalently, keep all points containing  $x$*

**example:** (disjoint union) Let  $A, B$  be rings, then the projection  $A \times B \rightarrow A$  corresponds to the monomorphism of affine schemes

$$\text{Spec } A \hookrightarrow \text{Spec } A \amalg \text{Spec } B \cong \text{Spec } A \times B$$

Categorically,  $A \times B$  is product in rings and  $\text{Spec } A \amalg \text{Spec } B$  is coproduct in affine schemes

**example:** (non-trivial fiber) Consider the ring map  $\phi : k[u] \rightarrow k[x, y]$  defined by  $u \mapsto y^2 - x$ . It induces a morphism of affine schemes

---

<sup>1</sup>meaning will be revealed later

<sup>2</sup>meaning will be revealed later

$$\begin{aligned}
\psi : \operatorname{Spec} k[x, y] &\rightarrow \operatorname{Spec} k[u] \\
(0) &\mapsto (0) \\
(x - a, y - b) &\mapsto (u - (b^2 - a)) \\
(y^2 - x) &\mapsto (u) \\
(f) &\mapsto (0)
\end{aligned}$$

for every  $a, b \in k$  and  $f$  is irreducible other than multiple of  $y^2 - x$ . (*check this carefully*).

**example:** (inclusion of open subschemes) Let  $A$  be a ring, and  $U \subseteq V \subseteq \operatorname{Spec} A$  be open sets. Then, the inclusion map of open affine subschemes corresponds to the canonical ring map between global sections.

$$\begin{aligned}
U &\hookrightarrow V \\
\mathcal{O}(U) &\leftarrow \mathcal{O}(V)
\end{aligned}$$

If  $U = D(T)$  and  $V = D(S)$ , then  $S \subseteq T$  and the canonical ring map is

$$\begin{aligned}
S^{-1}A = \mathcal{O}(V) &\rightarrow T^{-1}A = \mathcal{O}(U) \\
\frac{a}{s} &\mapsto \frac{a}{s}
\end{aligned}$$

In particular, the affine scheme map induced from  $\operatorname{Spec} A \rightarrow S^{-1}A = \operatorname{Spec} \mathcal{O}(V)$  is precisely  $V \hookrightarrow \operatorname{Spec} A$



## Chapter 2

# SCHEME

## 2.1 DEFINITION OF SCHEME AND MORPHISM OF SCHEMES

**Definition 2.1.1** (scheme, morphism of schemes)

A scheme is a locally ringed space  $(X, \mathcal{O})$  such that for every point  $x \in X$ , there is an open neighbourhood  $U_x$  containing  $x$  so that  $(U_x, \mathcal{O}|_{U_x})$  is an affine scheme. Morphism of schemes is morphism of locally ringed spaces.

We obtain the category of schemes, denoted by  $\text{Sch}$

**Definition 2.1.2** (scheme over  $S$ )

Let  $S$  be a scheme, the category  $(\text{Sch}/S)$  of schemes over  $S$  is the category where an object is a morphism of schemes  $X \rightarrow S$  and a morphism is a morphism of schemes  $X \rightarrow Y$  so that  $X \rightarrow S$  factors through  $Y$  by the map  $X \rightarrow Y$

$$\begin{array}{ccc} X & \xrightarrow{\quad} & Y \\ & \searrow \quad \swarrow & \\ & S & \end{array}$$

The collection of morphisms from  $X \rightarrow S$  into  $Y \rightarrow S$  is denoted by  $\text{Hom}_S(X, Y)$ . If  $S = \text{Spec } A$ , we also call scheme over  $S$  as scheme over  $A$

Since  $\text{Spec } \mathbb{Z}$  is a terminal object in the category of schemes, the category of schemes is canonically equivalent the category of schemes over  $\mathbb{Z}$  (*proof for  $\text{Spec } \mathbb{Z}$  is terminal in next section*)

## 2.2 GLUING OF MORPHISMS

**Definition 2.2.1** (open subscheme)

Let  $(X, \mathcal{O}_X)$  be a scheme and  $U \subseteq X$  be an open subset. Then, the locally ringed space  $(U, \mathcal{O}_X|_U)$  is a scheme and we call it an open subscheme of  $X$ . If the

### Lemma 2.2.2

Let  $X$  be a scheme and  $U, V$  be open affine subschemes of  $X$ . Then for every  $x \in U \cap V$ , there exists an open subscheme  $W \subseteq U \cap V$  containing  $x$  such that  $W$  is principal open in  $U$  as well as in  $V$ .

### Proposition 2.2.3 (gluing of morphisms)

Let  $X, Y$  be locally ringed spaces. Then  $U \mapsto \text{Hom}(U, Y)$  sending open subset  $U \subseteq X$  into the set of morphisms  $(U, \mathcal{O}_X|_U) \rightarrow (Y, \mathcal{O}_Y)$  is a sheaf of sets on  $X$ .

In other words, if  $X = \bigcup_i U_i$  is an open covering, then a family of morphisms  $U_i \rightarrow Y$  glues into a morphism  $X \rightarrow Y$  if and only if the morphisms coincide on intersections  $U_i \cap U_j$ . Moreover, in that case, the resulting morphism  $X \rightarrow Y$  is uniquely determined.

*TODO - proof idea - one can identify the morphism by map between stalks, it feels local hence glueable*

### Proposition 2.2.4

Let  $X$  be a locally ringed space and  $Y = \text{Spec } A$  an affine scheme. Then, the natural map

$$\begin{aligned} \text{Hom}(X, Y) &\rightarrow \text{Hom}(A, \Gamma(X)) \\ (f, f^\flat) &\mapsto f_Y^\flat \end{aligned}$$

is a bijection.

*Proof. (TODO - use lemma above to prove for the case  $X$  is a scheme, full proof in EGA new 1.6.3)* □

$\text{Spec } \mathbb{Z}$  being terminal object in the category of schemes follows this proposition. Moreover, when  $A = \Gamma(X)$ , there corresponds to  $\text{id}_{\Gamma(X)}$  a morphism

$$c_X : X \rightarrow \text{Spec } \Gamma(X)$$

which we call canonical.

## 2.3 SOME BASIC MORPHISMS

### 2.3.1 MORPHISM FROM AFFINE POINT

Let  $X$  be a scheme. Let  $x \in X$  and  $U \subseteq X$  be an open affine subscheme of  $x$ , let  $U = \text{Spec } A$ . Let  $\mathfrak{p} \in \text{Spec } A$  corresponds to  $x$ . Then, the natural map  $A \rightarrow A_{\mathfrak{p}}$  induces a morphism


$$j_x : \text{Spec } \mathcal{O}_{X,x} = \text{Spec } A_{\mathfrak{p}} \rightarrow \text{Spec } A = U \subseteq X$$

*(here, stalk at  $x \in X$  is the same as stalk at  $x \in X$  because stalk is defined as colimit, so it doesn't matter if we restrict to  $U$  or not - Görtz - Wedhorn didn't explain it clearly, so below is my attempt <sup>1</sup>)*

*proof for  $j_x : \text{Spec } \mathcal{O}_{X,x} \rightarrow X$  being independent of choice of  $U$ .* Let  $U$  and  $V$  be two open affine subschemes containing  $x$ , let  $U = \text{Spec } A$  and  $V = \text{Spec } B$ , let  $\mathfrak{p} \in \text{Spec } A$  and  $\mathfrak{q} \in \text{Spec } B$  correspond to  $x$ . Then  $A_{\mathfrak{p}} = B_{\mathfrak{q}}$  and we have two maps  $j_x^U : \text{Spec } \mathcal{O}_{X,x} \rightarrow U$  and  $j_x^V : \text{Spec } \mathcal{O}_{X,x} \rightarrow V$ . By definition,  $\mathcal{O}_{X,x}$  is the colimit of functor  $\mathcal{O}_X$  from the

<sup>1</sup>math.SE <https://math.stackexchange.com/q/5116709/700122>

category of open sets containing  $x$ . Since the contravariant functor  $\text{Spec}$  is fully faithful, it preserves limits,  $\text{Spec } \mathcal{O}_{X,x}$  is the limit of the functor  $\text{Spec } \mathcal{O}_X$ . By universal property of limit, let  $W = U \cap V$ , then  $j_x^U$  and  $j_x^V$  factor through a unique map  $\text{Spec } \mathcal{O}_{X,x} \rightarrow W$  into their intersection and inclusions of schemes  $W \hookrightarrow U$  and  $W \hookrightarrow V$



□


Image of  $j_x$  in  $X$  is the intersection of all open subsets containing  $x$  (consider the map  $A \rightarrow A_{\mathfrak{p}}$ , its kernel is  $\mathfrak{a} = \{a \in A : as = 0 \text{ for some } s \notin \mathfrak{p}\}$ . Hence, the image of  $j_x$  are those primes in  $A$  containing  $\mathfrak{a}$ , ...)

Let  $\kappa(x) = \mathcal{O}_{X,x}/\mathfrak{m}_x$  be the residue field at  $x$ , we obtain the map of schemes

$$i_x : \text{Spec } \kappa(x) \rightarrow \text{Spec } \mathcal{O}_{X,x} \rightarrow X$$

which is called canonical. The image point of  $i_x$  in  $X$  is  $x$ .

Now let  $k$  be a field and  $f : \text{Spec } k \rightarrow X$  be a map, let  $x$  be the image point on  $X$  of the singleton  $p \in \text{Spec } k$ . The map  $f$  induces a map on stalks  $f_p^\# : \mathcal{O}_{X,x} \rightarrow \mathcal{O}_{\text{Spec } k, p} = k$ . Since  $\mathcal{O}_{X,x}$  is local,  $\mathcal{O}_{X,x} \rightarrow k$  factors through  $\mathcal{O}_{X,x} \rightarrow \kappa(x)$ . Hence,  $\text{Spec } k \rightarrow \text{Spec } \mathcal{O}_{X,x}$  factors through  $\text{Spec } \kappa(x) \rightarrow \text{Spec } \mathcal{O}_{X,x}$ .



Moreover,  $f : \text{Spec } k \rightarrow X$  factors through  $i_x : \text{Spec } \kappa(x) \rightarrow X$  (this is very nice but GW didn't explain clearly why  $i_x$  has such a nice property - I guess what's outside a proof is informal)

### Proposition 2.3.1


There is a bijection of sets

$$\text{Hom}(\text{Spec } k, X) \rightarrow X \times \text{Hom}(\kappa(x), k)$$

## 2.3.2 MORPHISM INTO AFFINE POINT

(from Borchers lecture on gluing schemes)

Let  $k$  be a field and  $R$  be a ring, the map  $\text{Spec } R \rightarrow \text{Spec } k$  that factors through  $\text{Spec } k[x]$  corresponds to an element of  $R$ , that is, a regular function on the global section of  $\text{Spec } R$



Similar, any map from scheme  $X$  into  $\text{Spec } k$  that factors through  $\text{Spec } k[x]$  is a regular function on global section of  $X$  since  $X$  can be covered by open affine subschemes.

## 2.4 GLUING SCHEMES, DISJOINT UNION OF SCHEMES

### Definition 2.4.1 (gluing of schemes)

A gluing of schemes consists of the following data: given an index set  $I$

- for all  $i \in I$ , a scheme  $U_i$
- for all  $i, j \in I$ , an open subscheme  $U_{ij} \subseteq U_i$  such that  $U_{ii} = U_i$  for all  $i \in I$
- for all  $i, j \in I$ , an isomorphism  $\phi_{ji} : U_{ij} \rightarrow U_{ji}$  of schemes such that the cocycle condition holds

$$\phi_{kj}\phi_{ji} = \phi_{ki}$$

on  $U_{ij} \cap U_{ik}$  for all  $i, j, k \in I$  if  $\phi_{ji}(U_{ij} \cap U_{ik}) \subseteq U_{jk}$

For  $i = j = k$ , the cocycle condition implies that  $\phi_{ii} = \text{id}_{U_i}$  and for  $i = k$ ,  $\phi_{ij}^{-1} = \phi_{ji}$  and that  $\phi_{ji}$  is an isomorphism  $U_{ij} \cap U_{ik} \rightarrow U_{ji} \cap U_{jk}$

*(it looks like we can make the definition a bit stronger by imposing cocycle condition on those overlapped region where it makes sense, for example  $\phi_{ji}(U_{ij} \cap U_{ik}) \cap U_{jk}$ , but somehow the authors didn't do so, I don't understand now, will do later)*

### Proposition 2.4.2

Given a gluing of schemes, there exists a scheme  $X$  together with morphism  $\psi : U_i \rightarrow X$  such that

- for all  $i \in I$ , the map  $\psi_i$  is an isomorphism from  $U_i$  into an open subscheme of  $X$
- $\psi_j \phi_{ji} = \psi_i$  on  $U_{ij}$  for all  $i, j \in I$
- $X = \bigcup_{i \in I} \psi_i(U_i)$
- $\psi_i(U_i) \cap \psi_j(U_j) = \psi_i(U_{ij}) = \psi_j(U_{ji})$  for all  $i, j \in I$

Furthermore,  $(X, \psi_i)$  is unique up to unique isomorphism.

*(there's a similar result in differential manifold using sheaf. it seems like sheaf is the categorical notion tool for gluing, just like morphism is the categorical notion of mapping, (co)limit is the categorical notion of sequential limit)*

Gluing of morphisms implies the universal property of gluing schemes

**Remark 2.4.3** (universal property of gluing schemes)

If  $T$  is a scheme and for every  $i \in I$ , there is a map  $\xi_i : U_i \rightarrow T$  such that  $\xi_j \phi_{ji} = \xi_i$  on  $U_{ij}$  for all  $i, j \in I$ , then there exists a unique morphism  $\xi : X \rightarrow T$  with  $\xi \psi_i = \xi_i$  for every  $i \in I$

(GW requires each  $\xi_i$  is an isomorphism into its image but it's clearly not required)

## 2.4.1 SOME EXAMPLES OF GLUING SCHEMES

**example:** (gluing two schemes) When the index set  $I = \{1, 2\}$ , any two open subsets  $U_{12} \subseteq U_1$  and  $U_{21} \subseteq U_2$  and an isomorphism  $\phi : U_{12} \xrightarrow{\sim} U_{21}$  is a gluing of schemes. Let  $X$  be the glued scheme, for any open subset  $V \subseteq X$ , we have

$$\Gamma(V, \mathcal{O}_X) = \{(s_1, s_2) \in \Gamma(V \cap U_1, \mathcal{O}_{U_1}) \times \Gamma(V \cap U_2, \mathcal{O}_{U_2}) : \phi^b(s_2|_{V \cap U_{21}}) = s_1|_{V \cap U_{12}}\}$$

(this is just a pushout/colimit in schemes and pullback/limit in rings)

**example:** (affine line with double origin) Let  $k$  be a field, let  $U_1 = U_2 = \text{Spec } k[x]$ , let  $U_{12} = U_{21} = \text{Spec } k[x] - \{(x)\}$  and the gluing morphism  $U_{12} \rightarrow U_{21}$  the identity map. The global section is  $\Gamma(\mathcal{O}_X) = k[x]$  (exercise 3.26 -> showing  $X$  is not affine)

**example:** (projective space) Let  $R$  be a ring. Define the projective space  $\mathbb{P}_R^n$  over  $R$  by gluing  $n + 1$  copies of affine space  $\mathbb{A}_R^n$ , that is,  $U_i = \mathbb{A}_R^n$  for  $i = 0, \dots, n$ . For convenience, let

$$A^i = R[Y_0, \dots, \widehat{Y}_i, \dots, Y_n]$$

( $\widehat{Y}_i$  means  $Y_i$  is omitted) and we can view these rings as subring of  $R[X_0, \dots, X_n, X_0^{-1}, \dots, X_n^{-1}]$  by the inclusion  $Y_j \mapsto X_j X_i^{-1}$ , then  $U_i = \text{Spec } A^i$ . For each pair  $i \neq j$ , set  $U_{ij}$  to be the principal open set  $D(Y_j) = A^i[Y_j^{-1}]$ , let the gluing map  $\phi_{ji} : U_{ij} \rightarrow U_{ji}$  be induced by the equality  $A_{Y_j}^i \rightarrow A_{Y_i}^j$  in  $R[X_0, \dots, X_n, X_0^{-1}, \dots, X_n^{-1}]$  as follows:

$$\begin{array}{ccccc} A^i[Y_j^{-1}] = R[Y_0, \dots, \widehat{Y}_i, \dots, Y_n, Y_j^{-1}] & & & & A^j[Y_i^{-1}] = R[Y_0, \dots, \widehat{Y}_j, \dots, Y_n, Y_i^{-1}] \\ & \searrow & & \swarrow & \\ & & R[X_0, \dots, X_n, X_0^{-1}, \dots, X_n^{-1}] & & \\ & \nearrow & & \nwarrow & \\ Y_j^{-1} & & & & Y_i^{-1} \\ \downarrow & & & & \downarrow \\ X_j^{-1} X_i & & & & X_i^{-1} X_j \end{array}$$

This scheme is called projective space of relative dimension  $n$  over  $R$ . We consider the open subscheme  $U_i$  of  $\mathbb{P}_R^n$  and denoted them by  $D_+(X_i)$ . The canonical ring morphism  $R \rightarrow \Gamma(\mathbb{P}_R^n)$  is an isomorphism. Hence, projective space is not affine.

## 2.5 BASIC PROPERTIES OF SCHEME AND MORPHISM OF SCHEMES

### 2.5.1 ON LOCAL PROPERTIES

In commutative algebra, one often observes many properties called local property in which a property on ring  $A$  propagate into its prime ideals and maximal ideals. Below is an example

**Remark 2.5.1** (being zero is a local property)

Let  $A$  be a ring, let  $x \in A$ , then the following are equivalent

1.  $x = 0$  in  $A$
2.  $x = 0$  in  $A_{\mathfrak{p}}$  for every prime ideal  $\mathfrak{p}$
3.  $x = 0$  in  $A_{\mathfrak{m}}$  for every maximal  $\mathfrak{m}$

There is an equivalent notion of local property. Since GW did not discuss it, Vakil only briefly mentioned it, below is my attempt to properly define local property: A property  $P$  on some collection of objects is a subset of the collection, if an object  $X$  belong to the subset, we said  $X$  has property  $P$  or  $X$  is  $P$  or  $P(X)$

**Definition 2.5.2** (stalk-local)

Let  $X$  be a scheme and let  $P$  be a property on schemes.  $P$  is called stalk-local if there exists a property  $Q$  on stalks of schemes such that  $P(X) \iff Q(\mathcal{O}_{X,x})$  for every  $x \in X$

stalk-local property on schemes are the equivalence of local property on rings.

**Definition 2.5.3** (affine-local)

Let  $X$  be a scheme and let  $P$  be a property on schemes  $P$  is called affine-local if

1. (locality) if  $P(X)$ , then  $P(U)$  for every open affine subscheme  $U \subseteq X$
2. (gluability) if  $\{U_i\}_{i \in I}$  is an open affine cover of  $X$  and  $P(U_i)$  for every  $i \in I$ , then  $P(X)$

If we replace open affine subsets by open set, we say that  $P$  is open-local. However, the open affine sets generate the topology on  $X$ , it is easy to show that open-local and affine-local are equivalent. It is also clear that stalk-local implies affine-local.

Side note on sheaf: *(somehow I just randomly admire the beauty of sheaf)*

**Proposition 2.5.4** (sections are determined by germs)

Let  $\mathcal{F}$  be a sheaf on  $X$  and  $U \subseteq X$  be an open set, then the canonical map from sections to stalks is injective

$$\mathcal{F}(U) \rightarrow \prod_{x \in U} \mathcal{F}_x$$

This property has two important implications:

1. A sheaf  $\mathcal{F}$  on  $X$  is completely determined by its étale space  $E = \coprod_{x \in X} \mathcal{F}_x$ <sup>2</sup>
2. A section  $s \in \mathcal{F}(U)$  is zero if and only if  $s$  is zero on all stalks on  $U$

Back to local property, when  $X = \text{Spec } A$ , then being zero is stalk-local.

*(I will mumble something that I am not an expert in, so this is likely wrong)* Another subtle issue is these local definitions use higher order logic.

## 2.5.2 TOPOLOGICAL PROPERTIES

<sup>2</sup>topologized by the weakest topology generated by all compatible germs  $(s_x)_{x \in U}$  for some  $s \in \mathcal{F}(U)$

**Definition 2.5.5** (connected, quasi-compact, irreducible)

A scheme is called connected/quasi-compact/irreducible if the underlying topological space is connected/quasi-compact/irreducible.

All affine schemes are quasi-compact.

**Definition 2.5.6** (injective, surjective, bijective)

A morphism of schemes  $f : X \rightarrow Y$  is called injective/surjective/bijective if the underlying set function is injective/surjective/bijective.

**Definition 2.5.7** (open, closed, homeomorphism)

A morphism of schemes  $f : X \rightarrow Y$  is called open/closed/homeomorphism if the underlying continuous map is open/closed/homeomorphism.

## 2.5.3 NOETHERIAN SCHEMES

**Definition 2.5.8** (locally Noetherian, Noetherian)

A scheme  $X$  is called locally Noetherian if every open affine subscheme  $U \subseteq X$  is a Noetherian ring. A quasi-compact locally Noetherian scheme is called Noetherian.

Localization of Noetherian rings is Noetherian, hence a stalk of a locally Noetherian scheme is a Noetherian ring and every affine scheme of a Noetherian ring is Noetherian. However, being Noetherian scheme is not stalk-local, the converse statement is false.

**Proposition 2.5.9**

An affine scheme  $X$  is Noetherian if and only if it is a Noetherian ring.

**Lemma 2.5.10**

Let  $A$  be a ring and  $\text{Spec } A = \bigcup_{i \in I} D(f_i)$  be a finite open cover of  $\text{Spec } A$  by principal open subsets <sup>a</sup>. Let  $M$  be an  $A$ -module, then  $M$  is finitely generated if and only if for all  $i \in I$ , the localization  $M_{f_i}$  is a finitely generated  $A_{f_i}$ -module.

<sup>a</sup>that is, the ideal generated by  $f_i$  is the whole ring  $A$

**Remark 2.5.11** (Noetherian topological space)

A topological space  $X$  is called Noetherian if one of the following conditions satisfies:

1. Every nonempty collection of open sets has a maximal element
2. Decending chain condition for closed sets.

A finite covering  $X = \bigcup_{i \in I} X_i$  of Noetherian spaces is Noetherian.

Hence, the topological space of an Noetherian scheme is Noetherian but the converse statement is false.

**Remark 2.5.12** (subsets of Noetherian topological space)

Let  $X$  be a Noetherian topological space

1. Every subspace of  $X$  is Noetherian
2. Every open subset of  $X$  is quasi-compact
3. Every closed subset of  $X$  has only finitely many irreducible components.

**Proposition 2.5.13**

Let  $X$  be a (locally) Noetherian scheme, then every open subscheme  $U \subseteq X$  is (locally) Noetherian.

## 2.5.4 GENERIC POINTS

**Remark 2.5.14**

Let  $X$  be a topological space, a subspace  $Z \subseteq X$  is irreducible if and only if its closure is irreducible

**Remark 2.5.15**

Let  $X$  be a topological space and  $Z \subseteq X$ , a point  $z \in Z$  is called a generic point of  $Z$  if  $\overline{\{z\}} = Z$ .

As closure of an irreducible subset is irreducible, if a subset of  $X$  contains a generic point, it is irreducible. Moreover

**Proposition 2.5.16**

Let  $X$  be a scheme, the map

$$\begin{aligned} X &\rightarrow \{\text{closed irreducible subsets of } X\} \\ x &\mapsto \overline{\{x\}} \end{aligned}$$

is a bijection.

**Proposition 2.5.17**

Let  $f : X \rightarrow Y$  be an open morphism of schemes and  $Y$  be irreducible with generic point  $\eta$ . Then  $X$  is irreducible if and only if the fiber  $f^{-1}(\eta)$  is irreducible.

*Proof. (the proof in GW was too short, skipped many point-set topology things, this is a better version)*

It's clear that  $\overline{f^{-1}(\eta)} \subseteq f^{-1}(\overline{\{\eta\}})$  since  $f$  is continuous. Moreover, since  $f$  is open, any open set  $U \subseteq X$  is mapped into an open set  $f(U) \subseteq Y$ .  $\{\eta\}$  is dense in  $Y$ , hence  $\eta \in f(U)$ , so  $U \cap f^{-1}(\eta) \neq \emptyset$ .  $f^{-1}(\eta)$  cuts all open sets in  $X$ , so it is dense in  $X$ .  $\square$

Although schemes are never Hausdorff, they at least satisfy some weaker separation property.

**Proposition 2.5.18**

Let  $X$  be a scheme, then the underlying topological space  $X$  is Kolmogorov ( $T_0$ -space), that is, for any two distinct point  $x, y \in X$ , there exists an open subset of  $X$  containing exactly one of the points.



## 2.5.5 REDUCED SCHEME, INTEGRAL SCHEME, FUNCTION FIELD

A reduced ring is a ring with no nilpotent

**Definition 2.5.19** (reduced scheme, integral scheme)

A scheme is called reduced if every stalk is reduced. A scheme is called integral if it is reduced and irreducible.

*(from here, we diverted from GW a bit so that it is easier to internalize)*

By definition, being reduced is a stalk-local property, hence also an affine-local and open-local. Since being reduced ring is a local property, there for an affine scheme is reduced if and only if it is a reduced ring. One can characterized reduced scheme as follows:

**Remark 2.5.20**

A reduced scheme is characterized by the following two conditions:

1. being reduced scheme is an affine-local property
2. an affine scheme is reduced if and only if it is a reduced ring.

**Proposition 2.5.21**

An affine scheme is integral if and only if it is an integral domain.

*Proof. TODO*

□

**Proposition 2.5.22**

Let  $X$  be a scheme,  $X$  is integral if and only if the global sections of every nonempty open subscheme is an integral domain. If  $X$  is integral then every stalk is an integral domain. However, the converse does not hold, that is, being integral is not a stalk-local property.