



A Monte-Carlo investigation of the nematic-isotropic phase transition

Félix Bunel & Hadrien Vergnet


16/01/2017






Nematic phase:

- No positional order
- Orientational order



- 1 The order parameter
- 2 Numerical methods
 - Lebwohl Lasher Model
 - Monte-Carlo Algorithm
 - Equilibration
- 3 Nematic Isotropic Phase Transitions
 - Direct visualisation
 - Energy
 - Energy
 - Histograms
- 4 Influence of an Electric Field
 - Lebwohl Lasher Model
 - Critical point
 - Phase diagram

The Order Parameter




$$\mathbf{n} = \langle \mathbf{a} \rangle$$



$$S = \frac{3\langle (\mathbf{a} \cdot \mathbf{n})^2 \rangle - 1}{2}$$


Nematic phase : $S = 1$


Isotropic phase : $S = 0$




Size : $30 \times 30 \times 30$

$$E = -\epsilon \sum_{\langle i,j \rangle} \frac{3 \cos^2 \alpha_{i,j} - 1}{2}$$







- We select a random site




- We select a random site
- We compute the energy with the neighbor : E_{old}




- We select a random site
- We compute the energy with the neighbor : E_{old}
- We try to swap the chosen site



- We select a random site
- We compute the energy with the neighbor : E_{old}
- We try to swap the chosen site




- We select a random site
- We compute the energy with the neighbor : E_{old}
- We try to swap the chosen site
- We compute the energy with the neighbor : E_{new}



- We select a random site
- We compute the energy with the neighbor : E_{old}
- We try to swap the chosen site
- We compute the energy with the neighbor : E_{new}
- We accept the swap with a probability :


$$p = e^{-\frac{E_{old} - E_{new}}{k_B T}}$$




Equilibration at $T^* = 1$

Nematic Isotropic Phase Transitions


Direct visualisation



$$T^* = 0.05$$




$$T^* = 0.9$$



$$T^* = 1.5$$

Nematic Isotropic Phase Transitions


Energy



Energy and its variance.

Nematic Isotropic Phase Transitions





Energy




Order parameter and its variance.


Nematic Isotropic Phase Transitions

Energy Histograms






$$E = -\epsilon \sum_{\langle i,j \rangle} \frac{3 \cos^2 \alpha_{i,j} - 1}{2} - \epsilon \xi U^2 \sum_i \frac{3 \cos^2 \beta_i - 1}{2}$$



Energy for different electric field.



Transition temperature as a function of the electric field.

Conclusion

Merci pour votre attention